
Nucleotide sequence and organization of *Bacillus subtilis* RNA polymerase major sigma (σ^{43}) operon

Lin-Fa Wang and Roy H. Doi*

Department of Biochemistry and Biophysics, University of California, Davis, CA 95616, USA

Received 6 March 1986; Revised and Accepted 14 April 1986

ABSTRACT

The gene coding for *Bacillus subtilis* RNA polymerase major σ^{43} , *rpoD*, was cloned together with its neighboring genes in a 7 kb *EcoRI* fragment. The complete nucleotide sequence of a 5 kb fragment including the entire *rpoD* gene revealed the presence of two other genes preceding *rpoD* in the order *P23-dnaE-rpoD*. The *dnaE* codes for DNA primase while the function of *P23* remains unknown. The three genes reside in an operon that is similar in organization to the *E. coli* RNA polymerase major σ^{70} operon, which is composed of genes encoding small ribosome protein S21 (*rpsU*), DNA primase (*dnaG*), and RNA polymerase σ^{70} (*rpoD*). There is a relatively high degree of base and amino acid homology between the DNA primase and σ genes. The most significant differences between the two operons are observed in the molecular size of the first genes (*P23* and *rpsU*), the complete lack of amino acid homology between *P23* and S21, the molecular weights of the two *rpoD* genes, the size of the intercistronic region between the first two genes, and the regulatory elements of the operon.

INTRODUCTION

The existence of multiple RNA polymerase σ factors in *B. subtilis* has been well documented (1,2), but little is known about their genetic properties, the regulation of their synthesis, and the factors that govern their interactions with the RNA polymerase core. An analysis of their molecular organization and the parameters which regulate their genetic expression should provide a initial basis for determining their roles in the physiology of this Gram positive sporulating bacterium.

Our laboratory has been particularly interested in the study of the σ^{43} gene (*rpoD*), whose product is known to play a major role during vegetative growth, and the early stationary and sporulation phases (3). We have been able to clone (4),

genetically map (5), and sequence (6) the σ^{43} gene (rpoD), and show that its derived amino acid sequence had a very high degree of homology with that of the E.coli major σ^{70} (7). By genetic mapping (5) and DNA sequencing (8), we also showed that immediately upstream of the rpoD gene was located the dnaE gene, which encodes the B. subtilis DNA primase, whose product is very homologous to the E.coli dnaG DNA primase (9,10). No promoter region was observed in the intercistronic region between rpoD and dnaE, nor in the region immediately upstream of dnaE (6,8).

Recently, we have determined the nucleotide sequence of the region upstream of dnaE including the operon regulatory region, which provided support for our previous suggestion (6,8) that dnaE and rpoD were coordinately regulated with one or more unknown genes in an operon. The DNA sequence analysis of the region upstream of dnaE revealed an open reading frame capable of coding for a protein of molecular weight 22,540. The function of this protein is unknown, and hence the designation P23 is being used for this gene until a physiological role can be assigned to it.

In this paper we will discuss the similarities and differences of the structure and organization of the major sigma operons of B. subtilis and E. coli, the transcriptional and translational regulatory features of the operon, and the codon usage frequency encountered in the operon.

MATERIALS AND METHODS

Strains, Phages and Plasmids

E. coli JM101 was used as host for the sequencing phage vectors M13mp8, M13mp9, M13mpl0, and M13mpl1 (11,12), and the plasmid pCPS1 (5). E. coli BNN45 (13) was used to prepare the phage lysate of λ gtWES- σ 82 (4). Plasmid pSB was provided by Sui-Lam Wong (unpublished data).

DNA Manipulations

Standard procedures of Maniatis *et al.* (14) were followed exactly as described.

DNA Sequencing

DNA sequencing was conducted by the dideoxy chain

termination method of Sanger *et al.* (15) using the sequencing kit purchased from Amersham Corporation.

Computer Analysis

Routine analysis of DNA or protein sequences were carried out using either the Delaney (16) or the Pustell (17) program, while the homology search against the NBRF Data Bank was made using the Microgenie Sequence Analysis Program developed by Queen and Korn (18).

RESULTS

Nucleotide sequence of the Entire Operon

The nucleotide sequences and the sequencing strategies of *dnaE* and *rpoD* genes have been reported previously (6,8). The sequencing strategy for the upstream 1.5 kb fragment is shown in Fig. 1 (bottom) along with the physical map of the σ^{43} operon (upper). As indicated, the nucleotide sequence has been determined for both strands of virtually the entire region except for the 100 bp at the extreme 5' end. The sequence was determined across the junctions of all the restriction sites used for subcloning during sequencing, as well as for the *EcoRI* site between the *dnaE* and *rpoD* genes (not shown here). In our previous reports, the sequences for these two genes were determined separately (6,8). Although unlikely, the possibility existed that a small *EcoRI* fragment may have been left out during the subcloning of the *EcoRI* fragments into plasmids from the original phage λ gtWES- σ 82 (4,5). Therefore we sequenced the 0.9 kb *HindIII* fragment containing the *EcoRI* junction region, which was subcloned into M13mpl0 directly from λ gtWES- σ 82, and the possibility mentioned above has been experimentally excluded. Now, the entire *EcoRI-SphI* fragment has been sequenced, including all the junctions of restriction sites used for sequencing. The nucleotide sequence of the entire operon and its flanking regions, and the deduced amino acid sequence of each gene are given in Fig. 2 with the first base of the 5' end *EcoRI* site labeled as number 1.

Features of the First Gene of the Operon

When the sequence of the region upstream of *dnaE* was analyzed by computer, only one large open reading frame was




Figure 1. Sequencing strategy for the first gene and regulatory region of the σ^{43} operon. The upper part represents the physical location of the genes in the operon. The lower part indicates the restriction sites used for subcloning and sequencing. The bar indicates the location of the cryptic P23 protein. Abbreviations: H, *HpaII*; S, *Sau3A*.

discovered. But unlike the case for the other two genes in the operon, we could not identify any strong ribosomal binding site by sequence analysis within the open reading frame. We found instead several weak ones preceding the potential initiation codons ATG (855), TTG (930), ATG (951), and GTG (1,200), which were able to code for proteins of molecular weights 22,540, 19,734, 18,934 and 9,312, respectively. However, our previous maxicell data showed that a protein of molecular weight around 23,000 was encoded within the upstream region of *dnaE* (4,5), which corresponded very well with the largest open reading frame identified here by sequencing, i.e., the open reading frame starting from the initiation codon ATG at nt 855.

Although the ribosomal binding site was very weak as predicted from its calculated free energy of binding ($\Delta G = -9.2$ kcal/mol), our assignment was further strengthened by our recent protein fusion studies. When the N-terminal two thirds of the P23 was fused to the *E. coli* β -galactosidase in frame, a functional hybrid protein was expressed in both *E. coli* and *B. subtilis* with the expected size as determined by Western blot analysis using anti- β -galactosidase antibody (data not shown). Functional P23- β -galactosidase (P23-gal) fusion protein was expressed even when only the first 8 amino acid residues of P23

1184
 GCT GAT TTA TAT ATC GCA AAT CAC GTG AAA CCG GGA GAT ATT GTT GTG ACG CAG GAA ATC
 Ala Asp Leu Tyr Ile Ala Asn His Val Lys Pro Gly Asp Ile Val Val Thr Glu Asp Ile 1214

124
 GGA TTA GCA TCT CTG CTG TCG AAC GAT AAT GTC TCT GTT ATG TCG GAA ACA GGT GGT GGT
 Gly Leu Ala Ser Leu Leu Asn Arg Asn Val Ser Val Met Ser Glu Arg Gly Arg Leu 1274

1304
 TAC AAG GAA GAC ACG ATT GAT TTT GCC CTA GAG GGC GGT CAT TTT TCC GGC AAA CAA AGA
 Tyr Lys Glu Asp Thr Ile Asp Thr Ala Leu Glu Arg His Phe Ser Ser Lys Glu Arg 1334

1364
 AGA AAA GGC GTA TAT GCC AAA CGG CCT AAA AAA TTG AAT AAA GAA GAT CGA GAA CGA TTT
 Arg Lys Gly Val Tyr Ala Lys Gly Pro Lys Lys Leu Asn Lys Leu Asn Lys Glu Asp Arg Glu Arg Phe 1394

1424
 ATT ACA CTG CTG CAA AAA ATC CTG AAC GAT GAA GGG ATT TTG CAC TAA AGCATCGAATA
 Ile Thr Leu Leu Glu Lys Ile Leu Ser Asn Asp Glu Gly Ile Leu His End 1456

APRDNCGCG GAGTGTATA AG

1508
 ATG CGA AAT CCG ATA CCA GAT GAA ATT GCG GAT CAG GTG CAA AAG TCG GCA GAT ATC GGT
 Met Gly Asn Arg Ile Pro Asp Glu Ile Val Asp Glu Val Glu Lys Ser Ala Asp Ile Val 1538

1568
 GAA GTC ATA GGT GAT TAT GTT CAA TTA AAG AAG CAA GGC CGA AAC TAC TTT GGA CTC TGT
 Glu Val Ile Gly Asp Tyr Val Glu Leu Lys Lys Glu Gly Arg Asn Tyr Phe Gly Leu Cys 1598

1628
 GCT TTT CAT GGA GAA ACG ACG CCT TCG TCG TCC GTR TCG CCC GAC AAA CAG ATT TTT GAA
 Pro Phe His Gly Glu Ser Thr Pro Ser Phe Ser Val Ser Pro Asp Lys Glu Ile Phe His 1658

1688
 TGC TTT GGC TGC GGA CGC GGC AAT GTT TTC TCT TTT TTA AGG CAG ATG GAA GGC TAT
 Cys Phe Gly Cys Gly Ala Gly Gly Alan Val Phe Ser Phe Leu Arg Glu Met Glu Gly Tyr 1718

1748
 TCT TTT GCG GAG TCG GTT TCT CAC CTT GCT GAC AAA TAC CAA AAT GAT TTT CCA GAT GAT
 Ser Phe Ala Glu Ser Val Ser His Leu Ala Asp Lys Tyr Glu Ile Asp Phe Pro Asp Asp 1778

1808
 ATA CAA GTC CAT TCC GCA GCC CGC CCA GAG TCT TCT GGA GAA CAA AAA ATG GCT GAG GCA
 Ile Thr Val His Ser Gly Ala Arg Pro Glu Glu Ser Ser Gly Lys Met Ala Glu Ala 1838

1868
 CAT GAG CTC CTG AAG AAA TTT TAC CAT CAT TTG TTA ATA AAT ACA AAA GAA GGT CAA GAG
 His Glu Leu Leu Lys Lys Phe Tyr His His Leu Leu Ile Asn Thr Lys Glu Gly Glu 1898

1928
 GCA CTG GAT TAT CTG CTT TCT AGG GCG TTT AGC AAA GAG CTG ATT AAT GAA TTT CAG ATT
 Ala Leu Asp Tyr Leu Leu Ser Arg Gly Phe Thr Lys Glu Leu Ile Asn Glu Phe Glu Ile 1958

1988
 GGC TAT GCT CTT GAT TCT TCG GAC CTT ATC ACG AAA TTC CTT GTA AAG ACG GGA TTT AGT
 Gly Tyr Ala Leu Asp Ser Trp Asp Phe Ile Thr Lys Phe Leu Val Lys Arg Gly Phe Ser 2018

2048
 GAG GCG CAA ATG GAA AAA GCG GGT CTC CTC ATC AGA CCG GAA GGA GCG AAT TAT TTC
 Glu Ala Glu Met Glu Lys Ala Gly Leu Leu Ile Arg Arg Glu Asp Gly Ser Gly Tyr Phe 2078

diB.2

30
 GAAATTCAT GTTCAGATA TACGCACTAT CAGTAGAGT ATTTCATCTG CAAAGCCGA 80

70
 CGCGCGAAT ATCTGTTTAA CACTCGGTGT GCGGAAATC AGAGATAATT TGATAGCCGA 120

130
 ACCGGAAMA GCAAATGTTT TATATATGA TATTATCGC CCGTTGATTT ATAAATATGA 180

180
 AACAGCCAC GGTTHAACA GGAATATCA ACCGGGGCG GTGCGCCAGC TTGATGACGA 240

240
 TTAATTCAAA AACTGCGAGG CCATCGAGTT TCGAGTTAAA TACGATATATC GACGTGATCC 300

300
 AAGAGGAAAT TTAAAGACCTG ATATGCTTGT GATCGCGGT TCGAAGACGT CTAAACACCC 360

370
 GCTGTCTCA TATCTCCAC ACAACGGCT GAAGTTGCC AATGTTCCGA TTGTACCCGA 420

430
 GGTGATCG CCGGAGAAC TCTTTACCT TGATCCGAAA AATGATATCG GTTTAAGAT 480

490
 TAGCCCTGAT AACTGATATC ATATACGAAA AGAAGCTTTC AATATCATCC GCGTTAATCA 540

550
 TAAAGCGATT TATCGAATA TCAACAGAT CAAAGAGCA CTCGAGTATT TCGAAGAT 600

610
 TGTGATCG ATGCGTCCC AGGTTGTTCA TTTTTCAAAT AAAGCGTTG AGGAACACGC 660

670
 AATATATC CATCTCTCA AACCAAAA CAATATATC AGGACGCTCT ATCCGTGTT 720

730
 TTGCGCTGT CCAAAAGCA ATATGAAA ACAATAGAT CTATTGAGC TTTGTATAT 780

790
 AAAAAAAAA TGTATAAA TGATTAATT TAGGTTAAG GATCGTGA TAGCAATAA 840

850
 CATATATGCG TAG

914
 ATG TCA AGA ATT TCT CCC GGA AAT TTT TCG ACA AAT TCA TAT ACA TCC ACA ATA ATA AAG
 Met Ser Arg Ile Ser Pro Gly Asn Phe Ser Thr Asn Ser Tyr Thr Ser Thr Ile Ile Lys

974
 GAT GTC CGA TTT TCG TGC CTT TTA TCC ACG AGT TTA ATG GAG GGA TGG AGA ATT ACT CTT
 Asp Val Arg Phe Cys Leu Leu Leu Cys Arg Ser Leu Met Glu Gly Trp Arg Ile Thr Leu

1004
 CTT AAT GAA CAA GAA AAG ACG ATT TTT GTC GAT GCT GAT GCT TGT CCG GTA AAA GAT GAA
 Leu Asn Glu Glu Glu Lys Thr Ile Phe Val Asp Ala Asp Ala Cys Pro Val Lys Asp Glu

1064
 ATT TTA CAA ACA TCC GAT TAT GAA GTT CAA GTT CTT TTT GTC GCT TCA TTT GAA CAT
 Ile Leu Glu Thr Ala Ser Glu Tyr Glu Val Glu Val Leu Phe Val Ala Ser Phe Glu His

1124
 TAT CAG CTT TCC ACA ACG AAT GAA GAA TGG AAG TAT GTT GAT CCT CAT AAA GAA GCT
 Tyr Glu Leu Ser Arg Ser Asn Glu Glu Lys Trp Lys Tyr Val Asp Pro His Lys Glu Ala

p23

884
 ATG TCA AGA ATT TCT CCC GGA AAT TTT TCG ACA AAT TCA TAT ACA TCC ACA ATA ATA AAG
 Met Ser Arg Ile Ser Pro Gly Asn Phe Ser Thr Asn Ser Tyr Thr Ser Thr Ile Ile Lys

944
 GAT GTC CGA TTT TCG TGC CTT TTA TCC ACG AGT TTA ATG GAG GGA TGG AGA ATT ACT CTT
 Asp Val Arg Phe Cys Leu Leu Leu Cys Arg Ser Leu Met Glu Gly Trp Arg Ile Thr Leu

1128
 ATG CTT CAG GTT AAT CAA GAG CTT AGC GAA GCC GAG TTA TCA GAT TAT GTA AAA AAA
 Met Leu Glu Val Asn Glu Glu Leu Ser Asp Tyr Val Lys Lys Val

3128
 TTG AAT GAA AGA AAT TGG TCA ATG ATA AAA AAA GAA GCG GAA AGA GCC GAA GCA GAA
 Leu Asn Glu Arg Asn TTP Ser Met Ile Lys Glu Lys Glu Ala Glu Arg Ala Glu Ala Glu

3218
 AGG CAA AAA GAT TTT TTA AGA GCT GCT TCT TTS GCT CAA GAA ATC GTT ACA TTG AAC CGA
 Arg Glu Lys Asp Phe Leu Arg Ala Ala Ser Leu Ala Glu Glu Ile Val Thr Leu Asn Arg

3248
 TCT TTA AAA TAA
 Ser Leu Lys End

3300 3310 3320 3330 3340 3350
 CTGGAGACT GATGAGGAC ATTATTCGC AATGATTCCT TCGGAGAG CAANTAGATC 3350
 3360 3370 3380 3390 3400 3410
 GCTTACCTC ATCAATGATTT GTCATTCCTT PATTGCGACA TGTGTAAGG CAGTTCACAT 3410
 3420 3430 3440 3450 3460 3470
 AGAAGGCC TGAATGACC GAATAGATTT CATACGCTT ATAGATTCG TTGCAGCTT 3470
 3480
 TCGAAGGCG GATGCATA

3518
 ATG OCT GAT AAA CAA ACC CAC GAG ACA TTA ACA TTC GAC CAA GTA AAA GAA GAA TTA
 Met Ala Asp Lys Glu Thr His Glu Thr Glu Leu Thr Phe Asp Glu Val Lys Glu Glu Leu

3548
 ACA GAG TCT GGT AAA AAA CGT GCG GTT TTS ACA TAT GAA AAT GCT GAG COT ATG TCC
 Thr Glu Ser Gly Lys Lys Arg Gly Val Leu Thr Tyr Glu Glu Ile Ala Glu Arg Met Ser

3608
 AGC TTT GAA ATT GAA TCA GAC CAA ATG GAT GAG TAT TAT GAA TTT TTA GGT GAA CAA GGT
 Ser Phe Glu Ile Glu Ser Asp Glu Met Asp Glu Tyr Tyr Glu Phe Leu Glu Glu Glu

3638
 GTT GAA TTA ATT AGT GAG AAT GAA GAA ACA GAA GAT CCT AAT ATT CAG CAG CTT GCC AAA
 Val Glu Leu Ile Ser Glu Asn Glu Glu Thr Glu Asp Pro Asn Ile Glu Glu Leu Ala Lys

3658
 GCC GAA GAA GAA TTT GAC CTT AAT GAC GTA AGT GTA CCG CCT GGC GTT AAA ATC AAT GAG
 Ala Glu Glu Glu Phe Asp Leu Asn Asp Leu Ser Val Pro Gly Val Lys Ile Asn Asp

3758
 CCA GTT COT ATG TAT TTA AAG GAA GAT GGT CCG GTT AAC CTT CTT TCT GCA AAA GAA GAA
 Pro Val Arg Met Tyr TTA Lys Glu Ile Gly Arg Val Asn Leu Leu Ser Ala Lys Glu Glu

3818
 ATC GCC TAC GCT CAA TAG ATT GAA GAA GAT GAT GAT TAA GCG AGA TTG GCT GAA
 Ile Ala Tyr Ala Glu Lys Ile Glu Glu Gly Asp Glu Glu Ser Lys Arg Arg Leu Ala Glu

IPD

2138
 GAC CGC TTC AGA AAC CGT GTC ATG TTT CGC ATC CAT GAT CAT CAC GGG GCT GTT GTT GCT
 Asp Arg Phe Arg Asn Arg Val Met Phe Pro Ile His Asp His His Gly Ala Val Val Ala

2198
 TTC TCA GCG AGC GCT CTT GCG CAG CAG CTT AGC TAT AGC TAC ACT CTT GAA ACC CGC
 Phe Ser Gly Arg Ala Leu Gly Ser Glu Asn Pro Lys Tyr Met Asn Ser Pro Glu Thr Pro

2258
 CTC TTT CAT AAA AGC AAA CTG CTT TAC AAT TTT TAT AAG GCC GCG CTT CAT ATC AGA AAG
 Leu Phe His Lys Ser Lys Leu Leu Tyr Asn Phe Tyr Lys Ala Arg Leu His Ile Arg Lys

2318
 CAG GAA AGA GCA GTC TTA TTT GAA GCG TAT GAT GGC TAT AGC GCC GFA AGC TCG GAT
 Glu Glu Arg Ala Val Leu Phe Glu Gly Phe Ala Asp Val Tyr Thr Ala Val Ser Ser Asp

2378
 GTA AAG GAA AGC AFA GCC ACG ATG GGA ACG TCT CTT ACA GAT GAT CAT GTC AAG ATC CTG
 Val Lys Glu Ser Ile Ala Thr Met Gly Thr Ser Leu Thr Asp Asp His Val Lys Ile Leu

2438
 AGA AGA AAC GTC GAA GAA ATC ATT CTT TGC TAT GAC TCT GAT AAG GCC GGT TAT GAA GCG
 Arg Arg Asn Val Glu Glu Ile Ile Leu Cys Tyr Asp Ser Asp Lys Ala Gly Tyr Glu Ala

2498
 ACC TTA AAA GCT TCG GAG CTT CTG CAA AAA AAA GGC TCC AAA GTC AGA GTT GCA ATG ATT
 Thr Leu Lys Ala Ser Glu Leu Leu Glu Lys Lys Gly Cys Lys Val Arg Val Ala Met Ile

2558
 CCT GAC GGA TTG GAC CCT GAT GAT TAC ATC AAA AAA TTC GCG GGG GAA AAA TTT AAA AAC
 Pro Asp Gly Leu Asp Pro Asp Asp Tyr Ile Lys Lys Phe Gly Gly Glu Lys Phe Lys Asn

2618
 GAC ATT ATT GAC GCA AGT GTC ACC GTA ATG GCG TTC AAA ATG CAA TAT TTC CGA AAA GGA
 Asp Ile Ile Asp Ala Ser Val Thr Val Met Ala Phe Lys Met Glu Tyr Phe Arg Lys Gly

2678
 AAG AAC CTG TCC GAT GAA GGC GAC CCG CTA GCT TAC GCT AAT AAA GAC GTA CTG AAA GAA ATC
 Lys Asn Leu Ser Asp Glu Gly Asp Arg Leu Ala Tyr Ile Lys Asp Val Leu Lys Glu Ile

2738
 AGC ACG CTT TCA GGG TCT CTA GAG CAG GAA GTC TAT GTA AAG CAG CTT GCT TCA GAG TTT
 Ser Thr Leu Ser Gly Ser Leu Glu Glu Glu Val Tyr Val Lys Glu Leu Ala Ser Glu Phe

2798
 TCG CTT TCA CAG GAG TCT TTA ACT GAG CAG CTG TCT GCT TTC AGC AAG CAA AAC AAA CCT
 Ser Leu Ser Glu Glu Ser Leu Thr Glu Glu Leu Ser Val Phe Ser Lys Glu Asn Lys Pro

2858
 GCT GAC AAT AGC GGT GAA ACT AAA ACG CCG CCA GCG CAT CTG ACG ACA AAA GCA GCG CAA
 Ala Asp Asn Ser Gly Glu Thr Lys Thr Arg Arg Ala His Leu Thr Lys Ala Arg Glu

2918
 AAA COT TTG COT CCG GAT GAA AAT GCA GAA AGG CTG TTA CTC GCT CAC ATG CTT CGA
 Lys Arg Leu Arg Pro Ala Tyr Glu Asn Ala Glu Arg Leu Leu Leu Ala His Met Leu Arg

2978
 GAT CCG AGC GTC ATC AAA AAA CTG ATT GAC CCG GTA CCG TTT CAA TTT AAT ATT GAT GAG
 Asp Arg Ser Val Ile Lys Lys Val Ile Asp Arg Val Gly Phe Glu Phe Asn Ile Asp Glu

3038
 CAC CCG GCA TTA GCC OCT TAT CTT TAT GCT TTT TAT GAA GAG GAA GCC GAG CTG ACG COT
 His Arg Leu Ala Arg Pro Ala Tyr Glu Glu Glu Gly Ala Phe Tyr Glu Glu Glu Leu Thr Pro

3098
 CAG CAT CTG ATG GCC ACG GTG ACG GAT GAT ATA ACC GTC CTC TTG TCC GAT ATP TTA
 Glu His Leu Met Ala Arg Val Thr Asp Asp His Ile Ser Glu Leu Leu Ser Asp Ile Leu

4914 4924 4934 4944 4954 4964
 GCGCGAAGCA TCMAAAGCG GCGAAGGAAA TGAATAATGGAA CCGCTATT CCATTTTTCG
 4974 4984 4994 5004 5014 5024
 TGATCGCTGT TTTAGGATCG GGTCTAATCT TCTTTTATC AGTAAAGGCA CTTCTGACT
 5034 5044 5054 5064 5074 5084
 CTCGGAGGT TCGGACGGCA GCGAAGACCA AATCTGCTCA AAGAAGAT GCAAACTCTT
 5094 5104 5114 5124
 CACCAGAAGA AATTTACAG GCAAAATCGCA TCGCATCGCA TC

Figure 2. Nucleotide sequence of B. subtilis σ^{43} operon. The DNA sequence of the upper strand is given in the 5' to the 3' direction, numbered from nucleotide 1 at the 5' end EcoRI site. The predicted amino acid sequence for each open reading frame is given below the corresponding DNA sequence. Sequences for promoters, ribosomal binding sites, and terminator are underlined.

3938 3968
 GCG AAC CTG CCG CTT GTT GTC AGT ATC GCA AAA CCG TAT GTC GGA CCG GGT ATG CTG TTC
 Ala Asn Leu Arg Leu Val Val Ser Ile Ala Lys Arg Tyr Val Gly Arg Gly Met Leu Phe
 4028
 CTT GAT CTG ATC CAT GAA GGA AAC ATG CCG CTG ATG AAA GCC GGT GAA AAA TTT GAT TAT
 Leu Asp Leu Ile His Glu Asp Asn Met Gly Leu Met Lys Ala Val Glu Lys Phe Asp Tyr
 4088
 CCG AAA GGT TAT AAA TTA ACT GCT ACC TGG TCG ATC AGA CAG CGC ATT ACA CCG
 Arg Lys Gly Tyr Lys Phe Ser Thr Tyr Ala Thr Trp Trp Ile Arg Glu Ala Ile Thr Arg
 4148
 GGC ATT GCC GAT CAG GCG AGA CCG ATC CCG ATT CCC GTT CAT ATG GTT GAA ACC ATT AAT
 Ala Ile Ala Asp Glu Ala Arg Thr Ile Arg Ile Pro Val His Met Val Glu Thr Ile Asn
 4208
 AAA TTA ATC CGT GTG CAG CGT CAA TTA CTG CAA GAC TTA GGC AGA GAA CCA ACA CCT GAA
 Lys Leu Ile Arg Val Glu Arg Glu Leu Leu Glu Asp Leu Gly Arg Glu Pro Thr Pro Glu
 4268
 GAA ATT GCG GAA GAT ATG GAT TTA ACG CCT GAA AAA GTA CCG GAA ATC TTA AAG ATT GCT
 Glu Ile Ala Glu Asp Met Asp Leu Thr Pro Glu Lys Val Arg Glu Ile Leu Lys Ile Ala
 4328
 CAA GAG CCG GTA TCT CTG GAA ACA CCG ATC GGT GAA GAG GAT CAG TCG CAC CTT GGT GAT
 Glu Glu Pro Val Ser Leu Glu Thr Pro Ile Gly Glu Asp Asp His Leu Glu Asp
 4388
 TTC ATT GAA GAC CAA GAA GCA ACT TCA CCT TCT GAC CAC GCC GCA TAC GAG GTA TTG AAA
 Phe Ile Glu Asp Glu Ala Thr Ser Pro Ser Asp His Ala Ala Tyr Glu Leu Lys
 4448
 GAG CAG CTG GAA GAT GTG CTT GAT ACG TTA ACT GAT CGT GAA GAA MAT GTA TTG CGT CTT
 Glu Glu Leu Asp Val Leu Asp Thr Leu Thr Asp Arg Arg Glu Asp Val Leu Leu Lys
 4508
 CGA TTC GGT CTT GAT GAC GGC CGT ACA AGA ACA TTA GAA GAG GTC GGC AAA GTA TTT GGA
 Arg Phe Gly Leu Asp Asp Gly Arg Thr Arg Thr Leu Glu Val Gly Lys Val Phe Gly
 4538
 GTA ACG AGA GAG CGT ATT CGA CAA ATC GAA GCC AAA GCG TTC GCG AAA GTA ACA CAT CCT
 Val Thr Arg Glu Arg Ile Arg Glu Ile Glu Ala Lys Ala Leu Arg Lys Leu Arg His Pro
 4598
 ACG AGA AGT AAA CGT TTC AAA GAT TTC CTT GAA TAA
 Ser Arg Ser Lys Arg Lys Asp Phe Leu Glu End
 4634 4624 4634 4644 4654 4664
 GATGAGGCG CTCTGAGG TCGCTGCTC TTTTATTAA AGATATATG GATATATGC
 4674 4684 4694 4704 4714 4724
 CTTTATTTA CTGAAAATG ATGTCAATTC CAAATGACA TTCTGTGAA AATTTCAA
 4734 4744 4754 4764 4774 4784
 ATCTAATTC ATMTTCTA TTGTAGGCT ATACAAACA TTATCAATA GAATTAAG
 4794 4804 4814 4824 4834 4844
 GATATTAG ATTTAGCAT GTTCTATTT CAAATTTGTA TAAGTGTGA ATMAAAACT
 4854 4864 4874 4884 4894 4904
 TTTGTATAC ATCCATTTA CTTTTGTA AATATAGTA GAATTAGAG TTTTACATA

Table 1. Amino Acid Composition (mol%) Analysis

Group	<u>P23</u>	<u>dnaE</u>	<u>rpoD</u>	Ave. B.s. Proteins*
Small aliphatic (A+G)	10.2	13.3	11.2	15.0
Hydroxyl (S+T)	12.2	11.0	9.6	13.4
Acidic (D+E)	14.3	14.3	21.1	14.2
Acidic + acid amide (D+E+N+Q)	22.4	22.3	28.0	23.1
Basic (K+R+H)	14.3	16.7	16.0	14.8
Hydrophobic (L+I+V+M)	22.9	22.8	24.9	26.3
Aromatic (F+Y+W)	8.7	9.5	6.0	7.7
Charged (D+E+K+R+H)	28.6	31.0	37.1	29.0

* The amino acid composition of average B. subtilis proteins is calculated from 35 sequenced genes published up to 1985.

were fused to the 8th amino acid residue of β -galactosidase, which led us to the conclusion that ATG (855) was functionally active in vivo. Hence, we designated the first gene of the operon as P23 from these data and for the reason that its physiological function is still unknown.

The deduced amino acid sequence of P23 was examined for homology against the NBRF Protein Data Bank using the Microgenie Program (18), but no significant homology was found to any of the known proteins in the bank, indicating to us that P23 was not homologous with E. coli S21, the first gene in the σ^{70} operon (also their size difference is significant), and that P23 might be unique to B. subtilis, or that its counterpart in E. coli has not been characterized as yet. The latter case is a possibility, since a reasonable degree of homology has been found between many B. subtilis and E. coli proteins.

The deduced amino acid composition of P23 is shown in Table 1, together with those of DNA primase, σ^{43} and an average of B. subtilis proteins for comparison. One difference noted from σ^{43} , which is a highly acidic protein typical of most transcription factors (7,19,20), is that P23 is more like the average composition of B. subtilis proteins. Thus it is difficult to categorize this protein based on its amino acid

composition. We are currently raising antibody against P23 using a P23-gal fusion protein as antigen, hoping that this will provide us a tool to determine the location and possibly the function of P23 in B. subtilis.

Regulatory Features

Previously, we reported that there was no promoter activity detected within the intercistronic regions of the operon except for a weak heat shock promoter activity located at the C-terminal end of the dnaE gene (8). So we concluded that a promoter(s) should exist in front of P23 if the operon was composed of three genes as in the case of E.coli σ^{70} operon (10). By sequence analysis we did find at least two potential promoters with significant homology to the consensus sequence of B. subtilis σ^{43} promoters (1,2), which were then confirmed to function in vivo by fusing the 211 bp Sau3A fragment (609-821) to the subtilisin gene (aprA) in a promoter-probe plasmid pSB (Wong and Doi, unpublished data). These sequences are underlined in Fig. 2, and designated as P1 and P2. To our surprise, one additional promoter activity was detected when the 316 bp Sau3A fragment (829-1136) downstream of P1 and P2 was cloned in pSB. This promoter (P3) was temporally regulated in that it was not expressed until the culture reached the sporulation phase, while P1 and P2 were expressed efficiently mainly during growth. More detailed mapping and functional characterization of these promoters are in progress, and will be published elsewhere.

Earlier sequence analysis (6) allowed us to identify a sequence typical of rho independent terminator (21) immediately following the TAA stop codon of rpoD gene (underlined in Fig. 2). Recently, we confirmed its termination activity in vivo by subcloning the PvuII-AhaIII fragment (4394-4641) into a B. subtilis terminator-probe plasmid pST19 constructed in our laboratory (Wang and Doi, unpublished data). We were able to show that introduction of this 247 bp fragment reduced the activity of the indicator enzyme in the terminator probe (subtilisin, in this case) by more than 90% compared to the control (vector alone), indicating that this was a relatively strong terminator (data not shown).

Thus we have determined the presence of three genes in the

Table 2. Codon Usage of *E. subtilis* Sigma-43 Operon*

AA	Codon	P23	dnaE	rpoD	B.s.	E.c.	AA	Codon	P23	dnaE	rpoD	B.s.	E.c.
Phe	UUU	1.00	.73	.45	.67	.37	Tyr	UAU	.86	.74	.80	.62	.40
Phe	UUC	.00	.27	.55	.33	.63	Tyr	UAC	.14	.26	.20	.38	.60
Leu	UUA	.24	.20	.31	.23	.07	OCH	UAA	1.00	1.00	1.00	.57	.75
Leu	UUG	.19	.12	.15	.13	.09	AMB	UAG	.00	.00	.00	.06	.08
Leu	CUU	.29	.29	.28	.24	.07	His	CAU	.60	.78	.50	.68	.54
Leu	CUC	.00	.10	.00	.11	.07	His	CAC	.40	.22	.50	.32	.46
Leu	CUA	.05	.03	.08	.05	.02	Gln	CAA	.71	.53	.65	.56	.24
Leu	CUG	.24	.26	.18	.23	.68	Gln	CAG	.29	.47	.35	.44	.76
Ile	AUU	.62	.39	.54	.50	.36	Asn	AAU	.78	.58	.67	.53	.26
Ile	AUC	.23	.35	.40	.40	.61	Asn	AAC	.22	.42	.33	.47	.74
Ile	AUA	.15	.26	.00	.10	.03	Lys	AAA	.73	.73	.88	.75	.76
Met	AUG	1.00	1.00	1.00	1.00	1.00	Lys	AAG	.27	.27	.12	.25	.24
Val	GUU	.39	.29	.43	.20	.36	Asp	GAU	.83	.64	.58	.63	.46
Val	GUC	.23	.31	.14	.24	.15	Asp	GAC	.17	.36	.42	.37	.54
Val	GUA	.15	.26	.33	.23	.22	Glu	GAA	.81	.63	.77	.69	.73
Val	GUG	.23	.14	.10	.23	.27	Glu	GAG	.19	.37	.23	.31	.27
Ser	UCU	.20	.30	.31	.25	.23	Cys	UGU	.33	.20	.00	.50	.43
Ser	UCC	.27	.09	.06	.12	.27	Cys	UGC	.67	.80	.00	.50	.57
Ser	UCA	.20	.13	.13	.14	.07	OPL	UGA	.00	.00	.00	.37	.17
Ser	UCG	.20	.15	.06	.11	.11	Trp	UGG	1.00	1.00	1.00	1.00	1.00
Pro	CCU	.40	.53	.50	.33	.12	Arg	CGU	.15	.08	.33	.24	.56
Pro	CCC	.20	.07	.08	.09	.07	Arg	CGC	.00	.11	.17	.18	.36
Pro	CCA	.00	.20	.17	.18	.16	Arg	CGA	.23	.14	.07	.09	.03
Pro	CCG	.40	.20	.25	.40	.65	Arg	CGG	.00	.17	.17	.11	.03
Thr	ACU	.11	.10	.10	.17	.25	Ser	AGU	.07	.07	.31	.10	.06
Thr	ACC	.00	.14	.10	.14	.50	Ser	AGC	.07	.26	.13	.23	.26
Thr	ACA	.56	.29	.30	.42	.07	Arg	AGA	.54	.28	.27	.28	.01
Thr	ACG	.33	.47	.50	.27	.18	ARG	AGG	.08	.22	.00	.10	.01
Ala	GCU	.50	.37	.26	.26	.26	Gly	GGU	.10	.14	.53	.21	.48
Ala	GCC	.20	.26	.35	.19	.21	Gly	GGC	.30	.35	.32	.33	.39
Ala	GCA	.30	.22	.17	.31	.22	Gly	GGA	.40	.37	.16	.32	.05
Ala	GCG	.00	.15	.22	.24	.31	Gly	GGG	.20	.14	.00	.14	.08

- * 1) B.s. codon usage frequency of average *E. subtilis* proteins, compiled from 10,919 codons of 35 sequenced genes.
 2) E.c. codon usage frequency of average *E. coli* proteins, from reference 24.
 3) The number tabulated is the fraction usage of each codon compared with total for identical amino acid.

RNA polymerase major σ^{43} operon in the order P23, *dnaE* and *rpoD* and have physically and functionally located the promoter and terminator regions for the operon.

Codon Usage

It has been well established in *E. coli* that there is a correlation between expression level of a gene and its codon usage pattern. The more highly expressed genes show a very non-random pattern of codon usage, utilizing a restricted set of

codons which are recognized by major species of isoacceptor tRNAs, while genes which are expressed at very low levels show an almost random pattern of codon usage (22,23). The analysis of this kind of correlation in B. subtilis has been limited due to the small number of sequenced genes, and the lack of knowledge concerning the expression levels of these genes in vivo. Recently, the rapid advance of cloning and sequencing of B. subtilis genes have allowed us to compile a codon usage table for average B. subtilis proteins and compare this with the codon usage of each gene in the σ^{43} operon, and also with that of average E. coli proteins (24) (see Table 2). Comparison of the codon usages between B. subtilis and E. coli led us to the general conclusion that B. subtilis tends to more evenly or randomly distribute the codons for its amino acids than E. coli. Nevertheless, the rarely used codons in E. coli, CUA (Leu), AUA (Ile), CCC (Pro), AGG (Arg) and GGG (Gly), were also used least in B. subtilis, although the bias is not as dramatic as that in E. coli. When the usage frequencies of codons AUA, AGG and GGG in the three genes of the operon were carefully examined, striking differences were found. In rpoD, a relatively highly expressed gene in B. subtilis [2,000-10,000 molecules/cell during growth (6)] just as its counterpart in E. coli (10), these codons were not used at all, while in P23 and dnaE they were used quite frequently. Especially in dnaE, the usage frequencies for codons AUA and AGG were 0.26 and 0.22, which was much higher than those for the average B. subtilis proteins, which were 0.10 and 0.10, respectively. This preliminary analysis suggests to us that P23 and dnaE are expressed at lower levels than rpoD, which was not unexpected for dnaE since the DNA primase is required only in small amounts during DNA replication (25). Also consistent with this idea were the relative strengths of the ribosome binding sites for the three genes, which were found in an increasing order of $\Delta G^\circ = -9.2$ kcal/mol for P23, -13.8 kcal/mol for dnaE and -18.8 kcal/mol for rpoD.

Since we do not know the function of P23, we can only speculate that it may have some role in translation or act as a regulatory protein present in relatively low concentrations in the cell.




Figure 3. Schematic representation of the organization of *B. subtilis* (A) and *E. coli* (B) major σ operons. The number of base pairs is shown in parentheses for each structural unit. The organization of the *E. coli* σ^{70} operon is based on results from Burton *et al.* (10) and Lupski *et al.* (26). For simplicity, internal promoters and RNA processing sites are not shown.

Operon Organization

A comparison of the two major σ operons from *B. subtilis* and *E. coli* is illustrated in Fig. 3. Both operons contain three genes including the DNA primase and major σ genes, which are ordered in the same way. The operons are all under control of multiple promoters, indicating a complex transcription regulation system. However, significant differences do exist between them, of which the most striking is the first gene in the operon. P23 is more than twice as large as S21, and there is no sequence homology at all between them, while the middle gene products (DNA primases) are 31% homologous (8), and the last gene products (major σ factors) more than 50% homologous (6). Also different are the sizes of the intercistronic regions between the first and second genes, 33 bp in *B. subtilis* and 110 bp in *E. coli*, while that between the second and third genes are very similar in size (8,10). In *E. coli*, there is a mRNA processing site immediately following the *dnaG* gene (10), which was not found in *B. subtilis*.

DISCUSSION

We determined the nucleotide sequence of a 5 kb fragment in the *dnaE-rpoD* region of the *B. subtilis* chromosome and found three open reading frames transcribed in the same counterclockwise direction, two of which were identified as genes for the DNA primase and RNA polymerase major σ^{33} factor. The function of the first gene is still unknown. Discovery of the promoters in front of P23 and a terminator 3' to *rpoD*, and the absence of promoters in the intercistronic regions between these genes

provided strong evidence that P23, dnaE, and rpoD comprised a three gene operon. The operon was named RNA polymerase major sigma (σ^{43}) operon in analogy to that in E. coli.

The structure and organization of the σ^{43} operon resemble those of the E. coli σ^{70} operon except for the first gene. The size of P23 and its lack of amino acid sequence homology with S21 represent the most significant differences between the two operons at the molecular level. Since a reasonably high degree of homology exists between the DNA primase and σ genes, one might have expected some homology between the first genes P23 and rpsU. We are currently attempting to identify P23 by use of immunological and cell fractionation techniques to see whether P23 might be associated with the ribosome fraction of E. subtilis.

The regulation of the E. coli σ^{70} operon is very complex since it is an important operon controlling not only translation, but also DNA replication and transcription (10,26). One of its interesting features is the control mechanism(s) to keep the expression of the dnaG gene lower than its adjacent genes, rpsU and rpoD. At least four mechanisms have been proposed including an internal terminator between the first and second genes (10), a weak ribosomal binding site for dnaG (10), frequent use of rare codons (27), and a mRNA processing site between the second and third genes (10). Although no experimental data are available concerning the expression level of dnaE in vivo, a low expression is expected from its function, and its counterpart in E. coli. The results of our codon usage analysis and the comparison of ribosomal binding sites are also in good agreement with the notion that dnaE represents a weakly expressed gene, and rpoD a fairly highly expressed gene. However, besides the possible regulatory mechanisms at the translational level, it is very likely that there are also control mechanism(s) involved at the transcriptional level.

In general, E. subtilis requires a more stringent Shine-Dalgarno sequence for gene expression than E. coli (28,29). The calculated free energies of interaction of the Shine-Dalgarno regions of E. subtilis mRNAs with the 3' end of its 16s rRNA have an average value of -17 kcal/mol (30), contrasted with an

average of -11 kcal/mol for that in E. coli (31). However, the calculated free energy value for P23 gene, -9.2 kcal/mol, is far below that of the average for B. subtilis. Considering this and the codon analysis data, it is tempting to propose that P23 encodes a regulatory protein which is weakly expressed, but physiologically important. Also, since B. subtilis cells undergo differentiation and can form spores, it is possible that the cell may have evolved a unique regulation system that is absent in E. coli, and that P23 may be one of the members in that system. The possibility also exists that P23 encodes an unidentified component of the B. subtilis translation machinery which is absent or has not been identified as yet in E. coli, since it has been reported that sequences other than the Shine-Dalgarno region can affect the translation efficiency of a gene (29,32); it thus is possible that P23 might still be expressed efficiently in vivo. More experimental data are required before we can say anything conclusive about this cryptic gene.

ACKNOWLEDGEMENTS

We thank Sui-Lam Wong for use of plasmid pSB. This research was supported in part by National Institute of General Medical Sciences grant GM19673 and National Science Foundation grant PCM82-18304.

*To whom correspondence should be addressed

REFERENCES

1. Doi, R.H. (1982) Arch. Biochem. Biophys. 214, 772-781
2. Losick, R. and Pero, J. (1981) Cell 25, 582-584
3. Doi, R.H., Gitt, M., Wang, L.-F., Price, C.W. and Kawamura, F. (1984) In: Molecular Biology of Microbial differentiation (J.A. Hoch and P. Setlow, eds.), ppl47-161, American Society for Microbiology, Washington, D.C.
4. Price, C.W., Gitt, M.A. and Doi, R.H. (1983) Proc. Natl. Acad. Sci. USA 72, 1589-1593
5. Price, C.W. and Doi, R.H. (1985) Mol. Gen. Genet. 201,88-95
6. Gitt, M.A., Wang, L.-F. and Doi, R.H. (1985) J. Biol. Chem. 260, 7178-7185
7. Burton, Z., Burgess, R., Lin, J., Moore, D., Holder, S., and Gross, C. (1981) Nucl. Acids Res. 9, 2889-2903
8. Wang, L.-F., Price, C.W. and Doi, R.H. (1985) J. Biol. Chem. 260, 3368-3372
9. Smiley, B.L., Lupski, J.R., Svec, P.S., McMacken, R. and

-
- Godson, G.N. (1982) Proc. Natl. Acad. Sci. USA 79, 4550-4554
 10. Burton, Z.F., Gross, C.A., Watanabe, K.K. and Burgess, R.R. (1983) Cell 32, 335-349
 11. Messing, J., Crea, R. and Seeberg, P.H. (1981) Nucl. Acids Res. 9, 309-321
 12. Messing, J. and Vieira, J. (1982) Gene 19, 269-276
 13. Davis, R.W., Botstein, D. and Roth, J.R. (1980) Advanced Genetics: A Manual for Genetic Engineering, Cold Spring Harbor Laboratory, New York.
 14. Maniatis, R., Fritsch, E.f., and Sambrook, J. (1982) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, New York.
 15. Sanger, F., Nicklen, S. and Coulson, A.R. (1977) Proc. Natl. Acad. Sci. USA 74, 5463-5467
 16. Delaney, A.D. (1982) Nucl. Acids Res. 10, 61-67
 17. Pustell, J. and Kafatos, F. (1982) Nucl. Acids Res. 10, 51-59
 18. Queen, C. and Korn, L. (1984) Nucl. Acids Res. 12, 581-599
 19. Ishii, S., Ihara, M., Maekawa, T., Nakamura, Y., Uchida, H., and Imamoto, R. (1984) Nucl. Acids Res. 12, 3333-3342
 20. Hunt, T.P. and Magasanik, B. (1985) Proc. Natl. Acad. Sci. USA 82, 8453-8457
 21. Platt, T. and Bear, D.G. (1983) In: Gene Function in Prokaryotes (J. Beckwith, J. Davis, and J.A. Gallant, eds.), Cold Spring Harbor Laboratory, New York.
 22. Grantham, R., Gautier, C., Gouy, M., Jacobzone, M. and Mercier, R. (1981) Nucl. Acids Res. 9, r43-r74
 23. Ikemura, T. and Ozeki, H. (1982) Cold Spring Harbor Symp. Quant. Biol. 47, 1087-1097
 24. Alff-Steinberger, C. (1984) Nucl. Acids Res. 12, 2235-2241
 25. Kornberg, A. (1981) DNA Replication, Freeman & Co., San Francisco.
 26. Lupski, J.R., Smiley, B.L., Blattner, F.R. and Godson, G.N. (1982) Mol. Gen. Genet. 185, 120-128
 27. Konigsberg, W. and Godson, G.N. (1983) Proc. Natl. Acad. Sci. USA 80, 687-691
 28. McLaughlin, J.R., Murray, C.L. and Rabinowitz, J.C. (1981) J. Biol. Chem. 256, 11283-11291
 29. Band, L. and Henner, D.J. (1984) DNA 5, 17-21
 30. Murray, C.L. and Rabinowitz, J.c. (1982) J. Biol. Chem. 257, 1053-1062
 31. Gold, L., Pribnow, d., Schneider, T., Shinedling, S., Singer, B.S. and Stormo, G. (1981) Ann. Rev. Microbiol. 35, 365-403
 32. Stanssens, P., Remaut, E. and Fiers, W. (1985) Gene 36, 211-223