

Research Article

Numerical Investigation of the Time-Fractional Whitham–Broer–Kaup Equation Involving without Singular Kernel Operators

Kamsing Nonlaopon ¹, Muhammad Naeem ², Ahmed M. Zidan ^{3,4}, Rasool Shah ⁵,
Ahmed Alsanad ⁶ and Abdu Gumaei ^{6,7}

¹Department of Mathematics, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand

²Deanship of Joint First Year Umm Al-Qura University, Makkah, Saudi Arabia

³Department of Mathematics, College of Science, King Khalid University, Abha 9004, Saudi Arabia

⁴Department of Mathematics, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

⁵Department of Mathematics, Abdul Wali University Mardan, Mardan, Pakistan

⁶STC's Artificial Intelligent Chair, Department of Information Systems, College of Computer and Information Sciences, King Saud University, Riyadh 11451, Saudi Arabia

⁷Computer Science Department, Faculty of Applied Sciences, Taiz University, Taiz 6803, Yemen

Correspondence should be addressed to Rasool Shah; rasoolshah@awkum.edu.pk and Ahmed Alsanad; aaasanad@ksu.edu.sa

Received 6 May 2021; Revised 15 June 2021; Accepted 13 July 2021; Published 22 July 2021

Academic Editor: Muhammad Imran Asjad

Copyright © 2021 Kamsing Nonlaopon et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper aims to implement an analytical method, known as the Laplace homotopy perturbation transform technique, for the result of fractional-order Whitham–Broer–Kaup equations. The technique is a mixture of the Laplace transformation and homotopy perturbation technique. Fractional derivatives with Mittag-Leffler and exponential laws in sense of Caputo are considered. Moreover, this paper aims to show the Whitham–Broer–Kaup equations with both derivatives to see their difference in a real-world problem. The efficiency of both operators is confirmed by the outcome of the actual results of the Whitham–Broer–Kaup equations. Some problems have been presented to compare the solutions achieved with both fractional-order derivatives.

1. Introduction

In engineering and applied sciences and technology, fractional partial differential equations (FPDEs) containing nonlinearities define many phenomena, ranging from gravitation to dynamics. The nonlinear FPDEs are significant tools analyzed to model nonlinear dynamical behaviour in many areas such as plasma physics, mathematical biology, fluid dynamics, and solid-state physics. The widely held dynamical schemes can be denoted by an appropriate set of FPDEs. Moreover, it is well identified that FPDEs solve

mathematical models, such as Poincare conjecture and Calabi conjecture models [1–12].

It has been determined that the nonlinear development of shallow-water waves in fluid mechanics is described by a coupled system of Whitham–Broer–Kaup (WBK) equations [13]. Whitham [14], Broer [15], and Kaup [16] proposed the coupled scheme of the aforementioned equations. The aforementioned equations define shallow-water wave propagation with various spreading relations, as shown in [17]. The governing equations for the respective phenomena in classic order are provided by

$$\begin{aligned}
D_{\mathfrak{S}}^{\beta} \mu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S}) \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + g \frac{\partial \nu(\varphi, \mathfrak{S})}{\partial \varphi} &= 0, \\
D_{\mathfrak{S}}^{\beta} \nu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S}) \frac{\partial \nu(\varphi, \mathfrak{S})}{\partial \varphi} + \nu(\varphi, \mathfrak{S}) \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + p \frac{\partial^3 \mu(\varphi, \mathfrak{S})}{\partial \varphi^3} - g \frac{\partial^2 \nu(\varphi, \mathfrak{S})}{\partial \varphi^2} &= 0, \quad 0 < \beta \leq 1,
\end{aligned} \tag{1}$$

where $\mu(\varphi, \mathfrak{S})$ and $\nu(\varphi, \mathfrak{S})$ show the height and horizontal velocity that diverges from the liquid's equilibrium position, respectively, and pg are constant, signified in different diffusion powers. Over the last few decades, there has been a lot of research into solutions to such nonlinear PDEs. So, many researchers have created a variety of mathematical methods to investigate the analytical results of nonlinear PDEs. The HPM was used by Biazar and Khah [18] to solve the coupled schemes of the Burger and Brusselator problems. Amjad et al. [19] applied the solution of standard order coupled with fractional-order Whitham–Broer–Kaup equation by Laplace decomposition technique. Noor et al. [20] used the homotopy perturbation technique to investigate the results of much classical order of PDEs. Whitham–Broer–Kaup equations are used by other scholars who implemented several numerical techniques, such as residual power series technique [4], reduced differential transformation technique [21], Adomian decomposition technique [7], homotopy perturbation technique [22, 23], Lie symmetry analysis [24, 25], exp-function technique [26], G'/G -expansion technique [27], and homotopy analysis technique [28].

Fractional calculus (FC) is a new mathematical approach for describing models of nonlocal behaviour. Fractional derivatives have mathematically described many other physical problems in recent years; these representations have yielded excellent outcomes in the simulation of real-world issues. Some basic definitions of fractional operators were given by Riesz, Coimbra, Hadamard, Riemann–Liouville, Grunwald–Letnikov, Weyl, Caputo, Fabrizio, and Atangana Baleanu, among others [29–31]. To investigate the solutions of nonlinear FPDEs, some well-known techniques for finding actual results have been develop, for example, the homotopy perturbation transform technique [32, 33], the invariant subspace technique [34], the Hermite collocation technique [35], the q -homotopy analysis transform method [36], the optimal homotopy asymptotic technique [37], the homotopy analysis Sumudu transform technique [38], the Adomian decomposition technique [39], the Pade approximation and homotopy-Pade method [39], and the Sumudu transform series expansion technique [40]. The Laplace homotopy perturbation transform technique is a mixture of the homotopy perturbation technique introduced by Liao [41] and of the Laplace transformation [42].

The rest of the paper is arranged as follows. Section 2 discusses the basic definitions from fractional calculus. Section 3 is introduced to the fundamental methodology of HPTM. Sections 4 and 5 are the implementations of the techniques for the CF and AN operators. The conclusion of the work is written in Section 6.

2. Fractional Calculus

This section provides some fundamental concepts of fractional calculus.

Definition 1 (see [42]). The Liouville–Caputo operator is given by

$$D_{\mathfrak{S}}^{\beta} u(\varphi, \mathfrak{S}) = \frac{1}{\Gamma(n-\beta)} \int_0^{\mathfrak{S}} (\mathfrak{S}-\theta)^{n-1} u^{(n)}(\varphi, \theta) d\theta, \quad n-1 < \beta < n, \tag{2}$$

where $u^{(n)}(\varphi, \theta)$ is the derivative of integer n th order of $u(\varphi, \theta)$, $n \in \mathbb{N}$ and $n-1 < \beta \leq n$. If $0 < \beta \leq 1$, then we defined the Laplace transformation for the Caputo fractional derivative as follows:

$$\mathcal{L} \left[D_{\mathfrak{S}}^{\beta} u(\varphi, \mathfrak{S}) \right] (s) = s^{\beta} \mathcal{L} [u(\varphi, \mathfrak{S})] (s) - s^{\beta-1} [u(\varphi, 0)]. \tag{3}$$

Definition 2 (see [42]). The Caputo–Fabrizio operator (CF) is defined by

$$D_{\mathfrak{S}}^{\beta} u(\varphi, \mathfrak{S}) = \frac{(2-\beta)M(\beta)}{2(n-\beta)} \int_0^{\mathfrak{S}} \exp\left(-\beta \frac{(\mathfrak{S}-\theta)}{n-\beta}\right) u^{(n)}(\varphi, \theta) d\theta, \tag{4}$$

where $M(\beta)$ is a normalization form and $M(0) = M(1) = 1$. The exponential law is used as the nonsingular kernel in this fractional operator.

If $0 < \beta \leq 1$, then we define the Caputo–Fabrizio of Laplace transformation for the fractional derivative is given as

$$\mathcal{L} \left[D_{\mathfrak{S}}^{\beta} u(\varphi, \mathfrak{S}) \right] (s) = \left(\frac{s \mathcal{L} [u(\varphi, \mathfrak{S})] (s) - u(\varphi, 0)}{s + \beta(1-s)} \right). \tag{5}$$

Definition 3 (see [42]). The fractional generalized Mittag-Leffler law with the sense of Atangana–Baleanu operator is defined as follows:

$$D_{\mathfrak{S}}^{\beta} u(\varphi, \mathfrak{S}) = \frac{B(\beta)}{1-\beta} \int_0^{\mathfrak{S}} E_{\beta} \left[-\beta \frac{(\mathfrak{S}-\theta)^{\beta}}{n-\beta} \right] u^{(n)}(\varphi, \theta) d\theta, \tag{6}$$

where $B(\beta)$ is a normalization function with $B(0) = B(1) = 1$. The Mittag-Leffler law is used as a non-singular and nonlocal kernel in this fractional operator.

If $0 < \beta \leq 1$, then we express the Laplace transformation for the Atangana–Baleanu operator fractional derivative as

$$\mathcal{L}\left[D_{\mathfrak{S}}^{\beta}u(\varphi, \mathfrak{S})\right](s) = \left(\frac{s\mathcal{L}[u(\varphi, \mathfrak{S})](s) - u(\varphi, 0)}{s^{\beta}(1-\beta) + \beta}\right). \quad (7)$$

3. Implementation of the LHPTM for the Solution of Fractional Partial Differential Equation

The LHPTM is general methodology and can be written as follows.

The main procedure of this technique is defined as follows:

Step 1: let us consider the following equation:

$$\begin{aligned} \frac{\partial^{\beta}}{\partial \mathfrak{S}^{\beta}}F(\varphi, \mathfrak{S}) &= \Lambda(F(\varphi, \mathfrak{S})) + \Xi(F(\varphi, \mathfrak{S})) \\ &+ u(\varphi, \mathfrak{S}), \quad \beta = 1, 2, 3, \dots, \end{aligned} \quad (8)$$

under the initial condition

$$\begin{aligned} \frac{\partial^i}{\partial \mathfrak{S}^i}F(\varphi, 0) &= z_i(\varphi), \\ \frac{\partial^{\beta-1}}{\partial \mathfrak{S}^{\beta-1}}F(\varphi, 0) &= 0, \quad i = 0, 1, 2, 3, \dots, \beta - 2, \end{aligned} \quad (9)$$

where $u(\varphi, \mathfrak{S})$ is a known function, β is the order of the derivative, Λ represents a linear differential operator, and Ξ is the general nonlinear differential operator.

Step 2: using both sides Laplace transformation operator of (8), we obtain

$$\begin{aligned} \mathcal{L}[f(\varphi, \mathfrak{S})] &= s^{\beta}\mathcal{L}[\Lambda(F(\varphi, \mathfrak{S}))] + s^{\beta}\mathcal{L}[\Xi(F(\varphi, \mathfrak{S}))] \\ &+ s^{\beta}\mathcal{L}[u(\varphi, \mathfrak{S})], \end{aligned} \quad (10)$$

Laplace transformation is applied to Caputo–Fabrizio (5) and Atangana–Baleanu (7) operators.

Step 3: on both sides, using the inverse Laplace transformation of equation (10), we obtain

$$\begin{aligned} F(\varphi, \mathfrak{S}) &= H(\varphi, \mathfrak{S}) + \mathcal{L}^{-1}\left[s^{\beta}\mathcal{L}[\Lambda(F(\varphi, \mathfrak{S}))] \right. \\ &\left. + s^{\beta}\mathcal{L}[\Xi(F(\varphi, \mathfrak{S}))]\right]. \end{aligned} \quad (11)$$

Step 4: applying the homotopy producer, the result of the above equations in a series form is defined by

$$F(\varphi, \mathfrak{S}, k) = \sum_{n=0}^{\infty} k^n F_n(\varphi, \mathfrak{S}), \quad (12)$$

$$F(\varphi, \mathfrak{S}) = \lim_{k \rightarrow 1} F(\varphi, \mathfrak{S}, k),$$

and the nonlinear terms can be expressed as

$$\Xi F(\varphi, \mathfrak{S}) = \sum_{n=1}^{\infty} k^n h_n(f(\varphi, \mathfrak{S})), \quad (13)$$

where $k \in (0, 1]$ is an embedding parameter and $h_n(F(x, \mathfrak{S}))$ are He's polynomials that can be provided by

$$h_n(F_0, \dots, F_n) = \frac{1}{n!} \frac{\partial^n}{\partial k^n} \left[\Xi \left(\sum_{j=0}^n k^j F_j(\varphi, \mathfrak{S}) \right) \right], \quad n \in \mathbb{N}. \quad (14)$$

Finally, the LHPTM is achieved by coupling the decomposition technique which is defined by

$$\begin{aligned} \sum_{j=0}^{\infty} k^n F_n(\varphi, \mathfrak{S}) &= T(\varphi, \mathfrak{S}) + k \frac{1}{(m-1)!} \int_0^{\mathfrak{S}} (\mathfrak{S} - \tau)^{m-1} \\ &\cdot \left[u(\varphi, \mathfrak{S}) + \Lambda \left(\sum_{n=0}^{\infty} k^n F_n(\varphi, \mathfrak{S}) \right) + \sum_{n=0}^{\infty} k^n h_n(F(\varphi, \mathfrak{S})) \right] d\tau, \end{aligned} \quad (15)$$

where $T(\varphi, \mathfrak{S}) = H(\varphi, \mathfrak{S})$.

The terms, comparing with the same powers of k , produce results of many orders. The initial estimated of the approximation is $T(\varphi, \mathfrak{S})$, which is actually the Taylor series for the exact result of order β .

Using the aforementioned technique, we solved the fractional-order Whitham–Broer–Kaup equations in the Atangana–Baleanu and Caputo–Fabrizio senses using the LHPTM.

4. Implementation of Caputo–Fabrizio Operator

Example 1. Let us consider the coupled system of fractional-order WBKEs in the CFC sense:

$${}^{CF}D_{\mathfrak{S}}^{\beta}\mu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi} = 0,$$

$${}^{CF}D_{\mathfrak{S}}^{\beta}\nu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S})\frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi} + \nu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + 3\frac{\partial^3\mu(\varphi, \mathfrak{S})}{\partial\varphi^3} - \frac{\partial^2\nu(\varphi, \mathfrak{S})}{\partial\varphi^2} = 0, \quad 0 < \beta \leq 1, -1 < \mathfrak{S} \leq 1, -10 \leq \varphi \leq 10,$$
(16)

under the initial conditions,

$$\mu(\varphi, 0) = \frac{1}{2} - 8\tanh(-2\varphi),$$

$$\nu(\varphi, 0) = 16 - 16\tanh^2(-2\varphi).$$
(17)

Using the Laplace transformation to equation (16), we obtain

$$\frac{s\mathcal{L}[\mu(\varphi, \mathfrak{S})](s) - \mu(\varphi, 0)}{s + \beta(1-s)} = -\mathcal{L}\left[\mu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi}\right],$$

$$\frac{s\mathcal{L}[\nu(\varphi, \mathfrak{S})](s) - \nu(\varphi, 0)}{s + \beta(1-s)} = -\mathcal{L}\left[\mu(\varphi, \mathfrak{S})\frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi} + \nu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + 3\frac{\partial^3\mu(\varphi, \mathfrak{S})}{\partial\varphi^3} - \frac{\partial^2\nu(\varphi, \mathfrak{S})}{\partial\varphi^2}\right].$$
(18)

Simplify the above equation and use the initial conditions (17), and we obtain

$$\mathcal{L}[\mu(\varphi, \mathfrak{S})] = \frac{\mu(\varphi, 0)}{s} - \frac{s + \beta(1-s)}{s} \mathcal{L}\left[\mu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi}\right],$$

$$\mathcal{L}[\nu(\varphi, \mathfrak{S})] = \frac{\nu(\varphi, 0)}{s} - \frac{s + \beta(1-s)}{s} \mathcal{L}\left[\mu(\varphi, \mathfrak{S})\frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi} + \nu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + 3\frac{\partial^3\mu(\varphi, \mathfrak{S})}{\partial\varphi^3} - \frac{\partial^2\nu(\varphi, \mathfrak{S})}{\partial\varphi^2}\right].$$
(19)

The inverse Laplace transformation is implemented to (19), and we obtain

$$\mu(\varphi, \mathfrak{S}) = \mu(\varphi, 0) - \mathcal{L}^{-1}\left[\frac{s + \beta(1-s)}{s} \mathcal{L}\left\{\mu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + \frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi}\right\}\right],$$

$$\nu(\varphi, \mathfrak{S}) = \nu(\varphi, 0) - \mathcal{L}^{-1}\left[\frac{s + \beta(1-s)}{s} \mathcal{L}\left\{\mu(\varphi, \mathfrak{S})\frac{\partial\nu(\varphi, \mathfrak{S})}{\partial\varphi} + \nu(\varphi, \mathfrak{S})\frac{\partial\mu(\varphi, \mathfrak{S})}{\partial\varphi} + 3\frac{\partial^3\mu(\varphi, \mathfrak{S})}{\partial\varphi^3} - \frac{\partial^2\nu(\varphi, \mathfrak{S})}{\partial\varphi^2}\right\}\right].$$
(20)

The LHPTM is used in (20), and we obtain

$$\sum_{n=0}^{\infty} \mu_n(\varphi, \mathfrak{S}) = \mu(\varphi, 0) - p\mathcal{L}^{-1}\left[\frac{s + \beta(1-s)}{s} \mathcal{L}\left\{\sum_{n=0}^{\infty} p^n A_n + \frac{\partial}{\partial\varphi} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{S}) + \frac{\partial}{\partial\varphi} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{S})\right\}\right],$$

$$\sum_{n=0}^{\infty} \nu_n(\varphi, \mathfrak{S}) = \nu(\varphi, 0) - p\mathcal{L}^{-1}\left[\frac{s + \beta(1-s)}{s} \mathcal{L}\left\{\sum_{n=0}^{\infty} p^n B_n + \sum_{n=0}^{\infty} p^n C_n + 3\frac{\partial^3}{\partial\varphi^3} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{S}) - \frac{\partial^2}{\partial\varphi^2} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{S})\right\}\right].$$
(21)

The nonlinear can be found with the help of He's polynomial and can be defined as

$$\begin{aligned}
A_0 &= \mu_0 \frac{\partial \mu_0}{\partial \varphi}, \\
A_1 &= \mu_0 \frac{\partial \mu_1}{\partial \varphi} + \mu_1 \frac{\partial \mu_0}{\partial \varphi}, \\
B_0 &= \mu_0 \frac{\partial \nu_0}{\partial \beta}, \\
B_1 &= \mu_0 \frac{\partial \nu_1}{\partial \beta} + \mu_1 \frac{\partial \nu_0}{\partial \beta}, \\
C_0 &= \nu_0 \frac{\partial \mu_0}{\partial \varphi}, \\
C_1 &= \nu_0 \frac{\partial \mu_1}{\partial \varphi} + \nu_1 \frac{\partial \mu_0}{\partial \varphi}.
\end{aligned} \tag{22}$$

Comparing the coefficient of p , we have

$$\begin{aligned}
p^0: \mu_0(\varphi, \mathfrak{F}) &= \mu(\varphi, 0) = \frac{1}{2} - 8 \tanh(-2\varphi), \\
p^0: \nu_0(\varphi, \mathfrak{F}) &= \nu(\varphi, 0) = 16 - 16 \tanh^2(-2\varphi), \\
p^1: \mu_1(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ A_0 + \frac{\partial}{\partial \varphi} \mu_0(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_0(\varphi, \mathfrak{F}) \right\} \right] = -8 \sec^2 h^2(-2\varphi) [\beta \mathfrak{F} + (1-\beta)], \\
p^1: \nu_1(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ B_0 + C_0 + 3 \frac{\partial^3}{\partial \varphi^3} \mu_0(\varphi, \mathfrak{F}) - \frac{\partial^2}{\partial \varphi^2} \nu_0(\varphi, \mathfrak{F}) \right\} \right] \\
&= -32 \sec^2 h^2(-2\varphi) \tanh(-2\varphi) [\beta \mathfrak{F} + (1-\beta)], \\
p^2: \mu_2(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ A_1 + \frac{\partial}{\partial \varphi} \mu_1(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= -16 \sec^2 h^2(-2\varphi) \left(4 \sec^2 h^2(-2\varphi) - 8 \tanh^2(-2\varphi) + 3 \tanh(-2\varphi) \right) \left[(1-\beta)^2 + 2\beta(1-\beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right], \\
p^2: \nu_2(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ B_1 + C_1 + 3 \frac{\partial^3}{\partial \varphi^3} \mu_1(\varphi, \mathfrak{F}) - \frac{\partial^2}{\partial \varphi^2} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= -32 \sec^2 h^2(-2\varphi) \left\{ 40 \sec^2 h^2(-2\varphi) \tanh(-2\varphi) + 96 \tanh(-2\varphi) - 2 \tanh^2(-2\varphi) \right. \\
&\quad \left. - 32 \tanh^3(-2\varphi) - 25 \sec^2 h^2(-2\varphi) \right\} \left[(1-\beta)^2 + 2\beta(1-\beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right], \\
&\vdots
\end{aligned} \tag{23}$$

We can calculate few terms of (16) which can be written as

$$\mu(\varphi, \mathfrak{F}) = \mu_0(\varphi, \mathfrak{F}) + \mu_1(\varphi, \mathfrak{F}) + \mu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\nu(\varphi, \mathfrak{F}) = \nu_0(\varphi, \mathfrak{F}) + \nu_1(\varphi, \mathfrak{F}) + \nu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\begin{aligned} \mu(\varphi, \mathfrak{F}) &= \frac{1}{2} - 8 \tanh(-2\varphi) - 8 \sec h^2(-2\varphi) [\beta \mathfrak{F} + (1 - \beta)] - 16 \sec h^2(-2\varphi) \\ &\quad \times \left(4 \sec h^2(-2\varphi) - 8 \tanh^2(-2\varphi) + 3 \tanh(-2\varphi) \right) \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right] + \dots, \end{aligned} \quad (24)$$

$$\begin{aligned} \nu(\varphi, \mathfrak{F}) &= 16 - 16 \tanh^2(-2\varphi) - 8 \sec h^2(-2\varphi) [\beta \mathfrak{F} + (1 - \beta)] - 32 \sec h^2(-2\varphi) \{ 40 \sec h^2(-2\varphi) \tanh(-2\varphi) \\ &\quad + 96 \tanh(-2\varphi) - 2 \tanh^2(-2\varphi) - 32 \tanh^3(-2\varphi) - 25 \sec h^2(-2\varphi) \} \\ &\quad \times \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right] + \dots. \end{aligned}$$

The exact solution of (16) is

$$\begin{aligned} \mu(\varphi, \mathfrak{F}) &= \frac{1}{2} - 8 \tanh \left\{ -2 \left(\varphi - \frac{\mathfrak{F}}{2} \right) \right\}, \\ \nu(\varphi, \mathfrak{F}) &= 16 - 16 \tanh^2 \left\{ -2 \left(\varphi - \frac{\mathfrak{F}}{2} \right) \right\}. \end{aligned} \quad (25)$$

Figure 1 shows the actual and approximate solutions of $\mu(\varphi, \mathfrak{F})$ at $\beta = 1$, and Figure 2 shows the actual and

approximate solutions of $\nu(\varphi, \mathfrak{F})$ at $\beta = 1$. Figures 3 and 4 show that the first graph has a different fractional order with respect to φ and the second graph has a different fractional order with respect to \mathfrak{F} of Example 1. Tables 1 and 2 show the different fractional-order β of $\mu(\varphi, \mathfrak{F})$ and $\nu(\varphi, \mathfrak{F})$. Tables 3 and 4 show the comparisons with different methods.

Example 2. Let us consider the coupled system of fractional-order WBKEs in the CFC sense:

$${}^{CF}D_{\mathfrak{F}}^{\beta} \mu(\varphi, \mathfrak{F}) + \mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} = 0, \quad (26)$$

$${}^{CF}D_{\mathfrak{F}}^{\beta} \nu(\varphi, \mathfrak{F}) + \mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} = 0, \quad 0 < \beta \leq 1, 0 < \mathfrak{F} \leq 1, -100 \leq \varphi \leq 100,$$

under the initial conditions,

$$\begin{aligned} \mu(\varphi, 0) &= \xi - \kappa \coth[\kappa(\varphi + \theta)], \\ \nu(\varphi, 0) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)]. \end{aligned} \quad (27)$$

Using the Laplace transformation to equation (26), we obtain

$$\begin{aligned} \frac{s \mathcal{L}[\mu(\varphi, \mathfrak{F})](s) - \mu(\varphi, 0)}{s + \beta(1 - s)} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} \right], \\ \frac{s \mathcal{L}[\nu(\varphi, \mathfrak{F})](s) - \nu(\varphi, 0)}{s + \beta(1 - s)} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} \right]. \end{aligned} \quad (28)$$

Simplify the above equation and use the initial conditions (27), and we obtain




FIGURE 1: The exact and analytical solution with respect to the Caputo–Fabrizio operator of $\mu(\varphi, \mathfrak{Z})$ at $\beta = 1$ of Example 1.




FIGURE 2: The exact and analytical solution with respect to the Caputo–Fabrizio operator of $\nu(\varphi, \mathfrak{Z})$ at $\beta = 1$ of Example 1.




FIGURE 3: The different fractional-order solution with respect to the Caputo–Fabrizio operator of $\mu(\varphi, \mathfrak{I})$ at β of Example 1.




FIGURE 4: The different fractional-order solution with respect to the Caputo–Fabrizio operator of $\nu(\varphi, \mathfrak{I})$ at β of Example 1.

TABLE 1: HPTM for the approximate solution of Example 1.

(φ, \mathfrak{F})	$\mu(\varphi, \mathfrak{F})$ at $\beta = 0.5$	$\mu(\varphi, \mathfrak{F})$ at $\beta = 0.75$	$\mu(\varphi, \mathfrak{F})$ at $\beta = 1$	Exact solution
(0.1, 0.1)	-0.500817	-0.500795	-0.500782	-0.500782
(0.1, 0.3)	-0.500853	-0.500829	-0.50081	-0.50081
(0.1, 0.5)	-0.500878	-0.500857	-0.500837	-0.500837
(0.2, 0.1)	-0.49812	-0.498098	-0.498085	-0.498085
(0.2, 0.3)	-0.498154	-0.498131	-0.498112	-0.498112
(0.2, 0.5)	-0.498178	-0.498158	-0.498139	-0.498139
(0.3, 0.1)	-0.495491	-0.49547	-0.495458	-0.495458
(0.3, 0.3)	-0.495525	-0.495502	-0.495484	-0.495484
(0.3, 0.5)	-0.495548	-0.495529	-0.49551	-0.49551
(0.4, 0.1)	-0.49293	-0.492909	-0.492897	-0.492897
(0.4, 0.3)	-0.492963	-0.49294	-0.492922	-0.492922
(0.4, 0.5)	-0.492985	-0.492966	-0.492948	-0.492948
(0.5, 0.1)	-0.490433	-0.490413	-0.490401	-0.490401
(0.5, 0.3)	-0.490465	-0.490443	-0.490426	-0.490426
(0.5, 0.5)	-0.490487	-0.490469	-0.490451	-0.490451

TABLE 2: HPTM for the approximate solution of Example 1.

(φ, \mathfrak{F})	$\nu(\varphi, \mathfrak{F})$ at $\beta = 0.5$	$\nu(\varphi, \mathfrak{F})$ at $\beta = 0.75$	$\nu(\varphi, \mathfrak{F})$ at $\beta = 1$	Exact solution
(0.1, 0.1)	-0.0939215	-0.0939015	-0.09389	-0.09389
(0.1, 0.3)	-0.0939536	-0.0939319	-0.0939146	-0.0939146
(0.1, 0.5)	-0.0939757	-0.0939571	-0.0939391	-0.0939391
(0.2, 0.1)	-0.0915064	-0.091487	-0.0914759	-0.0914759
(0.2, 0.3)	-0.0915375	-0.0915165	-0.0914997	-0.0914997
(0.2, 0.5)	-0.0915589	-0.0915409	-0.0915235	-0.0915235
(0.3, 0.1)	-0.0891657	-0.0891469	-0.0891361	-0.0891361
(0.3, 0.3)	-0.0891958	-0.0891754	-0.0891592	-0.0891592
(0.3, 0.5)	-0.0892166	-0.0891992	-0.0891822	-0.0891822
(0.4, 0.1)	-0.0868965	-0.0868782	-0.0868678	-0.0868678
(0.4, 0.3)	-0.0869257	-0.0869059	-0.0868901	-0.08688901
(0.4, 0.5)	-0.0869458	-0.0869289	-0.0869125	-0.0869125
(0.5, 0.1)	-0.0846961	-0.0846784	-0.0846683	-0.0846683
(0.5, 0.3)	-0.0847244	-0.0847052	-0.0846899	-0.0846899
(0.5, 0.5)	-0.0847439	-0.0847275	-0.0847116	-0.0847116

$$\mathcal{L}[\mu(\varphi, \mathfrak{F})] = \frac{\mu(\varphi, 0)}{s} - \frac{s + \beta(1-s)}{s} \mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} \right], \quad (29)$$

$$\mathcal{L}[\nu(\varphi, \mathfrak{F})] = \frac{\nu(\varphi, 0)}{s} - \frac{s + \beta(1-s)}{s} \mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} \right].$$

The inverse Laplace transformation is implemented to (29), and we obtain

$$\begin{aligned} \mu(\varphi, \mathfrak{F}) &= \mu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} \right\} \right], \\ \nu(\varphi, \mathfrak{F}) &= \nu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} \right\} \right]. \end{aligned} \quad (30)$$

The LHPTM is used in (30), and we obtain

TABLE 3: HPTM for the approximate solution of Example 1.

(φ, \mathfrak{F})	Absolute error of ADM	Absolute error of VIM	Absolute error of OHAM	Absolute error HPTM
(0.1, 0.1)	1.04892×10^{-4}	1.23033×10^{-4}	1.07078×10^{-4}	1.67111×10^{-12}
(0.1, 0.3)	9.64474×10^{-5}	3.69597×10^{-4}	3.04565×10^{-4}	4.51196×10^{-11}
(0.1, 0.5)	8.88312×10^{-5}	6.16873×10^{-4}	4.81303×10^{-4}	2.08888×10^{-10}
(0.2, 0.1)	4.25408×10^{-4}	1.19869×10^{-4}	1.04388×10^{-4}	1.57879×10^{-12}
(0.2, 0.3)	3.91098×10^{-4}	3.60098×10^{-4}	2.97260×10^{-4}	4.26227×10^{-11}
(0.2, 0.5)	3.60161×10^{-4}	6.01006×10^{-4}	4.70138×10^{-4}	1.97328×10^{-10}
(0.3, 0.1)	9.71922×10^{-4}	1.16789×10^{-4}	1.01776×10^{-4}	1.49181×10^{-12}
(0.3, 0.3)	8.93309×10^{-4}	3.50866×10^{-4}	2.90150×10^{-4}	4.02799×10^{-11}
(0.3, 0.5)	8.22452×10^{-4}	5.85610×10^{-4}	4.59590×10^{-4}	1.86481×10^{-10}
(0.4, 0.1)	1.75596×10^{-3}	1.13829×10^{-4}	9.92418×10^{-5}	1.41043×10^{-12}
(0.4, 0.3)	1.61430×10^{-3}	3.41948×10^{-4}	2.83229×10^{-4}	3.80803×10^{-11}
(0.4, 0.5)	1.48578×10^{-3}	5.70710×10^{-4}	4.49118×10^{-4}	1.76298×10^{-10}
(0.5, 0.1)	2.79519×10^{-3}	1.10936×10^{-4}	9.67808×10^{-4}	1.33388×10^{-12}
(0.5, 0.3)	2.56714×10^{-3}	3.33274×10^{-4}	2.76492×10^{-4}	3.60145×10^{-11}
(0.5, 0.5)	2.36184×10^{-3}	5.56235×10^{-4}	4.38895×10^{-4}	1.66734×10^{-10}

TABLE 4: Comparison of different methods for the approximate solution of Example 1.

(φ, \mathfrak{F})	Absolute error of ADM	Absolute error of VIM	Absolute error of OHAM	Absolute error HPTM
(0.1, 0.1)	6.41419×10^{-3}	1.10430×10^{-4}	5.86860×10^{-5}	3.28081×10^{-12}
(0.1, 0.3)	5.99783×10^{-3}	3.31865×10^{-4}	3.04565×10^{-4}	8.85812×10^{-11}
(0.1, 0.5)	5.61507×10^{-3}	5.54071×10^{-4}	3.08812×10^{-4}	4.10099×10^{-10}
(0.2, 0.1)	1.33181×10^{-2}	1.07016×10^{-4}	5.56884×10^{-5}	3.07768×10^{-12}
(0.2, 0.3)	1.24441×10^{-2}	3.21601×10^{-4}	2.97260×10^{-4}	8.30963×10^{-11}
(0.2, 0.5)	1.16416×10^{-2}	5.36927×10^{-4}	2.92626×10^{-4}	3.84706×10^{-10}
(0.3, 0.1)	2.07641×10^{-2}	1.03737×10^{-4}	5.28609×10^{-5}	2.88849×10^{-12}
(0.3, 0.3)	1.93852×10^{-2}	3.11737×10^{-4}	2.90150×10^{-4}	7.79908×10^{-11}
(0.3, 0.5)	1.81209×10^{-2}	5.20447×10^{-4}	2.77382×10^{-4}	3.6107×10^{-10}
(0.4, 0.1)	2.88100×10^{-2}	1.00579×10^{-4}	5.01929×10^{-5}	2.71246×10^{-12}
(0.4, 0.3)	2.68724×10^{-2}	3.02245×10^{-4}	2.83229×10^{-4}	7.32356×10^{-11}
(0.4, 0.5)	2.50985×10^{-2}	5.04593×10^{-4}	2.63019×10^{-4}	3.39055×10^{-10}
(0.5, 0.1)	3.75193×10^{-2}	9.75385×10^{-5}	4.76741×10^{-5}	2.54828×10^{-12}
(0.5, 0.3)	3.49617×10^{-2}	2.93107×10^{-4}	2.76492×10^{-4}	6.88039×10^{-11}
(0.5, 0.5)	3.26239×10^{-2}	4.89335×10^{-4}	2.49480×10^{-4}	3.18537×10^{-10}

$$\sum_{n=0}^{\infty} \mu_n(\varphi, \mathfrak{F}) = \mu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n A_n + \frac{1}{2} \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \gamma_n(\varphi, \mathfrak{F}) \right\} \right], \tag{31}$$

$$\sum_{n=0}^{\infty} \nu_n(\varphi, \mathfrak{F}) = \nu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n B_n + \sum_{n=0}^{\infty} p^n C_n - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{F}) \right\} \right].$$

The nonlinear can be found with the help of He's polynomial and can be defined as

$$\begin{aligned}
A_0 &= \mu_0 \frac{\partial \mu_0}{\partial \varphi}, & \text{Comparing the coefficient of } p, \text{ we have} \\
A_1 &= \mu_0 \frac{\partial \mu_1}{\partial \varphi} + \mu_1 \frac{\partial \mu_0}{\partial \varphi}, \\
B_0 &= \mu_0 \frac{\partial \nu_0}{\partial \beta}, \\
B_1 &= \mu_0 \frac{\partial \nu_1}{\partial \beta} + \mu_1 \frac{\partial \nu_0}{\partial \beta}, \\
C_0 &= \nu_0 \frac{\partial \mu_0}{\partial \varphi}, \\
C_1 &= \nu_0 \frac{\partial \mu_1}{\partial \varphi} + \nu_1 \frac{\partial \mu_0}{\partial \varphi}.
\end{aligned} \tag{32}$$

$$p^0: \mu_0(\varphi, \mathfrak{F}) = \mu(\varphi, 0) = \xi - \kappa \coth[\kappa(\varphi + \theta)],$$

$$p^0: \nu_0(\varphi, \mathfrak{F}) = \nu(\varphi, 0) = -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)],$$

$$\begin{aligned}
p^1: \mu_1(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ A_0 + \frac{1}{2} \frac{\partial}{\partial \varphi} \mu_0(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_0(\varphi, \mathfrak{F}) \right\} \right] \\
&= -\xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] [\beta \mathfrak{F} + (1 - \beta)],
\end{aligned}$$

$$\begin{aligned}
p^1: \nu_1(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ B_0 + C_0 - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \nu_0(\varphi, \mathfrak{F}) \right\} \right] \\
&= -\xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \coth[\kappa(\varphi + \theta)] [\beta \mathfrak{F} + (1 - \beta)],
\end{aligned}$$

$$\begin{aligned}
p^2: \mu_2(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ A_1 + \frac{1}{2} \frac{\partial}{\partial \varphi} \mu_1(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= -\xi \kappa^4 \operatorname{cosech}^2[\kappa(\varphi + \theta)] (3 \coth^2([\kappa(\varphi + \theta)] - 1)) \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right] \\
&\quad + 2\xi^2 \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left(3\beta(-2\beta + 1 + \beta^2)\mathfrak{F} + \frac{\beta^3 \mathfrak{F}^3}{6} - \frac{3\beta^2(\beta - 1)\mathfrak{F}^2}{2} + 3\beta^2 - 3\beta + 1 - \beta^3 \right),
\end{aligned} \tag{33}$$

$$\begin{aligned}
p^2: \nu_2(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{s + \beta(1-s)}{s} \mathcal{L} \left\{ B_1 + C_1 - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= 2\xi \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left\{ \xi \kappa \operatorname{cosech}^2(3 \coth^2([\kappa(\varphi + \theta)] - 1)) + 2\xi \kappa \operatorname{cosech}^2 \coth^2([\kappa(\varphi + \theta)]) \right\} \\
&\quad \times \left(3\beta(-2\beta + 1 + \beta^2)\mathfrak{F} + \frac{\beta^3 \mathfrak{F}^3}{6} - \frac{3\beta^2(\beta - 1)\mathfrak{F}^2}{2} + 3\beta^2 - 3\beta + 1 - \beta^3 \right) \\
&\quad - 2\xi \coth(3 \operatorname{cosech}^2([\kappa(\varphi + \theta)] - 1)) \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{F} + \frac{\beta^2 \mathfrak{F}^2}{2} \right],
\end{aligned}$$

⋮

We can calculate few terms of (26) which can be written as

$$\mu(\varphi, \mathfrak{S}) = \mu_0(\varphi, \mathfrak{S}) + \mu_1(\varphi, \mathfrak{S}) + \mu_2(\varphi, \mathfrak{S}) + \dots,$$

$$\nu(\varphi, \mathfrak{S}) = \nu_0(\varphi, \mathfrak{S}) + \nu_1(\varphi, \mathfrak{S}) + \nu_2(\varphi, \mathfrak{S}) + \dots,$$

$$\begin{aligned} \mu(\varphi, \mathfrak{S}) &= \xi - \kappa \coth[\kappa(\varphi + \theta)] - \xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)][\beta \mathfrak{S} + (1 - \beta)] \\ &\quad - \xi \kappa^4 \operatorname{cosech}^2[\kappa(\varphi + \theta)](3 \coth^2([\kappa(\varphi + \theta)] - 1)) \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{S} + \frac{\beta^2 \mathfrak{S}^2}{2} \right] \\ &\quad + 2\xi^2 \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left(3\beta(-2\beta + 1 + \beta^2)\mathfrak{S} + \frac{\beta^3 \mathfrak{S}^3}{6} - \frac{3\beta^2(\beta - 1)\mathfrak{S}^2}{2} + 3\beta^2 - 3\beta + 1 - \beta^3 \right) + \dots, \end{aligned} \quad (34)$$

$$\begin{aligned} \nu(\varphi, \mathfrak{S}) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] - \xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \coth[\kappa(\varphi + \theta)][\beta \mathfrak{S} + (1 - \beta)] \\ &\quad + 2\xi \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left[\left\{ \xi \kappa \operatorname{cosech}^2(3 \coth^2([\kappa(\varphi + \theta)] - 1)) + 2\xi \kappa \operatorname{cosech}^2 \coth^2([\kappa(\varphi + \theta)]) \right\} \right. \\ &\quad \times \left. \left(3\beta(-2\beta + 1 + \beta^2)\mathfrak{S} + \frac{\beta^3 \mathfrak{S}^3}{6} - \frac{3\beta^2(\beta - 1)\mathfrak{S}^2}{2} + 3\beta^2 - 3\beta + 1 - \beta^3 \right) \right. \\ &\quad \left. - 2\xi \coth(3 \operatorname{cosech}^2([\kappa(\varphi + \theta)] - 1)) \left[(1 - \beta)^2 + 2\beta(1 - \beta)\mathfrak{S} + \frac{\beta^2 \mathfrak{S}^2}{2} \right] \right] + \dots \end{aligned}$$

The exact solution of (26) is

$$\begin{aligned} \mu(\varphi, \mathfrak{S}) &= \xi - \kappa \coth[\kappa(\varphi + \theta - \xi \mathfrak{S})], \\ \nu(\varphi, \mathfrak{S}) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta - \xi \mathfrak{S})]. \end{aligned} \quad (35)$$

5. Implementation of Atangana–Baleanu Operator

Example 3. Let us consider the coupled system of fractional-order WBKEs in the ABC sense:

$$\begin{aligned} {}^{AB}D_{\mathfrak{S}}^{\beta} \mu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S}) \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{S})}{\partial \varphi} &= 0, \\ {}^{AB}D_{\mathfrak{S}}^{\beta} \nu(\varphi, \mathfrak{S}) + \mu(\varphi, \mathfrak{S}) \frac{\partial \nu(\varphi, \mathfrak{S})}{\partial \varphi} + \nu(\varphi, \mathfrak{S}) \frac{\partial \mu(\varphi, \mathfrak{S})}{\partial \varphi} + 3 \frac{\partial^3 \mu(\varphi, \mathfrak{S})}{\partial \varphi^3} - \frac{\partial^2 \nu(\varphi, \mathfrak{S})}{\partial \varphi^2} &= 0, \quad 0 < \beta \leq 1, -1 < \mathfrak{S} \leq 1, -10 \leq \varphi \leq 10, \end{aligned} \quad (36)$$

under the initial conditions,

$$\mu(\varphi, 0) = \frac{1}{2} - 8 \tanh(-2\varphi), \quad (37)$$

$$\nu(\varphi, 0) = 16 - 16 \tanh^2(-2\varphi).$$

Using the Laplace transformation to (36), we obtain

$$\begin{aligned} \frac{B(\beta)}{1-\beta} \frac{s \mathcal{L}[\mu(\varphi, \mathfrak{T})](s) - s^{\beta-1} \mu(\varphi, 0)}{s + (\beta/(1-\beta))} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} \right], \\ \frac{B(\beta)}{1-\beta} \frac{s \mathcal{L}[\nu(\varphi, \mathfrak{T})](s) - s^{\beta-1} \nu(\varphi, 0)}{s + (\beta/(1-\beta))} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{T}) \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} + \nu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + 3 \frac{\partial^3 \mu(\varphi, \mathfrak{T})}{\partial \varphi^3} - \frac{\partial^2 \nu(\varphi, \mathfrak{T})}{\partial \varphi^2} \right]. \end{aligned} \quad (38)$$

Simplify the above equation and use the initial conditions (37), and we obtain

$$\begin{aligned} \mathcal{L}[\mu(\varphi, \mathfrak{T})] &= \frac{\mu(\varphi, 0)}{s} - \frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left[\mu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} \right], \\ \mathcal{L}[\nu(\varphi, \mathfrak{T})] &= \frac{\nu(\varphi, 0)}{s} - \frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left[\mu(\varphi, \mathfrak{T}) \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} + \nu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + 3 \frac{\partial^3 \mu(\varphi, \mathfrak{T})}{\partial \varphi^3} - \frac{\partial^2 \nu(\varphi, \mathfrak{T})}{\partial \varphi^2} \right]. \end{aligned} \quad (39)$$

The inverse Laplace transformation implement to (39), we obtain

$$\begin{aligned} \mu(\varphi, \mathfrak{T}) &= \mu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} \right\} \right], \\ \nu(\varphi, \mathfrak{T}) &= \nu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{T}) \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} + \nu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + 3 \frac{\partial^3 \mu(\varphi, \mathfrak{T})}{\partial \varphi^3} - \frac{\partial^2 \nu(\varphi, \mathfrak{T})}{\partial \varphi^2} \right\} \right]. \end{aligned} \quad (40)$$

The LHPTM is used in (40), and we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \mu_n(\varphi, \mathfrak{T}) &= \mu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n A_n + \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{T}) + \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{T}) \right\} \right], \\ \sum_{n=0}^{\infty} \nu_n(\varphi, \mathfrak{T}) &= \nu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n B_n + \sum_{n=0}^{\infty} p^n C_n + 3 \frac{\partial^3}{\partial \varphi^3} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{T}) - \frac{\partial^2}{\partial \varphi^2} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{T}) \right\} \right]. \end{aligned} \quad (41)$$

The nonlinear can be found with the help of He's polynomial and can be defined as

$$\begin{aligned}
A_0 &= \mu_0 \frac{\partial \mu_0}{\partial \varphi}, & \text{Comparing the coefficient of } p, \text{ we have} \\
A_1 &= \mu_0 \frac{\partial \mu_1}{\partial \varphi} + \mu_1 \frac{\partial \mu_0}{\partial \varphi}, \\
B_0 &= \mu_0 \frac{\partial \nu_0}{\partial \beta}, \\
B_1 &= \mu_0 \frac{\partial \nu_1}{\partial \beta} + \mu_1 \frac{\partial \nu_0}{\partial \beta}, \\
C_0 &= \nu_0 \frac{\partial \mu_0}{\partial \varphi}, \\
C_1 &= \nu_0 \frac{\partial \mu_1}{\partial \varphi} + \nu_1 \frac{\partial \mu_0}{\partial \varphi}.
\end{aligned} \tag{42}$$

$$p^0: \mu_0(\varphi, \mathfrak{F}) = \mu(\varphi, 0) = \frac{1}{2} - 8 \tanh(-2\varphi),$$

$$p^0: \nu_0(\varphi, \mathfrak{F}) = \nu(\varphi, 0) = 16 - 16 \tanh^2(-2\varphi),$$

$$\begin{aligned}
p^1: \mu_1(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ A_0 + \frac{\partial}{\partial \varphi} \mu_0(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_0(\varphi, \mathfrak{F}) \right\} \right] \\
&= -8 \sec h^2(-2\varphi) \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right],
\end{aligned}$$

$$\begin{aligned}
p^1: \nu_1(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ B_0 + C_0 + 3 \frac{\partial^3}{\partial \varphi^3} \mu_0(\varphi, \mathfrak{F}) - \frac{\partial^2}{\partial \varphi^2} \nu_0(\varphi, \mathfrak{F}) \right\} \right] \\
&= -32 \sec h^2(-2\varphi) \tanh(-2\varphi) \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right],
\end{aligned}$$

$$\begin{aligned}
p^2: \mu_2(\varphi, \mathfrak{F}) &= -\mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ A_1 + \frac{\partial}{\partial \varphi} \mu_1(\varphi, \mathfrak{F}) + \frac{\partial}{\partial \varphi} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= -16 \sec h^2(-2\varphi) (4 \sec h^2(-2\varphi) - 8 \tanh^2(-2\varphi) + 3 \tanh(-2\varphi)) \\
&\quad \times \frac{1}{B^2(\beta)} \left[(1-\beta)^2 + \frac{2\beta(1-\beta)\mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right],
\end{aligned} \tag{43}$$

$$\begin{aligned}
p^2: \nu_2(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ B_1 + C_1 + 3 \frac{\partial^3}{\partial \varphi^3} \mu_1(\varphi, \mathfrak{F}) - \frac{\partial^2}{\partial \varphi^2} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= -32 \sec h^2(-2\varphi) \{ 40 \sec h^2(-2\varphi) \tanh(-2\varphi) + 96 \tanh(-2\varphi) - 2 \tanh^2(-2\varphi) \\
&\quad - 32 \tanh^3(-2\varphi) - 25 \sec h^2(-2\varphi) \} \left[(1-\beta)^2 + \frac{2\beta(1-\beta)\mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right],
\end{aligned}$$

$$\vdots$$

We can calculate few terms of (36) which can be written as

$$\mu(\varphi, \mathfrak{F}) = \mu_0(\varphi, \mathfrak{F}) + \mu_1(\varphi, \mathfrak{F}) + \mu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\nu(\varphi, \mathfrak{F}) = \nu_0(\varphi, \mathfrak{F}) + \nu_1(\varphi, \mathfrak{F}) + \nu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\begin{aligned} \mu(\varphi, \mathfrak{F}) &= \frac{1}{2} - 8 \tanh(-2\varphi) - 8 \sec h^2(-2\varphi) \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right] - 16 \sec h^2(-2\varphi) \\ &\quad \left(4 \sec h^2(-2\varphi) - 8 \tanh^2(-2\varphi) + 3 \tanh(-2\varphi) \right) \frac{1}{B^2(\beta)} \left[(1-\beta)^2 + \frac{2\beta(1-\beta)\mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right] + \dots, \\ \nu(\varphi, \mathfrak{F}) &= 16 - 16 \tanh^2(-2\varphi) - 8 \sec h^2(-2\varphi) \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right] - 32 \sec h^2(-2\varphi) \\ &\quad \times \left\{ 40 \sec h^2(-2\varphi) \tanh(-2\varphi) + 96 \tanh(-2\varphi) - 2 \tanh^2(-2\varphi) - 32 \tanh^3(-2\varphi) - 25 \sec h^2(-2\varphi) \right\} \\ &\quad \times \left[(1-\beta)^2 + \frac{2\beta(1-\beta)\mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right] + \dots. \end{aligned} \quad (44)$$

The exact solution of (36) is

$$\begin{aligned} \mu(\varphi, \mathfrak{F}) &= \frac{1}{2} - 8 \tanh \left\{ -2 \left(\varphi - \frac{\mathfrak{F}}{2} \right) \right\}, \\ \nu(\varphi, \mathfrak{F}) &= 16 - 16 \tanh^2 \left\{ -2 \left(\varphi - \frac{\mathfrak{F}}{2} \right) \right\}. \end{aligned} \quad (45)$$

Figure 5 shows the actual and approximate solutions of $\mu(\varphi, \mathfrak{F})$ at $\beta = 1$, and Figure 6 shows the actual and

approximate solutions of $\nu(\varphi, \mathfrak{F})$ at $\beta = 1$. Figures 7 and 8 show that the first graph has a different fractional order with respect to φ and the second graph has a different fractional order with respect to \mathfrak{F} of Example 3.

Example 4. Let us consider the coupled system of fractional-order WBKEs in the ABC sense:

$$\begin{aligned} {}^{ABC}D_{\mathfrak{F}}^\beta \mu(\varphi, \mathfrak{F}) + \mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} &= 0, \\ {}^{ABC}D_{\mathfrak{F}}^\beta \nu(\varphi, \mathfrak{F}) + \mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} &= 0, \quad 0 < \beta \leq 1, -1 < \mathfrak{F} \leq 1, -10 \leq \varphi \leq 10, \end{aligned} \quad (46)$$

under the initial conditions,

$$\begin{aligned} \mu(\varphi, 0) &= \xi - \kappa \coth[\kappa(\varphi + \theta)], \\ \nu(\varphi, 0) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)]. \end{aligned} \quad (47)$$

Using the Laplace transformation to (46), we obtain

$$\begin{aligned} \frac{B(\beta)}{1-\beta} \frac{s \mathcal{L}[\mu(\varphi, \mathfrak{F})](s) - s^{\beta-1} \mu(\varphi, 0)}{s + (\beta/(1-\beta))} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} \right], \\ \frac{B(\beta)}{1-\beta} \frac{s \mathcal{L}[\nu(\varphi, \mathfrak{F})](s) - s^{\beta-1} \nu(\varphi, 0)}{s + (\beta/(1-\beta))} &= -\mathcal{L} \left[\mu(\varphi, \mathfrak{F}) \frac{\partial \nu(\varphi, \mathfrak{F})}{\partial \varphi} + \nu(\varphi, \mathfrak{F}) \frac{\partial \mu(\varphi, \mathfrak{F})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{F})}{\partial \varphi^2} \right]. \end{aligned} \quad (48)$$




FIGURE 5: The exact and analytical solution with respect to the Atangana–Baleanu operator of $\mu(\varphi, \mathfrak{Z})$ at $\beta = 1$ of Example 3.




FIGURE 6: The exact and analytical solution with respect to the Atangana–Baleanu operator of $\nu(\varphi, \mathfrak{Z})$ at $\beta = 1$ of Example 3.

Simplifying the above equation and using the initial conditions (47), we obtain

$$\begin{aligned} \mathcal{L}[\mu(\varphi, \mathfrak{Z})] &= \frac{\mu(\varphi, 0)}{s} - \frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left[\mu(\varphi, \mathfrak{Z}) \frac{\partial \mu(\varphi, \mathfrak{Z})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{Z})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{Z})}{\partial \varphi} \right], \\ \mathcal{L}[\nu(\varphi, \mathfrak{Z})] &= \frac{\nu(\varphi, 0)}{s} - \frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left[\mu(\varphi, \mathfrak{Z}) \frac{\partial \nu(\varphi, \mathfrak{Z})}{\partial \varphi} + \nu(\varphi, \mathfrak{Z}) \frac{\partial \mu(\varphi, \mathfrak{Z})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{Z})}{\partial \varphi^2} \right]. \end{aligned} \tag{49}$$

The inverse Laplace transformation is implemented to (49), and we obtain




FIGURE 7: The different fractional-order solution with respect to the Atangana–Baleanu operator of $\mu(\varphi, \mathfrak{T})$ at β of Example 3.




FIGURE 8: The different fractional-order solution with respect to the Atangana–Baleanu operator of $\nu(\varphi, \mathfrak{T})$ at β of Example 3.

$$\begin{aligned} \mu(\varphi, \mathfrak{T}) &= \mu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{1}{2} \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} + \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} \right\} \right], \\ \nu(\varphi, \mathfrak{T}) &= \nu(\varphi, 0) - \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \mu(\varphi, \mathfrak{T}) \frac{\partial \nu(\varphi, \mathfrak{T})}{\partial \varphi} + \nu(\varphi, \mathfrak{T}) \frac{\partial \mu(\varphi, \mathfrak{T})}{\partial \varphi} - \frac{1}{2} \frac{\partial^2 \nu(\varphi, \mathfrak{T})}{\partial \varphi^2} \right\} \right]. \end{aligned} \quad (50)$$

The LHPTM is used in (50), and we obtain

$$\begin{aligned} \sum_{n=0}^{\infty} \mu_n(\varphi, \mathfrak{S}) &= \mu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n A_n + \frac{1}{2} \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \mu_n(\varphi, \mathfrak{S}) + \frac{\partial}{\partial \varphi} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{S}) \right\} \right], \\ \sum_{n=0}^{\infty} \nu_n(\varphi, \mathfrak{S}) &= \nu(\varphi, 0) - p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ \sum_{n=0}^{\infty} p^n B_n + \sum_{n=0}^{\infty} p^n C_n - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \sum_{n=0}^{\infty} p^n \nu_n(\varphi, \mathfrak{S}) \right\} \right]. \end{aligned} \quad (51)$$

The nonlinear can be found with the help of He's polynomial and can be defined as

$$\begin{aligned} A_0 &= \mu_0 \frac{\partial \mu_0}{\partial \varphi}, \\ A_1 &= \mu_0 \frac{\partial \mu_1}{\partial \varphi} + \mu_1 \frac{\partial \mu_0}{\partial \varphi}, \\ B_0 &= \mu_0 \frac{\partial \nu_0}{\partial \beta}, \\ B_1 &= \mu_0 \frac{\partial \nu_1}{\partial \beta} + \mu_1 \frac{\partial \nu_0}{\partial \beta}, \\ C_0 &= \nu_0 \frac{\partial \mu_0}{\partial \varphi}, \\ C_1 &= \nu_0 \frac{\partial \mu_1}{\partial \varphi} + \nu_1 \frac{\partial \mu_0}{\partial \varphi}. \end{aligned} \quad (52)$$

Comparing the coefficient of p , we have

$$p^0: \mu_0(\varphi, \mathfrak{S}) = \mu(\varphi, 0) = \xi - \kappa \coth[\kappa(\varphi + \theta)],$$

$$p^0: \nu_0(\varphi, \mathfrak{S}) = \nu(\varphi, 0) = -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)],$$

$$\begin{aligned} p^1: \mu_1(\varphi, \mathfrak{S}) &= -\mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ A_0 + \frac{1}{2} \frac{\partial}{\partial \varphi} \mu_0(\varphi, \mathfrak{S}) + \frac{\partial}{\partial \varphi} \nu_0(\varphi, \mathfrak{S}) \right\} \right] \\ &= -\xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{S}^\beta}{\Gamma(\beta + 1)} + (1 - \beta) \right], \end{aligned}$$

$$\begin{aligned} p^1: \nu_1(\varphi, \mathfrak{S}) &= -p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ B_0 + C_0 - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \nu_0(\varphi, \mathfrak{S}) \right\} \right] \\ &= -\xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \coth[\kappa(\varphi + \theta)] \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{S}^\beta}{\Gamma(\beta + 1)} + (1 - \beta) \right], \end{aligned}$$

$$\begin{aligned} p^2: \mu_2(\varphi, \mathfrak{S}) &= -\mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ A_1 + \frac{1}{2} \frac{\partial}{\partial \varphi} \mu_1(\varphi, \mathfrak{S}) + \frac{\partial}{\partial \varphi} \nu_1(\varphi, \mathfrak{S}) \right\} \right] \\ &= -\xi \kappa^4 \operatorname{cosech}^2[\kappa(\varphi + \theta)] (3 \coth^2([\kappa(\varphi + \theta)] - 1)) \frac{1}{B^2} \left[(1 - \beta)^2 + \frac{2\beta(1 - \beta)\mathfrak{S}^\beta}{\Gamma(\beta + 1)} + \frac{\beta^2 \mathfrak{S}^{2\beta}}{\Gamma(2\beta + 1)} \right] \\ &\quad + 2\xi^2 \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \frac{1}{B^3} \left[(1 - \beta)^3 + \frac{3\beta(1 - \beta)^2 \mathfrak{S}^\beta}{\Gamma(2\beta + 1)} + \frac{3\beta^2(1 - \beta)\mathfrak{S}^{2\beta}}{\Gamma(2\beta + 1)} + \frac{\beta^3 \mathfrak{S}^{3\beta}}{\Gamma(3\beta + 1)} \right], \end{aligned}$$

$$\begin{aligned}
p^2: \nu_2(\varphi, \mathfrak{F}) &= -p \mathcal{L}^{-1} \left[\frac{(1-\beta)s^\beta + \beta}{B(\beta)s^\beta} \mathcal{L} \left\{ B_1 + C_1 - \frac{1}{2} \frac{\partial^2}{\partial \varphi^2} \nu_1(\varphi, \mathfrak{F}) \right\} \right] \\
&= 2\xi \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left\{ \xi \kappa \operatorname{cosech}^2(3 \coth^2([\kappa(\varphi + \theta)] - 1)) + 2\xi \kappa \operatorname{cosech}^2 \coth^2([\kappa(\varphi + \theta)]) \right\} \\
&\quad \times \frac{1}{B^3} \left[(1-\beta)^3 + \frac{3\beta(1-\beta)^2 \mathfrak{F}^\beta}{\Gamma(2\beta+1)} + \frac{3\beta^2(1-\beta) \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} + \frac{\beta^3 \mathfrak{F}^{3\beta}}{\Gamma(3\beta+1)} \right] \\
&\quad - 2\xi \coth(3 \operatorname{cosech}^2([\kappa(\varphi + \theta)] - 1)) \frac{1}{B^2} \left[(1-\beta)^2 + \frac{2\beta(1-\beta) \mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right], \\
&\quad \vdots
\end{aligned} \tag{53}$$

We can calculate few terms of (46) which can be written as

$$\mu(\varphi, \mathfrak{F}) = \mu_0(\varphi, \mathfrak{F}) + \mu_1(\varphi, \mathfrak{F}) + \mu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\nu(\varphi, \mathfrak{F}) = \nu_0(\varphi, \mathfrak{F}) + \nu_1(\varphi, \mathfrak{F}) + \nu_2(\varphi, \mathfrak{F}) + \dots,$$

$$\begin{aligned}
\mu(\varphi, \mathfrak{F}) &= \xi - \kappa \coth[\kappa(\varphi + \theta)] - \xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right] \\
&\quad - \xi \kappa^4 \operatorname{cosech}^2[\kappa(\varphi + \theta)] (3 \coth^2([\kappa(\varphi + \theta)] - 1)) \frac{1}{B^2} \left[(1-\beta)^2 + \frac{2\beta(1-\beta) \mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right] \\
&\quad + 2\xi^2 \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \frac{1}{B^3} \left[(1-\beta)^3 + \frac{3\beta(1-\beta)^2 \mathfrak{F}^\beta}{\Gamma(2\beta+1)} + \frac{3\beta^2(1-\beta) \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} + \frac{\beta^3 \mathfrak{F}^{3\beta}}{\Gamma(3\beta+1)} \right] + \dots,
\end{aligned} \tag{54}$$

$$\begin{aligned}
\nu(\varphi, \mathfrak{F}) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] - \xi \kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \coth[\kappa(\varphi + \theta)] \frac{1}{B(\beta)} \left[\frac{\beta \mathfrak{F}^\beta}{\Gamma(\beta+1)} + (1-\beta) \right] \\
&\quad + 2\xi \kappa^5 \operatorname{cosech}^2[\kappa(\varphi + \theta)] \left\{ \xi \kappa \operatorname{cosech}^2(3 \coth^2([\kappa(\varphi + \theta)] - 1)) + 2\xi \kappa \operatorname{cosech}^2 \coth^2([\kappa(\varphi + \theta)]) \right\} \\
&\quad \times \frac{1}{B^3} \left[(1-\beta)^3 + \frac{3\beta(1-\beta)^2 \mathfrak{F}^\beta}{\Gamma(2\beta+1)} + \frac{3\beta^2(1-\beta) \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} + \frac{\beta^3 \mathfrak{F}^{3\beta}}{\Gamma(3\beta+1)} \right] - 2\xi \coth(3 \operatorname{cosech}^2([\kappa(\varphi + \theta)] - 1)) \\
&\quad \times \frac{1}{B^2} \left[(1-\beta)^2 + \frac{2\beta(1-\beta) \mathfrak{F}^\beta}{\Gamma(\beta+1)} + \frac{\beta^2 \mathfrak{F}^{2\beta}}{\Gamma(2\beta+1)} \right] + \dots.
\end{aligned}$$

The exact solution of (46) is

$$\begin{aligned}
\mu(\varphi, \mathfrak{F}) &= \xi - \kappa \coth[\kappa(\varphi + \theta - \xi \mathfrak{F})], \\
\nu(\varphi, \mathfrak{F}) &= -\kappa^2 \operatorname{cosech}^2[\kappa(\varphi + \theta - \xi \mathfrak{F})].
\end{aligned} \tag{55}$$

6. Conclusions

In this paper, the LHPTM was considered to achieve an analytical result for the fractional-order Whitham–Broer–Kaup equations considering the Caputo–Fabrizio and

Atangana–Baleanu operators. Analytical results were obtained for different fractional order β . Both suggested operators have shown to be critical mathematical tools for scientists working in numerous fields of applied sciences. The polynomial expansion well-thought-out in the LHPTM permits to achieve an infinite series result for the Whitham–Broer–Kaup equations. This technique established a general system to achieve fractional-order models' analytical results, and the results are obtained in the series form, which converges quickly; the LHPTM is used for investigating other nonlinear fractional systems of partial differential equations.

Abbreviation

HPTM: Homotopy perturbation transform method
 IT: Laplace transform
 FPDEs: Fractional partial differential equations
 FC: Fractional calculus
 WBKEs: Whitham–Broer–Kaup equations
 ADM: Adomian decomposition method.

Data Availability

The numerical data used to support the findings of this study are included within the article.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this article.

Acknowledgments

One of the co-authors (A. M. Zidan) extend his appreciation to the Deanship of Scientific Research at King Khalid University, Abha 61413, Saudi Arabia, for funding this work through research groups program, under Grant no. R.G.P-2/142/42. The authors are grateful to the Deanship of Scientific Research, King Saud University, for funding through Vice Deanship of Scientific Research Chairs..

References

- [1] F. Xie, Z. Yan, and H. Zhang, "Explicit and exact traveling wave solutions of Whitham-Broer-Kaup shallow water equations," *Physics Letters A*, vol. 285, no. 1-2, pp. 76–80, 2001.
- [2] M. I. Asjad, M. Aleem, A. Ahmadian, S. Salahshour, M. Ferrara, and M. Ferrara, "New trends of fractional modeling and heat and mass transfer investigation of (SWCNTs and MWCNTs)-CMC based nanofluids flow over inclined plate with generalized boundary conditions," *Chinese Journal of Physics*, vol. 66, pp. 497–516, 2020.
- [3] A. A. Alderremy, K. M. Saad, S. Aly, D. Kumar, and J. Singh, "New models of fractional blood ethanol and two cell cubic autocatalator reaction equations," *Mathematical Methods in the Applied Sciences*, vol. 43, 2021.
- [4] L. Wang and X. Chen, "Approximate analytical solutions of time fractional Whitham-Broer-Kaup equations by a residual power series method," *Entropy*, vol. 17, no. 9, pp. 6519–6533, 2015.
- [5] M. Aleem, M. Imran Asjad, M. S. R. Chowdhury, and A. Hussanan, "Analysis of mathematical model of fractional viscous fluid through a vertical rectangular channel," *Chinese Journal of Physics*, vol. 61, pp. 336–350, 2019.
- [6] K. M. Saad, M. Alqhtani, and M. Alqhtani, "Numerical simulation of the fractal-fractional reaction diffusion equations with general nonlinear," *AIMS Mathematics*, vol. 6, no. 4, pp. 3788–3804, 2021.
- [7] S. M. El-Sayed and D. Kaya, "Exact and numerical traveling wave solutions of Whitham-Broer-Kaup equations," *Applied Mathematics and Computation*, vol. 167, no. 2, pp. 1339–1349, 2005.
- [8] N. A. Shah, I. Khan, M. Aleem, and M. A. Imran, "Influence of magnetic field on double convection problem of fractional viscous fluid over an exponentially moving vertical plate: new trends of Caputo time-fractional derivative model," *Advances in Mechanical Engineering*, vol. 11, no. 7, 2019.
- [9] K. M. Saad, M. Alqhtani, and J. F. Gómez-Aguilar, "Fractal-fractional study of the hepatitis C virus infection model," *Results in Physics*, vol. 19, Article ID 103555, 2020.
- [10] J. Ahmad, M. Mushtaq, and N. Sajjad, "Exact solution of whitham-broer-kaup shallow water wave equations," *Journal of Science and Arts*, vol. 15, p. 1, 2015.
- [11] M. Ahmad, M. A. Imran, M. Aleem, and I. Khan, "A comparative study and analysis of natural convection flow of MHD non-Newtonian fluid in the presence of heat source and first-order chemical reaction," *Journal of Thermal Analysis and Calorimetry*, vol. 137, no. 5, pp. 1783–1796, 2019.
- [12] H. M. Srivastava and K. M. Saad, "A comparative study of the fractional-order clock chemical model," *Mathematics*, vol. 8, no. 9, p. 1436, 2020.
- [13] B. A. Kupersmidt, "Mathematics of dispersive water waves," *Communications in Mathematical Physics*, vol. 99, no. 1, pp. 51–73, 1985.
- [14] G. B. Whitham, "Variational methods and applications to water waves," *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, vol. 299, no. 1456, pp. 6–25, 1967.
- [15] L. J. F. Broer, "Approximate equations for long water waves," *Applied Scientific Research*, vol. 31, no. 5, pp. 377–395, 1975.
- [16] D. J. Kaup, "A higher-order water-wave equation and the method for solving it," *Progress of Theoretical Physics*, vol. 54, no. 2, pp. 396–408, 1975.
- [17] D. D. Ganji, H. B. Rokni, M. G. Sfahani, and S. S. Ganji, "Approximate traveling wave solutions for coupled Whitham-Broer-Kaup shallow water," *Advances in Engineering Software*, vol. 41, no. 7-8, pp. 956–961, 2010.
- [18] J. Biazar and H. Aminikhah, "Study of convergence of homotopy perturbation method for systems of partial differential equations," *Computers & Mathematics with Applications*, vol. 58, no. 11-12, pp. 2221–2230, 2009.
- [19] A. Ali, K. Shah, and R. A. Khan, "Numerical treatment for traveling wave solutions of fractional Whitham-Broer-Kaup equations," *Alexandria Engineering Journal*, vol. 57, no. 3, pp. 1991–1998, 2018.
- [20] S. T. Mohyud-Din and M. A. Noor, "Homotopy perturbation method for solving partial differential equations," *Zeitschrift Fur Naturforschung A*, vol. 64, no. 3-4, pp. 157–170, 2009.
- [21] S. Saha Ray, "A novel method for travelling wave solutions of fractional Whitham-Broer-Kaup, fractional modified Bousinesq and fractional approximate long wave equations in shallow water," *Mathematical Methods in the Applied Sciences*, vol. 38, no. 7, pp. 1352–1368, 2015.
- [22] S. T. Mohyud-Din, A. Yıldırım, G. Demirli, and G. Demirli, "Traveling wave solutions of Whitham-Broer-Kaup equations by homotopy perturbation method," *Journal of King Saud University-Science*, vol. 22, no. 3, pp. 173–176, 2010.
- [23] A. Kadem and B. Dumitru, "On fractional coupled Whitham-Broer-Kaup equations," *Romanian Journal of Physics*, vol. 56, no. 5-6, pp. 629–635, 2011.
- [24] H. R. Ghehsareh, M. Ahmad, and Z. Ali, "Lie symmetry analysis and conservation laws for time fractional coupled Whitham-Broer-Kaup equations," *UPB Scientific Bulletin, Series A: Applied Mathematics and Physics*, vol. 80, no. 3, pp. 153–168, 2018.

- [25] Z. Zhang, X. Yong, and Y. Chen, "Symmetry analysis for Whitham-Broer-Kaup equations," *Journal of Nonlinear Mathematical Physics*, vol. 15, no. 4, pp. 383–397, 2008.
- [26] Y. Wang, Y.-F. Zhang, Z.-J. Liu, and M. Iqbal, "A fractional Whitham-Broer-Kaup equation and its possible application to tsunami prevention," *Thermal Science*, vol. 21, no. 4, pp. 1847–1855, 2017.
- [27] S. Arshed and M. Sadia, " G'/G -Expansion method: new traveling wave solutions for some nonlinear fractional partial differential equations," *Optical and Quantum Electronics*, vol. 50, no. 3, pp. 1–20, 2018.
- [28] A. Rani, Q. M. Ul-Hassan, M. Ashraf, K. Ayub, and M. Y. Khan, "A novel technique for solving nonlinear WBK equations of fractional-order," *Journal of Science and Arts*, vol. 18, no. 2, pp. 301–316, 2018.
- [29] K. M. Owolabi and A. Atangana, "Analysis and application of new fractional Adams-Bashforth scheme with Caputo-Fabrizio derivative," *Chaos, Solitons & Fractals*, vol. 105, pp. 111–119, 2017.
- [30] M. Naeem, A. M. Zidan, K. Nonlaopon, M. I. Syam, Z. Al-Zhour, and R. Shah, "A new analysis of fractional-order equal-width equations via novel techniques," *Symmetry*, vol. 13, no. 5, p. 886, 2021.
- [31] M. Caputo and M. Fabrizio, "A new definition of fractional derivative without singular kernel," *Progress in Fractional Differentiation and Applications*, vol. 1, no. 2, pp. 1–13, 2015.
- [32] R. K. Pandey and H. K. Mishra, "Homotopy analysis Sumudu transform method for time-fractional third order dispersive partial differential equation," *Advances in Computational Mathematics*, vol. 43, no. 2, pp. 365–383, 2017.
- [33] J. Vahidi, "The combined Laplace-homotopy analysis method for partial differential equations," *Journal of Mathematics and Computer Science*, vol. 16, pp. 88–102, 2016.
- [34] S. Choudhary and V. Daftardar-Gejji, "Invariant subspace method: a tool for solving fractional partial differential equations," *Fractional Calculus and Applied Analysis*, vol. 20, no. 2, pp. 477–493, 2017.
- [35] N. A. Pirim and F. Ayaz, "A new technique for solving fractional order systems: Hermite collocation method," *Applied Mathematics*, vol. 7, no. 18, pp. 2307–2323, 2016.
- [36] K. M. Saad, "Comparing the Caputo, Caputo-Fabrizio and Atangana-Baleanu derivative with fractional order: fractional cubic isothermal auto-catalytic chemical system," *The European Physical Journal Plus*, vol. 133, no. 3, pp. 1–12, 2018.
- [37] M. Hamarsheh, A. I. Ismail, and Z. Odibat, "An analytic solution for fractional order Riccati equations by using optimal homotopy asymptotic method," *Applied Mathematical Sciences*, vol. 10, no. 23, pp. 1131–1150, 2016.
- [38] S. Rathore, D. Kumar, J. Singh, and S. Gupta, "Homotopy analysis Sumudu transform method for nonlinear equations," *International Journal of Industrial Mathematics*, vol. 4, no. 4, pp. 301–314, 2012.
- [39] A. Ghorbani, "Beyond Adomian polynomials: He polynomials," *Chaos, Solitons & Fractals*, vol. 39, no. 3, pp. 1486–1492, 2009.
- [40] S. Abbasbandy and K. V. Elyas Shivanian, "A new approximate analytical technique for dual solutions of nonlinear differential equations arising in mixed convection heat transfer in a porous medium," *International Journal of Numerical Methods for Heat & Fluid Flow*, vol. 54, 2017.
- [41] S.-J. Liao, "An approximate solution technique not depending on small parameters: a special example," *International Journal of Non-linear Mechanics*, vol. 30, no. 3, pp. 371–380, 1995.
- [42] V. F. Morales-Delgado, J. F. Gómez-Aguilar, S. Kumar, and M. A. Taneco-Hernández, "Analytical solutions of the Keller-Segel chemotaxis model involving fractional operators without singular kernel," *The European Physical Journal Plus*, vol. 133, no. 5, p. 200, 2018.