

# Nutrient intakes of children aged 1–2 years as a function of milk consumption, cows' milk or growing-up milk

Jacques Ghisolfi<sup>1,\*†</sup>, Marc Fantino<sup>2</sup>, Dominique Turck<sup>3</sup>, Geneviève Potier de Courcy<sup>4</sup> and Michel Vidailhet<sup>5</sup>

<sup>1</sup>Department of Paediatrics, University Paul Sabatier, Toulouse, France: <sup>2</sup>Department of Physiology, University of Burgundy, Dijon, France: <sup>3</sup>Department of Paediatrics, University Lille 2, Lille, France:

<sup>4</sup>CNRS, UREN, Bobigny, France: <sup>5</sup>Department of Paediatrics, University Henri Poincaré, Nancy, France

Submitted 12 December 2011: Accepted 28 February 2012: First published online 6 July 2012

## Abstract

**Objective:** To evaluate the nutritional adequacy of diets in early childhood as a function of milk intake, cows' milk (CM) or growing-up milk (GUM).

**Design:** From a cross-sectional food consumption survey, two groups of children aged 1–2 years were defined: Group CM fed exclusively on CM  $\geq 250$  ml/d and Group GUM fed on GUM  $\geq 250$  ml/d. Proportions of children at risk of nutrient excess or insufficiency were estimated relative to the French Recommended Daily Allowances, Estimated Average Requirements or Adequate Intakes.

**Setting:** Parents participating in the survey were recruited from all regions of France by a polling organization. Distribution was adjusted to that of the French population.

**Subjects:** Sixty-three (Group CM) and fifty-five (Group GUM) children.

**Results:** Total energy and macronutrient intakes were similar in the two groups except protein intake of Group CM, which was much higher than the Recommended Daily Allowance and significantly higher than in Group GUM. A high percentage of children of Group CM had intake of linoleic acid (51%) and  $\alpha$ -linolenic acid (84%) below the lower limit of the Adequate Intake, and intake of Fe (59%) vitamin C (49%) and alimentary vitamin D (100%) less than the Estimated Average Requirement. Significant differences were observed in the proportions of children with a risk of dietary inadequacy between the two groups for all the mentioned nutrients ( $P < 0.001$ ). In Group GUM, this imbalance was only observed for vitamin D. Intake of foods other than milk and dairy products could not account for these discrepancies.

**Conclusions:** Consumption of CM ( $\geq 250$  ml/d) entails the risk of insufficiency in  $\alpha$ -linolenic acid, Fe, vitamin C and vitamin D. Use of GUM ( $\geq 250$  ml/d) significantly reduces the risk of insufficiencies in the mentioned nutrients.

**Keywords**  
Cows' milk  
Growing-up milk

With breast-feeding rarely being used for children after 1 year of age, breast milk substitutes are required to ensure the nutritional requirements of young children. In this regard, many child health professionals, not to mention many parents, consider cows' milk (CM) adequate and believe that a diet based on CM provides all nutritional requirements with the exception of Fe and vitamin D. For some 20 years now, an alternative to CM has been available. The so-called 'growing-up milks' (GUM) are not defined in any European Regulation or Directive, or in any Codex Alimentarius standard. GUM are intended for children after 1 year of age (up to 3 years of age in many European countries). They have a lower protein content than CM and are supplemented with several nutrients of interest including Fe and essential fatty acids (EFA), as well as vitamins D

and E. Directive 2009/39/EC of the European Parliament and of the Council of 6 May 2009 classifies these products as a 'foodstuff intended for particular nutritional uses' and states that 'the composition of the products shall be such that the products are appropriate to the nutritional use intended'<sup>(1)</sup>. However, although the commercialization of GUM continues to increase in many countries worldwide, particularly in Europe, their benefits are still a matter of debate. This controversy arises because the possible nutritional risks associated with the use of CM and the expected benefits from the use of GUM have not been clearly demonstrated. To gain further information on this important issue in paediatric nutrition, individual dietary data from a food consumption survey performed in France were retrospectively analysed to estimate the nutritional adequacy of the diets of children aged 1–2 years as a function of their milk intake, i.e. CM or GUM.

† Correspondence address: 28 Grand'rue, 31560 Cagnac, France.

\*Corresponding author: Email jacques.ghisolfi@nordnet.fr

The objectives of the present study were to answer the following two questions:

1. Does the use of CM in young children aged 1–2 years represent a risk of inadequate coverage of the nutritional requirements currently defined in France for this population?
2. If CM does not adequately meet these nutritional requirements, should GUM be used in preference by this population?

## Experimental methods


### Study sample

A cross-sectional food consumption survey involving 713 children aged from 15 d to 36 months was performed by the Department of Physiology of the University of Burgundy, Dijon, France. The results of this survey, sponsored by the Syndicat français des aliments de l'enfance (French Association of Baby Food Industries), were published in 2008<sup>(2)</sup>. According to French legislation, this type of nutritional survey does not need to be approved by an institutional review board. For the purpose of the present study, only data from 132 children aged 12–24 months (seventy-one boys) were taken into account. They were all full-term babies with birth weights of over 2500 g. At the time of the study, none of the children was breast-fed. The parents did not receive specific recommendations prior to the study, which reflects the daily food consumption of the children according to their parents' choices. Two groups were defined according to their type of milk intake. Group CM comprised the sixty-three children who received a daily intake of CM of at least 250 ml (70% as semi-skimmed milk) and who did not receive GUM or follow-on formula (FOF) or dairy products based on GUM or FOF (Fig. 1). Group GUM comprised

the fifty-five children who received a daily intake of GUM of at least 250 ml (Fig. 1). This minimal value of 250 ml/d for milk consumption was retained *a priori* since it corresponds to one daily bottle consumption.

### Data acquisition

Parents participating in the survey were recruited from all regions of France (excluding overseas territories) by the polling organization TNS/SOFRES<sup>®</sup>, according to a proportional sampling technique that took into account the population of each region, the age of the children, the professional status of the mother and the socio-economic level of the family. The distribution was adjusted to that of the general French population as defined by INSEE (Institut National de la Statistique et des Etudes Economiques/National Institute of Statistics and Economic Studies) on the basis of the 2002 census. None of the families in the study population was defined as economically precarious. With the consent of both parents, the usual caregiver of the child in the family environment was asked to record in the diary provided all foods and beverages ingested by the child over three consecutive days. The study period usually included one weekend day. The caregiver was instructed to note the time of each meal, the weight (or volume) of all the ingested foods and a precise description of them, including trade names, together with the methods of their preparation and detailed recipes for all home prepared foods. All foods and beverages were weighed on kitchen scales (accuracy  $\pm 1$  g), measured by mass or volume from the information on the packaging, or estimated from photographs of calibrated portions especially prepared for this purpose. Quantitative dietary records, together with the weights of any leftovers, were recorded in the diaries by the parents, then verified and if necessary clarified by especially trained researchers recruited by TNS/SOFRES.



**Fig. 1** Flow chart representing how Group CM and Group GUM were constituted (CM, cows' milk; GUM, growing-up milk; FOF, follow-on formula)

### **Analysis of food dairies**

The nutritional analysis was performed using tables that listed the compositions of 1260 foodstuffs in current use in home cooking in France ('current foods') and of all food items especially manufactured and intended for infants and young children (less than 3 years of age) available on the market in France at the time of the data acquisition (850 'baby foods'). The nutritional analysis of the 'current foods' was performed using a composition table provided by AFSSA (Agence Française de Sécurité Sanitaire des Aliments/French Food Safety Agency) and another composition table that was used for the SU.VI.MAX study (a French study on supplementation in vitamins and antioxidant minerals)<sup>(3)</sup>. The compositions of very few foods (less than ten) could not be determined and these were therefore deduced from the mean compositions of similar products. The compositions of all 'baby foods' were provided by the manufacturers. Dietary intake was calculated for each child with the especially developed software NUTRI 7<sup>®</sup>. Each diary was encoded and verified after data entry. Energy and nutrient intakes were calculated for each child by summing his/her reported consumption over the 3 d of the study. The child's intake was reduced to a daily rate. These data were then aggregated for all children in each study group (Group CM and Group GUM). Besides the total energy intake, the nutrients considered were: protein, lipids, total carbohydrates (excluding fibre), EFA (linoleic acid and  $\alpha$ -linolenic acid), Na, Ca, P, Mg, Zn, Fe, vitamins B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>5</sub>, B<sub>6</sub>, B<sub>9</sub>, B<sub>12</sub>, C, D (exclusively of food origin, referred to hereafter as 'alimentary vitamin D'), E (expressed as  $\alpha$ -tocopherol equivalents), total vitamin A (expressed as retinol equivalents), retinol and carotenoids (expressed as  $\beta$ -carotene equivalents). In France, children aged 1 to 2 years are frequently prescribed vitamin D and fluoride, rarely Fe and multivitamin supplements. These supplementary intakes were not taken into account in the current study because of the unreliable reporting of therapeutic compliance.

### **Analytical methods**

The means, errors and standard deviations of the daily energy and nutrient intakes were calculated for the two groups. They were compared with the French Recommended Daily Allowances (RDA), Estimated Average Requirements (EAR) and Adequate Intakes (AI)<sup>(4)</sup>. Because the weight of each infant was reported by the person filling out the diary and was not measured as part of the study, the daily intake was not calculated per unit body weight. The sex of the children was not considered. To express the variation in the intake of each nutrient, the first and third quartiles of intake are presented (with the 95% confidence intervals). The daily energy and nutrient intakes of the study groups were compared using Student's *t* test, and differences were considered significant at  $P < 0.05$ . The proportion of children with inadequate dietary intake in

each group was evaluated by estimating the percentage of individuals whose daily intake was less than the EAR (equivalent to 0.77 of the RDA) for the nutrient under consideration<sup>(5-7)</sup>. However, because there are no RDA for EFA, the lower limit of the AI<sup>(5)</sup> was used for these nutrients (linoleic acid and  $\alpha$ -linolenic acid). The 95% confidence intervals for these threshold values are indicated. To estimate the contribution to the global nutritional intake of various food groups, five categories of 'current foods' were distinguished: (i) milk; (ii) other dairy products; (iii) meat, fish and eggs; (iv) other foods (mainly cereals, fruits and vegetables); and (v) vegetable fats. Similarly, for foods specifically intended for infants younger than 3 years, three categories were defined: (i) GUM and GUM/FOF-based dairy products; (ii) meat and fish; and (iii) other 'baby foods'. The contribution of these food categories to each nutrient intake was expressed as daily amounts. The effect of variations in daily milk consumption on the intake of eight nutrients of interest (protein, Fe, linoleic acid,  $\alpha$ -linolenic acid, retinol, vitamin C, vitamin E and alimentary vitamin D) was evaluated by establishing a relationship between, on the one hand, either the daily consumption of CM and CM-based products or the daily consumption of GUM and GUM-based products (including FOF) and, on the other, the total daily intake of each of these eight nutrients, using the Pearson correlation coefficient. Statistical analysis was performed with the software Number Cruncher Statistical System version 2000 for Windows.

## **Results**

### **Group CM**

Mean age of the sixty-three children included in Group CM was 606 (SD 13) d (19.9 (SD 0.4) months). Their diet was characterized by (Tables 1 and 2): (i) a protein intake well above the French RDA<sup>(5)</sup>; (ii) a low lipid intake; (iii) intakes under the lower limit of the French AI for linoleic acid and  $\alpha$ -linolenic acid, and under the RDA for Fe, Zn, vitamin C, alimentary vitamin D and vitamin E. The large variations in energy and nutrient intakes were reflected in the large standard deviations and the values for the first and third quartiles (Table 1 and Fig. 2). These variations were associated primarily with the consumption of foods other than milk and dairy products (Table 3). A high proportion of these children consumed less than the lower limit of the AI for linoleic acid (51%) and  $\alpha$ -linolenic acid (84%). Their daily intake was less than the EAR for Fe (59%), Zn (56%), alimentary vitamin D (100%), vitamin E (94%) and vitamin C (49%; Table 2). Whereas CM and CM-based products represented 43% by weight of their total food consumption and 35% of their daily energy intake, they represented 45% of their daily intake of protein, 21% of Na, 17% of linoleic acid, 25% of  $\alpha$ -linolenic acid, 11% of Fe, 46% of Zn, 8% of vitamin C, 16% of vitamin E and 24% of alimentary vitamin D (Table 3). This population

**Table 1** Energy and nutrient daily intake distributions of French children aged 12–24 months consuming either CM or GUM

Energy or nutrient	AI* or RDA†	Group	Mean	SE	t Test	SD	First quartile	95% CI	Third quartile	95% CI
Mass of food ingested (g)		CM‡	1212	39		308	1020	882, 1085	1340	1240, 1537
		GUM§	1204	32		240	1054	943, 1104	1337	1261, 1466
TEI (kJ)		CM	4027	142		1125	3324	2906, 3468	4652	4234, 5421
		GUM	4052	79		584	3548	3399, 3697	4426	4270, 4605
Protein (g)	35.9*.a	CM	41.6	1.4	<i>P</i> < 0.001	10.9	33.3	31.2, 36.1	47.1	44.5, 54.5
		GUM	35.5	1.0		7.3	30.0	27.4, 32.0	39.7	37.1, 42.9
Lipids (g)	45.1*.b	CM	34.2	1.7		13.6	24.6	18.6, 27.6	40.8	37.4, 49.8
		GUM	37.0	1.0		7.1	31.5	27.2, 34.2	39.9	36.7, 44.2
Carbohydrate (g)	131.6*.c	CM	123	4.3		34.2	101	89, 107	141	132, 169
		GUM	125	2.9		21.5	112	101, 117	137	131, 143
Linoleic acid (g)	2.13*.d	CM	2.38	0.17	<i>P</i> < 0.001	1.38	1.58	1.08, 1.79	2.79	2.48, 3.71
		GUM	3.80	0.14		1.03	3.07	2.31, 3.36	4.49	4.08, 5.09
α-Linolenic acid (g)	0.43*.e	CM	0.33	0.03	<i>P</i> < 0.001	0.20	0.19	0.17, 0.24	0.38	0.33, 0.45
		GUM	0.51	0.02		0.13	0.42	0.35, 0.47	0.58	0.55, 0.63
Vitamin B <sub>1</sub> (mg)	0.4†	CM	0.76	0.03		0.24	0.59	0.51, 0.64	0.91	0.84, 1.02
		GUM	0.89	0.04		0.30	0.66	0.57, 0.76	1.03	0.95, 1.31
Vitamin B <sub>2</sub> (mg)	0.8†	CM	1.48	0.07		0.52	1.12	0.97, 1.28	1.71	1.49, 2.04
		GUM	1.44	0.05		0.35	1.22	1.04, 1.28	1.65	1.52, 1.93
Vitamin B <sub>3</sub> (niacin) (mg)	6†	CM	6.23	0.31	<i>P</i> < 0.001	2.44	4.65	4.16, 4.98	6.91	6.56, 8.8
		GUM	9.16	0.54		3.99	6.12	4.53, 6.94	12.19	10.3, 12.9
Vitamin B <sub>5</sub> (mg)	2.5†	CM	3.27	0.14		1.12	2.53	2.09, 2.77	3.97	3.38, 4.73
		GUM	3.11	0.13		0.98	2.32	1.93, 2.58	3.71	3.39, 4.28
Vitamin B <sub>6</sub> (mg)	0.6†	CM	0.91	0.04	<i>P</i> < 0.001	0.32	0.67	0.57, 0.78	1.07	0.97, 1.21
		GUM	1.21	0.06		0.43	0.81	0.70, 0.94	1.52	1.31, 1.69
Vitamin B <sub>9</sub> (μg)	100†	CM	126	6.2	<i>P</i> < 0.001	48.8	88.4	74, 98	155	140, 180
		GUM	162	6.8		50.6	123	98, 139	184	176, 231
Vitamin B <sub>12</sub> (μg)	0.8†	CM	3.24	0.51		4.03	2.00	1.64, 2.13	3.12	2.83, 3.72
		GUM	2.58	0.23		1.73	1.88	1.58, 1.98	2.84	2.53, 3.43
Retinol (μg)	400†	CM	392	121		961	139	117, 154	283	245, 384
		GUM	476	35		261	358	323, 380	502	487, 553
Carotenoids as β-carotene activity (μg)	–	CM	2377	249	<i>P</i> < 0.001	1977	819	486, 1109	3412	2778, 5003
		GUM	3858	344		2552	1925	975, 2521	5046	4262, 6643
Total vitamin A (μg retinol equivalents)	400†	CM	788	131	<i>P</i> < 0.001	1042	353	241, 420	895	665, 1128
		GUM	1119	75		558	773	542, 858	1363	1151, 1697
Vitamin C (mg)	60†	CM	52	3.7	<i>P</i> < 0.001	29.2	29.5	24, 33	69.3	59.0, 81.2
		GUM	82	4.1		30.5	58.0	50, 66	98.0	86.3, 117
Alimentary vitamin D (μg)	10†	CM	0.78	0.09	<i>P</i> < 0.001	0.69	0.32	0.22, 0.47	0.96	0.83, 1.36
		GUM	6.77	0.28		2.11	5.12	4.64, 5.31	7.76	6.88, 9.25
Vitamin E (mg)	6†	CM	2.75	0.14	<i>P</i> < 0.001	1.08	1.75	1.58, 2.17	3.56	3.11, 4.23
		GUM	6.17	0.22		1.63	4.74	4.52, 5.05	7.02	6.6, 7.99
P (mg)	360*	CM	779	30	<i>P</i> < 0.01	238	614	565, 652	871	792, 1115
		GUM	667	19		139.0	587	527, 603	738	681, 780
Ca (mg)	500†	CM	800	36		283	614	538, 670	962	834, 1039
		GUM	755	19		142	644	599, 703	826	781, 897
Mg (mg)	80†	CM	144	5.6	<i>P</i> < 0.001	45	112	101, 122	167	150, 196
		GUM	112	4.1		30	94	83, 101	124	114, 152

**Table 1** Continued

Energy or nutrient	AI* or RDA†	Group	Mean	SE	t Test	SD	First quartile	95% CI	Third quartile	95% CI
Na (mg)	920*	CM	1156	51		408	827	680, 961	1423	1238, 1602
		GUM	1023	44		327	809	690, 850	1146	1078, 1293
Fe (mg)	7†	CM	5.19	0.25	<i>P</i> < 0.001	1.96	3.65	3.05, 4.38	6.43	5.88, 7.47
		GUM	9.37	0.31		2.33	7.69	7.04, 8.23	10.44	9.74, 11.93
Zn (mg)	6†	CM	4.58	0.28	<i>P</i> < 0.001	2.19	3.12	2.55, 3.55	5.32	4.81, 7.00
		GUM	6.35	0.42		3.08	4.16	3.69, 4.62	7.40	6.06, 10.16

CM, cows' milk; GUM, growing-up milk; TEI, total energy intake.

\*15% of TEI; †40% of TEI; ‡55% of TEI; §lower limit of AI: 2% of TEI; ¶lower limit of AI: 0.4% of TEI (from Beaufrère *et al.*<sup>(4)</sup>).

\*AI: Adequate Intake (from Beaufrère *et al.*<sup>(4)</sup>).

†RDA: French Recommended Daily Allowance (from Beaufrère *et al.*<sup>(4)</sup>).

‡Group CM: children aged 12–24 months who received a daily intake of CM of at least 250 ml and who did not receive GUM or follow-on formula (FOF) or dairy products based on GUM or FOF (*n* 63).

§Group GUM: children aged 12–24 months who received a daily intake of GUM of at least 250 ml (*n* 55).

**Table 2** Percentage of individuals whose daily intakes were less than the AI or less than 0.77 of the RDA among French children aged 12–24 months consuming either CM or GUM

Nutrient	Linoleic acid	α-Linolenic acid	Fe	Zn	Retinol	Vitamin C	Vitamin D	Vitamin E
AI or RDA*	2.13 g	0.43 g¶	7 mg	6 mg	400 µg	60 mg	10 µg	6 mg
Threshold for a possible daily intake insufficiency†	2.13 g	0.43 g	5.4 mg	4.6 mg	308 µg	46 mg	7.7 µg	4.6 mg
Group CM ( <i>n</i> 63)‡								
No. of children with intake under the threshold for dietary inadequacy	32	53	37	35	51	31	63	59
Proportion (%)	51	84	59	56	81	50	100	94
95% CI	38, 63	75, 93	47, 71	43, 68	71, 91	37, 62	100, 100	88, 100
Group GUM ( <i>n</i> 55)§								
No. of children with intake under the threshold for dietary inadequacy	2	14	0	18	2	6	41	8
Proportion (%)	4	26	0	33	4	11	75	15
95% CI	0, 9	14, 37	0, 0	20, 45	0, 9	3, 19	63, 86	5, 24
Probability of a significant difference, within the two groups, in the proportion of children with a risk of dietary inadequacy ( <i>P</i> < 0.05)	<i>Z</i> = 5.6	<i>Z</i> = 6.4	<i>Z</i> = 6.8	<i>Z</i> = 2.4	<i>Z</i> = 8.4	<i>Z</i> = 4.5	<i>Z</i> = 4.2	<i>Z</i> = 8.7
	<i>P</i> < 0.001	<i>P</i> < 0.001	<i>P</i> < 0.001	<i>P</i> < 0.02	<i>P</i> < 0.001	<i>P</i> < 0.001	<i>P</i> < 0.001	<i>P</i> < 0.001

CM, cows' milk; GUM, growing-up milk.

\*AI or RDA: Adequate Intake calculated as the recommended contribution of the nutrient to total energy intake or Recommended Daily Allowance (from Beaufrère *et al.*<sup>(4)</sup>).


†AI or 77% of the French RDA.

‡Group CM: children aged 12–24 months who received a daily intake of CM of at least 250 ml and who did not receive GUM or follow-on formula (FOF) or dairy products based on GUM or FOF (*n* 63).

§Group GUM: children aged 12–24 months who received a daily intake of GUM of at least 250 ml (*n* 55).

||Lower limit of AI: 2% of observed total energy intake.

¶Lower limit of AI: 0.4% of observed total energy intake.



**Fig. 2** Effects of daily variation of milk product intake on nutritional daily intake of eight nutrients of interest (iron, protein, linoleic acid, α-linolenic acid, retinol, vitamin C, vitamin E, alimentary vitamin D) among French children aged 12–24 months. ●, Daily nutritional intake as a function of milk consumption by sixty-three children who received at least 250 ml of cows' milk (CM) daily, and no growing-up milk (GUM) or follow-on formula (FOF), or dairy products based on GUM or FOF (Group CM); ◆, daily nutritional intake as a function of milk consumption by fifty-five children who received at least 250 ml of GUM daily (Group GUM)

**Table 3** Contribution of various food groups to daily nutritional intake among French children aged 12–24 months consuming either CM or GUM

	Mass of food ingested (g)		Total energy intake (kJ)		Protein (g)		Lipids (g)		Carbohydrates (g)		Linoleic acid (g)		α-Linolenic acid (g)		Retinol (μg)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
<b>Current foods</b>																
<b>Milk</b>																
Group CM*	360	23.9	708	48.2	11.6	0.8	6.4	0.5	16.3	1.1	0.23	0.08	0.04	0.00	79.5	7.7
Group GUM†	26	9.9	50	18.8	0.8	0.3	0.4	0.2	1.2	0.5	0.01	0.00	0.00	0.00	5.1	2.0
<b>Other dairy products</b>																
Group CM	156	13.9	681	54.5	7.0	0.6	5.6	0.6	21.0	1.9	0.17	0.02	0.04	0.00	57.2	6.9
Group GUM	119	9.4	590	46.6	6.6	0.5	6.0	0.5	15.3	1.5	0.16	0.02	0.05	0.00	62.3	5.2
<b>Meat, fish and eggs</b>																
Group CM	31	2.8	256	24.3	7.2	0.7	3.2	0.3	0.9	0.2	0.30	0.04	0.03	0.00	175.4	121.5
Group GUM	22	3.1	187	26.6	5.2	0.7	2.5	0.4	0.4	0.2	0.19	0.03	0.02	0.00	42.9	34.4
<b>Other foods</b>																
Group CM	575	32.5	1959	135.7	13.9	1.0	15.1	1.4	69.3	4.6	1.24	0.13	0.17	0.02	335.4	37.4
Group GUM	436	27.6	1250	86.9	9.3	0.8	9.6	0.8	44.6	2.9	0.84	0.09	0.11	0.01	441.4	61.4
<b>Vegetable fats</b>																
Group CM	1.3	0.3	45.1	11.6	0.01	0.00	1.2	0.3	0.002	0.00	0.2	0.05	0.02	0.00	1.6	0.9
Group GUM	0.9	0.3	30.8	11.3	0.01	0.00	0.8	0.3	0.002	0.00	0.1	0.05	0.01	0.00	0.5	0.3
<b>Foods specially for young children</b>																
<b>GUM and GUM/FOF-based dairy products</b>																
Group CM	0	–	0	–	0	–	0	–	0	–	0	–	0	–	0	–
Group GUM	385	18	1139	44	9.0	0.3	11.1	0.4	33.3	1.6	1.94	0.09	0.27	0.02	288.4	13.1
<b>Meat and fish</b>																
Group CM	19	6	55	17	0.6	0.2	0.4	0.1	1.8	0.6	0.07	0.03	0.01	0.00	0.4	0.1
Group GUM	70	12	202	35	1.9	0.3	1.5	0.3	6.8	1.2	0.31	0.07	0.03	0.01	1.7	0.4
<b>Other 'baby foods'</b>																
Group CM	68	13	284	45	1.4	0.2	1.0	0.2	13.4	2.2	0.16	0.04	0.02	0.01	7.7	2.4
Group GUM	143	19	548	63	2.7	0.3	2.1	0.3	25.5	2.8	0.20	0.04	0.03	0.01	26.5	5.7
<b>Vitamin C (mg)</b>																
<b>Alimentary vitamin D (mg)</b>																
<b>Vitamin E (mg)</b>																
<b>Ca (mg)</b>																
<b>Mg (mg)</b>																
<b>Zn (mg)</b>																
<b>Na (mg)</b>																
<b>Fe (mg)</b>																
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
<b>Current foods</b>																
<b>Milk</b>																
Group CM	3.6	0.3	0.02	0.01	0.27	0.02	400	28	42.8	2.8	1.4	0.1	170	12	0.22	0.04
Group GUM	0.3	0.1	0.00	0.00	0.02	0.01	29	11	3.1	1.2	0.1	0.00	13	5.0	0.01	0.00
<b>Other dairy products</b>																
Group CM	0.4	0.0	0.17	0.02	0.16	0.02	182	17	17.0	1.5	0.5	0.1	71	7	0.35	0.04
Group GUM	0.3	0.0	0.16	0.01	0.17	0.01	135	11	12.4	1.1	0.5	0.00	50	4	0.30	0.06
<b>Meat, fish and eggs</b>																
Group CM	0.2	0.2	0.15	0.06	0.10	0.02	5.9	0.9	8.8	0.9	0.8	0.2	62	7	0.61	0.09
Group GUM	0.1	0.1	0.16	0.06	0.07	0.02	3.5	1.0	6.3	0.9	0.5	0.1	36	6	0.39	0.07
<b>Other foods</b>																
Group CM	34.6	3.1	0.26	0.03	1.46	0.12	188	15	66.9	4.5	1.8	0.2	774	56	3.07	0.22
Group GUM	27.4	2.7	0.20	0.03	1.10	0.11	143	13	49.7	3.9	1.3	0.1	588	54	2.14	0.19
<b>Vegetable fats</b>																
Group CM	0	–	0.02	0.01	0.21	0.05	0.1	0.02	0.0	0.0	0	–	0.4	0.2	0	–
Group GUM	0	–	0.02	0.01	0.13	0.04	0.1	0.04	0.0	0.0	0	–	0.6	0.4	0	–

Table 3 Continued

	Vitamin C (mg)		Alimentary vitamin D (mg)		Vitamin E (mg)		Ca (mg)		Mg (mg)		Zn (mg)		Na (mg)		Fe (mg)	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
Foods especially for young children																
GUM and GUM/FOF-based dairy products																
Group CM	0	—	0	—	0	—	0	—	0	—	0	—	0	—	0	—
Group GUM	28.3	1.3	5.75	0.26	3.57	0.20	382	15	30	2	3.9	0.4	156	8	4.97	0.19
Meat and fish																
Group CM	0.9	0.4	0.02	0.01	0.09	0.03	3	1	3	1	0.00	0.02	28	9	0.07	0.02
Group GUM	4.1	1.1	0.06	0.02	0.40	0.09	9	2	5	1	0.02	0.0	102	17	0.27	0.06
Other 'baby foods'																
Group CM	11.9	2.6	0.12	0.04	0.43	0.07	22	4	7	1	0.1	0.0	49	12	0.88	0.14
Group GUM	21.6	3.3	0.39	0.13	0.44	0.10	53	9	6	1	0.1	0.0	77	14	1.30	0.30

CM, cows' milk; GUM, growing-up milk; FOF, follow-on formula.

\*Group CM: children aged 12–24 months who received a daily intake of CM of at least 250 ml and who did not receive GUM or follow-on formula (FOF) or dairy products based on GUM or FOF (n 63).

†Group GUM: children aged 12–24 months who received a daily intake of GUM of at least 250 ml (n 55).

consumed few non-dairy manufactured foodstuffs specifically designed for children aged 1–3 years, which contributed minimally (less than 10%) to their total nutritional intake, except for Fe (18%), vitamin A (16%), alimentary vitamin D (19%), vitamin E (19%) and vitamin C (25%; Table 3). There was a significant relationship between the daily consumption of CM and CM-based products and daily protein intake ( $P < 0.001$ ). No relationship was found between the daily consumption of CM and CM-based products and linoleic acid,  $\alpha$ -linolenic acid, Fe, retinol, vitamin C, alimentary vitamin D or vitamin E (Fig. 2).

**Group GUM**

Mean age of the fifty-five children included in Group GUM was significantly lower than that of Group CM at 534 (SD 15) d (17.5 (SD 0.5) months;  $P = 0.003$ ). Their diet did not differ significantly from that of Group CM in terms of the total mass of food, energy, carbohydrates, lipids, Na, Ca, P and Mg (Table 1). Their diet was also characterized by a high protein intake, 35.5 (SD 7.3) g/d, of which an average of 5.2 g was in the form of 'current foods' (meat, eggs and fish) and 16.4 g in the form of milk products, which was nevertheless lower than that of Group CM (Table 3). Their lipid intake was low. Their intakes of EFA, Fe, Zn, vitamins B<sub>3</sub>, B<sub>6</sub>, B<sub>9</sub>, C and E, total vitamin A, alimentary vitamin D and  $\beta$ -carotene were significantly higher ( $P < 0.001$ ) than those of group CM (Table 1). Their consumption of GUM and GUM/FOF-based milk products represented 27% of their total daily intake of protein, 51% of linoleic acid, 52% of  $\alpha$ -linolenic acid, 96% of Fe, 85% of Zn, 26% of total vitamin A, 74% of retinol, 35% of vitamin C, 58% of vitamin E and 85% of alimentary vitamin D. The contribution of non-dairy manufactured foodstuffs specifically designed for children aged 1–3 years to their total nutritional intake was minimal (less than 10%), except for linoleic acid (13%),  $\alpha$ -linolenic acid (12%), Fe (17%), total vitamin A (24%), vitamin E (17%) and vitamin C (31%; Table 3). As in Group CM, the variations in their energy and nutrient intakes were pronounced, particularly for non-dairy products (Table 3 and Fig. 2). Compared with Group CM, a far smaller percentage of children in this group had a daily intake of  $\alpha$ -linolenic acid below the lower limit of the AI (26%) or a daily intake less than the EAR for Zn (33%), vitamin C (11%), alimentary vitamin D (75%) and vitamin E (15%; Table 2). There was a significant relationship between their daily consumption of GUM and their daily intake of Fe ( $P < 0.001$ ), linoleic acid and  $\alpha$ -linolenic acid ( $P < 0.001$ ), vitamin E ( $P < 0.001$ ), vitamin C ( $P < 0.05$ ) and alimentary vitamin D ( $P < 0.02$ ; Fig. 2). No relationship was found between the daily consumption of GUM and protein or retinol intake.

**Discussion**

Based on the expected effects of the composition of CM and GUM, it is commonly believed that feeding a young



child with CM can lead to a high protein intake and a deficient intake of linoleic and  $\alpha$ -linolenic acids, Fe, Zn and most vitamins. This analysis is not supported by convincing research data arising from plasma measurements of nutrients of interest, with the notable exceptions of Fe<sup>(8)</sup> and vitamin D<sup>(9,10)</sup>. Energy and macronutrient consumption was identical in the two groups in the present study with the exception of protein, for which the consumption was higher in Group CM, cows' milk being one possible but not exclusive causative factor. The main nutritional differences between the two groups concerned EFA, some trace elements and vitamins. In Group CM, consumption of EFA, Fe, Zn, vitamin C, alimentary vitamin D and vitamin E was below the French RDA or AI. As expected, given compositional differences between CM and GUM, consumption of these nutrients was higher in Group GUM and adequate as compared with the RDA or AI, except for alimentary vitamin D. Before concluding that these dietary inadequacies are attributable to the use of CM, it is also important to consider the role of the consumption of foods other than milk and dairy products. Our study shows that those foods represent a daily intake at least equivalent in protein and higher in EFA, Fe, Zn and many vitamins than that of a diet consisting solely of CM, and that their consumption is not sufficient to reach AI and RDA for the nutrients under consideration (Table 3). A food consumption survey performed in the USA in 2001 concerning 998 children aged 1–2 years reached similar conclusions. Eighty-five per cent of the study population was fed CM and 15% was fed formula. A high percentage of these children had intakes of protein, Zn and vitamin A above the RDA, and low intakes of fat, Fe and vitamin E<sup>(11)</sup>. These results and the results from the present study do not allow us to conclude that the health of these infants is at risk because of a dietary insufficiency in one or more nutrients. For any individual, an intake less than the EAR (or the lower limit of the AI for EFA) does not mean *per se* that that individual is effectively lacking input, especially because there are great variations in individual needs and in different food items.

Reports from food consumption surveys have well-known limitations: lack of homogeneity among the study populations, approximate reporting of dietary intakes, imprecision in food composition tables, and failure to take into account the bioavailability of nutrients in relation to diet and physiological status<sup>(12–14)</sup>. Long-term epidemiological studies with recurrent blood sampling would be the gold standard but are very hard to deal with due to ethical considerations, high financial burden and the large number of children lost to follow-up. Paediatric EAR and RDA are not generally supported by indisputable data, whether expressed in relation to the nutritional qualities of breast milk or evaluated by factorial methods with biomarkers or by extrapolation from defined adult dietary needs secondarily adapted to the height and weight of children<sup>(13)</sup>. At the present time, Fe and vitamin D are the only nutrients

for which the determination of requirements is based on reliable data. For EFA, Zn and vitamins C and E, it is not possible to infer that diets really constitute a risk for deficiencies in these nutrients, given our very poor knowledge of children's actual needs<sup>(14–16)</sup>. Taking into account the above-mentioned methodological drawbacks<sup>(17,18)</sup>, the only nutrients for which there is nowadays a higher risk of deficiency in relation to the consumption of CM are  $\alpha$ -linolenic acid, Fe, vitamin C and vitamin D.

The increased risk of dietary insufficiency in several nutrients in a population of children aged 1–2 years fed CM is the main scientific justification for proposing modifications to their diet. These may include changing the intake of foods other than milk and dairy products, suggesting nutritional supplements, or recommending specific milk products, i.e. GUM. The modification of children's diets with respect to non-milk foods can be effective. The correction of an insufficiency in EFA by increasing the consumption of vegetable oils, corn, rapeseed or soya (3–5 g/d) is more strongly indicated in these young children whose total lipid intake is less than 35% of total energy intake. An increased consumption of cereals, vegetables and fresh fruit (particularly citrus fruits) should correct any dietary insufficiency in vitamins, especially vitamin C. Conversely, correction of the dietary insufficiency in Fe by the increased consumption of meat products is not appropriate because it represents an additional daily protein intake of 10 g, leading to a very high total protein intake. The Committee on Nutrition of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) also emphasized that the only nutrients whose needs cannot be met by non-milk foods are Fe and vitamin D<sup>(19)</sup>. Dietary supplements, which are very much in use in the USA, even in young children<sup>(20)</sup>, can contribute to the correction of the insufficiencies observed in the paediatric population. In France, their use in young children, which seems to be increasing, is discouraged by most health professionals because they can lead to excessive micronutrient intake<sup>(20)</sup>, the consequences of which are unknown. GUM have now been available for about 20 years, their alleged purpose being to avoid the nutritional imbalance that may arise from giving children CM after 10–12 months of age. The present study confirms that the protein intake was lower in children fed GUM than in children fed CM, partly because the protein content of GUM is lower than that of CM, but predominantly because children in Group CM were fed more non-milk foods rich in protein. The present study also indicates that GUM plays a beneficial role by improving the intake of EFA, Fe and vitamins C and D in these children. The ingestion of at least 250 ml GUM/d supplies the RDA levels of the nutrients of interest, notably  $\alpha$ -linolenic acid and Fe, for the majority of children, with the exception of vitamin D.

There is no nutritional risk at the levels of use normally recommended for GUM (250–500 ml/d). Devaney *et al.* have

stressed the risk in this population of an excessive intake of Zn and vitamin A<sup>(11)</sup>. We saw no evidence of this with respect to Zn in either of our two study groups. However, Allen has reported that an excessive intake of Zn is very difficult to prove, given the lack of knowledge about the risk threshold for this micronutrient<sup>(13)</sup>. In our study, vitamin A intake exceeding the upper recommended limit for infants aged 1–3 years (600 or 800 µg/d depending on the reference source)<sup>(21,22)</sup> was observed in infants in Group GUM, but also in Group CM. Such an elevated intake cannot be explained by the retinol content of GUM. It seems to be associated, in a few cases, with an unusual intake of retinol (two children in the study displayed a very high intake of retinol, nearly 3000 µg/d, associated with the consumption of calf's liver during the 3 d study period) and above all carotenoids in non-milk foods. The same question can also be raised with respect to linoleic acid, Fe and vitamins B<sub>1</sub>, B<sub>3</sub>, and C, the intake of which was also elevated in several children. Carriquiry and Allen have demonstrated that this discussion about the excessive intake of most nutrients is at present fruitless because we lack knowledge of the risk thresholds, there are methodological uncertainties, and food consumption surveys report information derived from disciplined, regular meals, rather than from day-to-day life<sup>(6,13)</sup>.

## Conclusion

The results of the present study confirm that the daily consumption of ≥250 ml CM by young children aged 1–2 years partly increases the risk of a high protein intake and significantly increases the risk of an insufficient intake of nutrients of paramount importance, namely α-linolenic acid iron, vitamin C and vitamin D. The daily use of ≥250 ml GUM as a substitute for CM can prevent these nutritional risks, except for vitamin D. Randomized clinical trials comparing plasma levels of Fe, vitamin C, vitamin D and α-linolenic acid in young children fed CM or GUM are needed.

## Acknowledgements

The present study received no financial support from any baby food company or from the French Association of Baby Food Industries or any other public or private support. The declaration of interests of the authors is as follows: J.G., none; M.F., Lactalis (funding for the research unit of CREABio); D.T., Danone (funding for the university research lab), Mead Johnson, Nestlé, United Pharmaceuticals (refund for travel expenses or congress expenses); G.P.d.C., none; M.V., none. Author contributions: J.G. wrote the largest part of the paper and performed data analyses; M.F. organized the logistics of the study, reviewed data questionnaires and performed data analyses; D.T. and M.V. performed critical revision of the

paper for important intellectual content; G.P.d.C. assisted with interpretation of data.

## References

1. Council of Europe (2009) Directive 2009/39/EC of the European Parliament and of the Council of 6 May 2009 on foodstuffs intended for particular nutritional uses. *Official Journal of the European Union* 20.5.2009, L124/21–L124/29; available at [http://www.fsai.ie/uploadedFiles/Dir2009\\_39.pdf](http://www.fsai.ie/uploadedFiles/Dir2009_39.pdf)
2. Fantino M & Gourmet E (2008) Nutrient intakes in 2005 by non breast-fed French children of less than 36 months. *Arch Pediatr* **15**, 446–455.
3. SU.VI.MAX (2006) Food composition table. <http://www.lavoisier.fr/notice/fr281783.html> (accessed December 2011).
4. Beaufrère B, Briend A, Ghisolfi J *et al.* (2001) Nutrition Committee of the French Society of Paediatrics. Recommended dietary allowances for infants, children and adolescents. In *Recommended Dietary Allowances for the French Population*, 3rd ed., pp. 255–291 [A Martin, editor]. Paris: Editions TEC et DOC.
5. Agence Française de Sécurité Sanitaire des Aliments (2004) Specification for the selection of a Nutrient–Vector Food Pair. <http://www.afssa.fr/Documents/NUT-Ra-Alimentsvecteurs.pdf> (accessed December 2011).
6. Carriquiry AL (1999) Assessing the prevalence of nutrient adequacy. *Public Health Nutr* **2**, 23–33.
7. De Lauzon B, Volatier JL, Martin A *et al.* (2004) A Monte Carlo simulation to validate the EAR cut-point method for assessing the prevalence of nutrient inadequacy at the population level. *Public Health Nutr* **7**, 893–899.
8. Male C, Persson LA, Freeman V *et al.* (2001) Prevalence of iron deficiency in 12-mo-old infants from 11 European areas and influence of dietary factors on iron status (Euro-growth study). *Acta Paediatr* **90**, 492–498.
9. Gordon CM, Feldman HA, Sinclair L *et al.* (2008) Prevalence of vitamin D deficiency among healthy infants and toddlers. *Arch Pediatr Adolesc Med* **162**, 505–512.
10. Misra M, Pacaud D, Petryk A *et al.* (2008) Vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics* **122**, 398–417.
11. Devaney B, Ziegler P, Pac S *et al.* (2004) Nutrient intakes of infants and toddlers. *J Am Diet Assoc* **104**, Suppl. 1, S14–S121.
12. Bender DA (2003) Do we really know vitamin and mineral requirements for infants and children? *J R Soc Health Promot* **123**, 154–158.
13. Allen LH (2006) Setting dietary reference intakes for children. In *Dietary Reference Intakes Research Synthesis. Symposium Food and Nutrition Board*, pp. 95–105. Washington, DC: The National Academies Press.
14. Aggett PJ, Bresson JL, Haschke F *et al.* (1997) Recommended nutrient allowances (RDAs), recommended dietary intakes (RDIs), recommended nutrient intakes (RNIs) and population reference intakes (PRIs) are not 'recommended intakes'. *J Pediatr Gastroenterol Nutr* **25**, 236–241.
15. Krebs NF (2006) The use of zinc stable isotopes to inform the dietary reference intake process for infants and children. In *Dietary Reference Intakes Research Synthesis. Symposium Food and Nutrition Board*, pp. 105–112. Washington, DC: The National Academies Press.
16. Azais-Braesco V, Bruckert E, Durier P *et al.* (2001) Vitamin E. In *Recommended Dietary Allowances for the French Population*, 3rd ed., pp. 236–243 [A Martin, editor]. Paris: Editions TEC et DOC.

17. Carriquiry AL & Camano-Garcia G (2006) Evaluation of dietary intake data using the tolerable upper intake levels. *J Nutr* **136**, issue 2, 507S–513S.
18. Institute of Medicine (2000) *Dietary Reference Intakes: Applications in Dietary Assessment*. Washington, DC: The National Academies Press.
19. Agostoni C, Decsi T, Fewtrell M *et al.* (2008) Complementary feeding: a commentary by the EPSGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr* **46**, 99–110.
20. Briefel R, Hanson C, Fox MF *et al.* (2006) Feeding infants and toddlers study: do vitamin and mineral supplements contribute to nutrient adequacy or excess among US infants and toddlers? *J Am Diet Assoc* **106**, Suppl. 1, S52–S65.
21. Food and Nutrition Board, Institute of Medicine (2002) *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium and Zinc*. Washington DC: The National Academies Press.
22. Scientific Committee on Food (2002) Opinion of the Scientific Committee on Food on the tolerable upper intake level of preformed vitamin A (retinol and retinyl esters). [http://ec.europa.eu/food/fs/sc/scf/out145\\_en.pdf](http://ec.europa.eu/food/fs/sc/scf/out145_en.pdf) (accessed December 2011).