CrossMark

ERRATUM

Erratum to: ω-Amidase: an underappreciated, but important enzyme in L-glutamine and L-asparagine metabolism; relevance to sulfur and nitrogen metabolism, tumor biology and hyperammonemic diseases

Arthur J. L. Cooper[1] · Yevgeniya I. Shurubor[2,3] · Thambi Dorai[4] · John T. Pinto[1] ·
Elena P. Isakova[3] · Yulia I. Deryabina[3] · Travis T. Denton[5] · Boris F. Krasnikov[1,3]

Erratum to: Amino Acids DOI 10.1007/s00726-015-2061-7

In the original publication of the article, Fig. 1 and Acknowledgments were wrongly published. The correct Fig. 1 and acknowledgments are given below.

✉ Arthur J. L. Cooper
Arthur_Cooper@nymc.edu

[1] Department of Biochemistry and Molecular Biology, New York Medical College, Valhalla, NY 10595, USA

[2] Department of Neurology, The Brain and Mind Research Institute, Weill Cornell Medical College, New York, NY 10021, USA

[3] A.N. Bach Institute of Biochemistry, Russian Academy of Sciences, 33 Leninsky Prospekt, Moscow 119071, Russia

[4] Department of Urology, New York Medical College, Valhalla, NY 10595, USA

[5] Department of Pharmaceutical Sciences, College of Pharmacy, Washington State University, Spokane, WA 99210, USA

Fig. 1 Glutaminase II (**a**) and asparaginase II (**b**) pathways. Note the central importance of ω-amidase in hydrolyzing the α-keto acids derived from transamination of both L-glutamine (i.e., α-ketoglutaramate, KGM) and L-asparagine (α-ketosuccinamate, KSM). *Gln Ts* glutamine transaminases, *Asn Ts* asparagine transaminases