

Obesity in COVID-19: A Systematic Review and Meta-analysis

Jamie SY Ho, ^{*1}*MBBChir*, Daniel I Fernando, ^{*1}*MBBChir*, Mark Y Chan, ^{2,3}*MBBS*, Ching-Hui Sia, ^{2,3}*MBBS*

Abstract

Objective: Obesity has been shown to be associated with adverse outcomes in viral infections such as influenza, but previous studies on coronavirus disease 2019 (COVID-19) had mixed results. The aim of this systematic review is to investigate the relationship between COVID-19 and obesity.

Methods: We performed a systematic review and meta-analysis. A literature search of MEDLINE, EMBASE, Scopus, Web of Science, CENTRAL, OpenGrey and preprint servers medRxiv and bioRxiv was performed, with no restriction on language or date of publication. Primary outcomes of this study were intensive care unit (ICU) admission or critical disease, severe disease and mortality. Secondary outcome was a positive COVID-19 test. Meta-analysis was performed using OpenMeta-Analyst software, and heterogeneity was tested using Cochran's Q test and *I*² statistic. The study protocol was registered on PROSPERO (CRD42020184953).

Results: A total of 1,493 articles were identified and 61 studies on 270,241 patients were included. The pooled prevalence of obesity was 27.6% (95% confidence interval [CI] 22.0–33.2) in hospitalised patients. Obesity was not significantly associated with increased ICU admission or critical illness (odds ratio [OR] 1.25, 95% CI 0.99–1.58, *P*=0.062, *I*²=31.0) but was significantly associated with more severe disease (OR 3.13, 95% CI 1.41–6.92, *P*=0.005, *I*²=82.6), mortality (OR 1.36, 95% CI 1.09–1.69, *P*=0.006, *I*²=88.5) and a positive COVID-19 test (OR 1.50, 95% CI 1.25–1.81, *P*<0.001).

Conclusion: Obesity increased the risk of severe disease, mortality and infection with COVID-19. Higher body mass index was associated with ICU admission and critical disease. Patients who are obese may be more susceptible to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection, and infected patients should be monitored closely for adverse outcomes.

Ann Acad Med Singap 2020;49:996-1008

Keywords: Body mass index, coronavirus, intensive care, mortality, prognosis

Introduction

Coronavirus disease 2019 (COVID-19) is an ongoing global pandemic infection by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It is established that increasing age and comorbidities such as cardiovascular diseases are associated with risk of infection, more severe disease and adverse outcomes.¹

Obesity is an epidemic globally, causing more than 2.8 million deaths per year worldwide in 2019. The World Health Organization (WHO) defines obesity as a body mass index (BMI) of ≥ 30 , and overweight as a BMI of ≥ 25 . Other definitions of obesity include the WHO Asia-Pacific BMI guidelines, which defines obese as BMI ≥ 27.5 and overweight as BMI ≥ 23 .²

¹ School of Clinical Medicine, University of Cambridge, Cambridge, UK

² Department of Cardiology, National University Heart Centre, Singapore

³ Department of Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Singapore

Address for Correspondence: Dr Ching-Hui Sia, Department of Cardiology, National University Heart Centre, 5 Lower Kent Ridge Rd, Singapore 119074.

Email: ching_hui_sia@nuhs.edu.sg

*Joint first authors

Obese patients were at increased risk of hospitalisation, complications and death in the 2009 influenza A (H1N1) pandemic.³ There are suggestions that, similar to influenza, obesity may be a significant risk factor in COVID-19. Possible pathophysiology includes a chronic proinflammatory environment, reducing immunological response to infections, and altered dynamics of pulmonary ventilation with reduced diaphragmatic expansion and increased anatomical dead space.⁴ However, the evidence from reported cohort studies and case series has been mixed.^{5,6} In this comprehensive systematic review and meta-analysis of the current literature on obesity and COVID-19, we aimed to characterise the relationship between obesity and adverse prognostic outcomes.

Methods

The protocol for this review was registered and published on the International Prospective Register of Systematic Reviews (PROSPERO; CRD42020184953). A literature search of MEDLINE, EMBASE, Scopus, Web of Science, CENTRAL, OpenGrey and preprint servers medRxiv and bioRxiv was performed on 8 May 2020, using the search terms “coronavirus or COVID” or “SARS-COV-2” or similar terms to the pandemic, and “obesity or hyperphagia or overweight” or words to that effect. Additional articles were identified from hand searching of the reference lists of included studies. Inclusion criteria were clinical studies (1) that reported obesity prevalence or outcomes, and (2) that were performed on COVID-19 patients. All research study types, such as case series, cohort studies, longitudinal studies and randomised controlled trials, were included, with no restriction on publication date or language of publication. Articles discussing other infectious outbreaks and studies on animals or *in vitro* studies were excluded. Other systematic reviews, literature reviews, editorials and opinion articles were excluded, but references were screened for relevant articles. Articles from the Chinese WanFang and SinoMed databases were not searched owing to language limitations.

Titles and abstracts were screened independently by 2 researchers, and discrepancies were resolved through discussion or involvement of a third researcher. Full texts were identified, and data were extracted onto a standardised data extraction form by the 2 independent researchers. The extracted data included study type, patient characteristics, prevalence of obesity, and clinical outcomes such as severe disease, intensive care unit (ICU) admission and mortality. The quality of the included studies was assessed using the Newcastle-

Ottawa Scale for case series, cohort, cross-sectional and case-control studies.

The primary outcomes of this systematic review were prevalence of obesity in COVID-19 patients, and association of obesity with adverse outcomes such as ICU admission, critical illness, severe disease and mortality. The secondary outcome was a positive COVID-19 test.

Definitions

The definitions of terms including “obesity”, “severe disease” and “critical disease” used for each individual study were adopted in this meta-analysis. Both European and Asian definitions of obesity were used, for example, based on the geographical origin of the study. Critical illness was defined as any of the following: respiratory failure requiring invasive mechanical ventilation, shock, or any other organ failure, requiring intensive therapy unit (ITU) or ICU monitoring and treatment. Severe disease was defined according to the WHO interim guidance, and its definition was used by most study types that made the distinction between different severities of COVID-19.

For “positive COVID-19 test” results to be included in the analyses, COVID-19 needs to be diagnosed by reverse transcription-polymerase chain reaction (RT-PCR) testing using a nasopharyngeal swab.

Statistical analyses

A meta-analysis was performed for the primary and secondary outcomes using the OpenMeta-Analyst software. DerSimonian and Laird’s random effects model was employed to calculate the pooled odds ratio (OR) and 95% confidence interval (CI). Study heterogeneity was assessed using Cochran’s Q test and I^2 statistic. A P value of <0.05 was considered statistically significant.

Results

In total, 61 studies on 270,241 patients fit the inclusion criteria, including 38 cohort studies, 2 case-control studies, 8 cross-sectional studies and 13 case series (Fig. 1). The characteristics of the included studies are presented in Table 1. The quality of studies was generally moderate to good, with the exception of small case series that had a high risk of selection bias and lack of control group. A total of 46 studies were included in the meta-analysis, with any outcome analysed. The remaining 15 studies were descriptive studies including case series and case reports, and their reported patient characteristics are presented in Table 1. The prevalence of obesity was




Fig. 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram of the process of literature search, screening and inclusion of studies

27.6% (95% CI 22.0–33.2) across 31 cohort studies in the hospital setting,^{7,35} and univariate meta-regression model revealed no significant association with sample size (regression co-efficient <0.0001 , $P=0.921$). Only 31 cohort studies reported prevalence of obesity in the hospital setting and so were included with the pooled prevalence meta-analysis. On exclusion of 4 Chinese studies owing to ethnic differences in BMI,^{14,26,27,31} the prevalence of obesity was 31.1% (95% CI 25.1–37.2). The prevalence of obesity in Chinese studies alone was 5.4% (95% CI 1.9–8.9%).

ICU admission and critical illness

A total of 12 studies reported the outcome of ICU admission or critical illness, and comprised 9 cohort studies (8 retrospective, 1 prospective), 2 case series and 1 cross-sectional study, involving 10,314 patients with COVID-19.^{7,11,12,14–16,28,36–40} All studies were of moderate to good quality and were conducted in the hospital setting. The proportion of obese patients (BMI ≥ 30) in ICU or critical patients was compared with that of non-ICU hospitalised patients in 8 studies involving 9,869 patients.^{7,11,12,14–16,28,36} On meta-analysis, obesity was not significantly associated with increased ICU admission or critical illness (OR 1.25, 95% CI 0.99–1.58, $P=0.062$), with moderate heterogeneity ($Q=10.1$, $I^2=31.0$) (Fig. 2A).

Five studies involving 1,445 participants reported mean or median BMI of ICU patients and non-ICU

patients in the hospital setting (Fig. 2B).^{7,37–40} The ICU or critical illness group had a mean BMI of 28.7 (26.1–31.4) while the non-ICU group had a BMI of 25.2 (95% CI 22.0–28.5). The overall mean difference in BMI was 2.32 (95% CI 1.04–3.60, $P<0.001$), with substantial heterogeneity ($Q=13.0$, $I^2=69.2$). In particular, the study by Lau et al. showed a particularly large effect size and 95% CI, and the study focused on ICU admissions with recorded levels of serum 25-hydroxycholecalciferol in a single centre, which may not be representative of typical patients admitted to ICU for COVID-19.³⁷ There may also be ethnic differences across study populations, as the 2 studies from China^{39,40} reported lower BMIs than those from the US and Mexico,^{7,37,38} although the mean differences were found to be similar in magnitude.

Three studies found an increased risk of invasive mechanical ventilation in patients with obesity,^{18,21,34} including 1 study with a large cohort of 20,737 patients in the US.³⁴ A cohort study from France⁵ found that the risk of intubation increased in 124 ICU patients with BMI ≥ 35 (OR 7.36, 95% CI 1.63–33.14) but not in patients with BMI 30–35 (OR 3.45, 95% CI 0.83–14.31), which was similar to another cohort study involving 291 patients in France with BMI ≥ 35 (OR 6.24, 95% CI 2.30–16.93) and BMI 30–35 (OR 1.97, 95% CI 1.00–3.90). A case series of 2 obese patients reported perforation of the membranous trachea or cricoid membrane during intubation for acute

Table 1. Characteristics and quality assessment of included studies

Authors, Country, year	Study design	n	Setting	BMI, obesity (%)	Age (years)
Abou-Arab et al., ⁴¹ France, 2020	Case series	2	ICU	41, 34	59, 67
Argenziano et al., ⁷ US, 2020	Case series	1000	Hospital	Median 28.6 (25.2–33.1); ≥30, 48.3%	63 (50–75)
Auld et al., ⁶ US, 2020	Retrospective cohort	217	ICU	≥40, 9.7%	64 (54–73)
Barrasa et al., ⁸ Spain, 2020	Retrospective cohort	48	ICU	≥30, 48%; 30–40, 31%; >40, 14%	63 (51–75)
Bello-Chavolla et al., ⁹ Mexico, 2020	Retrospective cohort	15529	Hospital	≥30, 20.7%	46.6±15.5
Bhatraju et al., ^a US, 2020	Case series	24	Hospital	Mean 33.2±7.2	64±18
Borobia et al., ²⁰ Spain, 2020	Case series	2226	Hospital	≥30, 10.9%	61 (46–78)
Caussy et al., ⁵ France, 2020	Retrospective cohort	291	ICU	<25, 25.4%; 25–30, 41.6%; 30–35, 21.6%; >35, 11.3%	NR
Chaw et al., ^b Brunei, 2020	Cross-sectional	71	Community	≥30, 5.6%	33
Chen et al., ⁴³ China, 2020	Retrospective cohort	145	Hospital	Mean 23.2 (non-severe); mean 24.8 (severe)	47.5±14.6
Cummings et al., ¹⁰ US, 2020	Prospective cohort	1150	Hospital	Mean 30.8±7.7; ≥30, 46%; ≥35, 26%; ≥40, 13%	62 (51–72)
Dauchet et al., ^c France, 2020	Prospective cohort	187	Hospital	≥25, 43% of ICU; ≥30, 43% of ICU	NR
Docherty et al., ⁴⁷ UK, 2020	Prospective cohort	16749	Hospital		72 (57–82)
Du et al., ⁴⁶ China, 2020	Retrospective cohort	245	Hospital		55
Ebinger et al., ¹¹ US, 2020	Retrospective cohort	442	Hospital	≥30, 16%	52.7±19.7
Freedberg et al., ⁴ US, 2020	Retrospective cohort	1620	Hospital	Median 28.1 (24.9–32.6)	65 (52–77)
Gaibazzi et al., ²² US, 2020	Case series	279	Hospital	≥30, 16%	72 (60–80)
Garg et al., ²³ US, 2020	Retrospective cohort	178	Hospital	≥30, 48.3%	NR
Giacomelli et al., ¹² Italy, 2020	Prospective cohort	233	Hospital	≥30, 16.3%	61 (50–72)
Giorgi Rossi et al., ²⁵ Italy, 2020	Prospective cohort	2653	Hospital	≥30, 2.5%	<51y, 26.2%; 51–60y, 19.9%; 61–70y, 15.6%; >71y, 38.3%
Guo et al., ²⁶ China, 2020	Retrospective cohort	159	Hospital	≥30, 1.72%	71
Hadijadi et al., ²⁶ France, 2020	Case-control	50	NR	≥25, 10%; ≥30 excluded	55 (50–63)

BMI: body mass index; C: comparability; ICU: intensive care unit; NR: not reported

^aBhatraju PK, Ghassemi BJ, Nichols M, et al. COVID-19 in critically ill patients in the Seattle region – case series. *N Engl J Med* 2020;382(21):2012-22.

^bChaw L, Koh W, Jamaludin S, et al. Analysis of SARS-CoV-2 transmission in different settings, Brunei. *Emerg Infect Dis* 2020;26(11):2598-606.

^cDauchet L, Lambert M, Gauthier V, et al. ACE inhibitors, ATI receptor blockers and COVID-19: clinical epidemiology evidences for a continuation of treatments. The ACER-COVID study. *medRxiv* 2020. DOI: doi.org/10.1101/2020.04.28.20078071.

^dFreedberg DE, Congiario J, Wang TC, et al. Famotidine use is associated with improved clinical outcomes in hospitalized COVID-19 patients: a propensity score matched retrospective cohort study. *Gastroenterology* 2020;159:1129-31.e3.

Superscript numbers: refer to References

Table 1. Characteristics and quality assessment of included studies (Cont'd)

Authors, Country, year	Study design	n	Setting	BMI, obesity (%)	Age (years)
Ho et al., ¹³ UK, 2020	Retrospective cohort	340	Community	Mean 29.0±5.3; ≥25, 44.1%; ≥30, 34.1%	57.7±8.5
Hogan et al., ^e US, 2020	Cross-sectional	85	Hospital	≥30, 37.7%	55
Hu et al., ¹⁴ China, 2020	Retrospective cohort	323	Hospital	25–30, 16.1%; ≥30, 4%	61 (range 23–91)
Huang et al., ²⁷ China, 2020	Retrospective cohort	125	Hospital	>26, 5.6%	44.9±18.6
Kalligeros et al., ¹⁵ US, 2020	Retrospective cohort	103	Hospital	≥30, 47.5% (hospitalized), 56.8% (ICU), 65.5% (ventilated)	60 (52–70)
Kass et al., ^f US, 2020	Retrospective cohort	265	Hospital	Median 29.3	NR
Lau et al., ³⁷ US, 2020	Retrospective cohort	20	Hospital	Mean 31.4±9.3	65.2±16.2
Li et al., ^g China, 2020	Retrospective cohort	17	Hospital	Mean 25±4.7	45.1±12.8
Li et al., ^h US, 2020	Cross-sectional	NR	Community	NR	NR
Liao et al., ⁱ China, 2020	Retrospective cohort	46	Hospital	≥25, 37.0%	Range 10–35
Liao et al., ^j China, 2020	Retrospective cohort	539	Hospital	≥24, 50.6%; Median 24.0 (21.5–27.3)	50 (39–65)
Lighter et al., ²⁸ US, 2020	Retrospective cohort	3615	Hospital	30–34, 21%; >35, 16%	NR
Liu et al., ⁴⁴ China, 2020	Case series	30	Hospital	Mean 22.0±1.3 (non-severe); mean 27.0±2.5 (severe)	NR
Lochlainn et al., ^k UK, 2020	Retrospective cohort	171899	Community	Mean 27.1±5.9 (not hospitalised); mean 28.6±6.5 (hospitalised)	40.3±13.9
Mahévas et al., ²⁹ France, 2020	Retrospective cohort	181	Hospital	>30, 27.4%	60 (52–68)
Menter et al., ³⁰ Switzerland, 2020	Case series	21	Hospital	≥25, 48%; ≥30, 5%; ≥35, 5%; ≥40, 19%	76 (range 53–96)
Mercuro et al., ^l US, 2020	Retrospective cohort	90	Hospital	Mean 31.5±6.6	60.1±16.7
Nguyen et al., ^m Vietnam, 2020	Cross-sectional	3947	Hospital	≥25, 10.9%	44.4±17.0

^eHogan CA, Stevens BA, Sahoo MK, et al. High frequency of SARS-CoV-2 RNAemia and association with severe disease. *Clin Infect Dis* 2020;ciaa1054.

^fKass DA, Duggal P, Cingolani O. Obesity could shift severe COVID-19 disease to younger ages. *Lancet* 2020;395:1544-45.

^gLi AY, Hammah TC, Durbin JR, et al. Multivariate analysis of black race and environmental temperature on COVID-19 in the US. *Am J Med Sci* 2020;360:348-56.

^hLi J, Li S, Cai Y, et al. Epidemiological and clinical characteristics of 17 hospitalized patients with 2019 novel coronavirus infections outside Wuhan, China. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.02.11.20022053>.

ⁱLiao J, Fan S, Chen J, et al. Epidemiological and clinical characteristics of COVID-19 in adolescents and young adults. *Innovation* 2020;1:100001.

^jLiao X, Chen H, Wang B, et al. Critical care for severe COVID-19: a population-based study from a province with low case-fatality rate in China. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.03.22.20041277>.

^kLochlainn MN, Lee KA, Sudre CH, et al. Key predictors of attending hospital with COVID19: an association study from the COVID Symptom Tracker App in 2,618,948 individuals. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.25.20079251>.

^lMercuro NJ, Yen CF, Shim DJ, et al. Risk of QT interval prolongation associated with use of hydroxychloroquine with or without concomitant azithromycin among hospitalized patients testing positive for coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020:e201834.

^mNguyen HC, Nguyen MH, Do BN, et al. People with suspected COVID-19 symptoms were more likely depressed and had lower health-related quality of life: the potential benefit of health literacy. *J Clin Med* 2020;9:965.

Superscript numbers: refer to References

Table 1. Characteristics and quality assessment of included studies (Cont'd)

Authors, Country, year	Study design	n	Setting	BMI, obesity (%)	Age (years)
Peng et al., ³⁹ China, 2020	Retrospective cohort	112	Hospital	Median 22.0 (20.0–25.0)	62.0 (55.0–67.0)
Petrilli et al., ¹⁶ US, 2020	Cross-sectional	4103	Hospital	≥30, 26.8%	52 (36–65)
Piva et al., ¹⁷ Italy, 2020	Prospective cohort	33	ICU	Median 27.8 (27.0–32.1); ≥25, 58%; ≥30, 31%	64 (59–72)
Qi et al., ³¹ China, 2020	Case series	267	Hospital	≥30, 11.2%	48.0 (35.0–65.0)
Ramireddy et al., ^a US, 2020	Case series	98	Hospital	Mean 27.8±6.6	62.3±17.0
Rentsch et al., ⁴⁹ US, 2020	Retrospective cohort	585	Community	>30, 40.8%	66.1 (60.4–71.0)
Reyes Gil et al., ⁴⁸ US, 2020	Case series	217	Hospital	Median 29.0 (26.0–33.1) (died), median 29.5 (25.9–33.5) (discharged)	NR
Richardson et al., ³² US, 2020	Case series	5700	Hospital	>30, 41.7%; >35, 19%	63 (52–75)
Rodríguez-Cola et al., ²⁴ Spain, 2020	Prospective cohort	7	Hospital	≥30, 42.9%	68 (34–75)
Simonnet et al., ¹⁸ France, 2020	Retrospective cohort	124	ICU	>30, 47.6%; >35, 28.2%	60 (51–70)
Tahmasebi et al., ⁶ US, 2020	Cross-sectional	NR	NR	NR	NR
Vahidy et al., ¹⁹ US, 2020	Cross-sectional	754	Hospital, community	≥30, 7.7%	50.6±18.9
Valente-Acosta et al., ³⁸ Mexico, 2020	Retrospective case series	33	Hospital	Mean 26.9±4.1 (severe pneumonia); mean 30.2±6.2 (critical pneumonia); ≥25, 60% (severe pneumonia), 87.5% (critical pneumonia)	59.4±13.2 (severe pneumonia); 64.6±9.3 (critical pneumonia)
van der Voort et al., ⁵⁴ Netherlands, 2020	Cross-sectional	31	ICU	Mean 31 (range 24.8–48.4)	NR
Vuagnat et al., ³³ France, 2020	Prospective cohort	59	Hospital	≥30, 17%; median 26 (22–30)	58 (48–68)
Wang et al., ⁴⁵ US, 2020	Case series	6158	Hospital	Median 27.6 (24.1, 32.6) (hospitalised), median 27.6 (23.9, 32.5) (died); median 28.3 (24.8, 32.7) (discharged)	NR
Williamson et al., ^p UK, 2020	Retrospective cohort	5683	Community	25–30, 29.3%; 30–35, 20.5%; 35–40, 8.2%; ≥40, 4.5%	NR
Wollenstein-Betech et al., ³⁴ US, 2020	Retrospective cohort	20,737	Hospital	≥30, 33.7%	NR
Wu et al., ⁴⁰ China, 2020	Retrospective cohort	280	Hospital	Mean 24.1±3.0	43.1±19.0
Zheng et al., ⁴² China, 2020	Retrospective cohort	66	Hospital	Mean 26.5±3.9	47
Zao et al., ³⁵ US, 2020	Case-control	50	Hospital	≥30, 46%	61±15

^a Ramireddy A, Chugh H, Reinier K, et al. Experience with hydroxychloroquine and azithromycin in the coronavirus disease 2019 pandemic: implications for QT interval monitoring. *J Am Heart Assoc* 2020;9:e017144.z

⁶ Tahmasebi P, Shokri-Kuehni SMS, Sahimi M, et al. How do environmental, economic and health factors influence regional vulnerability to COVID-19? medRxiv 2020. DOI: <https://doi.org/10.1101/2020.04.09.20059659>.

^p Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;584:430-6. Superscript numbers: refer to References




Fig. 2. Meta-analysis of the proportion of COVID-19 patients with obesity, comparing (A) the prevalence in intensive care unit (ICU) or with critical illness, to that in hospitalised non-ICU patients; and (B) the mean body mass index in ICU patients to that in non-ICU patients. CI: confidence interval; I^2 : I-squared statistic of heterogeneity

respiratory distress syndrome, which was refractory to prone positioning and non-invasive ventilation.⁴¹

Severe disease

Severe disease was defined according to the WHO interim guidance, and the definition was very similar to that according to the National Health Commission of the People’s Republic of China. Severe disease was defined as respiratory distress or respiratory rate ≥ 30 /min, oxygen saturation $\leq 93\%$ at rest, or the ratio of arterial oxygen partial pressure to fraction of inspired oxygen (PaO_2/FiO_2) ≤ 300 mmHg. Eight studies involving 1,566 patients compared COVID-19 patients having severe disease with those having non-severe disease, including 6 retrospective cohort studies and 2 case series.^{14,27,31,36,40,42-44}

Of the 8 studies, 6 studies (1,111 patients) compared the proportion of obesity in patients having severe disease with that in patients having non-severe disease, and were of moderate to good quality.^{14,27,31,36,40,42} A meta-analysis found that obesity was significantly associated with more severe disease in hospitalised patients (OR 3.13, 95% CI 1.41–6.92, $P=0.005$) (Fig. 3). However, the studies had substantial and significant heterogeneity ($Q=28.8, I^2=82.6$), despite 5 of the 6 studies being performed in China. Differences in patient characteristics, investigations and treatment may have contributed to significant variability in these results. The pooled mean BMI was 25.6 (24.8–26.4) in the severe disease group and 23.0 (22.0–24.1) in the non-severe disease group of 3 studies.^{40,43,44} The overall mean




Fig. 3. Meta-analysis of the proportion of COVID-19 patients with obesity, comparing the prevalence in patients with severe disease to those with non-severe disease.

CI: confidence interval; I^2 : I-squared statistic of heterogeneity

difference in BMI was 2.37 (1.12–3.63, $P<0.001$), and there was substantial heterogeneity among the studies ($Q=6.9$, $I^2=71.0$). One study combined the severe disease group with the critical disease group.⁴⁰ The sensitivity analysis, which excluded this study by Wu et al., found that the proportion of obesity remained higher in severe disease (OR 4.89, 95% CI 1.39–17.18) than that in non-severe disease, with a mean difference in BMI of 3.02 (95% CI 0.50–5.54).

Notably, the study by Zheng et al. found a particularly large effect size (OR 72.3, 95% CI 9.1–573.7).⁴² This study only included patients with non-alcoholic fatty liver disease and may not be representative of unselected general hospital patient cohorts used in the other studies. A sensitivity analysis performed excluding this study found no significant effects on the results.

Mortality

The outcome of mortality was reported in 12 studies involving 45,768 patients, with 11 studies based on hospital inpatients^{9,10,12,20,22,25,39,45,46,47,48} and 1 study in the ICU setting.⁶ Overall, obesity was associated with increased risk of mortality (OR 1.33, 95% CI 1.07–1.66, $P=0.011$), with significant and substantial heterogeneity ($Q=95.8$, $I^2=88.5$) (Fig. 4). In the subgroup of studies performed in general hospitalised patients, mortality remained significantly associated with obesity (OR 1.36, 95% CI 1.09–1.69, $P=0.006$) with substantial heterogeneity ($Q=91.7$, $I^2=89.1$). The cohort study

performed by Auld et al.,⁶ which was the only study investigating the mortality rates in the ICU setting, found a smaller proportion of obese patients in those who died, in contrast to studies performed in the general hospital setting where non-survivors had a larger proportion of obese patients. Further research on obese patients in the ICU setting may be useful in clarifying their prognosis and outcomes in order to investigate the outlier results found in the study by Auld et al.

Peng et al.³⁹ showed that obese patients had an OR of 32.1 (95% CI 6.7–153.0) for mortality that was significantly larger than those in other studies. The study focused on COVID-19 patients with cardiovascular disease first and, within this subpopulation, obese patients were identified, whereas the other studies recruited unselected patients with severe or critical COVID-19. This differing population could explain the higher OR for mortality, given the compounding effect of both cardiovascular disease and obesity on mortality. Exclusion of this study did not have a significant effect on the results of the meta-analysis.

Positive test

Four studies involving 17,208 patients in the community and hospitals investigated the proportion of obese patients in patients who tested positive for COVID-19 and compared with that in patients who tested negative on RT-PCR of nasopharyngeal swab samples.^{9,13,19,49} The quality of these studies was moderate to good, and 3




Fig. 4. Meta-analysis of the proportion of COVID-19 patients with obesity, comparing the prevalence in patients with severe disease to those with non-severe disease. CI: confidence interval; I^2 : I-squared statistic of heterogeneity

were retrospective cohort studies while 1 was a cross-sectional study. Meta-analysis found that obesity was significantly associated with a positive COVID-19 test (OR 1.50, 95% CI 1.25–1.81, $P<0.001$), and there was significant heterogeneity among the studies ($Q=13.2$, $I^2=77.2$; Fig. 5). The study by Vahidy et al.¹⁹ focused on a population from a single health system in the US, in contrast to the use of national databases (e.g. the UK biobank¹⁹ or the Mexican national database⁹) in the other studies in this subgroup analysis.

Discussion

Obesity and cardiovascular diseases have been linked to viral infections, including influenza,^{3,50} and are risk factors for poor prognosis and severe disease. In this systematic review and meta-analysis, we found that obesity was significantly associated with a positive COVID-19 test, severe disease and mortality, and patients admitted to ICU or with critical disease

had significantly higher BMI. Many of the initial observational studies did not report obesity or BMI, and an earlier previous systematic review included 3 studies that concluded that obesity was a risk factor for severe disease and for requirement of advanced medical care.⁵¹ In this review, we further characterised the prognostic implications of obesity across a larger quantity of subjects and studies compared with previous systematic reviews.⁵²

The mechanisms for adverse outcomes in SARS-CoV-2 infection associated with obesity may be multifaceted. Mechanically, obesity is associated with reduced diaphragmatic expansion, increased anatomical dead space and increased difficulty of ventilation. Obesity is characterised by a state of chronic low-grade inflammation with elevated levels of proinflammatory cytokines such as interleukin-6, the levels of which were also observed to be elevated in non-survivors compared with survivors in COVID-19.⁵³ Leptin




Fig. 5. Meta-analysis of the proportion of COVID-19 patients with obesity, comparing the prevalence in patients who had a positive COVID-19 test to that in patients who tested negative. CI: confidence interval; I²: I-squared statistic of heterogeneity

promotes B cell maturation and inhibits anti-viral CD8⁺ T cell response. Leptin levels are increased in obesity, possibly reducing the effective immune response against viral infections.⁴ Intubated COVID-19 patients were shown to have an elevated leptin level compared with that of non-COVID-19 controls, despite similar BMI, suggesting that leptin may have a role in the pathogenesis of SARS-CoV-2 infection.⁵⁴ Adipose tissue also expresses high levels of angiotensin-converting enzyme-2 receptors, which mediate entry of SARS-CoV-2.

It is well-known that obesity is associated with diabetes mellitus, coronary artery disease and hypertension, forming the metabolic syndrome. In a cohort of 15,529 COVID-19 patients in Mexico, obese patients were more likely to have hypertension (33.7% versus 18.7%, *P*<0.001), diabetes (26.5% vs 16.2%, *P*<0.001) and cardiovascular disease (4.4% vs 2.6%, *P*<0.001).⁹ In multiple meta-analysis of comorbidities in COVID-19, hypertension, diabetes mellitus and cardiovascular disease were found to be associated with severe disease and ICU admission.⁵⁵

Increased BMI was significantly associated with ICU admission or critical illness, and obesity trended towards increasing risk, although not statistically significant. However, results from one study found that the proportion of obese patients who died in ICU was lower than that in survivors.⁶ This should be interpreted with caution owing to lack of data, and the lower mortality rate in obese patients in ICU may be related to the obesity paradox or selection bias of milder obese patients being admitted to ICU. A recent meta-analysis of 6 studies performed by Földi et al. found

that patients with obesity had increased risk of ICU admission (OR 1.21, 95% CI 1.002–1.46),⁵² which, combined with the analysis of 8 studies in this meta-analysis, suggests a relationship between obesity and ICU admission or critical disease.

The studies included in this review were significantly heterogeneous, suggesting that the true effect sizes were different among studies. Zheng et al. focused on a specific population of severe obesity disease (patients with metabolic associated fatty liver disease who are obese as opposed to obese patients alone) which, by definition, will have a compounding effect leading to an increased effect of obesity.⁴² This differing population could explain the outlier result when looking at mortality. The study by Auld et al.⁷ was the only one that specifically focused on an exclusive ITU population, whereas the other studies focused on an unselected general hospital population. Peng et al.³⁹ focused on cardiovascular disease patients with COVID-19, whereas most studies focused on unselected patients with severe or critical COVID-19. This differing population could explain the outlier result when looking at mortality. Additionally, cardiovascular comorbidities may confound the effects of obesity on the prognosis of COVID-19. Another cause may be the differences in ethnicity: the prevalence of obesity defined as BMI ≥30kg/m² was the lowest in Asians (9.8%) compared with those of the whites (22.0%), Latinos (33.6%) and African Americans (36.1%) in 42,935 adults in California, US.⁵⁶ Most studies adopted the WHO European definition of obesity, and future research should consider using the Asia-Pacific BMI classification in the Asian population.

Our meta-analysis also showed that obesity was significantly associated with a positive COVID-19 test (OR 1.50, 95% CI 1.25–1.81, $P < 0.001$). This finding adds to the evidence that obesity leads to increased infectivity of COVID-19 (positive tests as a proxy for infectivity). Most studies reported testing on symptomatic patients, which was the widely adopted method of testing during this stage of the pandemic, and it would be interesting to see studies of asymptomatic patients.⁵⁷ Recent large national studies utilising the UK primary care network⁵⁸ and the Information System for Research in Primary Care in Spain⁵⁹ found that obesity was associated with increased risk of positive COVID-19 testing. Reporting of obesity and positive test results may be a measure of how susceptible obese patients are to catching the virus, which could have implications on public health policies and healthcare utilisation.

Limitations of this study include the possibility of confounding factors and heterogeneity of the included studies. The heterogeneity in studies, mainly concentrating on study design, populations studied, geographical areas and definitions of COVID-19 severity used, will inevitably make it harder to draw definitive conclusions about the relationship between obesity and COVID-19 from across the literature. Only a subgroup of studies reported each of our identified primary and secondary outcomes, increasing the risk of type 1 and type 2 error in our conclusions. Seventeen articles from pre-print servers such as medRxiv that were not peer-reviewed but were included to ensure that the literature search was comprehensive, and the quality of these studies were assessed using validated tools. Publication bias was not assessed owing to the small number of studies in each meta-analysis.

Conclusion

Obesity was associated with an increased risk of severe disease, mortality and positive test for SARS-CoV-2 infection. Obesity trends towards higher risk of ICU admission and critical illness, and a higher BMI, rather than obesity, is associated with a statistically significant increase in risk of ICU admission and critical illness. Overall, this study suggests that obesity may be a marker of poor prognosis, and with the additional mechanistic challenges in ventilation, obese patients should be monitored closely and managed carefully. Compared with recently published systematic reviews and meta-analyses^{51,52} we analysed several important clinical outcomes (critical disease, severe disease and mortality) and comprehensively searched

the literature including pre-print texts and clinical trials for evidence, increasing the sample of studies from which to draw conclusions. Our findings support previous findings in the other systematic reviews. Future studies on COVID-19 patients should report prevalence of obesity and BMI in order to increase the understanding of the interaction between obesity and SARS-CoV-2 infection, and to inform the clinical management of these patients.

REFERENCES

1. Chew SY, Lee YS, Ghimiray D, et al. Characteristics and outcomes of COVID-19 patients with respiratory failure admitted to a “pandemic ready” intensive care unit – lessons from Singapore. *Ann Acad Med Singap* 2020;49:434-48.
2. WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004;363:157-63.
3. Morgan OW, Bramley A, Fowlkes A, et al. Morbid obesity as a risk factor for hospitalization and death due to 2009 pandemic influenza A(H1N1) disease. *PLoS One* 2010;5:e9694.
4. Michalakis K, Ilias I. SARS-CoV -2 infection and obesity: common inflammatory and metabolic aspects. *Diabetes Metab Syndr* 2020;14:469-71.
5. Caussy C, Pattou F, Wallet F, et al. Prevalence of obesity among adult inpatients with COVID-19 in France. *Lancet Diabetes Endocrinol* 2020;8:562-4.
6. Auld SC, Caridi-Scheible M, Blum JM, et al. ICU and ventilator mortality among critically ill adults with coronavirus disease 2019. *Crit Care Med* 2020;48:e799-e804.
7. Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020;369:m1996.
8. Barrasa H, Rello J, Tejada S, et al. SARS-CoV-2 in Spanish intensive care units: early experience with 15-day survival in Vitoria. *Anaesth Crit Care Pain Med* 2020;39:553-61.
9. Bello-Chavolla OY, Bahena-López JP, Antonio-Villa NE, et al. Predicting mortality due to SARS-CoV-2: a mechanistic score relating obesity and diabetes to COVID-19 outcomes in Mexico. *J Clin Endocrinol Metab* 2020;105:dga346.
10. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 2020;395:1763-70.
11. Ebinger JE, Achamallah N, Ji H, et al. Pre-existing traits associated with COVID-19 illness severity. *PLoS One* 2020;15:e0236240.
12. Giacomelli A, Ridolfo AL, Milazzo L, et al. 30-day mortality in patients hospitalized with COVID-19 during the first wave of the Italian epidemic: a prospective cohort study. *Pharmacol Res* 2020;158:104931.
13. Ho FK, Celis-Morales CA, Gray SR, et al. Modifiable and non-modifiable risk factors for COVID-19, and comparison to risk factors for influenza and pneumonia: results from a UK Biobank prospective cohort study. *BMJ Open* 2020;10:e040402.
14. Hu L, Chen S, Fu Y, et al. Risk factors associated with clinical outcomes in 323 coronavirus disease 2019 (COVID-19) hospitalized patients in Wuhan, China. *Clin Infect Dis* 2020;71:2089-98.
15. Kalligeros M, Shehadeh F, Mylona EK, et al. Association of obesity with disease severity among patients with coronavirus disease 2019. *Obesity (Silver Spring)* 2020;28:1200-4.

16. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. *BMJ* 2020;369:m1966.
17. Piva S, Filippini M, Turla F, et al. Clinical presentation and initial management critically ill patients with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in Brescia, Italy. *J Crit Care* 2020;58:29-33.
18. Simonnet A, Chetboun M, Poissy J, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity (Silver Spring)* 2020;28:1195-9.
19. Vahidy FS, Nicolas JC, Meeks JR, et al. Racial and ethnic disparities in SARS-CoV-2 pandemic: analysis of a COVID-19 observational registry for a diverse US metropolitan population. *BMJ Open* 2020;10:e039849.
20. Borobia AM, Carcas AJ, Arnalich F, et al. A cohort of patients with COVID-19 in a major teaching hospital in Europe. *J Clin Med* 2020;9:1733.
21. Caussy C, Wallet F, Laville M, et al. Obesity is associated with severe forms of COVID-19. *Obesity (Silver Spring)* 2020;28:1175.
22. Gaibazzi N, Martini C, Mattioli M, et al. Lung disease severity, coronary artery calcium, coronary inflammation and mortality in coronavirus disease 2019. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.05.01.20087114>.
23. Garg S, Kim L, Whitaker M, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 – COVID-NET, 14 states, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:458-64.
24. Rodríguez-Cola M, Jiménez-Velasco I, Gutiérrez-Henares F, et al. Clinical features of coronavirus disease 2019 (COVID-19) in a cohort of patients with disability due to spinal cord injury. *Spinal Cord Ser Cases* 2020;6:39.
25. Giorgi Rossi P, Marino M, Formisano D, et al. Characteristics and outcomes of a cohort of COVID-19 patients in the Province of Reggio Emilia, Italy. *PLoS One* 2020;15:e0238281.
26. Guo AX, Cui JJ, OuYang QY, et al. The clinical characteristics and mortal causes analysis of COVID-19 death patients. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.12.20062380>.
27. Huang H, Cai S, Li Y, et al. Prognostic factors for COVID-19 pneumonia progression to severe symptom based on the earlier clinical features: a retrospective analysis. *Front Med (Lausanne)* 2020;7:557453.
28. Lighter J, Phillips M, Hochman S, et al. Obesity in patients younger than 60 years is a risk factor for COVID-19 hospital admission. *Clin Infect Dis* 2020;71:896-7.
29. Mahévas M, Tran V-T, Roumier M et al. Clinical efficacy of hydroxychloroquine in patients with covid-19 pneumonia who require oxygen: observational comparative study using routine care data. *BMJ* 2020;369:m1844.
30. Menter T, Haslbauer JD, Nienhold R, et al. Post-mortem examination of COVID-19 patients reveals diffuse alveolar damage with severe capillary congestion and variegated findings of lungs and other organs suggesting vascular dysfunction. *Histopathology* 2020;77:198-209.
31. Qi D, Yan X, Tang X, et al. Epidemiological and clinical features of 2019-nCoV acute respiratory disease cases in Chongqing municipality, China: a retrospective, descriptive, multiple-center study. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.03.01.20029397>.
32. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020;323:2052-9.
33. Vuagnat P, Frelaut M, Ramtohl T, et al. COVID-19 in breast cancer patients: a cohort at the Institut Curie hospitals in the Paris area. *Breast Cancer Res* 2020;22:55.
34. Wollenstein-Betech S, Cassandras CG, Paschalidis IC. Personalized predictive models for symptomatic COVID-19 patients using basic preconditions: hospitalizations, mortality, and the need for an ICU or ventilator. *Int J Med Inform* 2020;142:104258.
35. Zuo Y, Yalavarthi S, Shi H, et al. Neutrophil extracellular traps (NETs) as markers of disease severity in COVID-19. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.09.20059626>.
36. Hadjadj J, Yatim N, Barnabei L, et al. Impaired type I interferon activity and exacerbated inflammatory responses in severe COVID-19 patients. *Science* 2020;369:718-24.
37. Lau FH, Majumder R, Torabi R, et al. Vitamin D insufficiency is prevalent in severe COVID-19. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.24.20075838>.
38. Valente-Acosta B, Hoyo-Ulloa I, Espinosa-Aguilar L, et al. COVID-19 severe pneumonia in Mexico City – first experience in a Mexican hospital. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.26.20080796>.
39. Peng YD, Meng K, Guan HQ, et al. Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV [in Chinese]. *Zhonghua Xin Xue Guan Bing Za Zhi* 2020;48:450-5.
40. Wu J, Li W, Shi X, et al. Early antiviral treatment contributes to alleviate the severity and improve the prognosis of patients with novel coronavirus disease (COVID-19). *J Intern Med* 2020;288:128-38.
41. Abou-Arab O, Huette P, Berna P, et al. Tracheal trauma after difficult airway management in morbidly obese patients with COVID-19. *Br J Anaesth* 2020;125:e168-e170.
42. Zheng KI, Gao F, Wang XB, et al. Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism* 2020;108:154244.
43. Chen Q, Zheng Z, Zhang C, et al. Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. *Infection* 2020;48:543-51.
44. Liu M, He P, Liu HG, et al. Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia [in Chinese]. *Zhonghua Jie He He Hu Xi Za Zhi* 2020;43:209-14.
45. Wang Z, Zheutlin A, Kao YH, et al. Hospitalized COVID-19 patients of the Mount Sinai Health System: a retrospective observational study using the electronic medical records. *BMJ Open* 2020;10:e040441.
46. Du X, Liu Y, Chen J, et al. Comparison of the clinical implications among two different nutritional indices in hospitalized patients with COVID-19. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.28.20082644>.
47. Docherty AB, Harrison EM, Green CA, et al. Features of 16,749 hospitalised UK patients with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.04.23.20076042>.
48. Reyes Gil M, Gonzalez-Lugo JD, Rahman S, et al. Correlation of coagulation parameters with clinical outcomes in Coronavirus-19 affected minorities in United States: observational cohort. *medRxiv* 2020. DOI: <https://doi.org/10.1101/2020.05.01.20087932>.
49. Rentsch CT, Kidwai-Khan F, Tate JP, et al. Covid-19 testing, hospital admission, and intensive care among 2,026,227 United

- States veterans aged 54–75 years. medRxiv 2020. DOI: <https://doi.org/10.1101/2020.04.09.20059964>.
50. Ho JSY, Tambyah PA, Sia CH. A call for vaccine against COVID-19: Implications for cardiovascular morbidity and healthcare utilization. *Cardiovasc Drugs Ther* 2020;34:585-7.
 51. Tamara A, Tahapary DL. Obesity as a predictor for a poor prognosis of COVID-19: a systematic review. *Diabetes Metab Syndr* 2020;14:655-9.
 52. Földi M, Farkas N, Kiss S, et al. Obesity is a risk factor for developing critical condition in COVID-19 patients: A systematic review and meta-analysis. *Obes Rev* 2020;21:e13095.
 53. Ruan Q, Yang K, Wang W, et al. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020;46:846-8.
 54. van der Voort PHJ, Moser J, Zandstra DF, et al. Leptin levels in SARS-CoV-2 infection related respiratory failure: A cross-sectional study and a pathophysiological framework on the role of fat tissue. *Heliyon* 2020;6:e04696.
 55. Ho JS, Tambyah PA, Ho AF, et al. Effect of coronavirus infection on the human heart: A scoping review. *Eur J Prev Cardiol* 2020;27:1136-48.
 56. Wang L, Southerland J, Wang K, et al. Ethnic differences in risk factors for obesity among adults in California, the United States. *J Obes* 2017;2017:2427483.
 57. Zhou JC, Hui TC, Tan CH, et al. Chest radiography in coronavirus disease 2019 (COVID-19): Correlation with clinical course. *Ann Acad Med Singap* 2020;49:456-61.
 58. de Lusignan S, Dorward J, Correa A, et al. Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: a cross-sectional study. *Lancet Infect Dis* 2020;20:1034-42.
 59. Burn E, Tebe C, Fernandez-Bertolin S, et al. The natural history of symptomatic COVID-19 in Catalonia, Spain: A multi-state model including 109,367 outpatient diagnoses, 18,019 hospitalisations, and 5,585 COVID-19 deaths among 5,627,520 people. medRxiv 2020. DOI: <https://doi.org/10.1101/2020.07.13.20152454>.
-