

## Observation of a Narrow Meson Decaying to $D_s^+ \pi^0 \gamma$ at a Mass of 2.458 GeV/ $c^2$

B. Aubert,<sup>1</sup> R. Barate,<sup>1</sup> D. Boutigny,<sup>1</sup> J.-M. Gaillard,<sup>1</sup> A. Hicheur,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> P. Robbe,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> A. Palano,<sup>2</sup> A. Pompili,<sup>2</sup> J. C. Chen,<sup>3</sup> N. D. Qi,<sup>3</sup> G. Rong,<sup>3</sup> P. Wang,<sup>3</sup> Y. S. Zhu,<sup>3</sup> G. Eigen,<sup>4</sup> I. Ofte,<sup>4</sup> B. Stugu,<sup>4</sup> G. S. Abrams,<sup>5</sup> A. W. Borgland,<sup>5</sup> A. B. Breon,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> E. Charles,<sup>5</sup> C. T. Day,<sup>5</sup> M. S. Gill,<sup>5</sup> A. V. Gritsan,<sup>5</sup> Y. Groysman,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> R. W. Kadel,<sup>5</sup> J. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> G. Kukartsev,<sup>5</sup> C. LeClerc,<sup>5</sup> M. E. Levi,<sup>5</sup> G. Lynch,<sup>5</sup> L. M. Mir,<sup>5</sup> P. J. Oddone,<sup>5</sup> T. J. Orimoto,<sup>5</sup> M. Pripstein,<sup>5</sup> N. A. Roe,<sup>5</sup> A. Romosan,<sup>5</sup> M. T. Ronan,<sup>5</sup> V. G. Shelkov,<sup>5</sup> A. V. Telnov,<sup>5</sup> W. A. Wenzel,<sup>5</sup> K. Ford,<sup>6</sup> T. J. Harrison,<sup>6</sup> C. M. Hawkes,<sup>6</sup> D. J. Knowles,<sup>6</sup> S. E. Morgan,<sup>6</sup> R. C. Penny,<sup>6</sup> A. T. Watson,<sup>6</sup> N. K. Watson,<sup>6</sup> K. Goetzen,<sup>7</sup> T. Held,<sup>7</sup> H. Koch,<sup>7</sup> B. Lewandowski,<sup>7</sup> M. Pelizaeus,<sup>7</sup> K. Peters,<sup>7</sup> H. Schmucker,<sup>7</sup> M. Steinke,<sup>7</sup> J. T. Boyd,<sup>8</sup> N. Chevalier,<sup>8</sup> W. N. Cottingham,<sup>8</sup> M. P. Kelly,<sup>8</sup> T. E. Latham,<sup>8</sup> C. Mackay,<sup>8</sup> F. F. Wilson,<sup>8</sup> K. Abe,<sup>9</sup> T. Cuhadar-Donszelmann,<sup>9</sup> C. Hearty,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> D. Thiessen,<sup>9</sup> P. Kyberd,<sup>10</sup> A. K. McKemey,<sup>10</sup> L. Teodorescu,<sup>10</sup> V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> V. B. Golubev,<sup>11</sup> V. N. Ivanchenko,<sup>11</sup> E. A. Kravchenko,<sup>11</sup> A. P. Onuchin,<sup>11</sup> S. I. Serednyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> E. P. Solodov,<sup>11</sup> A. N. Yushkov,<sup>11</sup> D. Best,<sup>12</sup> M. Bruinsma,<sup>12</sup> M. Chao,<sup>12</sup> D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> M. Mandelkern,<sup>12</sup> R. K. Mommsen,<sup>12</sup> W. Roethel,<sup>12</sup> D. P. Stoker,<sup>12</sup> C. Buchanan,<sup>13</sup> B. L. Hartfiel,<sup>13</sup> J. W. Gary,<sup>14</sup> J. Layter,<sup>14</sup> B. C. Shen,<sup>14</sup> K. Wang,<sup>14</sup> D. del Re,<sup>15</sup> H. K. Hadavand,<sup>15</sup> E. J. Hill,<sup>15</sup> D. B. MacFarlane,<sup>15</sup> H. P. Paar,<sup>15</sup> Sh. Rahatlou,<sup>15</sup> V. Sharma,<sup>15</sup> J. W. Berryhill,<sup>16</sup> C. Campagnari,<sup>16</sup> B. Dahmes,<sup>16</sup> N. Kuznetsova,<sup>16</sup> S. L. Levy,<sup>16</sup> O. Long,<sup>16</sup> A. Lu,<sup>16</sup> M. A. Mazur,<sup>16</sup> J. D. Richman,<sup>16</sup> W. Verkerke,<sup>16</sup> T. W. Beck,<sup>17</sup> J. Beringer,<sup>17</sup> A. M. Eisner,<sup>17</sup> C. A. Heusch,<sup>17</sup> W. S. Lockman,<sup>17</sup> T. Schalk,<sup>17</sup> R. E. Schmitz,<sup>17</sup> B. A. Schumm,<sup>17</sup> A. Seiden,<sup>17</sup> M. Turri,<sup>17</sup> W. Walkowiak,<sup>17</sup> D. C. Williams,<sup>17</sup> M. G. Wilson,<sup>17</sup> J. Albert,<sup>18</sup> E. Chen,<sup>18</sup> G. P. Dubois-Felsmann,<sup>18</sup> A. Dvoretzki,<sup>18</sup> R. J. Erwin,<sup>18</sup> D. G. Hitlin,<sup>18</sup> I. Narsky,<sup>18</sup> T. Piatenko,<sup>18</sup> F. C. Porter,<sup>18</sup> A. Ryd,<sup>18</sup> A. Samuel,<sup>18</sup> S. Yang,<sup>18</sup> S. Jayatilake,<sup>19</sup> G. Mancinelli,<sup>19</sup> B. T. Meadows,<sup>19</sup> M. D. Sokoloff,<sup>19</sup> T. Abe,<sup>20</sup> F. Blanc,<sup>20</sup> P. Bloom,<sup>20</sup> S. Chen,<sup>20</sup> P. J. Clark,<sup>20</sup> W. T. Ford,<sup>20</sup> U. Nauenberg,<sup>20</sup> A. Olivas,<sup>20</sup> P. Rankin,<sup>20</sup> J. Roy,<sup>20</sup> J. G. Smith,<sup>20</sup> W. C. van Hoek,<sup>20</sup> L. Zhang,<sup>20</sup> J. L. Harton,<sup>21</sup> T. Hu,<sup>21</sup> A. Soffer,<sup>21</sup> W. H. Toki,<sup>21</sup> R. J. Wilson,<sup>21</sup> J. Zhang,<sup>21</sup> D. Altenburg,<sup>22</sup> T. Brandt,<sup>22</sup> J. Brose,<sup>22</sup> T. Colberg,<sup>22</sup> M. Dickopp,<sup>22</sup> R. S. Dubitzky,<sup>22</sup> A. Hauke,<sup>22</sup> H. M. Lacker,<sup>22</sup> E. Maly,<sup>22</sup> R. Müller-Pfefferkorn,<sup>22</sup> R. Nogowski,<sup>22</sup> S. Otto,<sup>22</sup> J. Schubert,<sup>22</sup> K. R. Schubert,<sup>22</sup> R. Schwierz,<sup>22</sup> B. Spaan,<sup>22</sup> L. Wilden,<sup>22</sup> D. Bernard,<sup>23</sup> G. R. Bonneaud,<sup>23</sup> F. Brochard,<sup>23</sup> J. Cohen-Tanugi,<sup>23</sup> P. Grenier,<sup>23</sup> Ch. Thiebaux,<sup>23</sup> G. Vasileiadis,<sup>23</sup> M. Verderi,<sup>23</sup> A. Khan,<sup>24</sup> D. Lavin,<sup>24</sup> F. Muheim,<sup>24</sup> S. Playfer,<sup>24</sup> J. E. Swain,<sup>24</sup> M. Andreotti,<sup>25</sup> V. Azzolini,<sup>25</sup> D. Bettoni,<sup>25</sup> C. Bozzi,<sup>25</sup> R. Calabrese,<sup>25</sup> G. Cibinetto,<sup>25</sup> E. Luppi,<sup>25</sup> M. Negrini,<sup>25</sup> L. Piemontese,<sup>25</sup> A. Sarti,<sup>25</sup> E. Treadwell,<sup>26</sup> F. Anulli,<sup>27</sup> \* R. Baldini-Ferroli,<sup>27</sup> M. Biasini,<sup>27</sup> \* A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> D. Falciari,<sup>27</sup> G. Finocchiaro,<sup>27</sup> P. Patteri,<sup>27</sup> I. M. Peruzzi,<sup>27</sup> \* M. Piccolo,<sup>27</sup> M. Pioppi,<sup>27</sup> \* A. Zallo,<sup>27</sup> A. Buzzo,<sup>28</sup> R. Capra,<sup>28</sup> R. Contri,<sup>28</sup> G. Crosetti,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> S. Bailey,<sup>29</sup> M. Morii,<sup>29</sup> E. Won,<sup>29</sup> W. Bhimji,<sup>30</sup> D. A. Bowerman,<sup>30</sup> P. D. Dauncey,<sup>30</sup> U. Egede,<sup>30</sup> I. Eschrich,<sup>30</sup> J. R. Gaillard,<sup>30</sup> G. W. Morton,<sup>30</sup> J. A. Nash,<sup>30</sup> P. Sanders,<sup>30</sup> G. P. Taylor,<sup>30</sup> G. J. Grenier,<sup>31</sup> S.-J. Lee,<sup>31</sup> U. Mallik,<sup>31</sup> J. Cochran,<sup>32</sup> H. B. Crawley,<sup>32</sup> J. Lamsa,<sup>32</sup> W. T. Meyer,<sup>32</sup> S. Prell,<sup>32</sup> E. I. Rosenberg,<sup>32</sup> J. Yi,<sup>32</sup> M. Davier,<sup>33</sup> G. Grosdidier,<sup>33</sup> A. Höcker,<sup>33</sup> S. Laplace,<sup>33</sup> F. Le Diberder,<sup>33</sup> V. Lepeltier,<sup>33</sup> A. M. Lutz,<sup>33</sup> T. C. Petersen,<sup>33</sup> S. Plaszczynski,<sup>33</sup> M. H. Schune,<sup>33</sup> L. Tantot,<sup>33</sup> G. Wormser,<sup>33</sup> V. Brigljević,<sup>34</sup> C. H. Cheng,<sup>34</sup> D. J. Lange,<sup>34</sup> M. C. Simani,<sup>34</sup> D. M. Wright,<sup>34</sup> A. J. Bevan,<sup>35</sup> J. P. Coleman,<sup>35</sup> J. R. Fry,<sup>35</sup> E. Gabathuler,<sup>35</sup> R. Gamet,<sup>35</sup> M. Kay,<sup>35</sup> R. J. Parry,<sup>35</sup> D. J. Payne,<sup>35</sup> R. J. Sloane,<sup>35</sup> C. Touramanis,<sup>35</sup> J. J. Back,<sup>36</sup> P. F. Harrison,<sup>36</sup> H. W. Shorthouse,<sup>36</sup> P. B. Vidal,<sup>36</sup> C. L. Brown,<sup>37</sup> G. Cowan,<sup>37</sup> R. L. Flack,<sup>37</sup> H. U. Flaecher,<sup>37</sup> S. George,<sup>37</sup> M. G. Green,<sup>37</sup> A. Kurup,<sup>37</sup> C. E. Marker,<sup>37</sup> T. R. McMahon,<sup>37</sup> S. Ricciardi,<sup>37</sup> F. Salvatore,<sup>37</sup> G. Vaitsas,<sup>37</sup> M. A. Winter,<sup>37</sup> D. Brown,<sup>38</sup> C. L. Davis,<sup>38</sup> J. Allison,<sup>39</sup> N. R. Barlow,<sup>39</sup> R. J. Barlow,<sup>39</sup> P. A. Hart,<sup>39</sup> M. C. Hodgkinson,<sup>39</sup> F. Jackson,<sup>39</sup> G. D. Lafferty,<sup>39</sup> A. J. Lyon,<sup>39</sup> J. H. Weatherall,<sup>39</sup> J. C. Williams,<sup>39</sup> A. Farbin,<sup>40</sup> A. Jawahery,<sup>40</sup> D. Kovalskyi,<sup>40</sup> C. K. Lae,<sup>40</sup> V. Lillard,<sup>40</sup> D. A. Roberts,<sup>40</sup> G. Blaylock,<sup>41</sup> C. Dallapiccola,<sup>41</sup> K. T. Flood,<sup>41</sup> S. S. Hertzbach,<sup>41</sup> R. Kofler,<sup>41</sup> V. B. Koptchev,<sup>41</sup> T. B. Moore,<sup>41</sup> S. Saremi,<sup>41</sup> H. Staengle,<sup>41</sup> S. Willocq,<sup>41</sup> R. Cowan,<sup>42</sup> G. Sciolla,<sup>42</sup> F. Taylor,<sup>42</sup> R. K. Yamamoto,<sup>42</sup> D. J. J. Mangeol,<sup>43</sup> P. M. Patel,<sup>43</sup> S. H. Robertson,<sup>43</sup> A. Lazzaro,<sup>44</sup> F. Palombo,<sup>44</sup> J. M. Bauer,<sup>45</sup> L. Cremaldi,<sup>45</sup> V. Eschenburg,<sup>45</sup> R. Godang,<sup>45</sup> R. Kroeger,<sup>45</sup> J. Reidy,<sup>45</sup> D. A. Sanders,<sup>45</sup> D. J. Summers,<sup>45</sup> H. W. Zhao,<sup>45</sup> S. Brunet,<sup>46</sup> D. Cote-Ahern,<sup>46</sup> P. Taras,<sup>46</sup> H. Nicholson,<sup>47</sup> C. Cartaro,<sup>48</sup> N. Cavallo,<sup>48</sup> † G. De Nardo,<sup>48</sup> F. Fabozzi,<sup>48</sup> † C. Gatto,<sup>48</sup> L. Lista,<sup>48</sup> P. Paolucci,<sup>48</sup> D. Piccolo,<sup>48</sup> C. Sciacca,<sup>48</sup> M. A. Baak,<sup>49</sup> G. Raven,<sup>49</sup> J. M. LoSecco,<sup>50</sup> T. A. Gabriel,<sup>51</sup> B. Brau,<sup>52</sup> K. K. Gan,<sup>52</sup> K. Honscheid,<sup>52</sup> D. Hufnagel,<sup>52</sup> H. Kagan,<sup>52</sup> R. Kass,<sup>52</sup> T. Pulliam,<sup>52</sup> Q. K. Wong,<sup>52</sup> J. Brau,<sup>53</sup> R. Frey,<sup>53</sup> C. T. Potter,<sup>53</sup> N. B. Sinev,<sup>53</sup> D. Strom,<sup>53</sup> E. Torrence,<sup>53</sup> F. Colechia,<sup>54</sup> A. Dorigo,<sup>54</sup> F. Galeazzi,<sup>54</sup> M. Margoni,<sup>54</sup> M. Morandin,<sup>54</sup> M. Posocco,<sup>54</sup> M. Rotondo,<sup>54</sup>

F. Simonetto,<sup>54</sup> R. Stroili,<sup>54</sup> G. Tiozzo,<sup>54</sup> C. Voci,<sup>54</sup> M. Benayoun,<sup>55</sup> H. Briand,<sup>55</sup> J. Chauveau,<sup>55</sup> P. David,<sup>55</sup> Ch. de la Vaissière,<sup>55</sup> L. Del Buono,<sup>55</sup> O. Hamon,<sup>55</sup> M. J. J. John,<sup>55</sup> Ph. Leruste,<sup>55</sup> J. Ocariz,<sup>55</sup> M. Pivk,<sup>55</sup> L. Roos,<sup>55</sup> J. Stark,<sup>55</sup> S. T'Jampens,<sup>55</sup> G. Therin,<sup>55</sup> P. F. Manfredi,<sup>56</sup> V. Re,<sup>56</sup> P. K. Behera,<sup>57</sup> L. Gladney,<sup>57</sup> Q. H. Guo,<sup>57</sup> J. Panetta,<sup>57</sup> C. Angelini,<sup>58</sup> G. Batignani,<sup>58</sup> S. Bettarini,<sup>58</sup> M. Bondioli,<sup>58</sup> F. Bucci,<sup>58</sup> G. Calderini,<sup>58</sup> M. Carpinelli,<sup>58</sup> V. Del Gamba,<sup>58</sup> F. Forti,<sup>58</sup> M. A. Giorgi,<sup>58</sup> A. Lusiani,<sup>58</sup> G. Marchiori,<sup>58</sup> F. Martinez-Vidal,<sup>58, †</sup> M. Morganti,<sup>58</sup> N. Neri,<sup>58</sup> E. Paoloni,<sup>58</sup> M. Rama,<sup>58</sup> G. Rizzo,<sup>58</sup> F. Sandrelli,<sup>58</sup> J. Walsh,<sup>58</sup> M. Haire,<sup>59</sup> D. Judd,<sup>59</sup> K. Paick,<sup>59</sup> D. E. Wagoner,<sup>59</sup> N. Danielson,<sup>60</sup> P. Elmer,<sup>60</sup> C. Lu,<sup>60</sup> V. Miftakov,<sup>60</sup> J. Olsen,<sup>60</sup> A. J. S. Smith,<sup>60</sup> H. A. Tanaka,<sup>60</sup> E. W. Varnes,<sup>60</sup> F. Bellini,<sup>61</sup> G. Cavoto,<sup>60,61</sup> R. Faccini,<sup>61</sup> F. Ferrarotto,<sup>61</sup> F. Ferroni,<sup>61</sup> M. Gaspero,<sup>61</sup> M. A. Mazzoni,<sup>61</sup> S. Morganti,<sup>61</sup> M. Pierini,<sup>61</sup> G. Piredda,<sup>61</sup> F. Safai Tehrani,<sup>61</sup> C. Voena,<sup>61</sup> S. Christ,<sup>62</sup> G. Wagner,<sup>62</sup> R. Waldi,<sup>62</sup> T. Adye,<sup>63</sup> N. De Groot,<sup>63</sup> B. Franek,<sup>63</sup> N. I. Geddes,<sup>63</sup> G. P. Gopal,<sup>63</sup> E. O. Olaiya,<sup>63</sup> S. M. Xella,<sup>63</sup> R. Aleksan,<sup>64</sup> S. Emery,<sup>64</sup> A. Gaidot,<sup>64</sup> S. F. Ganzhur,<sup>64</sup> P.-F. Giraud,<sup>64</sup> G. Hamel de Monchenault,<sup>64</sup> W. Kozanecki,<sup>64</sup> M. Langer,<sup>64</sup> M. Legendre,<sup>64</sup> G. W. London,<sup>64</sup> B. Mayer,<sup>64</sup> G. Schott,<sup>64</sup> G. Vasseur,<sup>64</sup> Ch. Yeche,<sup>64</sup> M. Zito,<sup>64</sup> M. V. Purohit,<sup>65</sup> A. W. Weidemann,<sup>65</sup> F. X. Yumiceva,<sup>65</sup> D. Aston,<sup>66</sup> J. Bartelt,<sup>66</sup> R. Bartoldus,<sup>66</sup> N. Berger,<sup>66</sup> A. M. Boyarski,<sup>66</sup> O. L. Buchmueller,<sup>66</sup> M. R. Convery,<sup>66</sup> D. P. Coupal,<sup>66</sup> D. Dong,<sup>66</sup> J. Dorfan,<sup>66</sup> D. Dujmic,<sup>66</sup> W. Dunwoodie,<sup>66</sup> R. C. Field,<sup>66</sup> T. Glanzman,<sup>66</sup> S. J. Gowdy,<sup>66</sup> E. Grauges-Pous,<sup>66</sup> T. Hadig,<sup>66</sup> V. Halyo,<sup>66</sup> T. Hryn'ova,<sup>66</sup> W. R. Innes,<sup>66</sup> C. P. Jessop,<sup>66</sup> M. H. Kelsey,<sup>66</sup> P. Kim,<sup>66</sup> M. L. Kocian,<sup>66</sup> U. Langenegger,<sup>66</sup> D. W. G. S. Leith,<sup>66</sup> J. Libby,<sup>66</sup> S. Luitz,<sup>66</sup> V. Luth,<sup>66</sup> H. L. Lynch,<sup>66</sup> H. Marsiske,<sup>66</sup> R. Messner,<sup>66</sup> D. R. Muller,<sup>66</sup> C. P. O'Grady,<sup>66</sup> V. E. Ozcan,<sup>66</sup> A. Perazzo,<sup>66</sup> M. Perl,<sup>66</sup> S. Petrak,<sup>66</sup> B. N. Ratcliff,<sup>66</sup> A. Roodman,<sup>66</sup> A. A. Salnikov,<sup>66</sup> R. H. Schindler,<sup>66</sup> J. Schwiening,<sup>66</sup> G. Simi,<sup>66</sup> A. Snyder,<sup>66</sup> A. Soha,<sup>66</sup> J. Stelzer,<sup>66</sup> D. Su,<sup>66</sup> M. K. Sullivan,<sup>66</sup> J. Va'vra,<sup>66</sup> S. R. Wagner,<sup>66</sup> M. Weaver,<sup>66</sup> A. J. R. Weinstein,<sup>66</sup> W. J. Wisniewski,<sup>66</sup> D. H. Wright,<sup>66</sup> C. C. Young,<sup>66</sup> P. R. Burchat,<sup>67</sup> A. J. Edwards,<sup>67</sup> T. I. Meyer,<sup>67</sup> B. A. Petersen,<sup>67</sup> C. Roat,<sup>67</sup> M. Ahmed,<sup>68</sup> S. Ahmed,<sup>68</sup> M. S. Alam,<sup>68</sup> J. A. Ernst,<sup>68</sup> M. A. Saeed,<sup>68</sup> M. Saleem,<sup>68</sup> F. R. Wappler,<sup>68</sup> W. Bugg,<sup>69</sup> M. Krishnamurthy,<sup>69</sup> S. M. Spanier,<sup>69</sup> R. Eckmann,<sup>70</sup> H. Kim,<sup>70</sup> J. L. Ritchie,<sup>70</sup> R. F. Schwitters,<sup>70</sup> J. M. Izen,<sup>71</sup> I. Kitayama,<sup>71</sup> X. C. Lou,<sup>71</sup> S. Ye,<sup>71</sup> F. Bianchi,<sup>72</sup> M. Bona,<sup>72</sup> F. Gallo,<sup>72</sup> D. Gamba,<sup>72</sup> C. Borean,<sup>73</sup> L. Bosisio,<sup>73</sup> G. Della Ricca,<sup>73</sup> S. Dittongo,<sup>73</sup> S. Grancagnolo,<sup>73</sup> L. Lanceri,<sup>73</sup> P. Poropat,<sup>73, §</sup> L. Vitale,<sup>73</sup> G. Vuagnin,<sup>73</sup> R. S. Panvini,<sup>74</sup> Sw. Banerjee,<sup>75</sup> C. M. Brown,<sup>75</sup> D. Fortin,<sup>75</sup> P. D. Jackson,<sup>75</sup> R. Kowalewski,<sup>75</sup> J. M. Roney,<sup>75</sup> H. R. Band,<sup>76</sup> S. Dasu,<sup>76</sup> M. Datta,<sup>76</sup> A. M. Eichenbaum,<sup>76</sup> J. R. Johnson,<sup>76</sup> P. E. Kutter,<sup>76</sup> H. Li,<sup>76</sup> R. Liu,<sup>76</sup> F. Di Lodovico,<sup>76</sup> A. Mihalyi,<sup>76</sup> A. K. Mohapatra,<sup>76</sup> Y. Pan,<sup>76</sup> R. Prepost,<sup>76</sup> S. J. Sekula,<sup>76</sup> J. H. von Wimmersperg-Toeller,<sup>76</sup> J. Wu,<sup>76</sup> S. L. Wu,<sup>76</sup> Z. Yu,<sup>76</sup> and H. Neal<sup>77</sup>

(The BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

<sup>2</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>3</sup>Institute of High Energy Physics, Beijing 100039, China

<sup>4</sup>University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, CA 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, BC, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, CA 92697, USA

<sup>13</sup>University of California at Los Angeles, Los Angeles, CA 90024, USA

<sup>14</sup>University of California at Riverside, Riverside, CA 92521, USA

<sup>15</sup>University of California at San Diego, La Jolla, CA 92093, USA

<sup>16</sup>University of California at Santa Barbara, Santa Barbara, CA 93106, USA

<sup>17</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, CA 95064, USA

<sup>18</sup>California Institute of Technology, Pasadena, CA 91125, USA

<sup>19</sup>University of Cincinnati, Cincinnati, OH 45221, USA

<sup>20</sup>University of Colorado, Boulder, CO 80309, USA

<sup>21</sup>Colorado State University, Fort Collins, CO 80523, USA

<sup>22</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>23</sup>Ecole Polytechnique, LLR, F-91128 Palaiseau, France

<sup>24</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

<sup>25</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

<sup>26</sup>Florida A&M University, Tallahassee, FL 32307, USA

<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

- <sup>29</sup>Harvard University, Cambridge, MA 02138, USA  
<sup>30</sup>Imperial College London, London, SW7 2BW, United Kingdom  
<sup>31</sup>University of Iowa, Iowa City, IA 52242, USA  
<sup>32</sup>Iowa State University, Ames, IA 50011-3160, USA  
<sup>33</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France  
<sup>34</sup>Lawrence Livermore National Laboratory, Livermore, CA 94550, USA  
<sup>35</sup>University of Liverpool, Liverpool L69 3BX, United Kingdom  
<sup>36</sup>Queen Mary, University of London, E1 4NS, United Kingdom  
<sup>37</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>38</sup>University of Louisville, Louisville, KY 40292, USA  
<sup>39</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>40</sup>University of Maryland, College Park, MD 20742, USA  
<sup>41</sup>University of Massachusetts, Amherst, MA 01003, USA  
<sup>42</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, MA 02139, USA  
<sup>43</sup>McGill University, Montréal, QC, Canada H3A 2T8  
<sup>44</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>45</sup>University of Mississippi, University, MS 38677, USA  
<sup>46</sup>Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7  
<sup>47</sup>Mount Holyoke College, South Hadley, MA 01075, USA  
<sup>48</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy  
<sup>49</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands  
<sup>50</sup>University of Notre Dame, Notre Dame, IN 46556, USA  
<sup>51</sup>Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA  
<sup>52</sup>Ohio State University, Columbus, OH 43210, USA  
<sup>53</sup>University of Oregon, Eugene, OR 97403, USA  
<sup>54</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>55</sup>Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France  
<sup>56</sup>Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy  
<sup>57</sup>University of Pennsylvania, Philadelphia, PA 19104, USA  
<sup>58</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy  
<sup>59</sup>Prairie View A&M University, Prairie View, TX 77446, USA  
<sup>60</sup>Princeton University, Princeton, NJ 08544, USA  
<sup>61</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>62</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>63</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom  
<sup>64</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>65</sup>University of South Carolina, Columbia, SC 29208, USA  
<sup>66</sup>Stanford Linear Accelerator Center, Stanford, CA 94309, USA  
<sup>67</sup>Stanford University, Stanford, CA 94305-4060, USA  
<sup>68</sup>State Univ. of New York, Albany, NY 12222, USA  
<sup>69</sup>University of Tennessee, Knoxville, TN 37996, USA  
<sup>70</sup>University of Texas at Austin, Austin, TX 78712, USA  
<sup>71</sup>University of Texas at Dallas, Richardson, TX 75083, USA  
<sup>72</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>73</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>74</sup>Vanderbilt University, Nashville, TN 37235, USA  
<sup>75</sup>University of Victoria, Victoria, BC, Canada V8W 3P6  
<sup>76</sup>University of Wisconsin, Madison, WI 53706, USA  
<sup>77</sup>Yale University, New Haven, CT 06511, USA

(Dated: October 29, 2018)

A narrow state, which we label  $D_{sJ}(2458)^+$ , with mass  $2458.0 \pm 1.0$  (stat.)  $\pm 1.0$  (syst.)  $\text{MeV}/c^2$ , is observed in the inclusive  $D_s^+ \pi^0 \gamma$  mass distribution in  $91 \text{ fb}^{-1}$  of  $e^+e^-$  annihilation data recorded by the BABAR detector at the PEP-II asymmetric-energy  $e^+e^-$  storage ring. The observed width is consistent with the experimental resolution. The data favor decay through  $D_s^*(2112)^+ \pi^0$  rather than through  $D_{sJ}^*(2317)^+ \gamma$ . An analysis of  $D_s^+ \pi^0$  data accounting for the influence of the  $D_{sJ}(2458)^+$  produces a  $D_{sJ}^*(2317)^+$  mass of  $2317.3 \pm 0.4$  (stat.)  $\pm 0.8$  (syst.)  $\text{MeV}/c^2$ .

PACS numbers: 14.40.Lb, 13.25.Ft, 12.40.Yx

---

\*Also with Università di Perugia, Perugia, Italy

---

†Also with Università della Basilicata, Potenza, Italy

Interest in the spectrum of charmed mesons has been heightened by the discovery by this collaboration [1] of a narrow state, produced in  $e^+e^- \rightarrow c\bar{c}$  collisions at the PEP-II collider, decaying to  $D_s^+\pi^0$  [2], with mass  $2317 \text{ MeV}/c^2$ , approximately  $41 \text{ MeV}/c^2$  below the  $DK$  mass threshold. This state, the  $D_{s,J}^*(2317)^+$ , has been confirmed by CLEO [3] and Belle [4, 5]. Along with the  $D_{s,J}^*(2317)^+$ , we noted [1] the presence of a narrow peak in the  $D_s^+\pi^0\gamma$  mass distribution near  $2.46 \text{ GeV}/c^2$ . Because this signal is near the kinematic overlap of the  $D_{s,J}^*(2317)^+\gamma$  and  $D_s^*(2112)^+\pi^0$  systems, special attention is required to remove associated backgrounds and to distinguish between the two possible decay modes. Such an analysis is the subject of this paper.

This state near  $2.46 \text{ GeV}/c^2$  has been seen by CLEO [3] and Belle [4] in the inclusive  $D_s^+\pi^0\gamma$  mass spectrum and by Belle [5] in exclusively reconstructed  $B$  decays.

To investigate the  $D_s^+\pi^0\gamma$  spectrum, we study  $D_s^+$  candidates from  $e^+e^- \rightarrow c\bar{c}$  (at a center-of-mass energy near  $10.6 \text{ GeV}$ ) that decay to  $K^-K^+\pi^+$ . Particle identification is used to provide clean samples of charged  $K$  and  $\pi$  candidates, which are combined using a geometric fit to a common vertex. Backgrounds are suppressed by selecting decays to  $\bar{K}^{*0}K^+$  and  $\phi\pi^+$ . A description of this sample and additional details can be found elsewhere [1]. Events with  $1.954 < m(K^-K^+\pi^+) < 1.981 \text{ GeV}/c^2$  are taken as  $D_s^+$  candidates.

A candidate  $\pi^0$  is formed by constraining a pair of photons each with energy greater than  $100 \text{ MeV}$  to emanate from the intersection of the  $D_s^+$  trajectory with the beam envelope, performing a one-constraint fit to the  $\pi^0$  mass, and requiring a fit probability greater than 5%. A given event may yield several acceptable  $\pi^0$  candidates. We retain only those candidates for which neither photon belongs to another otherwise acceptable  $\pi^0$ .

Each  $D_s^+$  candidate is combined with all combinations of accompanying  $\pi^0$  candidates with momentum greater than  $300 \text{ MeV}/c$  and photon candidates of energy greater than  $100 \text{ MeV}$ . To suppress background, photons that belong to any  $\pi^0$  candidate are excluded and we require the momentum,  $p^*$ , of each  $D_s^+\pi^0\gamma$  combination in the  $e^+e^-$  center-of-mass frame to be greater than  $3.5 \text{ GeV}/c$ . The last requirement also removes any  $D_s^+\pi^0\gamma$  combination from  $B$  decay.

The  $D_s^+\pi^0\gamma$  invariant mass distribution is shown in Fig. 1a. A clear enhancement is observed near  $2.46 \text{ GeV}/c^2$ . The background underneath this peak is from several sources, which can be described in terms of




FIG. 1: (a) The mass distribution for all selected  $D_s^+\pi^0\gamma$  combinations. The shaded region is from  $D_s^+$  sidebands defined by  $1.912 < m(K^-K^+\pi^+) < 1.933 \text{ GeV}/c^2$  and  $1.999 < m(K^-K^+\pi^+) < 2.020 \text{ GeV}/c^2$ . (b) The value of  $\Delta m_\gamma$  versus  $\Delta m_{\pi^0}$  for all combinations. The horizontal lines delineate three ranges in  $\Delta m_\gamma$ . (c) The  $\Delta m_{\pi^0}$  mass distribution for the middle range of  $\Delta m_\gamma$  (points) and for the average of the upper and lower ranges (shaded histogram). (d) The difference between the two distributions shown in (c). The curve is the fit described in the text.

mass differences defined as

$$\Delta m_\gamma \equiv m(D_s^+\gamma) - m(D_s^+) \quad (1)$$

$$\Delta m_{\pi^0} \equiv m(D_s^+\gamma\pi^0) - m(D_s^+\gamma) \quad (2)$$

A scatter plot of the data is shown in Fig. 1b. Particular background patterns are visible:  $D_s^*(2112)^+ \rightarrow D_s^+\gamma$  decay combined with an unassociated  $\pi^0$ , which appears as a horizontal band, and  $D_{s,J}^*(2317)^+ \rightarrow D_s^+\pi^0$  decay combined with an unassociated  $\gamma$ , which appears as a band that is almost vertical.

To demonstrate the existence of a signal above these backgrounds, the upper histogram of Fig. 1c shows  $D_s^+\pi^0\gamma$  combinations in the  $D_s^*(2112)^+$  signal region, and the gray histogram, scaled to the area of the signal region, corresponds to the two  $D_s^*(2112)^+$  sidebands. We conclude that a signal for a state decaying to  $D_s^+\pi^0\gamma$  exists over a background resulting from  $D_{s,J}^*(2317)^+$  and an unassociated  $\gamma$ . This background peaks at a mass slightly higher than that of the signal. A Gaussian fit to the subtracted mass distribution (Fig. 1d) indicates a narrow signal at  $\Delta m_{\pi^0} = 346.2 \pm 0.9 \text{ MeV}/c^2$  (statistical error only).

The state corresponding to this signal, which we label  $D_{s,J}^*(2458)^+$ , may decay to  $D_s^+\pi^0\gamma$  through  $D_s^*(2112)^+\pi^0$

<sup>‡</sup>Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

<sup>§</sup>Deceased

or  $D_{sJ}^*(2317)^+\gamma$ . To disentangle these modes and reliably extract the parameters of the signal, we apply an unbinned maximum likelihood fit simultaneously to the  $D_s^+\pi^0\gamma$ ,  $D_s^+\pi^0$ , and  $D_s^+\gamma$  invariant masses of all  $D_s^+\pi^0\gamma$  combinations using the channel likelihood method [6]. This fit describes the probability density function of the two  $D_{sJ}(2458)^+$  decay channels as the product of a Gaussian shape in the  $D_s^+\pi^0\gamma$  mass distribution and a Gaussian shape projected into the  $D_s^+\pi^0$  or  $D_s^+\gamma$  mass axes, as appropriate. Because the daughter resonances are narrow, interference between the two  $D_{sJ}(2458)^+$  decay modes cannot be resolved, and so is ignored.

Sources of background in the  $D_s^+\pi^0\gamma$  spectrum included in the fit are purely combinatorial background ( $D_s^+$  meson combined with an unassociated  $\pi^0$  and  $\gamma$ ),  $D_s^*(2112)^+ \rightarrow D_s^+\gamma$  decay combined with an unassociated  $\pi^0$ , and  $D_{sJ}^*(2317)^+ \rightarrow D_s^+\pi^0$  decay combined with an unassociated  $\gamma$ . The fit also includes a contribution from  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  decay but with an unassociated  $\gamma$  replacing the  $\gamma$  from  $D_s^*(2112)^+$  decay. The fit determines the relative size of the background and signal contributions, the mass and width of the  $D_{sJ}(2458)^+$ , and the  $D_{sJ}^*(2317)^+$  mass.

The likelihood fit is validated using Monte Carlo (MC) simulation. This simulation includes  $e^+e^- \rightarrow c\bar{c}$  events and all known charm states and decays, including the  $D_{sJ}^*(2317)^+$  and the signal under study. The generated events were processed by a detailed detector simulation [7] and subjected to the same reconstruction and event-selection procedure as the data.

As shown in Fig. 2a, the fit provides a good description of the  $D_s^+\pi^0\gamma$  mass distribution observed in the data. The  $D_{sJ}(2458)^+$  signal for a particular decay mode can be isolated by calculating a weight for each  $D_s^+\pi^0\gamma$  combination proportional to the relative likelihood contributed by the decay mode of interest. Distributions of events so weighted can be compared to the likelihood function to validate the fit. This is shown in Figs. 2b and 2c. A  $\chi^2$  probability calculation gives 22%, 74% and 11% for fig. 2a, b and c respectively. The resulting yield of correctly reconstructed  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  ( $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+\gamma$ ) decays is  $195 \pm 26$  ( $0 \pm 23$ ), consistent with the fit shown in Fig. 1d. Excluding the  $D_{sJ}(2458)^+$  from the likelihood fit decreases the log likelihood by approximately 57, corresponding to a significance of more than 10 standard deviations. The fit yields a  $D_{sJ}(2458)^+$  mass of  $2458.0 \pm 1.0$  MeV/ $c^2$  with an rms width of  $8.5 \pm 1.0$  MeV/ $c^2$ .

The likelihood fit uses the shapes of the  $D_s^+\pi^0$  and  $D_s^+\gamma$  mass distributions to distinguish between the two possible decay modes,  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  and  $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+\gamma$ . These shapes are influenced by the kinematic constraints of  $D_{sJ}(2458)^+$  decay shown in Fig. 3a. Figs. 3b–3c show the sideband-subtracted  $D_s^+\pi^0$  and  $D_s^+\gamma$  mass projections compared with MC simulations of the two hypotheses (scaled to match the data yield). The  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  decay mode (solid histograms) produces a narrow  $D_s^+\gamma$




FIG. 2: Maximum likelihood fit results overlaid on the  $D_s^+\pi^0\gamma$  mass distribution with (a) no weights and after applying weights corresponding to (b) the decay  $D_s^*(2112)^+\pi^0$  and (c) the decay  $D_{sJ}^*(2317)^+\gamma$ . (d) The mass spectrum of  $D_s^+\pi^0$  combinations (with no  $\gamma$  requirement). The solid curve is the fit described in the text. The dashed and lower solid curves are the contributions from  $D_{sJ}(2458)^+$  decays and combinatorial background, respectively.

mass distribution and a wide  $D_s^+\pi^0$  mass distribution. In contrast, the  $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+\gamma$  decay mode (dashed histograms) produces a wide  $D_s^+\gamma$  mass distribution and a narrow  $D_s^+\pi^0$  mass distribution. Figures 3b and 3c show that the  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  hypothesis is in better agreement with the data.

Our previous measurement [1] of the  $D_{sJ}^*(2317)^+$  mass using the decay  $D_{sJ}^*(2317)^+ \rightarrow D_s^+\pi^0$  did not explicitly consider background from  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  decay. This background peaks in the  $D_s^+\pi^0$  mass spectrum just below the  $D_{sJ}^*(2317)^+$  mass. Shown in Fig. 2d is the  $D_s^+\pi^0$  invariant mass distribution for a sample of  $D_s^+$  candidates combined with all  $\pi^0$  candidates, with  $p^* > 3.5$  GeV/ $c$ . Superimposed on this distribution is a binned fit that includes the contribution from the  $D_{sJ}(2458)^+$  as estimated from MC simulation and a quadratic background function. The result is a  $D_{sJ}^*(2317)^+$  yield of  $1022 \pm 50$  events, a mass of  $2317.3 \pm 0.4$  MeV/ $c^2$ , and measured rms width  $7.3 \pm 0.2$  MeV/ $c^2$ . These results are an improvement over our earlier measurement [1].

We divide the sources of systematic uncertainty in the  $D_{sJ}(2458)^+$  and  $D_{sJ}^*(2317)^+$  mass values and production rates into three categories. The first category is associated with the fit procedure. Likelihood fits to MC samples that include samples of  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+\pi^0$  and  $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+\gamma$  decays correctly reproduce the given sample sizes within statistical errors. The average values of the fit results obtained using statisti-




FIG. 3: (a) The  $D_s^+ \gamma$  versus  $D_s^+ \pi^0$  mass distribution for all  $D_s^+ \pi^0 \gamma$  combinations. The decay of a zero-width  $D_{sJ}(2458)^+$  is kinematically restricted to the region between the two curves. (b) Sideband-subtracted  $D_s^+ \gamma$  mass distribution with MC simulation for (solid histogram)  $D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+ \pi^0$  and (dashed histogram)  $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+ \gamma$ . (c) A similar plot for the  $D_s^+ \pi^0$  mass distribution.

cally distinct MC samples corresponding to the measurements in the data are used to place limits on any fit bias.

We obtain the background distribution used in the likelihood from a random selection of  $D_s^+$ ,  $\pi^0$ , and  $\gamma$  candidates taken from the MC  $D_s^+ \pi^0 \gamma$  sample. To test our sensitivity to this distribution, various selection requirements are altered within reasonable bounds to provide alternate background samples for use in the fit. The resulting changes in yield and mass are used as the second category of systematic uncertainty.

Reconstruction of the decay sequences is the third source of systematic uncertainty. To evaluate the reliability of the MC determination of  $\pi^0$  efficiency and momentum calibration, we use control samples of  $K_S \rightarrow \pi^0 \pi^0$  and  $\tau \rightarrow \pi^0 X$ . On this basis, we assign a systematic uncertainty of  $\pm 5\%$  in  $\pi^0$  reconstruction efficiency and a relative  $\pm 1\%$  in  $\pi^0$  momentum bias. Similar studies for  $\gamma$  reconstruction reveal a systematic uncertainty of  $\pm 3\%$  in  $\gamma$  reconstruction efficiency and  $\pm 1\%$  in energy bias. Uncertainties in the  $D_s^+$  and  $D_s^*(2112)^+$  masses, taken from world averages [8], also contribute to the systematic uncertainty.

The resulting total systematic uncertainty in the  $D_{sJ}(2458)^+ [D_{sJ}^*(2317)^+]$  mass is  $\pm 1.0 [\pm 0.8] \text{ MeV}/c^2$ .

Using the yields from our fit and correcting for effi-

ciency, we estimate the relative production rate

$$R = \frac{\sigma(D_{sJ}(2458)^+) \mathcal{B}(D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+ \pi^0)}{\sigma(D_{sJ}^*(2317)^+) \mathcal{B}(D_{sJ}^*(2317)^+ \rightarrow D_s^+ \pi^0)} \quad (3)$$

to be  $0.25 \pm 0.03$  (stat.)  $\pm 0.03$  (syst.), requiring  $p^* > 3.5 \text{ GeV}/c$  for both states. We also estimate, at 95% C.L.,

$$\frac{\mathcal{B}(D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+ \gamma)}{\mathcal{B}(D_{sJ}(2458)^+ \rightarrow D_s^*(2112)^+ \pi^0)} < 0.22. \quad (4)$$

The observed rms width of the  $D_{sJ}(2458)^+$  is consistent with detector resolution, as determined by Monte Carlo studies. We conclude that the intrinsic width of the  $D_{sJ}(2458)^+$  is small ( $\Gamma \lesssim 10 \text{ MeV}/c^2$ ).

The mass of the  $D_{sJ}(2458)^+$  lies above  $DK$  and below  $D^*K$  thresholds. The narrow width and the isospin-violating decay to  $D_s^*(2112)^+ \pi^0$  indicate that decay to  $DK$  is forbidden and suggest an unnatural spin-parity assignment for the state. Belle has observed the decay  $D_{sJ}(2458)^+ \rightarrow D_s^+ \gamma$  in production from both  $c\bar{c}$  continuum [4] and  $B$  decay [5]. Such a decay rules out  $J = 0$  and favors a  $1^+$  interpretation. Decay distributions studied by Belle further support  $J = 1$  for  $D_{sJ}(2458)^+$  and also  $J^P = 0^+$  for  $D_{sJ}^*(2317)^+$ . The apparent absence of the decay  $D_{sJ}(2458)^+ \rightarrow D_{sJ}^*(2317)^+ \gamma$  may indicate that the electromagnetic decay mechanism cannot compete with  $D_s^*(2112)^+ \pi^0$ , which may be a strong, but isospin-violating, process resulting from  $\eta$ - $\pi^0$  mixing, as discussed by Cho and Wise [9].

Our measurement of the  $D_{sJ}(2458)^+$  mass ( $2458.0 \pm 1.4 \text{ MeV}/c^2$ , with combined statistical and systematic uncertainties) agrees with that obtained by Belle ( $2456.5 \pm 1.7 \text{ MeV}/c^2$ ) [4], but is two standard deviations smaller than that obtained by CLEO ( $2463.1 \pm 2.1 \text{ MeV}/c^2$ ) [3]. We obtain a relative yield ( $R = 0.25 \pm 0.04$ ) which agrees with that of Belle ( $0.26 \pm 0.08$ ). Both values are somewhat smaller than that reported by CLEO ( $0.44 \pm 0.13$ ). Our reanalysis of the  $D_{sJ}^*(2317)^+ \rightarrow D_s^+ \pi^0$  sample to account for background from the  $D_{sJ}(2458)^+$  gives a mass of  $2317.3 \pm 0.4$  (stat.)  $\pm 0.8$  (syst.)  $\text{MeV}/c^2$ , which remains consistent with results from CLEO [3] and Belle [4].

In summary, in  $91 \text{ fb}^{-1}$  of data collected from the BABAR experiment, we have observed a narrow state that decays to  $D_s^+ \pi^0 \gamma$  with a mass of  $2458.0 \pm 1.0$  (stat.)  $\pm 1.0$  (syst.)  $\text{MeV}/c^2$ . The only significant  $D_s^+ \pi^0 \gamma$  decay mode we observe is through  $D_s^*(2112)^+ \pi^0$ . We measure a mass and yield relative to the  $D_{sJ}^*(2317)^+$  similar to those measured by Belle though smaller than those reported by CLEO. The observed width is compatible with our mass resolution. After including the influence of this state, our new measurement of the  $D_{sJ}^*(2317)^+$  mass is  $2317.3 \pm 0.4$  (stat.)  $\pm 0.8$  (syst.)  $\text{MeV}/c^2$ .

### Acknowledgments

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy

Physics (China), the Commissariat à l’Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, and the Particle Physics and Astronomy Research Council (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

- 
- [1] B. Aubert et al. (*BABAR*), *Phys. Rev. Lett.* **90**, 242001 (2003).
  - [2] Inclusion of charge conjugate states is implied throughout this Letter.
  - [3] D. Besson et al. (CLEO), *Phys. Rev.* **D68**, 032002 (2003).
  - [4] Y. Mikami et al. (BELLE) (2003), hep-ex/0307052.
  - [5] P. Krokovny et al. (BELLE) (2003), *submitted to Phys. Rev. Lett.*, hep-ex/0308019.
  - [6] P. E. Condon and P. L. Cowell, *Phys. Rev.* **D9**, 2558 (1974).
  - [7] S. Agostinelli et al. (GEANT4), *Nucl. Instrum. Meth.* **A506**, 250 (2003).
  - [8] K. Hagiwara et al. (Particle Data Group), *Phys. Rev.* **D66**, 010001 (2002).
  - [9] P. L. Cho and M. B. Wise, *Phys. Rev.* **D49**, 6228 (1994).