

UC Irvine

UC Irvine Previously Published Works

Title

Observation of charmed-baryon production in e^+e^- annihilation

Permalink

<https://escholarship.org/uc/item/38k52110>

Journal

Physical Review Letters, 44(1)

ISSN

0031-9007

Authors

Abrams, GS
Alam, MS
Blocker, CA
et al.

Publication Date

1980

DOI

10.1103/PhysRevLett.44.10

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Observation of Charmed-Baryon Production in e^+e^- Annihilation

G. S. Abrams, M. S. Alam, C. A. Blocker, A. M. Boyarski, M. Breidenbach, D. L. Burke, W. C. Carithers, W. Chinowsky, M. W. Coles, S. Cooper, W. E. Dieterle, J. B. Dillon, J. Dorenbosch, J. M. Dorfan, M. W. Eaton, G. J. Feldman, M. E. B. Franklin, G. Gidal, G. Goldhaber, G. Hanson, K. G. Hayes, T. Himel, D. G. Hitlin,^(a) R. J. Hollebeek, W. R. Innes, J. A. Jaros, P. Jenni, A. D. Johnson, J. A. Kadyk, A. J. Lankford, R. R. Larsen, M. J. Longo,^(b) V. Lüth, R. E. Millikan, M. E. Nelson, C. Y. Pang, J. F. Patrick, M. L. Perl, B. Richter, A. Roussarie, J. J. Russell,^(c) D. L. Scharre, R. H. Schindler, R. F. Schwitters,^(d) J. L. Siegrist, J. Strait, H. Taureg, M. Tonutti,^(e) G. H. Trilling, E. N. Vella, R. A. Vidal, I. Videau, J. M. Weiss, and H. Zaccone^(f)
*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305, and
 Lawrence Berkeley Laboratory and Department of Physics,
 University of California, Berkeley, California 94720*

(Received 22 October 1979)

A peak in the $pK^-\pi^+$ and $\bar{p}K^+\pi^-$ invariant-mass spectra at 2.285 ± 0.006 GeV/ c^2 is observed, which is associated with the lowest-lying charmed baryon (Λ_c). A cross section times branching ratio of 0.037 ± 0.012 nb at $E_{c.m.} = 5.2$ GeV is measured with a substantial fraction of the events produced with an equal recoiling mass. New measurements of inclusive p and Λ cross sections are also presented, allowing an estimate of the branching ratio $B(\Lambda_c \rightarrow pK^-\pi^+) = 0.022 \pm 0.010$.

Our present understanding of the charmonium states and of charmed mesons leads us to expect the existence of weakly decaying charmed baryons.¹ Evidence for the production of such charmed baryons has been reported in neutrino interactions,² photon interactions,³ and p - p interactions.⁴ In an experiment with the Stanford Linear Accelerator Center-Lawrence Berkeley Laboratory Mark II detector at the SPEAR e^+e^- storage ring of the Stanford Linear Accelerator Center, we observe the $pK^-\pi^+$ and $\bar{p}K^+\pi^-$ decay modes of what appears to be the lowest-lying charmed baryon (Λ_c). As in the case of the charmed mesons, the baryon is associated with recoil masses equal to or greater than its observed mass. Together with a new measurement of the inclusive proton and Λ cross sections, reported here, we are able to estimate the absolute branching ratio of the Λ_c state to the $pK\pi$ channel.

The data sample used in this analysis represents an integrated luminosity of 9150 nb $^{-1}$. Of this 4000 nb $^{-1}$ was obtained during a scan of the region 4.5 – 6.0 GeV, and 5150 nb $^{-1}$ was obtained during a run at the center-of-mass energy 5.2 GeV. Additional data at other energies are used for the inclusive cross-section measurement.

Detailed descriptions of the Mark II detector have been given elsewhere.⁵ Particle identification is accomplished by the time-of-flight (TOF) system with a measured resolution for hadron tracks of $\sigma = 0.300$ nsec. This allows a 1σ separa-

tion for K - π at 1.35 -GeV/ c and for K - p at 2.0 -GeV/ c momentum. Each track is assigned weights $W(\pi)$, $W(K)$, and $W(p)$ corresponding to its possible identities. Each weight is proportional to the probability that if the particle has the assumed identity, its flight time would have the measured value, and the normalization is such that the sum of all weights corresponding to a given track is unity. The track is called a proton if its momentum is less than 2.0 GeV/ c and $W(p) > 0.5$. Kaons are identified for momenta less than 1.4 GeV/ c by $W(K) > 0.5$. All other tracks not positively identified as muons or electrons are called pions.

We are searching for charmed-baryon signals in various channels and report here results for the $pK\pi$ channel. To reduce the effects of beam-gas backgrounds, proton events which do not also contain an antiproton are required to have a net observed charge $< +1$. This additional requirement reduces the efficiency for protons relative to antiprotons by 21% as measured for the antiproton data sample.

Figure 1(a) shows the invariant-mass distribution for the $pK^-\pi^+$ combinations and charge conjugate $\bar{p}K^+\pi^-$ with the requirement that the mass recoiling against the $pK\pi$ system be greater than 2.2 GeV/ c^2 . A significant enhancement is observed at $M(pK\pi) = 2.285$ GeV/ c^2 in these channels which have the quantum numbers of the Cabibbo-favored weak decay of the Λ_c . Figure 1(b) shows the complementary mass distribution for




FIG. 1. The combined $pK^-\pi^+$ and $\bar{p}K^+\pi^-$ mass distribution (a) for recoil masses greater than $2.2 \text{ GeV}/c^2$, and (b) for recoil masses less than $2.2 \text{ GeV}/c^2$. (c) The $pK^+\pi^-$ and $pK^-\pi^+$ (and charge conjugate states) mass distribution for recoil masses greater than $2.2 \text{ GeV}/c^2$. (d) The beam-energy-constrained mass distribution for events with $pK^-\pi^+$ or $\bar{p}K^+\pi^-$ energy within 0.03 GeV of the beam energy.

those $pK\pi$ combinations with recoil mass less than $2.2 \text{ GeV}/c^2$. No enhancement is observed in Fig. 1(b), indicating that the observed signal is associated with an equal or larger recoil mass. In Fig. 1(c) we show the invariant-mass distribution for the channels $pK^+\pi^-$ and $pK^-\pi^+$ and their charge conjugates, with the same cuts as in Fig. 1(a). These channels have quantum numbers inconsistent with a Λ_c decay and do not exhibit any structure. The curve in Fig. 1(a) shows that the data are well fitted by a Gaussian error function plus a background shape determined from a fit to Fig. 1(c). The signal consists of 39 ± 8 events above a background of twenty events.

For those $pK\pi$ combinations with total measured energy within 0.03 GeV of the beam energy, Fig. 1(d) shows the beam-energy-constrained mass defined as $M_c = (E_{\text{beam}}^2 - p_{pK\pi}^2)^{1/2}$. The 10 ± 4 events in the peak near $M_c = 2.285 \text{ GeV}/c^2$ imply that the reaction $e^+e^- \rightarrow \Lambda_c \bar{\Lambda}_c$ is the source of $(26 \pm 11)\%$ of the observed $pK\pi$ signal.

The fit to Fig. 1(a) yields a mass of $2.286 \pm 0.007 \text{ GeV}/c^2$ and an rms width of $0.010 \text{ GeV}/c^2$. The quoted error includes a systematic component of $0.006 \text{ GeV}/c^2$ due to uncertainties in the magnetic field and the geometrical reconstruction. These error sources are checked by measurement of the $K^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^-\pi^+$ masses in the data sample. A shift in the mass of the observed $pK\pi$ signal of $0.003 \text{ GeV}/c^2$, for example, would require a change in the magnetic field which would displace these masses from their present agreement with nominal values by 1 standard deviation ($0.001 \text{ GeV}/c^2$ for M_{K^0} and $0.0025 \text{ GeV}/c^2$ for M_{D^0}). The mass determined from the spectrum of Fig. 1(d) is $2.284 \pm 0.008 \text{ GeV}/c^2$. This is consistent with the mass value determined above but is subject to different systematic errors⁶ and to a larger statistical error. From the combination of the two mass determinations our best estimate of the mass of the charmed baryon is then $2.285 \pm 0.006 \text{ GeV}/c^2$. As expected for the weak decay of a charmed baryon, the measured width agrees with the calculated detector resolution.

To determine a cross section, the detection efficiency has been calculated to be 0.13 ± 0.025 for the observed Λ_c momentum distribution. The 26 ± 7 $pK\pi$ signal events observed in the 5450-nb^{-1} integrated luminosity within 0.05 GeV of 5.2-GeV c.m. energy correspond to

$$\sigma(\Lambda_c + \bar{\Lambda}_c)B(\Lambda_c \rightarrow pK^-\pi^+) = 0.037 \pm 0.012 \text{ nb},$$

where $\sigma(\Lambda_c + \bar{\Lambda}_c)$ is defined as the inclusive cross section $[\sigma(\Lambda_c) + \sigma(\bar{\Lambda}_c)]$. The data are consistent with equal cross sections for both charge states.

To obtain an estimate of the total production of charmed baryons we have measured the inclusive cross sections for p and Λ from 3.52 to 7.40 GeV . Because of substantial beam-gas contamination in proton events, only antiprotons are used for these measurements. Two or more observed tracks are required and valid identification of antiprotons is ensured by a stricter TOF weight cut of 0.7 for momenta greater than $1.2 \text{ GeV}/c$. The observed \bar{p} sample includes contributions from weakly decaying hyperons. For Λ and $\bar{\Lambda}$ production, only multihadron events with three or more detected tracks are used. The Λ and $\bar{\Lambda}$ are identified from the invariant-mass distribution of all neutral $p\pi$ pairs identified by TOF. In addition, the decay products of the Λ and $\bar{\Lambda}$ are required to originate from a secondary vertex. To reduce beam-gas contamination to less than 4% , Λ (but not $\bar{\Lambda}$) events are required to have total observed

charge $< +1$. With these cuts, our Λ and $\bar{\Lambda}$ background subtractions are both $\lesssim 15\%$ at all energies.

The efficiency for detection of antiprotons is calculated from a Monte Carlo model which generates \bar{p} tracks with a momentum distribution corresponding to an invariant cross section⁷

$$E d^3\sigma/dp^3 \propto e^{-bE},$$

and which then chooses the other nucleon and a number of pions according to the remaining phase space. After adjustment of the slope parameter b and the mean particle multiplicity at each energy, this form gives a good description of the data. The overall detection efficiency for antiprotons is approximately 58% over the entire range 3.7–7.4 GeV.⁸ The efficiency for Λ and $\bar{\Lambda}$ detection is determined by the same Monte Carlo model with the parameters obtained from the antiprotons. The efficiency ranges from 10% at 3.67 GeV to 13% at 7.4 GeV, including the branching ratio for $\Lambda \rightarrow p\pi^-$. In this case also, the Monte Carlo calculation reproduces the observed Λ momentum and multiplicity distributions. As an additional check on our Λ efficiency calculations, we have verified that the ratio of single Λ 's and $\Lambda\bar{\Lambda}$ pairs detected in well-identified $\psi \rightarrow \Lambda\bar{\Lambda}$ events at $E_{c.m.} = 3.095$ GeV is correctly reproduced by the Monte Carlo simulation program.

Our results for inclusive production of p and Λ are presented in Fig. 2 as the ratio of the inclusive production cross section to the μ -pair cross section. Figure 2(a) shows $R(p+\bar{p}) \equiv 2\sigma(\bar{p})/\sigma_{\mu\mu}$ and Fig. 2(b) shows $R(\Lambda+\bar{\Lambda}) \equiv [\sigma(\Lambda) + \sigma(\bar{\Lambda})]/\sigma_{\mu\mu}$. The estimated overall systematic errors in R are $\pm 17\%$ and $\pm 27\%$ for the p and Λ , respectively, and are not shown in Fig. 2. These systematic errors are dominated by the model dependence of the Monte Carlo calculations, and are expected to vary slowly over our energy region.

We note that the rise previously reported⁹ in $R(p+\bar{p})$ and $R(\Lambda+\bar{\Lambda})$ is confirmed here with more precision. We observe for the first time clear steps in both $R(p+\bar{p})$ and $R(\Lambda+\bar{\Lambda})$ in the range of 4.5 to 5.2 GeV c.m. energy, although the R values probably continue to rise more slowly at higher energies. Within the quoted errors, our measurements of $R(p+\bar{p})$ are consistent with previous experiments.^{7,9} Our values of $R(\Lambda+\bar{\Lambda})$ are considerably higher than the previous measurements⁹ but have been obtained with larger solid angle, improved vertex reconstruction, and a more sophisticated efficiency calculation.¹⁰

The coincidence of the location of this step with the threshold for production of an object near the




FIG. 2. (a) $R(p+\bar{p})$ as a function of $E_{c.m.}$. (b) $R(\Lambda+\bar{\Lambda})$ as a function of $E_{c.m.}$. Errors are statistical only.

mass of the observed $pK\pi$ signal is consistent with its interpretation as the lowest-lying charmed baryon. The observed step sizes of $\Delta R(p+\bar{p}) = 0.31 \pm 0.06$ and $\Delta R(\Lambda+\bar{\Lambda}) = 0.10 \pm 0.03$ would indicate a Λ/p ratio for charmed-baryon decays of $(41 \pm 15)\%$ after explicitly removing protons which arise from Λ decay, but not from other weakly decaying strange baryons.

We can use our measurement of $R(p+\bar{p})$ and the measured σ_B for the $pK\pi$ signal at 5.2 GeV to estimate the absolute branching ratio for the $pK\pi$ decay mode. We make the following assumptions: (i) The observed step in $R(p+\bar{p})$ is due entirely to the onset of charmed-baryon pair production¹¹; (ii) all charmed baryons cascade down to the Λ_c state¹²; and (iii) the probability for a charmed baryon to give a proton (as opposed to a neutron) as a final product is 0.6 ± 0.1 .¹³ Using the relationship

$$\sigma(\Lambda_c + \bar{\Lambda}_c) = \frac{\Delta R(p+\bar{p})}{0.6} \sigma_{\mu\mu},$$

we find an inclusive cross section

$$\sigma(\Lambda_c + \bar{\Lambda}_c) = 1.7 \pm 0.4 \text{ nb}$$

at 5.2 GeV. Thus the cross section for producing pairs of charmed baryons is 0.85 nb. With these assumptions, the branching ratio is then estimated to be

$$B(\Lambda_c \rightarrow pK^-\pi^+) = (2.2 \pm 1.0)\%.$$

In conclusion, our observation of a narrow state with the proper quantum numbers, associated with equal and higher recoil masses, and at a mass compatible with the observed threshold in p and Λ production, argues for the interpretation of this state as the charmed baryon Λ_c .

This work was supported primarily by the U. S. Department of Energy under Contracts No. DE-AC03-765F00515, and No. W-7405-ENG-48. Support for individuals came from the listed institutions plus Deutsche Akademische Austausch Dienst, Bonn, Germany, and Ecole Polytechnique, France.

^(a)Present address: California Institute of Technology, Pasadena, Cal. 91125.

^(b)Present address: University of Michigan, Ann Arbor, Mich. 48104.

^(c)Present address: Southeastern Massachusetts University, N. Dartmouth, Mass. 02747.

^(d)Present address: Harvard University, Cambridge, Mass. 02138.

^(e)Present address: Universität Bonn, Bonn, Federal Republic of Germany.

^(f)Present address: Centre d'Etudes Nucléaires de Saclay, Gif-sur-Yvette, France.

¹A. De Rújula, H. Georgi, and S. L. Glashow, Phys. Rev. D **12**, 147 (1975), and M. K. Gaillard, B. W. Lee, and J. L. Rosner, Rev. Mod. Phys. **47**, 277 (1975).

²E. G. Cazzoli *et al.*, Phys. Rev. Lett. **34**, 1125 (1975); A. M. Cnops *et al.*, Phys. Rev. Lett. **42**, 197 (1979); C. Baltay *et al.*, Phys. Rev. Lett. **42**, 1721 (1979); C. Angelini *et al.*, CERN Report No. CERN-EP 79-44, 1979 (to be published).

³B. Knapp *et al.*, Phys. Rev. Lett. **37**, 882 (1976).

⁴D. Drijard *et al.*, Phys. Lett. **85B**, 452 (1979); K. L. Giboni *et al.*, Phys. Lett. **85B**, 437 (1979); W. Lockman *et al.*, Phys. Lett. **85B**, 443 (1979).

⁵G. S. Abrams *et al.*, Phys. Rev. Lett. **43**, 477 (1979); G. S. Abrams *et al.*, Phys. Rev. Lett. **43**, 481 (1979); G. S. Abrams *et al.*, Phys. Rev. Lett. **43**, 1555 (1979).

⁶An error which increases the magnitude of the momentum causes an increase in the directly calculated mass but a decrease in the mass calculated with the beam-energy constraint.

⁷R. Brandelik *et al.*, Nucl. Phys. B **148**, 189 (1979).

⁸Included in these efficiencies are corrections for low-momentum antiprotons (<0.3 GeV/c), which range out or are not tracked successfully (from +9% to +15%), corrections for antiprotons with momenta greater than 2.0 GeV/c (from 0% to +5%), and corrections for nuclear absorption (+8%), TOF electronic efficiency (+3%), and initial-state radiation (-15% to -5%). For the Λ efficiencies, the low-momentum (<0.5 GeV/c) correction is +20% to +33% and the nuclear-absorption corrections are +10% for Λ and +13% for $\bar{\Lambda}$.

⁹M. Piccolo *et al.*, Phys. Rev. Lett. **39**, 1503 (1977).

¹⁰An extensive Monte Carlo simulation of the detector, including the generation of drift-chamber coordinates, has been performed. The quoted efficiencies are derived from a complete reconstruction of Λ events so that tracking and vertex-formation inefficiencies are explicitly included.

¹¹At 5.2 GeV, the Λ_c can also be produced in association with a D meson via the final states $\Lambda_c \bar{D}^0 \bar{p}$ or $\Lambda_c D^- \bar{n}$. Approximately 20% of the Λ_c events have recoil masses above Dp threshold, providing an upper limit for such associated production in this sample.

¹²This neglects weak decays from all other charmed baryons, consistent with predictions and present experiments. See Refs. 1-3.

¹³This value is slightly model dependent. The quoted value as estimated from our measurement of $\Delta R(\Lambda + \bar{\Lambda})$ and $\Delta R(p + \bar{p})$ and a simple isospin statistical model.