

Observation of the decays $B^- \rightarrow D_s^{(*)+} K^- \pi^-$

The *BABAR* Collaboration

July 25, 2006

Abstract

We report first observations of the decays $B^- \rightarrow D_s^{(*)+} K^- \pi^-$, using 292 fb⁻¹ of data collected at the $\Upsilon(4S)$ resonance energy by the *BABAR* detector at the PEP-II e^+e^- collider. The branching fractions are measured to be $\mathcal{B}(B^- \rightarrow D_s^+ K^- \pi^-) = (1.88 \pm 0.13 \pm 0.41) \cdot 10^{-4}$ and $\mathcal{B}(B^- \rightarrow D_s^{*+} K^- \pi^-) = (1.84 \pm 0.19 \pm 0.40) \cdot 10^{-4}$.

Submitted to the 33rd International Conference on High-Energy Physics, ICHEP 06,
26 July—2 August 2006, Moscow, Russia.

Stanford Linear Accelerator Center, Stanford University, Stanford, CA 94309

Work supported in part by Department of Energy contract DE-AC02-76SF00515

The BABAR Collaboration,

B. Aubert, R. Barate, M. Bona, D. Boutigny, F. Couderc, Y. Karyotakis, J. P. Lees, V. Poireau,
V. Tisserand, A. Zghiche

*Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux,
France*

E. Grauges

Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

A. Palano

Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

J. C. Chen, N. D. Qi, G. Rong, P. Wang, Y. S. Zhu

Institute of High Energy Physics, Beijing 100039, China

G. Eigen, I. Ofte, B. Stugu

University of Bergen, Institute of Physics, N-5007 Bergen, Norway

G. S. Abrams, M. Battaglia, D. N. Brown, J. Button-Shafer, R. N. Cahn, E. Charles, M. S. Gill,
Y. Groysman, R. G. Jacobsen, J. A. Kadyk, L. T. Kerth, Yu. G. Kolomensky, G. Kukartsev, G. Lynch,
L. M. Mir, T. J. Orimoto, M. Pripstein, N. A. Roe, M. T. Ronan, W. A. Wenzel

Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

P. del Amo Sanchez, M. Barrett, K. E. Ford, A. J. Hart, T. J. Harrison, C. M. Hawkes, S. E. Morgan,
A. T. Watson

University of Birmingham, Birmingham, B15 2TT, United Kingdom

T. Held, H. Koch, B. Lewandowski, M. Pelizaeus, K. Peters, T. Schroeder, M. Steinke
Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

J. T. Boyd, J. P. Burke, W. N. Cottingham, D. Walker

University of Bristol, Bristol BS8 1TL, United Kingdom

D. J. Asgeirsson, T. Cuhadar-Donszelmann, B. G. Fulsom, C. Hearty, N. S. Knecht, T. S. Mattison,
J. A. McKenna

University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

A. Khan, P. Kyberd, M. Saleem, D. J. Sherwood, L. Teodorescu

Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

V. E. Blinov, A. D. Bukin, V. P. Druzhinin, V. B. Golubev, A. P. Onuchin, S. I. Serednyakov,
Yu. I. Skovpen, E. P. Solodov, K. Yu Todyshev

Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

D. S. Best, M. Bondioli, M. Bruinsma, M. Chao, S. Curry, I. Eschrich, D. Kirkby, A. J. Lankford, P. Lund,
M. Mandelkern, R. K. Mommsen, W. Roethel, D. P. Stoker

University of California at Irvine, Irvine, California 92697, USA

S. Abachi, C. Buchanan

University of California at Los Angeles, Los Angeles, California 90024, USA

S. D. Foulkes, J. W. Gary, O. Long, B. C. Shen, K. Wang, L. Zhang
University of California at Riverside, Riverside, California 92521, USA

H. K. Hadavand, E. J. Hill, H. P. Paar, S. Rahatlou, V. Sharma
University of California at San Diego, La Jolla, California 92093, USA

J. W. Berryhill, C. Campagnari, A. Cunha, B. Dahmes, T. M. Hong, D. Kovalskyi, J. D. Richman
University of California at Santa Barbara, Santa Barbara, California 93106, USA

T. W. Beck, A. M. Eisner, C. J. Flacco, C. A. Heusch, J. Kroseberg, W. S. Lockman, G. Nesom, T. Schalk,
B. A. Schumm, A. Seiden, P. Spradlin, D. C. Williams, M. G. Wilson
University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

J. Albert, E. Chen, A. Dvoretzkii, F. Fang, D. G. Hitlin, I. Narsky, T. Piatenko, F. C. Porter, A. Ryd,
A. Samuel
California Institute of Technology, Pasadena, California 91125, USA

G. Mancinelli, B. T. Meadows, K. Mishra, M. D. Sokoloff
University of Cincinnati, Cincinnati, Ohio 45221, USA

F. Blanc, P. C. Bloom, S. Chen, W. T. Ford, J. F. Hirschauer, A. Kreisel, M. Nagel, U. Nauenberg,
A. Olivas, W. O. Ruddick, J. G. Smith, K. A. Ulmer, S. R. Wagner, J. Zhang
University of Colorado, Boulder, Colorado 80309, USA

A. Chen, E. A. Eckhart, A. Soffer, W. H. Toki, R. J. Wilson, F. Winklmeier, Q. Zeng
Colorado State University, Fort Collins, Colorado 80523, USA

D. D. Altenburg, E. Feltresi, A. Hauke, H. Jasper, J. Merkel, A. Petzold, B. Spaan
Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

T. Brandt, V. Klose, H. M. Lacker, W. F. Mader, R. Nogowski, J. Schubert, K. R. Schubert, R. Schwierz,
J. E. Sundermann, A. Volk
Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

D. Bernard, G. R. Bonneaud, E. Latour, Ch. Thiebaux, M. Verderi
Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France

P. J. Clark, W. Gradl, F. Muheim, S. Playfer, A. I. Robertson, Y. Xie
University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

M. Andreotti, D. Bettoni, C. Bozzi, R. Calabrese, G. Cibinetto, E. Luppi, M. Negrini, A. Petrella,
L. Piemontese, E. Prencipe
Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

F. Anulli, R. Baldini-Ferroli, A. Calcaterra, R. de Sangro, G. Finocchiaro, S. Pacetti, P. Patteri,
I. M. Peruzzi,¹ M. Piccolo, M. Rama, A. Zallo
Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

¹Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

A. Buzzo, R. Capra, R. Contri, M. Lo Vetere, M. M. Macri, M. R. Monge, S. Passaggio, C. Patrignani,
E. Robutti, A. Santroni, S. Tosi

Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

G. Brandenburg, K. S. Chaisanguanthum, M. Morii, J. Wu

Harvard University, Cambridge, Massachusetts 02138, USA

R. S. Dubitzky, J. Marks, S. Schenk, U. Uwer

Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany

D. J. Bard, W. Bhimji, D. A. Bowerman, P. D. Dauncey, U. Egede, R. L. Flack, J. A. Nash,
M. B. Nikolich, W. Panduro Vazquez

Imperial College London, London, SW7 2AZ, United Kingdom

P. K. Behera, X. Chai, M. J. Charles, U. Mallik, N. T. Meyer, V. Ziegler

University of Iowa, Iowa City, Iowa 52242, USA

J. Cochran, H. B. Crawley, L. Dong, V. Eyges, W. T. Meyer, S. Prell, E. I. Rosenberg, A. E. Rubin

Iowa State University, Ames, Iowa 50011-3160, USA

A. V. Gritsan

Johns Hopkins University, Baltimore, Maryland 21218, USA

A. G. Denig, M. Fritsch, G. Schott

Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany

N. Arnaud, M. Davier, G. Grosdidier, A. Höcker, F. Le Diberder, V. Lepeltier, A. M. Lutz, A. Oyanguren,
S. Pruvot, S. Rodier, P. Roudeau, M. H. Schune, A. Stocchi, W. F. Wang, G. Wormser

*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique
d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France*

C. H. Cheng, D. J. Lange, D. M. Wright

Lawrence Livermore National Laboratory, Livermore, California 94550, USA

C. A. Chavez, I. J. Forster, J. R. Fry, E. Gabathuler, R. Gamet, K. A. George, D. E. Hutchcroft,
D. J. Payne, K. C. Schofield, C. Touramanis

University of Liverpool, Liverpool L69 7ZE, United Kingdom

A. J. Bevan, F. Di Lodovico, W. Menges, R. Sacco

Queen Mary, University of London, E1 4NS, United Kingdom

G. Cowan, H. U. Flaecher, D. A. Hopkins, P. S. Jackson, T. R. McMahon, S. Ricciardi, F. Salvatore,
A. C. Wren

*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United
Kingdom*

D. N. Brown, C. L. Davis

University of Louisville, Louisville, Kentucky 40292, USA

J. Allison, N. R. Barlow, R. J. Barlow, Y. M. Chia, C. L. Edgar, G. D. Lafferty, M. T. Naisbit,
J. C. Williams, J. I. Yi

University of Manchester, Manchester M13 9PL, United Kingdom

C. Chen, W. D. Hulsbergen, A. Jawahery, C. K. Lae, D. A. Roberts, G. Simi

University of Maryland, College Park, Maryland 20742, USA

G. Blaylock, C. Dallapiccola, S. S. Hertzbach, X. Li, T. B. Moore, S. Saremi, H. Staengle

University of Massachusetts, Amherst, Massachusetts 01003, USA

R. Cowan, G. Sciolla, S. J. Sekula, M. Spitznagel, F. Taylor, R. K. Yamamoto

*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139,
USA*

H. Kim, S. E. McLachlin, P. M. Patel, S. H. Robertson

McGill University, Montréal, Québec, Canada H3A 2T8

A. Lazzaro, V. Lombardo, F. Palombo

Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy

J. M. Bauer, L. Cremaldi, V. Eschenburg, R. Godang, R. Kroeger, D. A. Sanders, D. J. Summers,
H. W. Zhao

University of Mississippi, University, Mississippi 38677, USA

S. Brunet, D. Côté, M. Simard, P. Taras, F. B. Viaud

Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7

H. Nicholson

Mount Holyoke College, South Hadley, Massachusetts 01075, USA

N. Cavallo,² G. De Nardo, F. Fabozzi,³ C. Gatto, L. Lista, D. Monorchio, P. Paolucci, D. Piccolo,
C. Sciacca

Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy

M. A. Baak, G. Raven, H. L. Snoek

*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The
Netherlands*

C. P. Jessop, J. M. LoSecco

University of Notre Dame, Notre Dame, Indiana 46556, USA

T. Allmendinger, G. Benelli, L. A. Corwin, K. K. Gan, K. Honscheid, D. Hufnagel, P. D. Jackson,
H. Kagan, R. Kass, A. M. Rahimi, J. J. Regensburger, R. Ter-Antonyan, Q. K. Wong

Ohio State University, Columbus, Ohio 43210, USA

N. L. Blount, J. Brau, R. Frey, O. Igonkina, J. A. Kolb, M. Lu, R. Rahmat, N. B. Sinev, D. Strom,
J. Strube, E. Torrence

University of Oregon, Eugene, Oregon 97403, USA

²Also with Università della Basilicata, Potenza, Italy

³Also with Università della Basilicata, Potenza, Italy

A. Gaz, M. Margoni, M. Morandin, A. Pompili, M. Posocco, M. Rotondo, F. Simonetto, R. Stroili, C. Voci
Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy

M. Benayoun, H. Briand, J. Chauveau, P. David, L. Del Buono, Ch. de la Vaissière, O. Hamon,
B. L. Hartfiel, M. J. J. John, Ph. Leruste, J. Malcès, J. Ocariz, L. Roos, G. Therin
*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie
Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*

L. Gladney, J. Panetta
University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

M. Biasini, R. Covarelli
Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy

C. Angelini, G. Batignani, S. Bettarini, F. Bucci, G. Calderini, M. Carpinelli, R. Cenci, F. Forti,
M. A. Giorgi, A. Lusiani, G. Marchiori, M. A. Mazur, M. Morganti, N. Neri, E. Paoloni, G. Rizzo,
J. J. Walsh
Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy

M. Haire, D. Judd, D. E. Wagoner
Prairie View A&M University, Prairie View, Texas 77446, USA

J. Biesiada, N. Danielson, P. Elmer, Y. P. Lau, C. Lu, J. Olsen, A. J. S. Smith, A. V. Telnov
Princeton University, Princeton, New Jersey 08544, USA

F. Bellini, G. Cavoto, A. D’Orazio, D. del Re, E. Di Marco, R. Faccini, F. Ferrarotto, F. Ferroni,
M. Gaspero, L. Li Gioi, M. A. Mazzoni, S. Morganti, G. Piredda, F. Polci, F. Safai Tehrani, C. Voena
Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy

M. Ebert, H. Schröder, R. Waldi
Universität Rostock, D-18051 Rostock, Germany

T. Adye, N. De Groot, B. Franek, E. O. Olaiya, F. F. Wilson
Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom

R. Aleksan, S. Emery, A. Gaidot, S. F. Ganzhur, G. Hamel de Monchenault, W. Kozanecki, M. Legendre,
G. Vasseur, Ch. Yèche, M. Zito
DSM/Daphnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France

X. R. Chen, H. Liu, W. Park, M. V. Purohit, J. R. Wilson
University of South Carolina, Columbia, South Carolina 29208, USA

M. T. Allen, D. Aston, R. Bartoldus, P. Bechtle, N. Berger, R. Claus, J. P. Coleman, M. R. Convery,
M. Cristinziani, J. C. Dingfelder, J. Dorfan, G. P. Dubois-Felsmann, D. Dujmic, W. Dunwoodie,
R. C. Field, T. Glanzman, S. J. Gowdy, M. T. Graham, P. Grenier,⁴ V. Halyo, C. Hast, T. Hryn’ova,
W. R. Innes, M. H. Kelsey, P. Kim, D. W. G. S. Leith, S. Li, S. Luitz, V. Luth, H. L. Lynch,
D. B. MacFarlane, H. Marsiske, R. Messner, D. R. Muller, C. P. O’Grady, V. E. Ozcan, A. Perazzo,
M. Perl, T. Pulliam, B. N. Ratcliff, A. Roodman, A. A. Salnikov, R. H. Schindler, J. Schwiening,
A. Snyder, J. Stelzer, D. Su, M. K. Sullivan, K. Suzuki, S. K. Swain, J. M. Thompson, J. Va’vra, N. van

⁴Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France

Bakel, M. Weaver, A. J. R. Weinstein, W. J. Wisniewski, M. Wittgen, D. H. Wright, A. K. Yarritu, K. Yi,
C. C. Young

Stanford Linear Accelerator Center, Stanford, California 94309, USA

P. R. Burchat, A. J. Edwards, S. A. Majewski, B. A. Petersen, C. Roat, L. Wilden

Stanford University, Stanford, California 94305-4060, USA

S. Ahmed, M. S. Alam, R. Bula, J. A. Ernst, V. Jain, B. Pan, M. A. Saeed, F. R. Wappler, S. B. Zain

State University of New York, Albany, New York 12222, USA

W. Bugg, M. Krishnamurthy, S. M. Spanier

University of Tennessee, Knoxville, Tennessee 37996, USA

R. Eckmann, J. L. Ritchie, A. Satpathy, C. J. Schilling, R. F. Schwitters

University of Texas at Austin, Austin, Texas 78712, USA

J. M. Izen, X. C. Lou, S. Ye

University of Texas at Dallas, Richardson, Texas 75083, USA

F. Bianchi, F. Gallo, D. Gamba

Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy

M. Bomben, L. Bosisio, C. Cartaro, F. Cossutti, G. Della Ricca, S. Dittongo, L. Lanceri, L. Vitale

Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy

V. Azzolini, N. Lopez-March, F. Martinez-Vidal

IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain

Sw. Banerjee, B. Bhuyan, C. M. Brown, D. Fortin, K. Hamano, R. Kowalewski, I. M. Nugent, J. M. Roney,
R. J. Sobie

University of Victoria, Victoria, British Columbia, Canada V8W 3P6

J. J. Back, P. F. Harrison, T. E. Latham, G. B. Mohanty, M. Pappagallo

Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom

H. R. Band, X. Chen, B. Cheng, S. Dasu, M. Datta, K. T. Flood, J. J. Hollar, P. E. Kutter, B. Mellado,
A. Mihalyi, Y. Pan, M. Pierini, R. Prepost, S. L. Wu, Z. Yu

University of Wisconsin, Madison, Wisconsin 53706, USA

H. Neal

Yale University, New Haven, Connecticut 06511, USA

1 INTRODUCTION

First evidence for so-called inclusive *flavor correlated production* of D_s^+ in B^- decays was reported recently [1] with a branching fraction of $\mathcal{B}(B^- \rightarrow D_s^+ X) = (1.2 \pm 0.4)\%$ [2]. These decays are mediated by a $b \rightarrow c$ quark transition and require at least three final state particles, including the production of an $s\bar{s}$ pair from the vacuum ($s\bar{s}$ “popping”). An example for a three-body B^- decay with a D_s^+ in the final state is $B^- \rightarrow D_s^+ K^- \pi^-$. The corresponding \bar{B}^0 decay is $\bar{B}^0 \rightarrow D_s^+ \bar{K}^0 \pi^-$. The Feynman diagram for $B^- \rightarrow D_s^{(*)+} K^- \pi^-$ decays is shown in Fig. 1. In case of $\bar{B}^0 \rightarrow D_s^+ \bar{K}^0 \pi^-$, an additional contribution from a W -exchange diagram with $s\bar{s}$ and $d\bar{d}$ popping may exist. If we




Figure 1: Feynman diagram for $B^- \rightarrow D_s^+ K^- \pi^-$.

replace the π^- in Fig. 1, which comes from the hadronization of the W^- boson with a K^- , we get the Cabibbo-suppressed decays $B^- \rightarrow D_s^+ K^- K^-$ and $\bar{B}^0 \rightarrow D_s^+ \bar{K}^0 K^-$. It is interesting to note that the final state $D_s^+ \bar{K}^0 K^-$ can also be reached from a B^0 decay. In this case the decay is mediated by a $b \rightarrow u$ quark transition, but the W hadronization is not Cabibbo-suppressed. Thus a \bar{B}^0 can either decay directly to $D_s^+ \bar{K}^0 K^-$ or via $B^0 \bar{B}^0$ mixing followed by $B^0 \rightarrow D_s^+ \bar{K}^0 K^-$. The interference between the two decay amplitudes for decay with and without $B^0 \bar{B}^0$ mixing leads to a time-dependent CP -asymmetry that is sensitive to $\sin(2\beta + \gamma)$. In case the contribution from the higher D^{**} resonances decaying into $D_s^+ \bar{K}$ turns out to be large, it may also be interesting to measure the resonant parameters independently from the analysis using $B \rightarrow \bar{D} \pi \pi$ decays [3].

No exclusive $B^- \rightarrow D_s^{(*)+} X$ or $\bar{B}^0 \rightarrow D_s^{(*)+} X$ decay mode has hitherto been observed. Limits on the branching fractions from the analyses by other experiments are listed in Table 1. In this paper we report the first measurement of the decay modes $B^- \rightarrow D_s^{(*)+} K^- \pi^-$.

2 THE BABAR DETECTOR AND DATASET

The analysis uses a sample of approximately 292 fb^{-1} , which corresponds to about 324 million $\Upsilon(4S)$ decays into $B\bar{B}$ pairs collected with the *BABAR* detector at the PEP-II [4] asymmetric-energy B -factory. The *BABAR* detector is described elsewhere [5] and only the components crucial to this analysis are summarized here. Charged particle tracking is provided by a five-layer silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH). For charged-particle identification,

Table 1: Upper limits from ARGUS [6] and CLEO [7] on $B^- \rightarrow D_s^{(*)+} K^- \pi^-$ branching fractions.

Experiment	Decay Mode	Upper limit (@90% C.L.)
ARGUS	$B^- \rightarrow D_s^+ K^- \pi^-$	8×10^{-4}
	$B^- \rightarrow D_s^{*+} K^- \pi^-$	12×10^{-4}
CLEO	$B^- \rightarrow D_s^+ K^- \pi^-$	5×10^{-4}
	$B^- \rightarrow D_s^{*+} K^- \pi^-$	6.8×10^{-4}

ionization energy loss (dE/dx) in the DCH and SVT, and Cherenkov radiation detected in a ring-imaging device are used. Photons are identified and measured using a thallium-doped CsI-crystal electromagnetic calorimeter. These systems are located inside a 1.5 T solenoidal superconducting magnet. We use GEANT4 [8] software to simulate interactions of particles traversing the *BABAR* detector, taking into account the varying detector conditions and beam backgrounds.

3 ANALYSIS METHOD

The optimal selection criteria as well as the probability density distributions of selection variables are determined by a blind analysis based on Monte Carlo (MC) simulation of both signal and background. For the calculation of the expected signal yield we assume $\mathcal{B}(B^- \rightarrow D_s^{(*)+} K^- \pi^-)$ to be 10^{-4} (*i.e.* about 10% of the measured $\mathcal{B}(B^- \rightarrow D^+ \pi^- \pi^-)$ [3]). We use MC samples of our signal modes and, to simulate background, inclusive samples of $B^+ B^-$ (784 fb^{-1}), $B^0 \bar{B}^0$ (774 fb^{-1}), $c\bar{c}$ (247 fb^{-1}), and $q\bar{q}$, $q = u, d, s$ (246 fb^{-1}). In addition, we use large samples of simulated events of rare background modes which have final states similar to the signal. We have verified that our MC correctly describes the data by comparing distributions of various selection variables.

Candidates for D_s^+ mesons are reconstructed in the modes $D_s^+ \rightarrow \phi \pi^+$, $\bar{K}^{*0} K^+$, and $K_s^0 K^+$, with $\phi \rightarrow K^+ K^-$, $\bar{K}^{*0} \rightarrow K^- \pi^+$ and $K_s^0 \rightarrow \pi^+ \pi^-$. The K_s^0 candidates are reconstructed from two oppositely-charged tracks, that come from a common vertex displaced from the $e^+ e^-$ interaction point. We require the significance of this displacement (measured flight distance divided by an estimated error) to exceed 2. All other tracks are required to originate less than 1.5 cm away from the $e^+ e^-$ interaction point in the transverse plane and less than 10 cm along the beam axis. Charged kaon candidates must satisfy kaon identification criteria that are typically around 92% efficient, depending on momentum and polar angle, and have a pion misidentification rate at the 5% level. The $\phi \rightarrow K^+ K^-$, $\bar{K}^{*0} \rightarrow K^- \pi^+$ and $K_s^0 \rightarrow \pi^+ \pi^-$ candidates are required to have invariant masses close to their nominal masses (we require the absolute differences between their measured masses and the nominal values [9] to be in the range $\pm 15 \text{ MeV}$, $\pm 50 \text{ MeV}$ and $\pm 10 \text{ MeV}$, respectively). The polarizations of the \bar{K}^{*0} and ϕ mesons in the D_s^+ decays are employed to reject backgrounds through the use of the helicity angle θ_H , defined as the angle between the K^- momentum vector and the direction of flight of the D_s^+ in the \bar{K}^{*0} or ϕ rest frame. The \bar{K}^{*0} and ϕ candidates are required to have $|\cos \theta_H|$ greater than 0.5.

The D_s^{*+} candidates are reconstructed in the mode $D_s^{*+} \rightarrow D_s^+ \gamma$. The photons are accepted if their energy is greater than 100 MeV. The D_s^+ and D_s^{*+} candidates are required to have invariant

masses in the interval $[-10, 10]$ MeV/ c^2 (for D_s^+) and $[-15, 10]$ MeV/ c^2 (for D_s^{*+}) from their nominal values [9] (the D_s^+ mass resolution is around 6 MeV/ c^2 , and the asymmetric mass cut on D_s^{*+} has an efficiency of about 90%). All D_s^+ candidates are mass-constrained. The invariant mass of the D_s^{*+} is calculated after a mass constraint on the daughter D_s^+ has been applied. Subsequently, all D_s^{*+} candidates are subjected to a mass-constrained fit.

We also require that photons from D_s^{*+} are inconsistent with π^0 hypothesis when combined with any other photon having an energy greater than 150 MeV in the event (the π^0 veto window is ± 10 MeV/ c^2). Finally, the B^- meson candidates are formed using the reconstructed combinations of $D_s^+ K^- \pi^-$ and $D_s^{*+} K^- \pi^-$.

The background from continuum $q\bar{q}$ production (where $q = u, d, s, c$) is suppressed based on the event topology. We calculate the angle (θ_T) between the thrust axis of the B meson candidate and the thrust axis of all other particles in the event in the center-of-mass frame (c.m.). In this frame, $B\bar{B}$ pairs are produced approximately at rest and have a uniform $\cos\theta_T$ distribution. In contrast, $q\bar{q}$ pairs are produced in the c.m. frame with high momenta, which results in a $|\cos\theta_T|$ distribution peaking at 1. $|\cos\theta_T|$ is required to be smaller than 0.8. In addition, the ratio of the second and zeroth order Fox-Wolfram moments [10] must be less than 0.3.

We extract the signal using the kinematical variables $m_{\text{ES}} = \sqrt{E_b^{*2} - (\sum_i \mathbf{p}_i^*)^2}$ and $\Delta E = \sum_i \sqrt{m_i^2 + \mathbf{p}_i^{*2}} - E_b^*$, where E_b^* is the beam energy in the c.m. frame, \mathbf{p}_i^* is the c.m. momentum of the daughter particle i of the B^- meson candidate, and m_i is the mass hypothesis for particle i . For signal events, m_{ES} peaks at the B^- meson mass with a resolution of about 2.6 MeV/ c^2 and ΔE peaks near zero with a resolution of 13 MeV, indicating that the B^- candidate has a total energy consistent with the beam energy in the c.m. frame. The B^- candidates are required to have $|\Delta E| < 25$ MeV (around 2σ of the signal ΔE resolution) and $m_{\text{ES}} > 5.2$ GeV/ c^2 .

The fraction of events with multiple B^- candidates is estimated using the MC simulation and found to be around 3% for $D_s^+ K^- \pi^-$ and 9% for $D_s^{*+} K^- \pi^-$ combinations. In each event with more than one B^- candidate that passes the selection requirements, we select the one with the lowest $|\Delta E|$ value.

After all selection criteria are applied, we estimate the B^- reconstruction efficiencies, excluding the subsequent branching fractions (see Table 2).

Table 2: Reconstruction efficiencies for $B^- \rightarrow D_s^{(*)+} K^- \pi^-$ decays (excluding the subsequent branching fractions).

Decay mode	$D_s^+ \rightarrow \phi\pi^+$	$D_s^+ \rightarrow \bar{K}^{*0}K^+$	$D_s^+ \rightarrow K_S^0 K^+$
$B^- \rightarrow D_s^+ K^- \pi^-$	11.0%	7.0%	10.0%
$B^- \rightarrow D_s^{*+} K^- \pi^-$	5.3%	3.4%	4.8%

Background events that pass these selection criteria are represented by approximately equal amounts of $q\bar{q}$ continuum and $B\bar{B}$ events. We parametrize their m_{ES} distribution by a threshold function [11]:

$$f(m_{\text{ES}}) \sim m_{\text{ES}} \sqrt{1 - x^2} \exp[-\xi(1 - x^2)],$$

where $x = 2m_{ES}/\sqrt{s}$, \sqrt{s} is the total energy of the beams in their center of mass frame, and ξ is the fit parameter.

A study using simulated B^0 and B^+ decays shows that some background events with distributions in m_{ES} and in ΔE peaking near the signal region are expected in reconstructed $B^- \rightarrow D_s^+ K^- \pi^-$ candidates due to charmless and charmonium B^- decays with the same set of particles in the final state. For $B^- \rightarrow D_s^{*+} K^- \pi^-$, no background of this kind is expected, due to the presence of the γ , which suppresses charmless and charmonium decay contributions. The peaking contribution is evaluated using the data by reconstructing “ $D_s^{(*)+}$ ” $K^- \pi^-$ combinations, where “ D_s^+ ” candidates are selected from $[\pm 40, \pm 25]$ MeV sidebands around the D_s^+ nominal mass. In this procedure, we use the same selection requirements, as for the signal, except that “ D_s^+ ” candidates are not mass constrained. The resulting m_{ES} spectra are shown in Figure 2. We fit the distributions using an extended unbinned maximum likelihood (ML) fit with a sum of a Gaussian (with a width and central value fixed from the MC simulation) and a threshold function $f(m_{ES})$ with the floating shape and normalization (see detailed expression of the likelihood function is Section 5). The fit yields 34 ± 12 events in the “signal” m_{ES} peak for “ D_s^+ ” $K^- \pi^-$ and 3 ± 7 for “ D_s^{*+} ” $K^- \pi^-$. Since the sideband interval is 1.5 times larger than the D_s^+ mass region used for signal selection, this translates into 23 ± 8 peaking background events expected in $B^- \rightarrow D_s^+ K^- \pi^-$.




Figure 2: m_{ES} spectra for the data “ D_s^+ ” $K^- \pi^-$ (left) and “ D_s^{*+} ” $K^- \pi^-$ (right) combinations with no mass constraint applied on “ D_s^+ ” candidates, using D_s^+ mass sidebands $[\pm 25, \pm 40]$ MeV (1.5 times the signal interval).

We also study cross-feed between the signal modes and other decays with final states similar to our signal modes, including $D_s^{(*)+} K^- K^-$. The cut on ΔE of the B^- candidates effectively suppresses the cross-feed contributions, which do not exceed 2% of the reconstructed signal after all the selection criteria are applied.

4 SYSTEMATIC STUDIES

The summary of the systematic uncertainties is presented in Table 3. The total relative systematic error is estimated to be 22% for each B^- decay mode, with the largest contribution coming from the D_s^+ branching fractions uncertainty (15%). Other significant sources of systematic errors are found to be due to the difference between the selection efficiency in MC events and in the data

(estimated using the control mode $B^- \rightarrow D_s^- D^0$, $D^0 \rightarrow K^- \pi^+$), and also due to the efficiency dependence on the $D_s^{(*)+} K^-$ invariant mass and its potential effect if the resonant contribution is present.

Table 3: Summary of relative systematic errors (in %) for $B^- \rightarrow D_s^{(*)+} K^- \pi^-$ decays.

Source	$B^- \rightarrow D_s^+ K^- \pi^-$	$B^- \rightarrow D_s^{*+} K^- \pi^-$
B counting	1.1	1.1
MC statistics	0.8	1.4
Tracking	5	5
Particle identification efficiency	4	4
K_S^0 efficiency	0.5	0.5
γ (from $D_s^{*+} \rightarrow D_s^+ \gamma$) efficiency	–	2
\mathcal{B} of sub-decays	15	15
Peaking background contribution	6	3
Cross-feed contribution	1	2
Selection efficiency, Data/MC	12	12
Signal and background shape uncertainty	3	3
$M(D_s^{(*)-} K^+)$ efficiency dependence	7	9
Total	22	22

5 RESULTS

Figure 3 shows the m_{ES} distributions for the reconstructed candidates $B^- \rightarrow D_s^+ K^- \pi^-$ and $B^- \rightarrow D_s^{*+} K^- \pi^-$. For each mode, we perform an extended unbinned ML fit to the m_{ES} distributions using the candidates from all D_s^+ decay modes combined. We fit the m_{ES} distributions with the sum of the function $f(m_{\text{ES}})$ characterizing the combinatorial background and a Gaussian function to describe the signal. The mean and width of the Gaussian function, the threshold shape parameter ξ , and the numbers of signal (n_{sig}) and background (n_{bkg}) events are free parameters of the fit. The likelihood function is given by:

$$\mathcal{L} = \frac{e^{-(n_{\text{sig}}+n_{\text{bkg}})}}{N!} \prod_{i=1}^N (n_{\text{sig}} P_i^{\text{sig}} + n_{\text{bkg}} P_i^{\text{bkg}}),$$

where P_i^{sig} and P_i^{bkg} are the probability density functions for the signal and background, N is the total number of events in the fit and i is the index over all events in the fit.

The fit yields 393 ± 25 events in the $B^- \rightarrow D_s^+ K^- \pi^-$ mode. Taking into account the estimated peaking background contribution, we obtain 370 ± 26 signal events for $B^- \rightarrow D_s^+ K^- \pi^-$. The




Figure 3: m_{ES} spectra for the $B^- \rightarrow D_s^+ K^- \pi^-$ (left) and $B^- \rightarrow D_s^{*+} K^- \pi^-$ (right) using the data. Solid curves show the fit results, as explained in the text. Dashed lines in the signal regions correspond to the background components of the fit.

number of $B^- \rightarrow D_s^{*+} K^- \pi^-$ signal events from the fit is 164 ± 17 (no peaking contribution is subtracted in this mode as it was estimated to be consistent with 0). We also fit signal yields in each of the D_s^+ modes, fixing the width and central values of the signal Gaussians to that of the combined fit, letting the background level and shape float. The ratio of the signal yields between submodes is consistent with the expectations from MC.

The total signal yield in each B^- decay mode is calculated as a sum over D_s^+ modes ($i = \phi\pi^+, \bar{K}^{*0}K^+, K_s^0K^+$) and is related to the B^- branching fraction \mathcal{B} using the following expression:

$$n_{sig} = \mathcal{B} \cdot N_{B\bar{B}} \cdot \sum_i \mathcal{B}_i \cdot \epsilon_i,$$

where $N_{B\bar{B}}$ is the number of produced $B\bar{B}$ pairs, \mathcal{B}_i is the product of the intermediate branching ratios and ϵ_i is the reconstruction efficiency (from Table 2). As an input to the calculation, we used branching fraction numbers from [9] and [12]. The relative systematic uncertainties are converted into absolute numbers using the measured central values. The results are:

$$\mathcal{B}(B^- \rightarrow D_s^+ K^- \pi^-) = (1.88 \pm 0.13 \pm 0.41) \cdot 10^{-4}$$

$$\mathcal{B}(B^- \rightarrow D_s^{*+} K^- \pi^-) = (1.84 \pm 0.19 \pm 0.40) \cdot 10^{-4}$$

In summary, two decay modes of charged B mesons are observed for the first time. The significance of the observation is 14.2σ for $B^- \rightarrow D_s^+ K^- \pi^-$ and 9.6σ for $B^- \rightarrow D_s^{*+} K^- \pi^-$.

6 ACKNOWLEDGMENTS

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and

National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy Physics (China), the Commissariat à l’Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, Ministerio de Educación y Ciencia (Spain), and the Particle Physics and Astronomy Research Council (United Kingdom). Individuals have received support from the Marie-Curie IEF program (European Union) and the A. P. Sloan Foundation.

References

- [1] BaBar Collaboration, B. Aubert *et al.*, [hep-ex/0606026](#).
- [2] Charge conjugate reactions are implicitly included, throughout this paper.
- [3] Belle Collaboration, K. Abe *et al.*, *Phys. Rev. D* **69**, 112002 (2004).
- [4] PEP-II Conceptual Design Report, SLAC-0418 (1993).
- [5] The *BABAR* Collaboration, B. Aubert *et al.*, *Nucl. Instrum. Methods* **A479**, 1-116 (2002).
- [6] ARGUS Collaboration, H. Albrecht *et al.*, *Zeitschrift für Physik* **C60**, 11 (1993).
- [7] A. H. Cho, “*A study of lower vertex production of D_s mesons in B meson decays*”, Cornell University, Ph.D thesis (unpublished).
- [8] Geant4 Collaboration, S. Agostinelli *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [9] Particle Data Group, S. Eidelman *et al.*, *Phys. Lett.* **B**, 592 (2004).
- [10] G.C. Fox and S. Wolfram, *Phys. Rev. Lett.* **41**, 1581 (1978).
- [11] ARGUS Collaboration, H. Albrecht *et al.*, *Z. Phys. C* **48**, 543 (1990).
- [12] BaBar Collaboration, B. Aubert *et al.*, *Phys. Rev. D* **71**, 091104 (2005).