

**Observed atlantification of the Barents Sea causes the Polar Front to limit the expansion of winter sea ice**

Barton, Benjamin; Lenn, Yueng-Djern; Lique, Camille

Journal of Physical Oceanography

DOI:
[10.1175/JPO-D-18-0003.1](https://doi.org/10.1175/JPO-D-18-0003.1)

Published: 15/08/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):
Barton, B., Lenn, Y-D., & Lique, C. (2018). Observed atlantification of the Barents Sea causes the Polar Front to limit the expansion of winter sea ice. *Journal of Physical Oceanography*.
<https://doi.org/10.1175/JPO-D-18-0003.1>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Observed atlantification of the Barents Sea causes the Polar Front to limit**
2 **the expansion of winter sea ice**

3 Benjamin I. Barton^{*}, Yueng-Djern Lenn[†] and Camille Lique[‡]

4 *Laboratoire d'Océanographie Physique et Spatiale, UMR6523, CNRS-Ifremer-UBO-IRD, Brest,*
5 *France,*

6 **Corresponding author address: Benjamin I. Barton, Laboratoire d'Océanographie Physique et*
7 *Spatiale (LOPS), IUEM, Rue Dumont d'Urville, 29280 Plouzané, France.*

8 E-mail: benjamin.barton@univ-brest.fr

9 [†]Ocean Sciences, Bangor University, Bangor, UK.

10 [‡]Laboratoire d'Océanographie Physique et Spatiale, UMR6523, CNRS-Ifremer-UBO-IRD, Brest,
11 France

ABSTRACT

12 Barents Sea Water (BSW) is formed from Atlantic Water that is cooled
13 through atmospheric heat loss and freshened through seasonal sea ice melt.
14 In the eastern Barents Sea, the BSW and fresher, colder Arctic Water meet
15 at the surface along the Polar Front (PF). Despite its importance in setting
16 the northern limit of BSW ventilation, the PF has been poorly-documented,
17 mostly eluding detection by observational surveys that avoid seasonal sea ice.
18 In this study, satellite sea surface temperature (SST) observations are used
19 in addition to a temperature and salinity climatology to examine the location
20 and structure of the PF, and characterise its variability over the period 1985 –
21 2016. It is shown that the PF is independent of the position of the sea ice
22 edge and is a shelf slope current constrained by potential vorticity. The main
23 driver of interannual variability in SST is the variability of the Atlantic Water
24 temperature, which has significantly increased since 2005. The SST gradient
25 associated with the PF has also increased after 2005, preventing sea ice from
26 extending south of the front during winter in recent years. The disappearance
27 of fresh, seasonal sea ice melt south of the PF has led to a significant increase
28 in BSW salinity and density. As BSW forms the majority of Arctic Interme-
29 diate Water, changes to BSW properties may have far-reaching impacts for
30 Arctic Ocean circulation and climate.

31 **1. Introduction**

32 The Arctic has been predicted to be sea-ice-free in summer by the middle of the twenty-first cen-
33 tury (Wang and Overland 2012; Snape 2013; Notz and Stroeve 2016). This follows an Arctic-wide
34 decline in sea ice extent over recent decades (Screen and Simmonds 2010). The Barents Sea alone
35 has seen a 50% reduction in annual sea ice area between 1998 and 2008 (Årthun et al. 2012), asso-
36 ciated with a strong sea ice decline in all seasons including winter (Onarheim and Årthun 2017).
37 Seasonal sea ice extent variations are very predictable in the Barents Sea compared with other
38 parts of the Arctic (Sigmond et al. 2016). For instance, Day et al. (2014) found significant correla-
39 tions between Arctic sea ice extent in May, and Barents Sea sea surface temperature (SST) for the
40 same month, as well as with SST in the preceding December. The variability of the Barents Sea
41 ice edge location has also been associated with atmospheric circulation (Sorteberg and Kvingedal
42 2006), and ice exported from the Arctic to the Barents Sea due to local winds (Koenigk et al. 2009;
43 Kwok 2009). On longer time scales, the reduction in annual and winter Barents Sea sea ice area is
44 thought to be driven by an increase in the heat transport into the Barents Sea due to the combined
45 increase in advection and temperature of Atlantic Water (AW) (Årthun et al. 2012; Onarheim et al.
46 2015). AW temperature and salinity in the Barents Sea are also varying on multidecadal timescales
47 (Levitus et al. 2009; Smedsrud et al. 2013), making it challenging to distinguish between long term
48 trend and natural variability.

49 Along with Fram Strait, the Barents Sea Opening (BSO) is a gateway for the warm and salty AW
50 (defined in Table 1) entering the Arctic Ocean and its marginal seas (Figure 1). The branch of AW
51 entering through the BSO transits the Barents Sea, where it is modified en route, forming Barents
52 Sea Water (BSW, Table 1) (Schauer et al. 2002; Harris et al. 1998). This transformation into BSW
53 is driven mainly by surface interactions with the atmosphere resulting in winter convection and

54 entrainment of freshwater. Heat is lost from the ocean through turbulent heat flux and longwave
55 radiation (Long and Perrie 2017), while freshwater input mostly comes from seasonal sea ice
56 import and rivers (Ellingsen et al. 2009). Thus the length and location of the pathway along which
57 AW flows determines to what extent its properties are modified by surface fluxes, sea ice and rivers
58 before it enters the Arctic Basin. The Barents Sea bathymetry is known to strongly influence the
59 path of AW inflow (Loeng 1991) (Figure 1). Part of the AW inflow crosses Murmansk Rise, south
60 of Central Bank, into the Central Basin (Skagseth 2008; Ingvaldsen 2005). The Central Basin
61 acts as a reservoir for AW until it loses enough buoyancy to propagate northwards below ArW as
62 BSW. As a result, the water column is stratified in the northern Barents Sea, with the upper 100 m
63 occupied by relatively fresh and cold Arctic Water (ArW, Table 1) and the lower layer occupied
64 by BSW (Harris et al. 1998; Lind and Ingvaldsen 2012).

65 *In situ* observations in the western Barents Sea have revealed that the surface expression of the
66 front separating AW from ArW follows isobaths in the range 150 - 275 m (Gawarkiewicz and
67 Plueddemann 1995; Harris et al. 1998; Våge et al. 2014; Fer and Drinkwater 2014). In the eastern
68 Barents Sea, the northern front (referred to as the Polar Front - PF hereafter) is defined as the
69 location where BSW meets ArW but its geographic position is poorly defined (Oziel et al. 2016).
70 The PF is a water mass boundary and therefore should have an SST signature. This PF should be
71 distinguished from another nearby SST front (hereafter thermal-surface front) that is also expected
72 to be present in the surface layer of the northern Barents Sea following the sea ice edge, due to the
73 transition from freezing-point water to ice-free water. In the range of temperatures and salinities
74 of BSW and ArW, salinity and temperature tend to contribute equally to the determination of
75 density (Parsons et al. 1996; Våge et al. 2014). Thus, both surface temperature and surface salinity
76 contribute to the PF's surface density gradient, suggesting that the variability of the PF position
77 can be influenced by other processes than just the position of the sea ice edge.

78 BSW exits the Barents Sea, entering the Arctic Ocean mainly through St. Anna Trough (Rudels
79 et al. 2000; Smedsrud et al. 2013). In the Arctic Ocean, BSW is entrained into Arctic Intermediate
80 Water (AIW), accounting for 50 - 80% of the volume of AIW (Schauer et al. 1997; Maslowski et al.
81 2004). AIW is ultimately exported to the North Atlantic through Fram Strait and in turn contributes
82 to the deeper branch of the Atlantic Meridional Overturning Circulation (AMOC) (Aagaard et al.
83 1985; Fahrbach et al. 2001). There is some debate in the literature about the extent to which
84 BSW properties at the exit of the Barents Sea are preserved into the Arctic Ocean and beyond.
85 Observations have revealed that some mixing of BSW occurs on continental slopes and within the
86 Arctic Ocean (Shapiro 2003; Rudels et al. 2015) but model results of Lique et al. (2010) show
87 modifications to BSW properties within the Arctic Ocean are small compared to the modification
88 within the Barents Sea. In either case, the properties with which BSW exits the Barents Sea are
89 important as they pre-condition it for the target depth at which it may settle and mix with ambient
90 water masses within the Arctic Basin. Anomalies in BSW density have been traced to Denmark
91 Strait suggesting far-reaching impacts from processes occurring in the Barents Sea (Karcher et al.
92 2011).

93 It has been hypothesised by Aagaard and Woodgate (2001) that a prolonged reduction in the
94 fresh, melt water input from seasonal sea ice into BSW could cause a modification of the BSW
95 properties, and in turn induce a warming and salinification of AIW. This hypothesis overlooks
96 the role the PF could play in determining whether the meltwater is entrained into BSW or ArW
97 and discounts the influence of changes in other water masses in the Barents Sea. Indeed, both
98 the transport and the temperature of AW circulating in the Barents Sea have increased in recent
99 decades (Årthun et al. 2012), resulting in a reduction in winter sea ice area through a decrease
100 in wind-driven sea ice advection and delayed winter refreezing (Lien et al. 2017). Thus, winter
101 sea ice extent trends are consistent with the emerging evidence of ongoing atlantification (i.e. the

102 increased influence of AW resulting in a warming and salinification) of the Barents Sea (Reigstad
103 et al. 2002; Oziel et al. 2017) and Arctic Ocean (Polyakov et al. 2017). This makes it important to
104 quantify the role that Barents Sea ice trends play on BSW properties.

105 The goal of this study is to investigate the variability of SST in order to characterise the PFs
106 location in the eastern Barents Sea, determine how this compares to the seasonal sea ice edge and
107 what the implications for BSW formation are given the documented sea ice loss and atlantifica-
108 tion of the Barents Sea? To that aim, we use a combination of the new high-resolution, 32-year
109 OSTIA SST dataset, satellite observations of sea ice concentration and 3D optimally interpolated
110 temperature and salinity products.

111 The methods and tools are presented in Section 2. In order to identify forcings on the formation
112 of BSW, in Section 3, the mechanisms that cause variability in SST on seasonal and to multidecadal
113 time-scales in the Barents Sea are explored. In Section 4, SST is used to pinpoint the surface
114 expression of the PF, and determine whether the winter sea ice edge has become bound by it. In
115 Section 5, the results of Section 3 and Section 4 are brought together and the consequences of a
116 regime shift for BSW properties are discussed. Conclusions are presented in Section 6.

117 **2. Data and Methods**

118 *a. Datasets*

119 This study makes use of satellite SST and sea ice concentration data from the OSTIA
120 project spanning January 1985 to December 2016 (Donlon et al. (2012); downloaded from ma-
121 rine.copernicus.eu portal). This dataset is optimally interpolated from multiple satellite sensors
122 together with *in situ* observations onto a 0.05° grid (1.5 x 5.6 km for Barents Sea) at a daily fre-
123 quency. The feature resolution is 10 km and the accuracy of the daily data is ~ 0.57 K (Donlon

124 et al. 2012). At the current spatial and temporal resolution, the satellite SST data used in this
125 study can not yet resolve mesoscale variability (with a characteristic scale of only a few kilome-
126 ters) in the Barents Sea. Sea ice extent in the Barents Sea is computed from the OSTIA sea ice
127 concentration. The sea ice edge is defined as the 15% contour the sea ice concentration.

128 Bathymetry is taken from the GEBCO 2014 30 arcsecond resolution dataset (Weatherall et al.
129 (2015); GEBCO_2014 Grid, version 20150318, www.gebco.net). In the Barents Sea, it corre-
130 sponds to a resolution of 0.2 km in longitude and 0.9 km in latitude. We also use fields of surface
131 air temperature (SAT; corresponding to temperature at 2 m above surface) and sea level pressure
132 (SLP) from the ECMWF ERA-Interim reanalysis (Berrisford et al. (2011); www.ecmwf.int). This
133 dataset is provided on a 0.75° grid (84 x 16 km for Barents Sea) with 3-hourly temporal resolution,
134 averaged into monthly means.

135 Observations from the Fugløy-Bear Island section along 20.0° E in the BSO (red line, Fig-
136 ure 1) are used to characterize the variations of the AW inflow temperature and salinity (Larsen
137 et al. 2016). This dataset is available through the ICES portal (ocean.ices.dk/iroc) and corresponds
138 to hydrographic profiles, collected six times a year, used for the period January 1985 to October
139 2015. The time series presented here is averaged over the 50 - 200 m depth range and between
140 71.5° N and 73.5° N (Ingvaldsen et al. 2003), and is thus representative of the subsurface temper-
141 ature and salinity variability. We also use observations from the Kola section (available through
142 www.pinro.ru), extending from 73.0° N to 74.0° N along 33.5° E (orange line, Figure 1), as a
143 proxy for the AW temperature in the central Barents Sea. Along the section, Conductivity Tem-
144 perature Depth (CTD) profiles have been collected between 7 to 9 times per year and we use the
145 subset from January 1985 to December 2015 (Bochkov 1982; Ingvaldsen et al. 2003). We consider
146 again the depth range between 50 and 200 m (i.e the sub-surface), which is below the depth of the

147 summer mixed layer and is the depth range over which the core of AW enters the Barents Sea
148 (Ingvaldsen 2005).

149 In order to examine the variability and long-term trends over the wider region than just
150 these two sections, temperature and salinity fields from the EN4 dataset are analysed (EN4.2.0,
151 www.metoffice.gov.uk/hadobs/en4). EN4 comprises *in situ* ship CTD profile data and Argo float
152 data optimal-interpolated on a 1° , monthly z-grid with 42 levels (Gouretski and Reseghetti 2010).
153 Data used in this analysis are from the January 1985 to December 2016 period. Between 1985 and
154 2015, there is a minimum of 2 profiles of temperature and salinity per year in the northeastern Bar-
155 ents Sea. It should be noted that there is a summer bias in this dataset based on when most of ship
156 based profiles were collected. To accommodate the variability in profile sampling the uncertainty
157 values provided in EN4 are considered throughout this study (Good et al. 2013).

158 Additional temperature and salinity fields are retrieved from the MIMOC ocean climatology
159 project (www.pmel.noaa.gov/mimoc), which optimally interpolates *in situ* ship CTD profiles and
160 Argo float data onto a 0.5° σ -grid followed by an 81 level z-grid (Schmidtke et al. 2013). The
161 monthly climatology is weighted to be representative of 2007 to 2011. MIMOC data was included
162 in this study as its higher spatial resolution allows a better description of the 3D structure of the
163 front than EN4 data, although it does not provide information on the interannual variations of the
164 fields.

165 *b. Methods*

166 The Barents Sea SST seasonal climatology is calculated from the OSTIA data. To resolve the PF,
167 the magnitude of the gradient in SST for both the latitude and longitude directions are calculated
168 using the equation: $|\nabla T_{(x,y)}| = \sqrt{(\partial T / \partial x)^2 + (\partial T / \partial y)^2}$.

169 In order to characterise the temporal and spatial variability in SST over the Barents Sea, em-
170 pirical orthogonal functions (EOF) are calculated using the singular value decomposition (SVD)
171 method (Thomson and Emery 2014). EOF analysis extracts the main mode of SST interannual
172 variability, providing us with a spatial pattern and an associated time-varying index referred to as
173 the principal component (PC). The area selected for EOF analysis covers the Barents Sea (10° E –
174 65° E and 68° N – 80° N, see green box, Figure 1). Prior to the EOF decomposition several steps
175 were taken. These are: (1) data points within 28 km (5 grid cells) of land were removed as well
176 as the Kara Sea and any isolated inlets with restricted connectivity to the Barents Sea that would
177 be unrepresentative of the variability in the Barents Sea; (2) the seasonal cycle was then removed
178 from the SST monthly means in each grid cell by applying a 12-month running mean to the data;
179 (3) the mean SST at each grid cell was then removed, (4) finally SST in each grid cell was divided
180 by its respective standard deviation. We also compute correlations between the PC and other fields
181 which were also subject to a 12-month running mean.

182 A 2-tailed Welch’s t-test is used to estimate the significance of a difference between two given
183 time periods, while a 2-tailed Student’s t-test is used for the significance of linear trends in monthly
184 SST. For estimating the 95%-level significance of correlations, a 2-tailed Student’s t-test is used
185 and an appropriate reduction in degrees of freedom associated with a 12-month running mean is
186 accounted for.

187 In this study, the criteria used to define the different water masses mostly follows previous
188 definitions found in the literature (e.g. Loeng (1991); Table 1). The main adjustment made to
189 existing definitions is the minimum density set for the BSW definition ($\sigma > 27.85 \text{ kg m}^{-3}$); it
190 ensures that we reject the warm and fresh surface water that is not dense enough to sink into the
191 Arctic Ocean. Note that our results are mostly insensitive to the exact definition of the different
192 water masses. For the EN4 and MIMOC datasets, potential density is determined using TEOS-

193 10 (McDougall et al. 2012). Practical salinity and potential temperature are also estimated and
194 presented throughout to allow direct comparison to results found in the literature.

195 In order to quantify the changes of the BSW properties over time, we estimate the mean BSW
196 temperature and salinity from EN4 data within a domain in the north eastern Barents Sea (Northern
197 Basin, 44° E – 54° E and 76.5° N – 78.5° N, see cyan-dashed box, Figure 1). We only consider
198 the depth range 100 – 300 m, as in this region, BSW is isolated from the atmosphere by the ArW
199 layer, inhibiting further modification before BSW reaches the Arctic Basin. ArW properties are
200 defined in the 0 – 100 m layer within the same region from EN4. Surface BSW properties south
201 of the PF are defined from EN4 in the Central Basin (40° E – 50° E and 74.5° N – 76.5° N, see
202 yellow-dashed box, Figure 1)

203 **3. Seasonal and Interannual Variability of Sea Surface Temperature in the Barents Sea**

204 In this section, we characterize the temporal and spatial variations of SST over the Barents Sea.
205 SST, by which the surface expression of PF is defined in Section 4, is representative of air-sea
206 interactions that are key to the formation of BSW. We first examine the seasonal cycle because this
207 has been suggested, from model analysis, to play an important role in BSW formation (Årthun
208 et al. 2011; Dmitrenko et al. 2014). When averaged over the Barents Sea domain (see green
209 box in Figure 1), the amplitude of the SST seasonal cycle amounts to 1.69° C, with minimum
210 and maximum occurring in April and July, respectively. This value is large when compared to
211 the standard deviation of the mean SST once the seasonal cycle is removed, which amounts to
212 0.41° C. Clearly, SST is dominated by seasonal variability. The annual winter reduction in SST is
213 key to the formation of BSW through heat loss and given this is an annual event suggests a link
214 between BSW and the 1 – 2.5 year residence time of AW within the Barents Sea (Smedsrud et al.
215 2010; Årthun et al. 2011).

216 Maps of seasonal mean SST, over the period 2005 – 2016 are shown in Figure 3a-d. It reveals
217 a pool of warm AW in the southwestern Barents Sea with a tongue of AW in Central Basin.
218 This warm AW tongue is intensified in winter and spring but present throughout the year. In
219 the southwestern Barents Sea, SST increases from 4 °C in spring to 8 °C in summer. In the
220 remainder of the Barents Sea, the SST also increases by 4 °C between spring and summer but
221 approaches -1.8 °C in the spring due to the presence of sea ice (Figure 4a-d). The sea ice edge also
222 shows strong seasonality, retreating to the northern margins of the Barents Sea in summer, while
223 advancing towards Central Bank from the north and the south-east in winter. As discussed later in
224 this section, the long term trend in SST changes in 2005, posing the question of a possible change
225 of SST seasonal cycle across the full period considered. The most striking difference between
226 the 1985 – 2005 (Figure 3) and the 2005 – 2016 (Figure 4) time periods is the location of the
227 sea ice edge, with appreciably larger areas of open water post-2005 in all the seasons. This is
228 accompanied by changes in SST where the seasonal sea ice has retreated.

229 This seasonality is primarily driven by the seasonal cycle of the net surface heat flux with a
230 contribution from AW heat transport (Ding et al. 2016; Smedsrud et al. 2010). In the northern
231 Barents Sea, seasonal surface heat fluxes roughly balance over a year. In contrast there is a net
232 heat flux from ocean to atmosphere in the southern Barents Sea, suggesting the importance of heat
233 brought here by AW for the formation of BSW (Smedsrud et al. 2010).

234 To examine SST variability on interannual and longer time-scales, the seasonal cycle is first
235 removed and EOF analysis is performed (see Section 2 for methodology). The trend is not removed
236 as this could be related to multidecadal variability discussed later in this section. The first mode
237 (EOF 1) of variability in SST explains 72.9% of the variance. As the second mode explains less
238 than 10% of the variance, we only discuss EOF 1. The spatial pattern of EOF 1 is a positive
239 anomaly across the full Barents Sea (Figure 5a). PC 1 has a periodicity of 6 to 10 years but

240 also exhibits multidecadal variability (Figure 5c). PC 1 is strongly correlated with the interannual
241 variations of SAT over the Barents Sea where SAT leads by 2 months (Figure 5b). Regressing
242 PC 1 on the SAT fields reveals an area of significant positive correlation over the Arctic Ocean,
243 eastern Arctic shelf seas and northern Russia. Lag correlations with AW temperature show AW
244 leads SST by 2 to 4 months. PC 1 is significantly correlated with the variation of AW temperature
245 at the Kola section ($r = 0.89$, lag = -2 months, Figure 5d) and the Fugløy-Bear Island section
246 ($r = 0.80$, lag = -4 months, Figure 5d). PC 1 is also anti-correlated with the variations of the sea
247 ice extent in the Barents Sea ($r = -0.93$, lag = 1 month, Figure 5e).

248 These correlations suggest that, when mode 1 is in positive phase, SST is warm in the Barents
249 Sea, the sub-surface AW temperature is warmer than average, sea ice extent is low and SAT is
250 warmer than average. A mechanism that could explain this mode is an increase in the temperature
251 of the AW inflow to the Barents Sea, which would in turn reduce sea ice extent in the Barents Sea,
252 both acting to increase AW heat loss to the atmosphere (Smedsrud et al. 2010) and resulting in
253 warmer SAT. During a positive phase of this mode, both the increase of oceanic heat lost and the
254 decrease of the sea ice extent will most likely affect the formation of BSW, as discussed in more
255 detail in Section 5.

256 We could not find a significant correlation between PC 1 and SLP variations across the Barents
257 Sea. This is at odds with the results of Herbaut et al. (2015), which suggested a link between
258 the variations sea ice (and thus SST) and SLP. The different results could be due to the different
259 periods considered as they only considered the variations up to 2004.

260 In summary, our lagged correlation analysis is consistent with heat carried in the AW inflow
261 gradually influencing both SAT and BSW SST as it propagates from Fugloya-Bear Island section
262 to the Kola section and onwards to the interior Barents Sea where SAT can feed heat back to SST.
263 Our results suggest AW inflow temperature may be at least as important as SAT in setting the

264 Barents Sea SST. Indeed, this BSW-SST-forcing mechanism is supported by the conclusions of
265 Smedsrud et al. (2010) who found that AW heat input has a bigger impact on SST variability than
266 SAT forcing. The mechanism proposed here is also consistent with the results of Schlichtholz
267 and Houssais (2011) who found that the temperature of recirculating AW exiting the Barents Sea
268 through the BSO was driven by SAT within the Barents Sea.

269 We now examine the SST multidecadal variability. We find a significant positive linear trend
270 of up to $0.05\text{ }^{\circ}\text{C year}^{-1}$ in the western Barents Sea for the period from 1985 to 2004 (pre-2005;
271 Figure 2b). Post-2005 (2005 to 2016) however, the SST in the western Barents Sea stabilises, such
272 that the trend becomes insignificant here while a positive trend of roughly $0.10\text{ }^{\circ}\text{C year}^{-1}$ arises in
273 most of the eastern Barents Sea (Figure 2c). A positive trend is also found in the analysis of Singh
274 et al. (2013) for the time period 2002 to 2010. The shift in SST trend since 2005 is consistent
275 with the results of Herbaut et al. (2015), who found a significant reduction of both the mean and
276 variance of the winter sea ice concentration after 2005. The positive trend in the eastern Barents
277 Sea coincides with an increase in AW temperatures observed at the Kola section (Figure 5e). As
278 AW temperature at the Kola section is correlated with PC 1, this suggests mode 1 also captured
279 part of the variability at multidecadal or longer time scales. As suggested by Smedsrud et al.
280 (2010), an increase in AW heat transport would manifest in an expansion of a warm heat anomaly
281 in the Barents Sea basin resulting in an increase in the surface area in which heat loss takes place.
282 The change in trend across the eastern Barents Sea could represent the expansion of this surface
283 area.

284 Although the SST dataset is limited to 1985 onwards, there are other datasets which have been
285 used to address longer term variability in the Barents Sea. A 16–20 year and 30–50 year timescale
286 fluctuation was found in ~ 100 year observational datasets of both sea ice concentration and SLP
287 (Venegas and Mysak 2000). These timescales are too long to be fully resolved in our analysis

288 period, so we can not fully distinguish between long term trend and natural variations occurring
289 on these timescales. Yet, the results of Venegas and Mysak (2000) suggest that the sea ice extent
290 variations on the 16–20 year timescale are likely linked with SLP anomalies. Our time period of
291 32 years should capture some variability at the 16–20 year time period which could be manifested
292 as the change in temperature occurring in 2005. However, as our EOF 1 is not driven by SLP
293 variations, we hypothesis that the change occurring in 2005 is likely the manifestation of a regime
294 shift rather than natural variability causing SLP to become decoupled across this time period. This
295 hypothesis is also supported by the analysis of the observed sea ice extent from 1850 onward by
296 Onarheim and Årthun (2017), who found that the winter sea ice extent is at its lowest level since
297 1990. This is discussed in relation to long-term trends in Section 5.

298 **4. The Polar Front's Constraint on the Sea Ice Edge**

299 The magnitude of the 2D gradient in SST shows the surface manifestation of fronts in the Barents
300 Sea (Figure 3e-h). Starting in the west, a front follows Spitsbergen Bank but then bifurcates at
301 Central Bank and splits into two branches (Figure 3e), in agreement with the results of Oziel et al.
302 (2016). The southern branch of this front (referred to hereafter as the Barents Sea Front) follows
303 the western side of Central Bank southward, dividing the Barents Sea into an AW-influenced
304 western region and a BSW-influenced eastern region. The Barents Sea Front is most prominent
305 during winter and spring (Figure 3e,h) and has been discussed in greater detail by Oziel et al.
306 (2016, 2017).

307 Further to the north, the PF divides the eastern Barents Sea into an ArW-influenced northern
308 region and a BSW-influenced southern region. Our results show the PF to be a persistent feature
309 following the ~ 220 m isobath throughout the year, although Oziel et al. (2016) found that the
310 PF was positioned further north than the present analysis with no fixed position. Their analysis

311 was limited by the dataset used, comprising temperature and salinity *in situ* profiles collected in
312 the Barents Sea, which captures only the sub-surface expression of the front in the 50 to 100 m
313 depth range. SST observations reveal that the PF pathway on the east side of the Barents Sea
314 follows the southern sides of Great Bank and Ludlov Saddle eastward to Novaya Zemlya Bank
315 (Figure 3e-h). At Novaya Zemlya Bank, the PF extends northward along Novaya Zemlya Bank to
316 78° N. It should be noted that a second, weaker thermal-surface front exists in the SST data due to
317 the transition from freezing ice-covered water to warmer ice-free water. The thermal-surface front
318 does move with the sea ice edge and sometimes coincides with the more permanent PF.

319 Previous studies have investigated several aspects of the PF (Våge et al. 2014; Oziel et al. 2016)
320 but the dynamics controlling it are still poorly pinned down. Here we present some evidences that
321 the PF is controlled by potential vorticity constraints. Within the Barents Sea, the PF is closely tied
322 to the 220 m isobath (Figures 3 and 4), which is located on a steep slope separating the northern and
323 southern Barents Sea (Figure 1). Potential vorticity constraints usually force currents to flow along
324 topographic contours rather than across them (Taylor 1917; Proudman 1916). Planetary potential
325 vorticity (q) can be estimated by the equation $q = f/h$, where f is the coriolis parameter and h is the
326 depth. The planetary potential vorticity contours in the Barents Sea follow closely the bathymetry
327 contours as f is roughly constant in the region. In the case of a basin with a shallower northern
328 outflow depth than inflow i.e. a ridge, an idealised model with potential vorticity constraints drives
329 anticyclonic/clockwise circulation around the basin and eastward along the ridge in the northern
330 hemisphere (Yang and Price 2000). This is consistent with the path of the PF we resolved by
331 the OSTIA SST (Figure 3), as well as the eastward flow found in velocity observations on the
332 southwestern slope of Great Bank (Våge et al. 2014) and simulations showing eastward flow along
333 the southern slope of Great Bank (Slagstad and McClimans 2005; Lind and Ingvaldsen 2012).

334 Following Pratt (2004), additional evidence that the PF is constrained by potential vorticity can
335 be provided by estimating the Froude number associated with the flow across the ridge towards
336 the eastern boundary (i.e Novaya Zemlya Bank in our case). The Froude number is given by
337 $F = u/(g'd)^{1/2}$, where u is current speed, g' is reduced gravity and d is depth of the layer at
338 the ridge. Here we take $u = 0.2 \text{ m s}^{-1}$ (based on observations by Våge et al. (2014), assuming
339 current speed is constant along the ridge), and values for g' and d are calculated from MIMOC
340 data (Figure 7), obtaining a Froude number of 0.3. Following the argument developed by Pratt
341 (2004) and given that in our case the height of the ridge occupies roughly $1/3$ of the water column,
342 a Froude number greater than 0.2 suggests that the Great Bank–Ludlov Saddle ridge imposes a
343 hydraulic control on the flow associated with the PF, providing further evidence that the PF is
344 constrained by potential vorticity.

345 We next examine time variations of the PF, in relation to the position of the sea ice edge over
346 time. According to Smedsrud et al. (2010), the PF sets the limit on surface area available for winter
347 heat loss over the Barents Sea. Logically, the PF may also play a role in determining the volume of
348 summer freshwater input from sea ice melt water. Thus the interplay between the eastern Barents
349 Sea PF and mobile sea ice edge mediates the properties of BSW that will be carried into the Arctic
350 as AIW. A comparison of SST gradients and sea ice concentration shows that the sea ice edge
351 follows the PF in both the eastern and western Barents Sea during winter and spring from 2005
352 to 2016 (Figure 3a-d) but this was not the case before 2005 (Figure 4). Steele and Ermold (2015)
353 suggest that during the expansion and retreat of seasonal sea ice, the edge loiters at fronts where
354 there is a gradient in temperature inhibiting further expansion. This then implies that the expansion
355 of sea ice south of the PF before 2005 could be consistent with cooler SST or stronger northerly
356 winds enabling greater transport of the mobile sea ice pack across the PF enabling it to loiter closer
357 to the Barents Sea Front.

358 We then focus on the interannual variability of the PF and its relationship with the sea ice edge
359 (Figures 6). To perform this analysis, the SST gradient is calculated meridionally and these gra-
360 dients are averaged zonally within the box shown as a blue-dashed line on Figure 1. Zonally-
361 averaged SST gradients on a given day are normalized by the daily standard deviation of the
362 gradient in the same analysis box (Figure 1), in order to remove to potential large effect of the
363 strong seasonality and interannual variability in the intensity of the SST gradient and sea ice ex-
364 tent. Figures 6(a) shows that the PF is persistent in its location throughout the majority of the year.
365 Between 1985 and 2004, the PF was covered by sea ice for parts of winter and spring but held
366 position at 76.5° N, rather than moving south with the advancing winter sea ice edge as previously
367 thought (Smedsrud et al. 2010). As expected, there is also a thermal-surface front at the position
368 of the sea ice edge to the north of the PF in summer, but the PF is always present as a stronger and
369 more persistent front at 76.5° N along the 220 m bathymetry contour.

370 A change in the location of the winter sea ice edge relative to the position of the PF is also
371 evident on decadal timescales (Figure 6). Unlike in the pre-2005 period, since 2005, the winter
372 sea ice edge has been unable to sustain a southwards breach of the PF for more than a few days
373 (Figure 6b). We define a region in the Barents Sea between the PF to the north and the Coastal
374 Water front to the south shown by the dark-blue box in Figure 1, within which sea ice melt can
375 be entrained into BSW. The change in 2005 has reduced the mean seasonal change in sea ice
376 area in this region, from $77\,000\text{ km}^2$ between 1985 and 2004 to $8\,700\text{ km}^2$ between 2005 and
377 2016. This provides useful information in efforts to predict the location of the winter sea ice in
378 the Barents Sea (examples of predictions include Onarheim and Årthun (2017); Sigmond et al.
379 (2016); Nakanowatari et al. (2014)). This is important because changes in sea ice conditions in
380 the Barents Sea have been linked to widespread, anomalous atmospheric conditions over northern
381 continents (Petoukhov and Semenov 2010; Yang et al. 2016).

382 At the same time, while remaining fixed to topography, the mean SST gradient across the PF
383 increases significantly from $0.011 \pm 10^{-4} \text{ }^\circ\text{C km}^{-1}$ pre-2005 to $0.015 \pm 10^{-4} \text{ }^\circ\text{C km}^{-1}$ in post-
384 2005 (Figure 6c). This steepening in the PF SST gradient coincides with a significant increase in
385 AW temperature at the Kola section from $3.1 \pm 0.05 \text{ }^\circ\text{C}$ in the pre-2005 period to $4.0 \pm 0.05 \text{ }^\circ\text{C}$ in
386 the post-2005 period. Given that the SST north of the PF is changing at a slower rate than south of
387 the PF (Figure 2), the intensification of the PF can then be mainly attributed to the increase in AW
388 temperature in the Barents Sea. One important consequence of the increase in AW temperature is
389 that the heat content on the southern side of the front prevents sea ice from accumulating. A link
390 between changes in sea ice and AW temperature has been discussed by Smedsrud et al. (2013)
391 but not in relation to the PF. We assess this result in relation to trend and long-term variability in
392 Section 5.

393 In addition to the changes found in the southern side of the front, discussed above, changes in
394 the properties of the ArW north of the PF could also occur. To the northeast of Svalbard where the
395 AW lies close to the surface, Ivanov et al. (2016) have suggested that a positive feedback could
396 exist between entrainment of warm AW and reduced midwinter sea ice thickness, due to a decrease
397 of the stratification driven by change in salinity. The mean ArW properties from EN4 pre-2005
398 were $T = -1.15 \pm 0.04 \text{ }^\circ\text{C}$, $S = 34.463 \pm 0.014$; while post-2005 they were $T = -0.76 \pm 0.06 \text{ }^\circ\text{C}$,
399 $S = 34.569 \pm 0.022$ (Figure 1 shows the cyan-dashed box selected for ArW properties north of the
400 PF). This significant increase in temperature and salinity could be caused by a similar process to
401 the one described by Ivanov et al. (2016).

402 The mean surface BSW properties pre-2005 were $T = -0.22 \pm 0.03 \text{ }^\circ\text{C}$, $S = 34.828 \pm 0.009$;
403 while post-2005 they significantly increased to $T = 0.50 \pm 0.05 \text{ }^\circ\text{C}$ $S = 34.943 \pm 0.013$ (Figure 1
404 shows the yellow-dashed box selected for surface BSW properties south of the PF). The salinity
405 increase is comparable for the surface BSW and ArW within the error bounds, but the increase in

406 the temperature of surface BSW is almost double the change in ArW temperature over the same
407 period. The result on ArW density and surface BSW density is an increase of $0.071 \pm 0.017 \text{ kg m}^{-3}$
408 and $0.054 \pm 0.009 \text{ kg m}^{-3}$, respectively, indicating a decrease in the density gradient across the
409 PF after 2005. This suggests that the steepening of the temperature gradient and weakening of the
410 density gradient across the PF in the eastern Barents Sea are primarily driven by changes occurring
411 in the southern side of the PF.

412 Transect data through the eastern Barents Sea (Figure 7) show the SST gradient across the PF is
413 the surface expression of a vertically-coherent front. In both the EN4 and MIMOC climatologies,
414 the PF is present near 76.5° N as a negative south-north temperature gradient over the depth range
415 $0 - 100 \text{ m}$, and a similar sub-surface salinity gradient. The PF is a transition between the southern
416 region that is temperature-stratified (α -ocean) and the northern region that is salinity-stratified (β -
417 ocean) (Carmack 2007). Here, α is the coefficient of thermal expansion and β is the coefficient
418 of haline contraction. This makes the PF an important transition zone where the contribution to
419 density from temperature and salinity can be in balance. Note the presence of water that is too fresh
420 to fit the BSW definition and too warm to fit the ArW definition between 77° N and 78° N over 0 -
421 50 m (Figure 7b,d). This water mass sits on the mixing line between BSW and ArW (Figure 7h),
422 suggesting that mixing between BSW and ArW occurs at the front. Previous studies based on
423 observations in the western Barents Sea have revealed the presence of interleaving between BSW
424 and ArW along the PF that could enhance mixing (Parsons et al. 1996; Våge et al. 2014; Fer and
425 Drinkwater 2014).

426 On the northern side of the transect, the ArW layer (Table 1) is present in the MIMOC data over
427 the depth range $0 - 100 \text{ m}$ at 80° N and extends down to 50 m at 77° N . In EN4, the ArW layer
428 extends to a deeper depth of 150 m at 80° N and 100 m at 77° N . The main difference between
429 the EN4 climatology and the MIMOC climatology is the 1° C cooler temperature of BSW in

430 the EN4 than in MIMOC (Figure 7c,d). This may represent a change in BSW temperature over
431 time given that the MIMOC climatology is weighted to be characteristic of 2007 – 2011 whereas
432 the EN4 climatology is an average over the period 1985 – 2016. Regardless of the difference
433 in temperature between the two datasets, BSW occupies roughly the same area (black dots in
434 Figure 7e,f). As BSW is denser than ArW (Figure 7g,h), it sits below ArW north of the PF at
435 76.5° N. From Ludlov Saddle, BSW flows eastward and exits the Barents Sea through St. Anna
436 Trough in a layer below ArW (Schauer et al. 2002). As the Central Basin is the source of BSW
437 (Oziel et al. 2016), this suggests BSW propagates northwards of the PF either by subducting below
438 ArW or by undergoing modification at the surface due to fast-mixing processes in the the upper
439 portion of BSW that Rudels et al. (1996) has hypothesized occurs during winter heat loss.

440 **5. Atlantification of the Barents Sea and implications for Barents Sea Water**

441 As a consequence of the intensification of the PF since 2005, it now forms a persistent barrier
442 to the formation and export of sea ice south of the PF (Figure 6). Having identified the forcing on
443 BSW in Section 3, here we discuss the possible implications of the barrier imposed by the PF on
444 the properties of the BSW exiting the Barents Sea:

- 445 1. The northern limit of the surface area available for AW winter heat loss has become fixed to
446 the location of the PF. The sub-surface EN4-averaged BSW temperature has warmed from
447 -0.51 ± 0.03 °C to -0.13 ± 0.03 °C when comparing the pre-2005 and post-2005 periods (Fig-
448 ure 6c, averaged over 100 – 300 m in the cyan-dashed box in Figure 1). The increase of the
449 temperature at the Kola section between the same two periods is more than twice as large
450 (0.9 °C). The observed reduction in Barents Sea ice extent has resulted in an increase of
451 the surface heat flux from the ocean to the atmosphere (Long and Perrie 2017; Årthun et al.

452 2012), likely explaining the different rate of temperature increase between the BSW and the
453 Kola section.

454 Before 2005, the expansion and retreat of sea ice in the eastern Barents Sea buffered BSW
455 properties against changes in AW temperatures (Smedsrud et al. 2010), but our analysis sug-
456 gests that this buffering capacity has reduced since 2005, enabling the temperature increase
457 of BSW in recent years visible on Figure 6d. Such a temperature change requires that most of
458 the AW heat is lost to the atmosphere in the ice-free southern Barents Sea (which is consistent
459 with the results of Smedsrud et al. (2010)) and that the heat lost by BSW through mixing with
460 ArW north of the PF is small. While Lind et al. (2016) have pointed out that mixing between
461 ArW and BSW can exist, in particular during years with lower sea ice cover, the heat lost
462 through that process is most likely much smaller than the heat lost to the atmosphere south of
463 the PF.

- 464 2. The reduction of sea ice south of the PF reduces the seasonal freshwater input to BSW asso-
465 ciated with local sea ice melt. Based on their model simulations, Ellingsen et al. (2009) found
466 that between 1979 and 2007, melt water from imported sea ice contributed 0.02 Sv of fresh-
467 water on average. This is enough to decrease the mean salinity of their simulated 1.1 Sv AW
468 inflow (salinity 35.1) to salinity of 34.4. However, in their study, Ellingsen et al. (2009) does
469 not account for the PF's role in partitioning sea ice meltwater between BSW and ArW, and
470 considers that the input of sea ice meltwater takes place entirely south of the PF, and thus can
471 convert AW into ArW. Here we revisit their calculation, taking into account the partitioning
472 of meltwater at the PF.

473 To calculate the meltwater input south of the PF before 2005, we assume that the sea ice found
474 south of the front was 1 m thick, which is a typical thickness for first-year ice in the Barents

475 Sea (Ellingsen et al. 2009; Smedsrud et al. 2010). In contrast to Ellingsen et al. (2009), we
476 only consider the box that contains the area south of the PF and north of the Coastal Water
477 front shown by the dark-blue box on Figure 1, and assume that the AW is not modified before
478 it enters that box. Within this box, the reduction in sea ice area south of the PF by 68 300 km²
479 between the pre and post-2005 periods (Figure 6) corresponds to a 0.0022 Sv reduction in
480 the freshwater input south of the PF after 2005 when the sea-ice is no longer present. This is
481 assumed to mix ubiquitously into BSW.

482 To calculate the dilution of AW by sea ice melt, we estimate the volume to be diluted by
483 comparing the AW inflow to the BSW outflow. Following Gammelsrød et al. (2009), we
484 assume a BSW transport leaving the Barents Sea between Novaya Zemlya and Franz-Josef
485 Land of 1.25 Sv (observed transport scaled up by the difference between virtual current meters
486 and modeled, whole-section transport). For comparison the net annual observed AW inflow
487 through the BSO is 1.1 – 1.2 Sv (Skagseth 2008; Ingvaldsen et al. 2004) (excluding transport
488 associated with Norwegian Coastal Current). This implies that there is no net storage of
489 BSW in the Barents Sea, such that the volume of AW to be diluted is $V_{AW} = 1.1$ Sv (note, a
490 change of AW volume transport across our time period cannot be estimated from the available
491 observations).

492 The salinity of inflowing AW should also be taken into account when calculating a change
493 in BSW salinity. As shown on Figure 5d, the mean properties of AW at the Fugløya–Bear
494 Island section for 1985 to 2005 were $T = 5.44 \pm 0.06$ °C, $S = 35.067 \pm 0.003$ and for 2005
495 to 2016 they were $T = 6.08 \pm 0.07$ °C, $S = 35.120 \pm 0.005$ (the changes between the two
496 periods are significant). Using these different salinity values and considering that the input
497 of freshwater south of the PF vanishes after 2005, we perform a simple dilution calculation,

498 following the equation: $C = [M_{AW} + M_{FW}] / [V_{AW} + V_{FW}]$, where C is the concentration of
499 salt, M is the mass of salt and V is the volume, AW is Atlantic Water and FW is fresh sea
500 ice meltwater. We also assume a constant salinity value of 3 for first-year sea ice (Ellingsen
501 et al. 2009), and constant net AW inflow (BSW outflow) of 1.1 Sv (1.25 Sv) (Skagseth 2008;
502 Gammelsrød et al. 2009). Based on this framework, pre-2005 the mean input of 0.0022 Sv
503 of freshwater results in a reduction of -0.063 (-0.056) of the BSW salinity, while post-2005,
504 the BSW salinity would equal the AW salinity which additionally increased by 0.053 across
505 this time period. Our dilution calculation predicts a change of BSW salinity by ~ 0.11 , which
506 is in broad agreement with the significant increase of BSW salinity estimated from the EN4
507 dataset (from 34.844 ± 0.003 to 34.900 ± 0.002 , Figure 6c). This suggests that the increase in
508 BSW salinity is likely a combination of the change in sea ice area and the change in inflowing
509 AW salinity.

510 When comparing the mean BSW temperature over the two periods in EN4, it increases by
511 $0.38 \text{ }^\circ\text{C}$, which is about a half of the $0.8 \text{ }^\circ\text{C}$ required to compensate density changes arising
512 from the 0.056 mean salinity increase observed. These changes in temperature and salinity
513 have led to a significant increase of BSW density from $1029.092 \pm 0.002 \text{ kg m}^{-3}$ pre-2005 to
514 $1029.116 \pm 0.002 \text{ kg m}^{-3}$ post-2005.

515 The 0.024 kg m^{-3} increase in BSW density between the two periods has to be compared against
516 the gain in density resulting from the transformation of AW to BSW. Pre-2005, the density trans-
517 formation amounted to $\sim 0.33 \text{ kg m}^{-3}$, a combination of $5.9 \text{ }^\circ\text{C}$ decrease and 0.23 salinity decrease
518 (based on AW properties at the BSO). This means a further 8% density change in BSW relative to
519 the pre-2005 era.

520 Our comparison of the two periods (pre and post 2005) suggests that a regime shift occurred
521 in 2005. Yet, one needs to remember that there is well-known multidecadal variability affecting
522 SLP, sea ice concentration, SAT and AW temperature (Venegas and Mysak 2000; Smedsrud et al.
523 2013; Levitus et al. 2009; Ingvaldsen et al. 2003). Variability at a 30–50 year frequency is thought
524 to be driven by the Atlantic Multidecadal Oscillation, suggesting that long-term variations in the
525 Barents Sea are driven by large-scale fluctuations (Levitus et al. 2009). These variations are also
526 affecting the formation, properties and volume of BSW on similar timescales (Årthun et al. 2011).
527 Analysis by Onarheim and Årthun (2017) of an observed time-series of winter sea ice extent from
528 1850 to 2017 in the Barents Sea complemented by analysis of climate simulations also emphasises
529 the existence of variations with a 50 year periodicity. However, their results show winter sea ice
530 extent in the Barents Sea has been lower since the 1990 than in the the rest of the time period and
531 that there is an unprecedented negative trend in the last 30 years that has less than 5% probability
532 of occurring in all preindustrial simulations. This suggests that winter sea ice in the Barents Sea
533 has most likely not been inhibited by the PF during 1850 to 2005. Further evidence comes from
534 the observations by Smedsrud et al. (2013), suggesting that Arctic SAT and AW temperature at the
535 Kola section were both greater after ~ 2000 than at any time from the last century.

536 **6. Conclusion**

537 The goal of this study was to investigate how changes and feedbacks between sea ice and the PF
538 in the Barents Sea may have affected BSW properties over the past decades. We have identified
539 and located the PF in the eastern Barents Sea using satellite SST observations, a feature that has
540 been obscured by seasonal sea ice between 1985 and 2004. While a summer mixed layer and
541 seasonal front does form in association with the melt of seasonal sea ice, as is the case in other
542 regions (Dewey et al. 2017), the PF persists throughout the year as a front with steeper gradients

543 in salinity and temperature in the eastern Barents Sea at 76.5° N, running parallel to the 220 m
544 isobath (Figure 3). The PF is a potential vorticity-constrained, shelf slope current at the steep
545 ridge formed by Great Bank and Ludlov Saddle. Since 2005, the sea ice is inhibited in its winter
546 southward extent by the increase in temperature gradients across the PF, a change most likely
547 driven by an increase in AW temperature.

548 Our results provide new evidence that, in addition to the natural multidecadal variability, the
549 Barents Sea is currently undergoing atlantification, with the corresponding temperature and salin-
550 ity increases catalysed by the observed PF constraint on the sea ice edge. The loss of winter sea
551 ice south of the front represents a loss of freshwater input to BSW, a water mass which makes
552 up 50 – 80% of AIW. As the stationary PF, rather than the mobile sea ice edge, has become the
553 limiting factor controlling the northern boundary of the surface area available for AW cooling in
554 winter, the buffering effect to BSW temperature from the variations of sea ice conditions has de-
555 creased. Observations show a change in BSW properties over the same time period resulting in
556 denser BSW, which could in turn result in a deeper settling depth of BSW once exported to the
557 Arctic Basin through St. Anna Trough (Dmitrenko et al. 2015), with potential far-reaching impacts
558 for the dense water outflow through Fram Strait (Lique et al. 2010; Moat et al. 2014) or the density
559 of the Denmark Strait overflow (Karcher et al. 2011), both of which are important for the deeper
560 branch of the AMOC.

561 *Acknowledgments.* This project was funded through the joint UK-France PhD program by
562 DGA/Dstl, and overseen by Carole Nahum and Timothy Clarke. Data from the Fugløyå–Bear
563 Island section are provided courtesy of Institute of Marine Research, Norway.

564 **References**

- 565 Aagaard, K., J. H. Swift, and E. C. Carmack, 1985: Thermohaline circulation in the Arc-
566 tic Mediterranean Seas. *Journal of Geophysical Research*, **90 (C3)**, 4833, doi:10.1029/
567 JC090iC03p04833, URL <http://doi.wiley.com/10.1029/JC090iC03p04833>.
- 568 Aagaard, K., and R. A. Woodgate, 2001: Some thoughts on the freezing and melting of sea ice
569 and their effects on the ocean. *Ocean Modelling*, **3 (1-2)**, 127–135, doi:10.1016/S1463-5003(01)
570 00005-1.
- 571 Årthun, M., T. Eldevik, L. H. Smedsrud, Ø. Skagseth, and R. B. Ingvaldsen, 2012: Quantifying the
572 influence of atlantic heat on barents sea ice variability and retreat. *Journal of Climate*, **25 (13)**,
573 4736–4743, doi:10.1175/JCLI-D-11-00466.1.
- 574 Årthun, M., R. B. Ingvaldsen, L. H. Smedsrud, and C. Schrum, 2011: Dense water formation
575 and circulation in the Barents Sea. *Deep-Sea Research Part I: Oceanographic Research Papers*,
576 **58 (8)**, 801–817, doi:10.1016/j.dsr.2011.06.001.
- 577 Berrisford, P., D. Dee, K. Fielding, M. Fuentes, P. Kallberg, S. Kobayashi, and S. Uppala, 2011:
578 The ERA-Interim Archive: Version 2.0. *ERA report series*, **2 (1)**, 1–16.
- 579 Bochkov, Y. A., 1982: Water temperature in the 0-200 m layer in the Kola-Meridian section in the
580 Barents Sea, 1900-1981. *Trudy PINRO, Murmansk*, **46**, 113–122.
- 581 Carmack, E. C., 2007: The alpha/beta ocean distinction: A perspective on freshwater fluxes,
582 convection, nutrients and productivity in high-latitude seas. *Deep-Sea Research Part II: Topical
583 Studies in Oceanography*, **54 (23-26)**, 2578–2598, doi:10.1016/j.dsr2.2007.08.018.

- 584 Day, J. J., S. Tietsche, and E. Hawkins, 2014: Pan-arctic and regional sea ice predictabil-
585 ity: Initialization month dependence. *Journal of Climate*, **27** (12), 4371–4390, doi:10.1175/
586 JCLI-D-13-00614.1.
- 587 Dewey, S. R., J. H. Morison, and J. Zhang, 2017: An Edge-Referenced Surface Fresh Layer in the
588 Beaufort Sea Seasonal Ice Zone. *Journal of Physical Oceanography*, **47** (5), 1125–1144, doi:
589 10.1175/JPO-D-16-0158.1, URL <http://journals.ametsoc.org/doi/10.1175/JPO-D-16-0158.1>.
- 590 Ding, Y., J. A. Carton, G. A. Chepurin, M. Steele, and S. Hakkinen, 2016: Seasonal heat and fresh-
591 water cycles in the Arctic Ocean in CMIP5 coupled models. *Journal of Geophysical Research :
592 Oceans*, **121**, 2043–2057, doi:10.1002/2015JC011534.Received.
- 593 Dmitrenko, I. A., and Coauthors, 2014: Heat loss from the Atlantic water layer in the northern Kara
594 Sea: Causes and consequences. *Ocean Science*, **10** (4), 719–730, doi:10.5194/os-10-719-2014.
- 595 Dmitrenko, I. A., and Coauthors, 2015: Atlantic water flow into the Arctic Ocean through the St.
596 Anna Trough in the northern Kara Sea. *Journal of Geophysical Research: Oceans*, **120**, 5158–
597 5178, doi:10.1002/2015JC010969.
- 598 Donlon, C. J., M. Martin, J. Stark, J. Roberts-Jones, E. Fiedler, and W. Wimmer, 2012: The
599 Operational Sea Surface Temperature and Sea Ice Analysis (OSTIA) system. *Remote Sensing
600 of Environment*, **116**, 140–158, doi:10.1016/j.rse.2010.10.017, URL [http://dx.doi.org/10.1016/
601 j.rse.2010.10.017](http://dx.doi.org/10.1016/j.rse.2010.10.017).
- 602 Ellingsen, I., D. Slagstad, and A. Sundfjord, 2009: Modification of water masses in the Barents
603 Sea and its coupling to ice dynamics: A model study. *Ocean Dynamics*, **59** (6), 1095–1108,
604 doi:10.1007/s10236-009-0230-5.

605 Fahrbach, E., J. Meincke, S. Österhus, G. Rohardt, U. Schauer, V. Tverberg, J. Verduin, and R. a.
606 Woodgate, 2001: Direct measurements of heat and mass transports through the Fram Strait.
607 *Polar Research*, **20** (2), 217–224, doi:10.1111/j.1751-8369.2001.tb00059.x.

608 Fer, I., and K. Drinkwater, 2014: Mixing in the Barents Sea Polar Front near Hopen in spring.
609 *Journal of Marine Systems*, **130**, 206–218, doi:10.1016/j.jmarsys.2012.01.005, URL [http://dx.](http://dx.doi.org/10.1016/j.jmarsys.2012.01.005)
610 [doi.org/10.1016/j.jmarsys.2012.01.005](http://dx.doi.org/10.1016/j.jmarsys.2012.01.005).

611 Gammelsrød, T., Ø. Leikvin, V. Lien, W. P. Budgell, H. Loeng, and W. Maslowski, 2009: Mass and
612 heat transports in the NE Barents Sea: Observations and models. *Journal of Marine Systems*,
613 **75** (1-2), 56–69, doi:10.1016/j.jmarsys.2008.07.010, URL [http://dx.doi.org/10.1016/j.jmarsys.](http://dx.doi.org/10.1016/j.jmarsys.2008.07.010)
614 [2008.07.010](http://dx.doi.org/10.1016/j.jmarsys.2008.07.010).

615 Gawarkiewicz, G., and A. J. Plueddemann, 1995: Topographic control of thermohaline frontal
616 structure in the Barents Sea Polar Front on the south flank of Spitsbergen Bank. *Journal of*
617 *Geophysical Research*, **100** (C3), 4509–4524.

618 Good, S. A., M. J. Martin, and N. A. Rayner, 2013: EN4: Quality controlled ocean tempera-
619 ture and salinity profiles and monthly objective analyses with uncertainty estimates. *Journal of*
620 *Geophysical Research: Oceans*, **118** (12), 6704–6716, doi:10.1002/2013JC009067.

621 Gouretski, V., and F. Reseghetti, 2010: On depth and temperature biases in bathythermograph
622 data: Development of a new correction scheme based on analysis of a global ocean database.
623 *Deep-Sea Research Part I: Oceanographic Research Papers*, **57** (6), 812–833, doi:10.1016/j.
624 [dsr.2010.03.011](http://dx.doi.org/10.1016/j.dsr.2010.03.011), URL <http://dx.doi.org/10.1016/j.dsr.2010.03.011>.

625 Harris, C. L., A. J. Plueddemann, and G. G. Gawarkiewicz, 1998: Water mass distribution and
626 polar front structure in the western Barents Sea. *Journal of Geophysical Research*, **103** (C2),

627 2905–2917.

628 Herbaut, C., M.-N. Houssais, S. Close, and A.-C. Blaizot, 2015: Two wind-driven modes of win-
629 ter sea ice variability in the Barents Sea. *Deep-Sea Research Part I: Oceanographic Research*
630 *Papers*, **106**, 97–115, doi:10.1016/j.dsr.2015.10.005, URL [http://dx.doi.org/10.1016/j.dsr.2015.](http://dx.doi.org/10.1016/j.dsr.2015.10.005)
631 10.005.

632 Ingvaldsen, R. B., 2005: Width of the North Cape Current and location of the Polar Front in the
633 western Barents Sea. *Geophysical Research Letters*, **32 (16)**, 1–4, doi:10.1029/2005GL023440.

634 Ingvaldsen, R. B., L. Asplin, and H. Loeng, 2004: The seasonal cycle in the Atlantic transport to
635 the Barents Sea during the years 1997–2001. *Continental Shelf Research*, **24 (9)**, 1015–1032,
636 doi:10.1016/j.csr.2004.02.011.

637 Ingvaldsen, R. B., H. Loeng, O. Geir, and A. Bjorn, 2003: Climate variability in the Barents Sea
638 during the 20th century with a focus on the 1990s. *ICES Journal of Marine Science*, 160–
639 168, URL [http://www.ices.dk/sites/pub/PublicationReports/MarineScienceSymposia/Phase2/](http://www.ices.dk/sites/pub/PublicationReports/MarineScienceSymposia/Phase2/ICESMarineScienceSymposia-Volume219-2003-Part20of75.pdf)
640 [ICESMarineScienceSymposia-Volume219-2003-Part20of75.pdf](http://www.ices.dk/sites/pub/PublicationReports/MarineScienceSymposia/Phase2/ICESMarineScienceSymposia-Volume219-2003-Part20of75.pdf).

641 Ivanov, V., V. Alexeev, N. V. Koldunov, I. Repina, A. B. Sandø, L. H. Smedsrud, and A. Smirnov,
642 2016: Arctic Ocean Heat Impact on Regional Ice Decay: A Suggested Positive Feed-
643 back. *Journal of Physical Oceanography*, **46 (5)**, 1437–1456, doi:10.1175/JPO-D-15-0144.
644 1, URL <http://journals.ametsoc.org/doi/abs/10.1175/JPO-D-15-0144.1>{\%}5Cn[http://journals.](http://journals.ametsoc.org/doi/10.1175/JPO-D-15-0144.1)
645 [ametsoc.org/doi/10.1175/JPO-D-15-0144.1](http://journals.ametsoc.org/doi/10.1175/JPO-D-15-0144.1).

646 Karcher, M., A. Beszczynska-Möller, F. Kauker, R. Gerdes, S. Heyen, B. Rudels, and U. Schauer,
647 2011: Arctic Ocean warming and its consequences for the Denmark Strait overflow. *Journal of*
648 *Geophysical Research: Oceans*, **116 (C02037)**, 1–10, doi:10.1029/2010JC006265.

- 649 Koenigk, T., U. Mikolajewicz, J. H. Jungclaus, and A. Kroll, 2009: Sea ice in the Barents Sea:
650 Seasonal to interannual variability and climate feedbacks in a global coupled model. *Climate*
651 *Dynamics*, **32 (7-8)**, 1119–1138, doi:10.1007/s00382-008-0450-2.
- 652 Kwok, R., 2009: Outflow of Arctic Ocean Sea Ice into the Greenland and Barents Seas: 19792007.
653 *Journal of Climate*, **22 (9)**, 2438–2457, doi:10.1175/2008JCLI2819.1.
- 654 Larsen, K. M. H., C. Gonzalez-Pola, P. Fratantoni, A. Beszczynska-Möller, and S. L. E. Hughes,
655 2016: ICES Report on Ocean Climate 2015. *ICES Cooperative Research Report*, **331 (May)**,
656 1–79.
- 657 Levitus, S., G. Matishov, D. Seidov, and I. Smolyar, 2009: Barents Sea multidecadal variability.
658 *Geophysical Research Letters*, **36 (19)**, 1–5, doi:10.1029/2009GL039847.
- 659 Lien, V. S., P. Schlichtholz, Ø. Skagseth, and F. B. Vikebø, 2017: Wind-Driven Atlantic Water
660 Flow as a Direct Mode for Reduced Barents Sea Ice Cover. *Journal of Climate*, **30 (2)**, 803–
661 812, doi:10.1175/JCLI-D-16-0025.1.
- 662 Lind, S., and R. B. Ingvaldsen, 2012: Variability and impacts of Atlantic Water entering the Bar-
663 ents Sea from the north. *Deep-Sea Research Part I: Oceanographic Research Papers*, **62**, 70–88,
664 doi:10.1016/j.dsr.2011.12.007.
- 665 Lind, S., R. B. Ingvaldsen, and T. Furevik, 2016: Arctic layer salinity controls heat loss from deep
666 Atlantic layer in seasonally ice-covered areas of the Barents Sea. *Geophysical Research Letters*,
667 **43 (10)**, 5233–5242, doi:10.1002/2016GL068421.
- 668 Lique, C., A. M. Treguier, B. Blanke, and N. Grima, 2010: On the origins of water masses exported
669 along both sides of Greenland: A Lagrangian model analysis. *Journal of Geophysical Research:*
670 *Oceans*, **115 (5)**, 1–20, doi:10.1029/2009JC005316.

- 671 Loeng, H., 1991: Features of the physical oceanographic conditions of the Barents Sea. *Polar*
672 *Research*, **10** (1), 5–18, doi:10.1111/j.1751-8369.1991.tb00630.x.
- 673 Long, Z., and W. Perrie, 2017: Changes in Ocean Temperature in the Barents Sea in the Twenty-
674 First Century. *Journal of Climate*, **30** (15), 5901–5921, doi:10.1175/JCLI-D-16-0415.1, URL
675 <http://journals.ametsoc.org/doi/10.1175/JCLI-D-16-0415.1>.
- 676 Maslowski, W., D. Marble, W. Walczowski, U. Schauer, J. L. Clement, and A. J. Semtner, 2004:
677 On climatological mass, heat, and salt transports through the Barents Sea and Fram Strait from
678 a pan-Arctic coupled ice-ocean model simulation. *Journal of Geophysical Research*, **109** (C3),
679 C03 032, doi:10.1029/2001JC001039.
- 680 McDougall, T. J., D. R. Jackett, F. J. Millero, R. Pawlowicz, and P. M. Barker, 2012: A
681 global algorithm for estimating Absolute Salinity. *Ocean Science*, **8** (6), 1123–1134, doi:
682 [10.5194/os-8-1123-2012](https://doi.org/10.5194/os-8-1123-2012).
- 683 Moat, B., S. Josey, and B. Sinhu, 2014: Impact of Barents Sea winter air-sea exchanges on Fram
684 Strait dense water transport. *Journal of Geophysical Research: Oceans*, **119** (2), 1009–1021,
685 doi:10.1002/2013JC009220.Received.
- 686 Nakanowatari, T., K. Sato, and J. Inoue, 2014: Predictability of the barents sea ice in early winter:
687 Remote effects of oceanic and atmospheric thermal conditions from the North Atlantic. *Journal*
688 *of Climate*, **27** (23), 8884–8901, doi:10.1175/JCLI-D-14-00125.1.
- 689 Notz, D., and J. Stroeve, 2016: Observed Arctic sea-ice loss directly follows anthropogenic CO2
690 emission. *Science*, **354** (6313), 747–750.

- 691 Onarheim, I. H., and M. Årthun, 2017: Toward an ice-free Barents Sea. *Geophysical Re-*
692 *search Letters*, 8387–8395, doi:10.1002/2017GL074304, URL <http://doi.wiley.com/10.1002/>
693 2017GL074304.
- 694 Onarheim, I. H., T. Eldevik, M. Årthun, R. B. Ingvaldsen, and L. H. Smedsrud, 2015: Skillful
695 prediction of Barents Sea ice cover. *Geophysical Research Letters*, **42**, 5364–5371, doi:10.1002/
696 2015GL064359.Abstract.
- 697 Oziel, L., J. Sirven, and J. C. Gascard, 2016: The Barents Sea frontal zones and water masses
698 variability (1980-2011). *Ocean Science*, **12** (1), 169–184, doi:10.5194/os-12-169-2016.
- 699 Oziel, L., and Coauthors, 2017: Role for Atlantic inflows and sea ice loss on shifting phyto-
700 plankton blooms in the Barents Sea. *Journal of Geophysical Research: Oceans*, **122**, 1–19,
701 doi:10.1002/2016JC012582.Received.
- 702 Parsons, A. R., R. H. Bourke, R. D. Muench, C.-S. Chiu, J. F. Lynch, J. H. Miller, A. J. Pluedde-
703 mann, and R. Pawlowicz, 1996: The Barents Sea Polar Front in summer. *Journal of Geophysi-*
704 *cal Research*, **101** (C6), 14 201–14 221, doi:10.1029/96JC00119, URL [http://doi.wiley.com/10.](http://doi.wiley.com/10.1029/96JC00119)
705 1029/96JC00119.
- 706 Petoukhov, V., and V. A. Semenov, 2010: A link between reduced Barents-Kara sea ice and
707 cold winter extremes over northern continents. *Journal of Geophysical Research Atmospheres*,
708 **115** (21), 1–11, doi:10.1029/2009JD013568.
- 709 Polyakov, I. V., and Coauthors, 2017: Greater role for Atlantic inflows on sea-ice loss in the
710 Eurasian Basin of the Arctic Ocean. *Science*, **291** (April), 285–291, doi:10.1126/science.
711 aai8204.

- 712 Pratt, L. J., 2004: Recent progress on understanding the effects of rotation in models of sea straits.
713 *Deep-Sea Research Part II: Topical Studies in Oceanography*, **51 (4-5)**, 351–369, doi:10.1016/
714 j.dsr2.2003.06.005.
- 715 Proudman, J., 1916: On the motion of solids in a liquid possessing vorticity. *Proceedings of the*
716 *Royal Society London A*, **92**, 408–424, doi:10.1103/RevModPhys.4.87, 0511310.
- 717 Reigstad, M., P. Wassmann, C. Wexels Riser, S. Øygarden, and F. Rey, 2002: Variations in hydrog-
718 raphy, nutrients and chlorophyll a in the marginal ice-zone and the central Barents Sea. *Journal*
719 *of Marine Systems*, **38 (1-2)**, 9–29, doi:10.1016/S0924-7963(02)00167-7.
- 720 Rudels, B., L. G. Anderson, and E. P. Jones, 1996: Formation and evolution of the surface mixed
721 layer and halocline of the Arctic Ocean. *Journal of Geophysical Research*, **101**, 8807–8821,
722 doi:10.1029/96JC00143.
- 723 Rudels, B., M. Korhonen, U. Schauer, S. Pisarev, B. Rabe, and A. Wisotzki, 2015: Circulation
724 and transformation of Atlantic water in the Eurasian Basin and the contribution of the Fram
725 Strait inflow branch to the Arctic Ocean heat budget. *Progress in Oceanography*, **132**, 128–152,
726 doi:10.1016/j.pocean.2014.04.003, URL <http://dx.doi.org/10.1016/j.pocean.2014.04.003>.
- 727 Rudels, B., R. D. Muench, J. Gunn, U. Schauer, and H. J. Friedrich, 2000: Evolution of the Arctic
728 Ocean boundary current north of the Siberian shelves. *Journal of Marine Systems*, **25 (1)**, 77–99,
729 doi:10.1016/S0924-7963(00)00009-9.
- 730 Schauer, U., H. Loeng, B. Rudels, V. K. Ozhigin, and W. Dieck, 2002: Atlantic Water flow through
731 the Barents and Kara Seas. *Deep-Sea Research Part I: Oceanographic Research Papers*, **49 (12)**,
732 2281–2298, doi:10.1016/S0967-0637(02)00125-5.

733 Schauer, U., R. D. Muench, B. Rudels, and L. Timokhov, 1997: Impact of eastern Arctic shelf
734 waters on the Nansen Basin intermediate layers. *Journal of Geophysical Research*, **102 (C2)**,
735 3371, doi:10.1029/96JC03366.

736 Schlichtholz, P., and M. N. Houssais, 2011: Forcing of oceanic heat anomalies by air-sea in-
737 teractions in the Nordic Seas area. *Journal of Geophysical Research: Oceans*, **116 (1)**, 1–22,
738 doi:10.1029/2009JC005944.

739 Schmidtko, S., G. C. Johnson, and J. M. Lyman, 2013: MIMOC: A global monthly isopycnal
740 upper-ocean climatology with mixed layers. *Journal of Geophysical Research: Oceans*, **118 (4)**,
741 1658–1672, doi:10.1002/jgrc.20122.

742 Screen, J. A., and I. Simmonds, 2010: The central role of diminishing sea ice in recent Arc-
743 tic temperature amplification. *Nature*, **464 (7293)**, 1334–1337, doi:10.1038/nature09051, URL
744 <http://dx.doi.org/10.1038/nature09051>.

745 Shapiro, G. I., 2003: Dense water cascading off the continental shelf. *Journal of Geophysical*
746 *Research*, **108 (C12)**, 3390, doi:10.1029/2002JC001610, URL [http://doi.wiley.com/10.1029/](http://doi.wiley.com/10.1029/2002JC001610)
747 [2002JC001610](http://doi.wiley.com/10.1029/2002JC001610).

748 Sigmond, M., M. C. Reader, G. M. Flato, W. J. Merryfield, and A. Tivy, 2016: Skillful seasonal
749 forecasts of Arctic sea ice retreat and advance dates in a dynamical forecast system. *Geophysical*
750 *Research Letters*, **43 (24)**, 12,457–12,465, doi:10.1002/2016GL071396.

751 Singh, R. K., M. Maheshwari, S. R. Oza, and R. Kumar, 2013: Long-term variability in Arctic sea
752 surface temperatures. *Polar Science*, **7 (3-4)**, 233–240, doi:10.1016/j.polar.2013.10.003, URL
753 <http://dx.doi.org/10.1016/j.polar.2013.10.003>.

- 754 Skagseth, Ø., 2008: Recirculation of Atlantic Water in the western Barents Sea. *Geophysical*
755 *Research Letters*, **35** (11), 1–5, doi:10.1029/2008GL033785.
- 756 Slagstad, D., and T. A. McClimans, 2005: Modeling the ecosystem dynamics of the Barents sea
757 including the marginal ice zone: I. Physical and chemical oceanography. *Journal of Marine*
758 *Systems*, **58** (1-2), 1–18, doi:10.1016/j.jmarsys.2005.05.005.
- 759 Smedsrud, L. H., R. Ingvaldsen, J. E. Ø. Nilsen, and Ø. Skagseth, 2010: Heat in the Bar-
760 ents Sea: transport, storage, and surface fluxes. *Ocean Science*, **6** (1), 219–234, doi:10.5194/
761 os-6-219-2010.
- 762 Smedsrud, L. H., and Coauthors, 2013: The role of the Barents Sea in the Arctic climate system.
763 *Reviews of Geophysics*, **51** (3), 415–449, doi:10.1002/rog.20017.
- 764 Snape, T., 2013: Decline of Arctic Sea Ice: Evaluation and weighting of CMIP5 projections. *Jour-*
765 *nal of Geophysical Research : Atmospheres*, **119** (2), 546–554, doi:10.1002/2013JD020593.
766 Received.
- 767 Sorteberg, A., and B. Kvingedal, 2006: Atmospheric forcing on the Barents Sea winter ice extent.
768 *Journal of Climate*, **19** (19), 4772–4784, doi:10.1175/JCLI3885.1.
- 769 Steele, M., and W. Ermold, 2015: Loitering of the retreating sea ice edge in the Arctic Seas.
770 *Journal of Geophysical Research: Oceans*, **120**, 7699–7721, doi:10.1002/2015JC010969.
- 771 Taylor, G. I., 1917: Motion of solids in fluids when the flow is not irrotational. *Proceedings of the*
772 *Royal Society London A*, **93**, 92–113.
- 773 Thomson, R. E., and W. J. Emery, 2014: *Data analysis methods in physical oceanography*. 3rd
774 ed., Newnes.

- 775 Våge, S., S. L. Basedow, K. S. Tande, and M. Zhou, 2014: Physical structure of the Barents
776 Sea Polar Front near Storbanken in August 2007. *Journal of Marine Systems*, **130** (August
777 **2007**), 256–262, doi:10.1016/j.jmarsys.2011.11.019, URL [http://dx.doi.org/10.1016/j.jmarsys.](http://dx.doi.org/10.1016/j.jmarsys.2011.11.019)
778 2011.11.019.
- 779 Venegas, S. A., and L. A. Mysak, 2000: Is there a dominant timescale of natural climate vari-
780 ability in the Arctic? *Journal of Climate*, **13** (19), 3412–3434, doi:10.1175/1520-0442(2000)
781 013<3412:ITADTO>2.0.CO;2.
- 782 Wang, M., and J. E. Overland, 2012: A sea ice free summer Arctic within 30 years: An update
783 from CMIP5 models. *Geophysical Research Letters*, **39** (17), 2–6, doi:10.1029/2012GL052868.
- 784 Weatherall, P., and Coauthors, 2015: A new digital bathymetric model of the world’s oceans. *Earth*
785 *and Space Science*, **2**, 331–345, doi:10.1002/2015EA000107.Received.
- 786 Yang, J., and J. F. Price, 2000: Water-mass formation and potential vorticity balance in
787 an abyssal ocean circulation. *Journal of Marine Research*, **58**, 789–808, doi:10.1357/
788 002224000321358918.
- 789 Yang, X. Y., X. Yuan, and M. Ting, 2016: Dynamical link between the Barents-Kara sea ice and
790 the arctic oscillation. *Journal of Climate*, **29** (14), 5103–5122, doi:10.1175/JCLI-D-15-0669.1.

791 **LIST OF TABLES**

792 **Table 1.** Definitions of the water masses present in the Barents Sea used in this study,
793 along with definitions used in previous studies. Note that Barents Sea Water
794 can be referred to as Modified Atlantic Water in literature. 38


795 TABLE 1. Definitions of the water masses present in the Barents Sea used in this study, along with definitions
 796 used in previous studies. Note that Barents Sea Water can be referred to as Modified Atlantic Water in literature.

Water Mass	Source	Temperature	Salinity	Density
Atlantic Water (AW)	Present Study	T >3.0	S >35.0	
	Oziel et al. (2016)	T >3.0	S >34.8	
	Loeng (1991)	T >3.0	S >35.0	
Arctic Water (ArW)	Present Study	T <0.0	S <34.7	
	Oziel et al. (2016)	T <0.0	S <34.7	
	Loeng (1991)	T <0.0	34.3 < S < 34.8	
Coastal Water (CW)	Present Study	T >2.0	S <34.7	
	Oziel et al. (2016)	T >3.0	S <34.4	
	Loeng (1991)	T >2.0	S <34.7	
Barents Sea Water (BSW)	Present Study	T <2.0	S >34.7	σ >27.85
	Schauer et al. (2002)			σ >27.85
	Oziel et al. (2016)	T <2.0	S >34.8	σ >27.8
	Loeng (1991)	-1.5 < T < 2.0	34.7 < S < 35.0	


LIST OF FIGURES

- 798 **Fig. 1.** Bathymetry of the Barents Sea. The different lines and box indicate the area used for EOF
799 analysis of SST (green box), the region used for Hovmoller analysis (blue-dashed box), the
800 cross-front transect (light-blue line), the area selected for calculating the contribution of sea
801 ice to AW/BSW (dark-blue box), the area selected for 100 – 300 m BSW properties from
802 EN4 data and 0 – 100 m ArW properties from EN4 data (cyan-dashed box), the area selected
803 for 0 – 100 m surface BSW properties from EN4 data south of the PF (yellow-dashed line),
804 the the Kola section (orange line) and the Fugløya–Bear Island section (red line). 41
- 805 **Fig. 2.** (a) Mean SST across the Barents Sea with a 12-month running mean (blue line). The linear
806 trend for the periods 1985 to 2004 and 2005 to 2016 are shown (green lines). Trend in SST
807 for the periods (b) 1985 to 2004 and (c) 2005 to 2016. Note that a different colour scale is
808 used in the two panels. Trends are significant at a level of 95% in un-hatched areas. The
809 black line indicates the 220 m isobath. 42
- 810 **Fig. 3.** SST seasonal climatology from 2005 to 2016 for (a) spring (March, April and May), (b)
811 summer (June, July and August), (c) autumn (September, October and November) and (d)
812 winter (December, January and February), respectively. Gradient in SST seasonal climatol-
813 ogy from 2005 to 2016 for (e) spring, (f) summer, (g) autumn and (h) winter, respectively.
814 The sea ice edge is defined by 15% sea ice concentration (white line) and the black line
815 indicates the 220 m isobath. 43
- 816 **Fig. 4.** As Figure 3 but for the seasonal climatology from 1985 to 2004. 44
- 817 **Fig. 5.** (a) Spatial pattern of first EOF mode of SST variability. The black line indicates the 220 m
818 isobath. (b) Regression of PC 1 with SAT. Maximum correlation (r-value) is shown in the
819 bottom left hand corner and the location of the maximum correlation is shown by a black
820 cross. Hatched areas are not significant at the 95%-level. (c) Time series of PC 1. (d) Time
821 series of AW temperature at the Kola section (blue line, 12-month running mean applied) and
822 Fugløya–Bear Island (FB) section (green line, 12-month running mean applied). (e) Time
823 series of sea ice extent in the Barents Sea (12-month running mean applied). Correlations
824 between each variable and PC 1 are indicated. 45
- 825 **Fig. 6.** (a) Magnitude of the meridional gradient in zonally-averaged SST between 35° E and 50° E
826 (blue-dashed box on Figure 1) and Polar Front location (dashed line). The magnitude is
827 normalized on a daily basis by its standard deviation to show the changes in the position
828 of the front over time. Note that changes in intensity over time cannot be deduced from
829 (a). (b) Latitude of the sea ice edge for the same region. (c) mean SST gradient between
830 76.3° N and 76.7° N before normalization (blue line, 12-month running mean applied) and
831 AW temperature from the Kola section (green line, 12-month running mean applied, section
832 marked in Figure 1). Gaps indicate missing data and sea ice coverage for AW and the SST
833 gradient respectively. (d) BSW salinity (blue line) and temperature (green line) between 100
834 and 300 m from the EN4 data, averaged in the cyan-dashed box on Figure 1. Uncertainty
835 values for EN4 data are shown by the shaded areas. Dashed green and blue lines in (c) and
836 (d) show the respective means for 1985 – 2004 and 2005 – 2016. 46
- 837 **Fig. 7.** Eastern Barents Sea transect at 44° E (shown in Figure 1) from the MIMOC and EN4 cli-
838 matology during winter (December, January, February), EN4 is averaged over 1985 – 2016.
839 The Polar Front is marked by the black triangle. (a,b) Salinity, (c,d) temperature and (e,f)
840 potential density. White areas in (a-f) indicate grid cell with no data, black points show the
841 grid cells containing BSW, and the EN4 sub-section (black dashed area) used to produce the
842 BSW temperature and salinities (Figure 6). (g,h) T-S diagrams showing the different water


843 masses present in (a-f). The color indicates the latitude of the profile. The green dotted area
844 in (g,h) shows the limits of the BSW definitions, and AW, ArW and CW water masses are
845 indicated (see Table 1 for their definitions) 47



846 FIG. 1. Bathymetry of the Barents Sea. The different lines and box indicate the area used for EOF analysis of
 847 SST (green box), the region used for Hovmoller analysis (blue-dashed box), the cross-front transect (light-blue
 848 line), the area selected for calculating the contribution of sea ice to AW/BSW (dark-blue box), the area selected
 849 for 100 – 300 m BSW properties from EN4 data and 0 – 100 m ArW properties from EN4 data (cyan-dashed
 850 box), the area selected for 0 – 100 m surface BSW properties from EN4 data south of the PF (yellow-dashed
 851 line), the the Kola section (orange line) and the Fugløy-Bear Island section (red line).



852 FIG. 2. (a) Mean SST across the Barents Sea with a 12-month running mean (blue line). The linear trend for
 853 the periods 1985 to 2004 and 2005 to 2016 are shown (green lines). Trend in SST for the periods (b) 1985 to
 854 2004 and (c) 2005 to 2016. Note that a different colour scale is used in the two panels. Trends are significant at
 855 a level of 95% in un-hatched areas. The black line indicates the 220 m isobath.



856 FIG. 3. SST seasonal climatology from 2005 to 2016 for (a) spring (March, April and May), (b) summer
 857 (June, July and August), (c) autumn (September, October and November) and (d) winter (December, January
 858 and February), respectively. Gradient in SST seasonal climatology from 2005 to 2016 for (e) spring, (f) summer,
 859 (g) autumn and (h) winter, respectively. The sea ice edge is defined by 15% sea ice concentration (white line)
 860 and the black line indicates the 220 m isobath.







FIG. 4. As Figure 3 but for the seasonal climatology from 1985 to 2004.



861 FIG. 5. (a) Spatial pattern of first EOF mode of SST variability. The black line indicates the 220 m isobath.
 862 (b) Regression of PC 1 with SAT. Maximum correlation (r-value) is shown in the bottom left hand corner and
 863 the location of the maximum correlation is shown by a black cross. Hatched areas are not significant at the
 864 95%-level. (c) Time series of PC 1. (d) Time series of AW temperature at the Kola section (blue line, 12-month
 865 running mean applied) and Fugløya–Bear Island (FB) section (green line, 12-month running mean applied). (e)
 866 Time series of sea ice extent in the Barents Sea (12-month running mean applied). Correlations between each
 867 variable and PC 1 are indicated.



868 FIG. 6. (a) Magnitude of the meridional gradient in zonally-averaged SST between 35° E and 50° E (blue-
 869 dashed box on Figure 1) and Polar Front location (dashed line). The magnitude is normalized on a daily basis
 870 by its standard deviation to show the changes in the position of the front over time. Note that changes in
 871 intensity over time cannot be deduced from (a). (b) Latitude of the sea ice edge for the same region. (c) mean
 872 SST gradient between 76.3° N and 76.7° N before normalization (blue line, 12-month running mean applied)
 873 and AW temperature from the Kola section (green line, 12-month running mean applied, section marked in
 874 Figure 1). Gaps indicate missing data and sea ice coverage for AW and the SST gradient respectively. (d) BSW
 875 salinity (blue line) and temperature (green line) between 100 and 300 m from the EN4 data, averaged in the
 876 cyan-dashed box on Figure 1. Uncertainty values for EN4 data are shown by the shaded areas. Dashed green
 877 and blue lines in (c) and (d) show the respective means for 1985 – 2004 and 2005 – 2016.



878 FIG. 7. Eastern Barents Sea transect at 44° E (shown in Figure 1) from the MIMOC and EN4 climatology
 879 during winter (December, January, February), EN4 is averaged over 1985 – 2016. The Polar Front is marked
 880 by the black triangle. (a,b) Salinity, (c,d) temperature and (e,f) potential density. White areas in (a-f) indicate
 881 grid cell with no data, black points show the grid cells containing BSW, and the EN4 sub-section (black dashed
 882 area) used to produce the BSW temperature and salinities (Figure 6). (g,h) T-S diagrams showing the different
 883 water masses present in (a-f). The color indicates the latitude of the profile. The green dotted area in (g,h)
 884 shows the limits of the BSW definitions, and AW, ArW and CW water masses are indicated (see Table 1 for their
 885 definitions)