
Commentary

Occupational lung disease in the South African mining industry: Research and policy implementation

Jill Murray^{a,b,*}, Tony Davies^b, and David Rees^{a,b}

^aNational Institute for Occupational Health (NIOH), National Health Laboratory Service, PO Box 4788, Johannesburg 2000, South Africa.
E-mail: jill.murray@nioh.nhls.ac.za

^bSchool of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa.

*Corresponding author.

Abstract South African miners face an epidemic of occupational lung diseases. Despite a plethora of research on the mining industry, and the gold mining industry in particular, research impact (including disease surveillance) on policy implementation and occupational health systems performance lags. We describe the gold mining environment, and research on silicosis, tuberculosis, HIV and AIDS, and compensation for occupational disease including initiatives to influence policy and thus reduce dust levels and disease. As these have been largely unsuccessful, we identify possible impediments, some common to other low- and middle-income countries, to the translation of research findings and policy initiatives into effective interventions.

Journal of Public Health Policy (2011) 32, S65–S79. doi:10.1057/jphp.2011.25

Keywords: occupational health; silicosis; tuberculosis; HIV/AIDS; compensation; policy research

Introduction

One hundred and twenty years after the discovery of gold on the Witwatersrand in 1886, mining remains a major force in the South African economy. Despite policy and legislative reform, partly informed by extensive research, 16 years into a democratic South Africa, gold miners in particular face an epidemic of occupational lung diseases. This provides an occupational health context in which to examine the impact of research on policy implementation and occupational

health systems performance. This article outlines the gold mining environment, presents disease burdens, describes initiatives to reduce disease, and concludes with factors that may have impeded effective interventions, some common to other countries.

The Gold Mining Industry

Although well below the half million employed in the 1990s, South African gold mines still employ approximately 160 000 people.¹ The industry is characterized by male cross-border and internal rural-urban migrants who leave their families and live mostly in single-sex mining compounds, returning home for variable periods.² Until 1975, work contracts were temporary and short, but labour arrangements stabilized thereafter and duration of service increased.^{3,4} These social circumstances and stabilization have contributed to serious inter-related epidemics of silicosis (silica dust-induced scarring of the lung),⁵ tuberculosis (TB), and HIV infection in miners, in surrounding communities and in labor-sending regions.² Post-apartheid reforms deracialized legislation, but race remains an important determinant of occupation, salary, housing, and disease burden.

Surveillance and Research

In South Africa, surveillance of occupational exposures and diseases is weak notwithstanding efforts of the Department of Mineral Resources to maintain registers such as the South African Mining Occupational Diseases database, initiated in 1998.⁶ The Pathology Automation System (PATHAUT), an electronic database of approximately 100 000 autopsies of deceased miners, dating back to 1975, is a striking exception. Although provision for compensation for occupational lung diseases to current and former miners and their families in life and after death has been in place since 1911, the Occupational Diseases in Mines and Works Act of 1973 catalysed the development of the database by the South African Medical Research Council.⁷ PATHAUT provides a rich data-source for research and monitoring disease trends.^{8,9} Universities and research entities, some partially funded by the state or from levies on industry administered by the state, filled gaps in data, on black miners in particular, in the 1990s and 2000s, showing rates of disease previously unappreciated. Silicosis demonstrates these aspects



well. Importantly, the proportion of black gold miners found to have silicosis at autopsy increased from 3 per cent in 1975 to 32 per cent in 2007. Key studies describing silicosis prevalence in gold miners (Table 1) show substantial disease burdens into the late 2000s, with no evidence that interventions have reduced these to acceptable levels.

Biologic and social factors combine to create a 'perfect storm' for the interaction among silicosis, TB, and HIV. Thus, integrated, multifaceted remedies are required to control these epidemics. Silicosis substantially increases the risk of TB^{19–22} to a magnitude similar to that of HIV infection.²³ Importantly, silica exposure is associated with TB even in the absence of silicosis^{20,24} and the increased risk is lifelong.²⁴ Risk factors such as migrancy²⁵ and single-sex compounds increase high-risk sexual behaviour,^{26,27} and thus HIV rates, which are close to 30 per cent among these miners.²⁸ The TB risks of silicosis and HIV infection combine multiplicatively.²³ Consequently, the highest recorded rates of TB worldwide have been reported in South African gold miners.²⁹ Mortality from TB is higher than that from mine accidents.^{30,31} The prevalence of TB in gold miners has increased from 806 per 100 000 in 1991 to 3821 in 2004.³² HIV prevalence rose from less than 1 per cent in 1987³³ to 27 per cent in 2000.²⁸ More recent data, after the roll out of antiretroviral treatment in 2003, are not available.

There is a large body of research on the control of TB in South African mines that addresses the use of isoniazid therapy to prevent TB in people with HIV;^{34–36} the spread of TB in the mining environment;^{37–41} and TB case finding.^{42–44} Studies have evaluated clinical management,^{45–47} and delayed, missed, or inaccurate diagnosis of TB in life^{28,48} that have added to TB transmission. Drug resistance^{37,49,50} and recurrence of TB^{41,51–53} have also complicated control. Researchers and occupational health practitioners have developed materials to assist mine health-care workers with TB management.^{54,55} The Mine Health and Safety Council, a tripartite body formed under the Mine Health and Safety Act of 1996, and comprising representatives from the Department of Mineral Resources, the mining industry, and trade unions, funded and published the work. Neither the Council nor any other organization has formally measured the uptake of the materials but anecdotal evidence suggests that they are not widely used.

The migrant labour system has weakened incentives to control dust and disease by externalizing costs of disease, moving them away from the gold mining industry to communities and the State.^{2,56–61}



Table 1: Silicosis prevalence in gold miners, 1940–2007

Authors	Study design	Study period	Study population	Mean employment (years)	Diagnostic tool	Proportion (%)
Hnizdo and Sluis-Cremer ¹⁰	Cohort	1940–1991	Deceased white gold miners	23.5	Chest X-rays	14.0
Murray et al ¹¹	Trend analysis	1975–1991	Deceased black gold miners	4.8	Autopsy	9.3–12.8
Murray and Hnizdo ¹²	Cohort	1940–2003	Deceased white gold miners	23.5	Autopsy	51.6
Steen et al ¹³	Cross-sectional	1994	Living black ex-miners	15.5	Chest X-rays	26.6–31.0
Trapido et al ¹⁴	Cross-sectional	1996	Living black ex-miners	12.2	Chest X-rays	22.0–36.0
Meel ¹⁵	Cross-sectional	1997–1999	Living black ex-miners	—	Chest X-rays	34
Churchyard et al ¹⁶	Cross-sectional	2000–2001	Employed black miners	21.8	Chest X-rays	18.3–19.9
Girdler-Brown et al ¹⁷	Cross-sectional	1999	Living black ex-miners	25.6	Chest X-rays	24.6
Park et al ¹⁸	Cohort	1999–2000	Living black ex-miners	26.1	Chest X-rays	27
Nelson et al ⁴	Trend analysis	1975–2007	Deceased black gold miners	13.4	Autopsy	3–32



South Africa requires multifaceted public policies to address this negative impact of migrancy that is common to many low and middle-income countries. Bilateral agreements between countries to provide industry-funded health services for former miners in labour-sending areas is one example. Another is patient-retained medical records to improve continuity of care for migrant workers.⁶² Accompanying this are failures of the occupational disease compensation system. Barriers to compensation are considerable and the majority of qualifying claimants have not received awards (Table 2), thus reducing the substantial financial incentive to control dust that would be brought

Table 2: Studies of occupational disease compensation processes and outcomes in miners employed on South African mines

<i>Study</i>	<i>Context</i>	<i>Study period</i>	<i>Outcome</i>
Steen <i>et al</i> ¹³	304 former gold miners living in Thamanga, Botswana	1994	Very few miners with occupational lung disease had been compensated (proportion not specified)
Trapido <i>et al</i> ¹⁴	238 former gold miners living in Eastern Cape, South Africa	1996	62% of those eligible not compensated. Only 2.5% fully compensated
Murray <i>et al</i> ⁶⁴	All 2530 miners who came to autopsy	1999	446 cases (19%) had occupational lung disease not identified and submitted for compensation in life, or more severe disease than had been compensated in life. 31/446 (7%) of families had received benefits by April 2001
Roberts ⁶⁵	205 former miners, Eastern Cape, South Africa	2008	175/205 (85%) reported not receiving the statutory medical examination when leaving the mine (which is partly to identify compensable disease). 203/205 (99%) did not know of the Compensation Act and its benefits
Maiphethlo <i>et al</i> ⁶⁶	84 former mine workers diagnosed with silicosis at Groote Schuur Hospital, Cape Town, and submitted for compensation	1993–2005	17/84 (20%) recorded as having received compensation. Median time from submission of medical records to receipt of award was 51 months (range 22–84 months)

about by compensation payments and hence increased levies on mines. Trapido *et al*⁶³ estimated the unpaid occupational lung disease compensation for gold miners to be 10 billion South African rands (US\$1.4 billion; \$1 = R7) in 1996, closer to 20 billion rands (US\$2.9 billion) today. Even in poor countries, investments in compensation systems should bring greater returns in reduced disease burdens and their attendant costs to the public health system, provided that compensation costs are borne by industry.

Policy, Legislative and Service Initiatives

Although the history of disease reduction initiatives dates back to the early 1900s, these cannot be seen in isolation from South Africa's apartheid history of racial discrimination.⁶⁷ We describe attempts to achieve control of silica dust, reduced impact of migrancy, improved housing, and better TB management and compensation systems necessary to reduce disease burdens. In 1973, the Occupational Diseases in Mines and Works Act (ODMWA) provided compensation for occupational diseases for miners, but benefits were substantially higher for white miners than other groups. The 1993 Amendment to ODMWA established racial parity in compensation benefits but access to benefits is unsatisfactory (Table 2). In Table 3 we summarize key initiatives in the last two decades and comment on their outcomes.

The 1994 Leon Commission of Inquiry into Safety and Health in the Mining Industry, whose recommendations were informed by evidence presented by the National Union of Mineworkers, researchers, and activists, has had far-reaching consequences. The inquiry led to promulgation of a modern and comprehensive Mine Health and Safety Act with ancillary tripartite structures (representation of labour, government, and industry), such as the Mine Health and Safety Council (MHSC), to promote, develop, and oversee policy, standards, and encourage relevant research. However, neither the state, academics, nor trade unions have evaluated the impact of this law on practice.

Government has not implemented its long-standing policy of improving compensation benefits for miners (under the auspices of the Department of Health), in line with those available to non-mining workers (under the auspices of the Department of Labour), probably because of the increased and unfunded costs.



Numerous initiatives have sought to address living conditions of miners, but progress has been slow.⁶⁸ A substantial body of applied research to reduce silica dust and disease^{69,70} coincides with widely published targets (milestones),⁷¹ the establishment of task teams to disseminate good practice, and campaigns by stakeholders. There is, however, limited evidence of sustained reductions in silica dust levels.^{72,73} Translation of research into practice has been patchy; for example, MHSC-sponsored research-based materials to raise awareness and skills among practitioners, miners, health and safety representatives, and managers⁷¹ have not been disseminated by the MHSC, mining houses, or trade unions.

Efforts of mine medical services to control TB in the mines have had limited impact.⁶² Although the mining industry has made substantial progress in developing HIV and AIDS programmes for miners,⁷⁴ there is little evidence of declining HIV rates. Despite research and numerous policy initiatives, South Africa's mining industry falls short of compliance with health and safety legal requirements, as demonstrated by the Department of Health,⁷³ the AIDS and Rights Alliance for Southern Africa,⁷⁵ and a Presidential Audit of compliance⁷⁶ reports.

Barriers to Translating Research into Action

The mining industry, the state, trade unions, and academics have done scant formal critical analysis of factors that have impeded effective interventions to reduce disease.⁷⁷ In the absence of this analysis, we have identified financial, social, and organizational factors with face-validity that might explain much of the failure to implement translation of research and policy into action (Table 4). Many of these factors could apply to other countries.

Conclusion

Scientific studies have motivated and informed many sound policy decisions and legislative reforms, and have identified strategies for strengthening practice and health systems. South Africa has adopted some recommendations, but very few studies seem to have fostered sustained remedial action. There are clear gaps between research, policy, and implementation in occupational health practice in the South African mining industry.

Table 3: Policy, legislative and service initiatives affecting miners in South Africa: 1993–2010

<i>Year</i>	<i>Initiative</i>	<i>Primary responsibility</i>	<i>Comment</i>
1990s to present	Substantial body of research initiated and sponsored by Mine Health and Safety Council, notably the Silicosis Control Programme, to improve measurement and control of dust and awareness of the diseases and the roles of workers, management, and practitioners	Department of Mineral Resources	Important findings widely disseminated but scant evaluation of impact on practice and limited availability of key research-based tools
1993	Amendment to Occupational Diseases in Mines and Works Act to provide uniform compensation benefits, irrespective of race	Department of Health	Did not achieve purpose because of failure to award benefits to the majority of qualifying claimants, and to make benefit examinations accessible to former miners in the labour-sending areas
1993 to present	HIV/AIDS research and numerous tripartite engagements and summits. Widespread implementation at large mines of comprehensive programmes on HIV and AIDS, including prevention, treatment, community outreach, and evaluation	Department of Mineral Resources, Department of Health, academics, mining industry, trade unions	Consensus on principles and policies achieved and national and industry plans formulated. Business case for action widely accepted. Little evidence of declining HIV rates
1994	Leon Commission of Inquiry into Safety and Health in the Mining Industry to recommend measures to improve mine health and safety	Presidential Inquiry	Swift incorporation of many recommendations into law, but no published analysis of impact of these regulatory reforms
1995	Mine Health and Safety Act with subsequent amendments, regulations, codes of practice, guidelines, and standards	Department of Mineral Resources	Modern comprehensive Act. Impact not formally evaluated but probably limited by relative focus on accident prevention, poor data on exposures and occupational lung diseases, and weaknesses in enforcement of dust control and measures to reduce lung disease



1999	Task-force to integrate the occupational health and compensation systems in the country, partly to improve compensation benefits for miners	Department of Labour	Separate systems for mining and non-mining industries persist. Contested domains of authority and unfunded liabilities may explain lack of progress
2005	Social Contract for Rapid Housing Delivery partly to reduce single-sex compounds	Department of Housing	Some progress but 51% of gold miners still housed in shared single-sex dwellings in 2007
2009	Housing and Living Conditions Standard for the South African Minerals Industry to provide standards and tools for monitoring and evaluation	Department of Mineral Resources	Promulgated, but not yet implemented
2003	Formulation of milestones (targets) to reduce silica dust and eliminate new cases of silicosis in mining	Department of Mineral Resources	Extensively publicized and endorsed; task teams established to promote attainment. Insufficient data to evaluate progress
2003	Guidance Note for Occupational Medicine Practitioners: Tuberculosis Control Programmes. Detailed guide to assist employers in preparing a comprehensive TB control programme	Department of Mineral Resources	Excellent recommendations but, to date, no evaluation of implementation published and Guidance Note not updated as stipulated
2007	Tuberculosis Strategic Plan for South Africa 2007–2011 outlines key role of mining and importance of dust control and improved living conditions in TB control	Department of Health	Weak intersectoral collaboration among government departments limited impact

Table 4: Factors influencing translation of research findings and policy initiatives into effective interventions to reduce disease

<i>Factor</i>	<i>Comment</i>
<i>Financial and other resources</i>	
Unknown cost of disease reduction	Mining industry, government departments, or other agencies have not done any cost analysis, but costs perceived to be high
Compensation system weaknesses	Low compensation costs limit the financial incentive for the mining industry to reduce dust and disease
Declining industry	Government departments may redirect their attention to growing industries
Competing demands for limited resources	Other public health concerns of the Department of Health and the mining industry may override occupational disease, eg the rapid escalation, scale, and the complexity of the HIV epidemic overtook existing infrastructure, policy formulation, and programme development capacity
<i>Social</i>	
High rates of migrancy	Integral to southern African economies but poorly developed government solutions to reduce impact as a disease-driver
Unequal power relations among social partners	Trade unions under-resourced (finance, knowledge, etc) to contribute adequately to self-regulatory tripartite model of Mine Health and Safety Act
Limited civil society activism	Weak occupational health and safety non-governmental organizations
<i>Organizational</i>	
Fragmented occupational health system	Domain-contestation among Departments of Health, Labour, and Mineral Resources inhibits formulation of integrated national policies and appropriate resource allocation
Weak inspectorate in the Department of Mineral Resources	Capacity directed to safety issues
Poor monitoring and evaluation	Inadequate information on dust levels and disease burdens, and evaluation of enterprise-level programmes and impact of policies/legislation has prevented identification of effective interventions. Government policies do not identify agencies responsible for monitoring and evaluation and/or allocate appropriate resources
Focus on accident prevention	Immediacy of injuries and traumatic deaths reduces focus on the more insidious impact of disease mortality, by all role players

Mining is growing in southern Africa and in many low- and middle-income countries. Other countries could avoid the high levels of occupational diseases experienced in South Africa through an enhanced



understanding of the implications of the failure to use research evidence. Key messages to other countries are to: monitor dust and disease levels reliably, evaluate the impact of policy and regulatory reforms, and define the roles and responsibilities of individual government departments and other agencies clearly.

About the Authors

Jill Murray (MD) is an associate professor in the School of Public Health at the University of the Witwatersrand, Johannesburg, and is Head of the Pathology Division at the National Institute for Occupational Health (NIOH), South Africa.

David Rees (MD, PhD) is Professor of Occupational Health at the University of the Witwatersrand, Johannesburg, and is Head of the Occupational Medicine Division at the NIOH, South Africa. He was Director of the NIOH from 1998 to 2003.

Tony Davies (MD) is Professor Emeritus at the University of the Witwatersrand, Johannesburg, and was Director of the NIOH from 1985 to 2002.

References

1. Chamber of Mines. (2009) *Facts and Figures*. Johannesburg: Chamber of Mines of South Africa.
2. Rees, D., Murray, J., Nelson, G. and Sonnenberg, P. (2009) Oscillating migration and the epidemics of silicosis, tuberculosis, and HIV infection in South African gold miners. *American Journal of Industrial Medicine* 53(4): 398–404.
3. Leger, J.P. (1992) Occupational diseases in South African mines – A neglected epidemic? *South African Medical Journal* 81(4): 197–201.
4. Nelson, G., Girdler-Brown, B., Ndlovu, N. and Murray, J. (2010) Three decades of silicosis: Disease trends at autopsy in South African gold miners. *Environmental Health Perspectives* 118(3): 421–426.
5. Rees, D. and Murray, J. (2007) Silica, silicosis and tuberculosis. *International Journal of Tuberculosis and Lung Disease* 11(5): 474–484.
6. Torres, N.G., Boyce, J.E., Barnes, D.F., White, N. and du Plessis, A. (1998) The development of an occupational diseases database, to be managed by the Department of Minerals and Energy, to record morbidity and mortality of occupational diseases in the South African mining industry (GEN 505). Johannesburg: Safety in Mines Research Advisory Committee, <http://www.mhsc.org.za/dmdocuments/Reports/GEN/gen505/Gen505.pdf>, accessed 20 December 2010.
7. Soskolne, C.L., Goldstein, B., Haffajee, I.A. and Cowling, G. (1976) An integrated cardio-respiratory pathology information system. *South African Medical Journal* 50(45): 1832–1836.

8. Ndlovu, N., Murray, J., Davies, T. and Nelson, G. (2010) Pathology Division Surveillance Report: Demographic Data and Disease Rates for January to December 2009. Johannesburg: National Institute for Occupational Health. National Health Laboratory Services, Johannesburg. NIOH Report 3/2010, ISSN 1812-7681.
9. National Institute for Occupational Health. (2009) Fact Sheet: Mining industry related burden of disease. Johannesburg: National Institute for Occupational Health, National Health Laboratory Service, http://www.nioh.ac.za/docs/fact_sheets/Mining.pdf, accessed 20 December 2010.
10. Hnizdo, E. and Sluis-Cremer, G.K. (1993) Risk of silicosis in a cohort of white South African gold miners. *American Journal of Industrial Medicine* 24(4): 447-457.
11. Murray, J., Kielkowski, D. and Reid, P. (1996) Occupational disease trends in black South African gold miners. An autopsy-based study. *American Journal of Respiratory and Critical Care Medicine* 153(2): 706-710.
12. Murray, J. and Hnizdo, E. (2005) *Development of silicosis in a cohort of South African gold miners – radiological and autopsy-based study*, Presented at the 10th International Conference on Occupational Respiratory Disease, Beijing, China.
13. Steen, T.W. *et al* (1997) Prevalence of occupational lung disease among Botswana men formerly employed in the South African mining industry. *Occupational and Environmental Medicine* 54(1): 19-26.
14. Trapido, A.S. *et al* (1998) Prevalence of occupational lung disease in a random sample of former mineworkers, Libode District, Eastern Cape Province, South Africa. *American Journal of Industrial Medicine* 34(4): 305-313.
15. Meel, B.L. (2002) Patterns of lung diseases in former mine workers of the former Republic of the Transkei: An X-ray-based study. *International Journal of Occupational and Environmental Health* 8(2): 105-110.
16. Churchyard, G.J. *et al* (2004) Silicosis prevalence and exposure-response relations in South African goldminers. *Occupational and Environmental Medicine* 61(10): 811-816.
17. Girdler-Brown, B.V., White, N.W., Ehrlich, R.I. and Churchyard, G.J. (2008) The burden of silicosis, pulmonary tuberculosis and COPD among former Basotho goldminers. *American Journal of Industrial Medicine* 51(9): 640-647.
18. Park, H.H., Girdler-Brown, B.V., Churchyard, G.J., White, N.W. and Ehrlich, R.I. (2009) Incidence of tuberculosis and HIV and progression of silicosis and lung function impairment among former Basotho gold miners. *American Journal of Industrial Medicine* 52(12): 901-908.
19. Cowie, R.L. (1994) The epidemiology of tuberculosis in gold miners with silicosis. *American Journal of Respiratory and Critical Care Medicine* 150: 1460-1462.
20. teWaterNaude, J.M. *et al* (2006) Tuberculosis and silica exposure in South African gold miners. *Occupational and Environmental Medicine* 63(3): 187-192.
21. Corbett, E.L. *et al* (1999) Risk factors for pulmonary mycobacterial disease in South African gold miners. A case-control study. *American Journal of Respiratory and Critical Care Medicine* 159(1): 94-99.
22. Stuckler, D., Basu, S., McKee, M. and Lurie, M. (2011) Mining and risk of tuberculosis in sub-Saharan Africa. *American Journal of Public Health* 101(3): 524-530.
23. Corbett, E.L. *et al* (2000) HIV infection and silicosis: The impact of two potent risk factors on the incidence of mycobacterial disease in South African miners. *AIDS* 14(17): 2759-2768.
24. Hnizdo, E. and Murray, J. (1998) Risk of pulmonary tuberculosis relative to silicosis and exposure to silica dust in South African gold miners. *Occupational and Environmental Medicine* 55(7): 496-502.
25. Lurie, M.N. *et al* (2003) The impact of migration on HIV-1 transmission in South Africa: A study of migrant and non-migrant men and their partners. *Sexually Transmitted Diseases* 30(2): 149-156.



26. Hargrove, J. (2008) Migration, mines and mores: The HIV epidemic in southern Africa. *South African Journal of Science* 104: 53–61.
27. Campbell, C. (1997) Migrancy, masculine identities and AIDS: The psychosocial context of HIV transmission on the South African gold mines. *Social Science & Medicine* 45(2): 273–281.
28. Corbett, E.L. *et al* (2004) Human immunodeficiency virus and the prevalence of undiagnosed tuberculosis in African gold miners. *American Journal of Respiratory and Critical Care Medicine* 170(6): 673–679.
29. Department of Health. (2008) *Tuberculosis Strategic Plan for South Africa, 2007–2011*. Pretoria, South Africa: Department of Health, Republic of South Africa.
30. Murray, J., Sonnenberg, P., Nelson, G., Bester, A., Shearer, S. and Glynn, J.R. (2007) Cause of death and presence of respiratory disease at autopsy in an HIV-1 seroconversion cohort of southern African gold miners. *AIDS* 21(Suppl 6): S97–S104.
31. Murray, J. *et al* (2005) Effect of HIV on work-related injury rates in South African gold miners. *AIDS* 19(17): 2019–2024.
32. Glynn, J.R., Murray, J., Bester, A., Nelson, G., Shearer, S. and Sonnenberg, P. (2008) Effects of duration of HIV infection and secondary tuberculosis transmission on tuberculosis incidence in the South African gold mines. *AIDS* 22(14): 1859–1867.
33. Brink, B. and Clausen, L. (1987) The acquired immune deficiency syndrome. *Proceedings of the Mine Medical Officers' Association* 63: 10–17.
34. van Halsema, C.L. *et al* (2010) Tuberculosis outcomes and drug susceptibility in individuals exposed to isoniazid preventive therapy in a high HIV prevalence setting. *AIDS* 24(7): 1051–1055.
35. Churchyard, G.J. *et al* (2003) Efficacy of secondary isoniazid preventive therapy among HIV-infected Southern Africans: Time to change policy? *AIDS* 17(14): 2063–2070.
36. Grant, A.D. *et al* (2005) Effect of routine isoniazid preventive therapy on tuberculosis incidence among HIV-infected men in South Africa: A novel randomized incremental recruitment study. *Journal of the American Medical Association* 293(22): 2719–2725.
37. Calver, A.D. *et al* (2010) Emergence of increased resistance and extensively drug-resistant tuberculosis despite treatment adherence, South Africa. *Emerging Infectious Diseases* 16(2): 264–271.
38. Corbett, E.L. *et al* (2003) Stable incidence rates of tuberculosis (TB) among human immunodeficiency virus (HIV)-negative South African gold miners during a decade of epidemic HIV-associated TB. *Journal of Infectious Diseases* 188(8): 1156–1163.
39. Godfrey-Faussett, P. *et al* (2000) Tuberculosis control and molecular epidemiology in a South African gold-mining community. *Lancet* 356(9235): 1066–1071.
40. Sonnenberg, P., Glynn, J.R., Fielding, K., Murray, J., Godfrey-Faussett, P. and Shearer, S. (2004) HIV and pulmonary tuberculosis: The impact goes beyond those infected with HIV. *AIDS* 18(4): 657–662.
41. Sonnenberg, P., Murray, J., Glynn, J.R., Shearer, S., Kambashi, B. and Godfrey-Faussett, P. (2001) HIV-1 and recurrence, relapse, and reinfection of tuberculosis after cure: A cohort study in South African mineworkers. *Lancet* 358(9294): 1687–1693.
42. Girdler-Brown, B.V., Murray, J.K.W., Thooe, S. and Godfrey-Faussett, P. (2004) *Development of Sensitive Tools for Active Case Finding of Tuberculosis (Phase 1) (SIM 03-08-02)*. Johannesburg: Safety in Mines Research Advisory Committee 2004.
43. Lowe, R.E. and Murray, J. (1992) Tuberculosis case finding. *Proceedings of the Mine Medical Officers' Association of South Africa* 65: 31–38.
44. Lewis, J.J. *et al* (2009) HIV infection does not affect active case finding of tuberculosis in South African gold miners. *American Journal of Respiratory and Critical Care Medicine* 180(12): 1271–1278.

45. Mqoqi, N.P., Churchyard, G.A., Kleinschmidt, I. and Williams, B. (1997) Attendance versus compliance with tuberculosis treatment in an occupational setting – A pilot study. *South African Medical Journal* 87(11): 1517–1521.
46. Sonnenberg, P., Ross, M.H., Shearer, S.C. and Murray, J. (1998) The effect of dosage cards on compliance with directly observed tuberculosis therapy in hospital. *International Journal of Tuberculosis and Lung Disease* 2(2): 168–171.
47. Murray, J., Sonnenberg, P., Shearer, S.C. and Godfrey-Faussett, P. (1999) Human immunodeficiency virus and the outcome of treatment for new and recurrent pulmonary tuberculosis in African patients. *American Journal of Respiratory and Critical Care Medicine* 159(3): 733–740.
48. Murray, J., Back, P., Coetzee, L. and Lowe, J.P. (2000) Clinico-Pathological Study to reduce the rate of missed and misdiagnosis of Pulmonary Tuberculosis in the South African Mining Industry. Final Report. HEALTH 611: SIMRAC, Braamfontein.
49. Churchyard, G.J., Corbett, E.L., Kleinschmidt, I., Mulder, D. and De Cock, K.M. (2000) Drug-resistant tuberculosis in South African gold miners: Incidence and associated factors. *International Journal of Tuberculosis and Lung Disease* 4(5): 433–440.
50. Murray, J., Sonnenberg, P., Shearer, S. and Godfrey-Faussett, P. (2000) Drug-resistant pulmonary tuberculosis in a cohort of southern African gold-miners with a high prevalence of HIV infection. *South African Medical Journal* 90(4): 381–386.
51. Mallory, K.F., Churchyard, G.J., Kleinschmidt, I., De Cock, K.M. and Corbett, E.L. (2000) The impact of HIV infection on recurrence of tuberculosis in South African gold miners. *International Journal of Tuberculosis and Lung Disease* 4(5): 455–462.
52. Charalambous, S. et al (2008) Contribution of reinfection to recurrent tuberculosis in South African gold miners. *International Journal of Tuberculosis and Lung Disease* 12(8): 942–948.
53. Glynn, J.R., Murray, J., Bester, A., Nelson, G., Shearer, S. and Sonnenberg, P. (2010) High rates of recurrence in HIV-infected and HIV-uninfected patients with tuberculosis. *Journal of Infectious Diseases* 201(5): 704–711.
54. Murray, J., Wong, M.L., Hopley, M.J. and Lowe, P.J. (2002) Process-based performance review for the diagnosis of pulmonary tuberculosis. Technology transfer of SIMRAC project Health 611 to enhance clinical performance.
55. Page-Shipp, L., Murray, J., Churchyard, G., Girdler-Brown, B., Sonnenberg, P. and Chicksen, E. (2010) *Development of a TB Programme Review Tool for the South African Mining Industry*, February 2010 Revision Johannesburg: Mine Health and Safety Council.
56. Calver, A. (2008) Miners' compensation: Who cares? *Labour Bulletin* 32(4): 26–28.
57. Davies, J.C. (2001) Silicosis and tuberculosis among South African goldminers – An overview of recent studies and current issues. *South African Medical Journal* 91(7): 562–566.
58. Gonsalves, G. and Akugizibwe, P. (2008) Migrants and TB in Southern Africa: Reaching across borders. *Labour Bulletin* 32(4): 29–31.
59. Grainger, L. (2010) Silicosis compensation editorial. *Occupational Health Southern Africa* 16(2): 2.
60. Trapido, A. and Goode, R. (1999) Polluters should pay. *South African Labour Bulletin* 23: 53–58.
61. White, N. (1997) Dust related diseases in former miners-the ODMWA legacy. *Occupational Health South Africa* 3(4): 20–24.
62. Basu, S., Stuckler, D., Gonsalves, G. and Lurie, M. (2009) The production of consumption: Addressing the impact of mineral mining on tuberculosis in southern Africa. *Globalization and Health* 5: 11.
63. Trapido, A., Goode, R. and White, N. (1998) Costs of occupational lung disease in South African gold mining. *Journal of Mineral Policy, Business and Environment* 13(2): 26–33.



64. Murray, J., Coetzee, L., Back, P., Banyini, A. and Ross, M. (2002) Analysis of occupational lung disease identified at autopsy and compensated in the South African mining industry. *Occupational Health South Africa* 8: 3–5.
65. Roberts, J. (2009) *The Hidden Epidemic amongst Former Miners: Silicosis, Tuberculosis and the Occupational Diseases in Mines and Works Act in the Eastern Cape, South Africa*. Westville, South Africa: Health Systems Trust.
66. Maiphethlo, L. (2010) Claims experience of former gold miners with silicosis – A clinic series. *Occupational Health Southern Africa* 16(2): 10–16.
67. Murray, J. and Rees, D. (2003) Tuberculosis and silicosis in South African gold mining. *Occupational Safety, Health and Development* 5: 2–10.
68. AngloGold Ashanti. (2007) *Country Report South Africa West Wits operations: AngloGold Ashanti*, http://www.anglogold.co.za/NR/rdonlyres/576A5C42-3524-49EE-A4A2-9A4D161572Do0/west_wits.pdf, accessed 28 April 2011.
69. Banyini, A. (2006) Simrac 03-06-03. *Occupational Health Southern Africa* 12(1): 18–21.
70. Stanton, D. (2006) Best practices on silicosis prevention. *Occupational Health Southern Africa* 12(1): 13.
71. Rees, D., Murray, J. and Ingham, F. (2009) Silicosis Elimination Programme (Track C). Silicosis elimination awareness for persons affected by mining operations in South Africa (SIM 03 06 03). Johannesburg: Mine Health and Safety Council.
72. Kemsley, D.M. (2008) *Respirable Dust and Quartz Exposure of Rock Drill Operators in Two Free State Gold Mines*. Johannesburg: University of the Witwatersrand.
73. Department of Health, South Africa. (2010) *Report on Tuberculosis in the Mining Industry*. Pretoria, South Africa: Department of Health.
74. Brink, B. and Pienaar, J. (2007) Business and HIV/AIDS: the case of Anglo American. *AIDS* 21(Suppl 3): S79–S84.
75. AIDS and Rights Alliance for Southern Africa (ARASA). (2008) *The Mining Sector, Tuberculosis and Migrant Labour in Southern Africa: Policy and Programmatic Interventions for the Cross-Border Control of Tuberculosis between Lesotho and South Africa, Focusing on Miners, Ex-Miners and Their Families*, http://arasa.info/sites/default/files/pub_ARASA_Mines_TB_and_Southern_Africa.pdf, accessed 28 April 2011.
76. Department of Mineral Resources. (2008) *Presidential Audit on Health and Safety in the Mining Industry*. Pretoria, South Africa: Department of Mineral Resources.
77. Stuckler, D., Basu, S. and McKee, M. (2010) Governance of mining, HIV and tuberculosis in Southern Africa. *Global Health Governance* IV(1), http://www.ghgj.org/Stuckler_final.pdf, accessed 28 April 2011.