

OCCURRENCE OF *SERGENTOMYA (PARROTOMYIA) BARRAUDI* (SINTON, 1929) IN MACAU

ALVES-PIRES C.*, RAMOS H.C.**, RIBEIRO H.* & EASTON E.R.***

Summary:

The presence of phlebotomine sandflies (*Sergentomyia barraudi*) in Macau, a Territory in Southern China under Portuguese administration, is here recorded for the first time. Nineteen specimens (14 males and 5 females) were examined for the most important features which were shown to agree with known descriptions of the species.

KEY WORDS : phlebotomine sandflies, *Sergentomyia barraudi*, Macau, distribution.

Résumé : PRÉSENCE DE *SERGENTOMYA BARRAUDI* (SINTON, 1929) À MACAO

Les auteurs signalent pour la première fois la présence des phlébotomes à Macao, Territoire du Sud de la Chine sous administration portugaise. Dix-neuf adultes (14 mâles et 5 femelles) ont été capturés dans une grotte artificielle en septembre 1994. Les caractères observés sont conformes à la description de l'espèce *Sergentomyia barraudi* (Sinton, 1929).

MOTS CLÉS : phlébotomes, *Sergentomyia barraudi*, Macao, distribution.

INTRODUCTION

S. barraudi was described by Sinton (1929) and later redescribed by Raynal & Gaschen (1934, 1935), from adults caught in India and Vietnam, respectively.

This species has a wide distribution in the Oriental Region (Lewis, 1978).

In Macau, a Territory in Southern China presently under Portuguese administration, phlebotomine sandflies have not yet been recorded, although *S. barraudi* and *S. indica* are known in near by Hong Kong (Lewis, 1978; Leng & Zhang, 1993), while more than forty species are known in China (Leng & Zhang, 1993).

It is presumed that *S. barraudi* is not responsible for the transmission of leishmaniases (Killick-Kendrick, 1990).

The wings, head and abdomen were cut, separated from the body, mounted on a slide in Berlese's medium (Léger *et al.*, 1983), and observed in light microscopy under Nomarsky contrast.

The identification was based on keys of Lewis (1978) and Lane (1993), and the descriptions by Sinton (1929), Raynal & Gaschen (1934, 1935), Parrot & Clastrier (1952), and Cates & Lien (1970).

The terminology used in this paper is the same as that of Lewis (1978).

RESULTS

The 5 females had : antenna 3 > A4 + A5, A3 = 181.4 ± 11.8 µm (mean ± SD) and A4 + A5 = 176.8 ± 4.6 µm (mean ± SD); cibarium with about 60 teeth and a bifid pigmented patch at the tip (Fig. 1a); pharynx with lamp-glass shape; smooth and elliptical spermathecae (Fig. 1b) with a length of 67.5 ± 2.5 µm (mean ± SD) and a width of 25.4 ± 1.3 µm (mean ± SD).

In the 14 males, we observed that antenna 3 ≥ A4 + A5, A3 = 215.6 ± 12.2 µm (mean ± SD) and A4 + A5 = 211.1 ± 10.5 µm (mean ± SD); style with four apical spines and four times as long as thick (Fig. 1c); cibarium with about 16 teeth; aedeagus slender and with rounded tip, not sclerotized (Fig. 1d), and hooked paramere (Fig. 1e).

MATERIAL AND METHODS

Nineteen adults (14 males and 5 females) were caught with an electric aspirator by the last three authors in an artificial cave, in Taipa's island - "trilho da Taipa" - (Macau), and preserved in 70 % ethanol.

* Disciplina de Entomologia Médica, Instituto de Higiene e Medicina Tropical, R. da Junqueira 96, P-1300 Lisboa.

** Centro de Zoologia, Instituto de Investigação Científica Tropical, R. da Junqueira 14, P-1300 Lisboa.

*** Universidade de Macau, PO Box 3001, Macau.




Fig. 1. — *Sergentomyia barraudi* from Macau; a. : cibarium, b. : spermathecae, c. : style, d. : aedeagus, e. : paremere.

DISCUSSION AND CONCLUSION

FEMALES

The presence of an antennal segment A3 longer than A4 + A5, and the value of the ratio between length and width of the spermathecae (2.7) were observed by Sinton (1929), Raynal & Gaschen (1934) and Cates & Lien (1970).

We found about 60 cibarial teeth in our five females, a little more (40-50) than observed by Sinton (*op. cit.*), Raynal & Gaschen (1934) and Parrot & Clastrier (1952), but similar to the values cited by Lewis (1978). The lamp-glass shaped pharynx and a vein R1 shorter than R2 + R3 in the wings, are similar to the observations of Sinton (1929), Raynal & Gaschen (1934) and Cates & Lien (1970).

The bifid shape of the pigmented patch of cibarium in the females, characteristic of this species (Sinton, 1929), was present in our five females.

MALES

The style with four apical spines, with length about four times the width, the hooked paramere, and a cibarium with about 16 teeth, as seen in our 14 males, are also characteristic of *Sergentomyia barraudi* (Sinton, 1929; Raynal & Gaschen, 1934 and Cates & Lien, 1970).

ACKNOWLEDGEMENT

We are grateful to Pr. R. Killick-Kendrick, Imperial College at Silwood Park, for critical examination of the manuscript.

REFERENCES

- CATES M.D. & LIEN J.C. The *Phlebotomus* of Taiwan. *Journal of Medical Entomology*, 1970, 7, 529-543.
- KILICK-KENDRICK R. Phlebotomine vectors of the leishmaniases: a review. *Medical and Veterinary Entomology*, 1990, 4, 1-24.
- LANE R.P. Sandflies (Phlebotominae). In: *Medical Insects and Arachnids*. Lane R.P. & Crosskey R.W. (eds). Published by Chapman & Hall, Londres, 1993, 78-119.
- LÉGER N., PESSON B., MADULO-LEBLOND G. & ABONNENC E. Sur la différenciation des femelles du sous-genre *Larroussius* Nitzulescu, 1931 (Diptera: Phlebotominae) de la région méditerranéenne. *Annales de Parasitologie Humaine et Comparée*, 1983, 58, 611-623.
- LENG Y.J. & ZHANG L.M. Check list and geographical distribution of phlebotomine sandflies. *Annals of Tropical Medicine and Parasitology*, 1993, 87, 83-94.

LEWIS D.J. The phlebotominae sandflies (Diptera: Psychodidae) of the Oriental Region. *Bulletin of the British Museum (Natural History), Entomology series*, 1978, 45, 121-209.

PARROT L. & CLASTRIER J. Notes sur les phlébotomes. LXV. - Phlébotomes d'Indochine. *Archives de l'Institut Pasteur d'Algérie*, 1952, 30, 153-171.

RAYNAL J. & GASCHEN H. Sur les phlébotomes d'Indochine. I. Présence de *Plebotomus barraudi* ♀ au Tonkin. *Bulletin de la Société de Pathologie Exotique*, 1934, 27, 559-563.

RAYNAL J. & GASCHEN H. Sur les phlébotomes d'Indochine. V. Présence de *Plebotomus barraudi*, Sinton 1929, dans le Haut-Bassin du Fleuve-Rouge et description de *Plebotomus barraudi* ♂. *Bulletin de la Société de Pathologie Exotique*, 1935, 28, 117-118.

SINTON J.A. Notes on some Indian species of the genus *Phlebotomus*. Part XXIV. *Phlebotomus barraudi* n. sp. *Indian Journal of Medical Research*, 1929, 16, 716-724.

Reçu le 3 octobre 1995

Accepté le 11 décembre 1995