

Odd Crossing Number and Crossing Number Are Not the Same

Michael J. Pelsmajer · Marcus Schaefer ·
Daniel Štefankovič

Abstract The *crossing number* of a graph is the minimum number of edge intersections in a plane drawing of a graph, where each intersection is counted separately. If instead we count the number of pairs of edges that intersect an odd number of times, we obtain the *odd crossing number*. We show that there is a graph for which these two concepts differ, answering a well-known open question on crossing numbers. To derive the result we study drawings of maps (graphs with rotation systems).

1 A Confusion of Crossing Numbers

Intuitively, the crossing number of a graph is the smallest number of edge crossings in any plane drawing of the graph. As it turns out, this definition leaves room for interpretation, depending on how we answer the questions: what is a drawing, what is a crossing, and how do we count crossings? The papers by Pach and Tóth [7] and Székely [9] discuss the historical development of various interpretations and definitions—often implicit—of the crossing number concept.

A *drawing* D of a graph G is a mapping of the vertices and edges of G to the Euclidean plane, associating a distinct point with each vertex, and a simple plane curve with each edge so that the ends of an edge map to the endpoints of the corresponding curve. For simplicity, we also require that

M.J. Pelsmajer (✉)

Department of Applied Mathematics, Illinois Institute of Technology, Chicago, IL 60616, USA
e-mail: pelsmajer@iit.edu

M. Schaefer

Department of Computer Science, DePaul University, Chicago, IL 60604, USA
e-mail: mschaefer@cs.depaul.edu

D. Štefankovič

Computer Science Department, University of Rochester, Rochester, NY 14627-0226, USA
e-mail: stefanko@cs.rochester.edu

- A curve does not contain any endpoints of other curves in its interior
- Two curves do not touch (that is, intersect without crossing), and
- No more than two curves intersect in a point (other than at a shared endpoint)

In such a drawing the intersection of the interiors of two curves is called a *crossing*. Note that by the restrictions we placed on a drawing, crossings do not involve endpoints, and at most two curves can intersect in a crossing. We often identify a drawing with the graph it represents. For a drawing D of a graph G in the plane we define

- $\text{cr}(D)$ - the total number of crossings in D
- $\text{pcr}(D)$ - the number of pairs of edges which cross at least once; and
- $\text{ocr}(D)$ - the number of pairs of edges which cross an odd number of times

Remark 1 For any drawing D , we have $\text{ocr}(D) \leq \text{pcr}(D) \leq \text{cr}(D)$.

We let $\text{cr}(G) = \min \text{cr}(D)$, where the minimum is taken over all drawings D of G in the plane. We define $\text{ocr}(G)$ and $\text{pcr}(G)$ analogously.

Remark 2 For any graph G , we have $\text{ocr}(G) \leq \text{pcr}(G) \leq \text{cr}(G)$.

The question (first asked by Pach and Tóth [7]) is whether the inequalities are actually equalities.¹ Pach [6] called this “perhaps the most exciting open problem in the area.” The only evidence for equality is an old theorem by Chojnacki, which was later rediscovered by Tutte—and the absence of any counterexamples.

Theorem 1.1 (Chojnacki [4], Tutte [10]) *If $\text{ocr}(G) = 0$, then $\text{cr}(G) = 0$.*²

In this paper we will construct a simple example of a graph with $\text{ocr}(G) < \text{pcr}(G) = \text{cr}(G)$. We derive this example from studying what we call weighted maps on the annulus. Section 2 introduces the notion of weighted maps on arbitrary surfaces and gives a counterexample to $\text{ocr}(M) = \text{pcr}(M)$ for maps on the annulus. In Section 3 we continue the study of crossing numbers for weighted maps, proving in particular that $\text{cr}(M) \leq c_n \cdot \text{ocr}(M)$ for maps on a plane with n holes. One of the difficulties in dealing with the crossing number is that it is **NP**-complete [2]. In Section 4 we show that the crossing number can be computed in polynomial time for maps on the annulus. Finally, in Section 5 we show how to translate the map counterexample from Section 2 into an infinite family of simple graphs for which $\text{ocr}(G) < \text{pcr}(G)$.


2 Map Crossing Numbers

A *weighted map* M is a surface S and a set $P = \{(a_1, b_1), \dots, (a_m, b_m)\}$ of pairs of distinct points on ∂S with positive weights w_1, \dots, w_m . A *realization* R of the map

¹Doug West lists the problem on his page of open problems in graph theory [12]. Dan Archdeacon even conjectured that equality holds [1].

²In fact they proved something stronger, namely that in any drawing of a non-planar graph there are two non-adjacent edges that cross an odd number of times. Also see [8].

Fig. 1 Optimal drawings: pcr and cr (above left), ocr (above right)



$M = (S, P)$ is a set of m properly embedded arcs $\gamma_1, \dots, \gamma_m$ in S where γ_i connects a_i and b_i .³

Let

$$\begin{aligned} \text{cr}(R) &= \sum_{1 \leq k < \ell \leq m} \iota(\gamma_k, \gamma_\ell) w_k w_\ell, \\ \text{pcr}(R) &= \sum_{1 \leq k < \ell \leq m} [\iota(\gamma_k, \gamma_\ell) > 0] w_k w_\ell, \\ \text{ocr}(R) &= \sum_{1 \leq k < \ell \leq m} [\iota(\gamma_k, \gamma_\ell) \equiv 1 \pmod{2}] w_k w_\ell, \end{aligned}$$

where $\iota(\gamma, \gamma')$ is the geometric intersection number of γ and γ' and $[x]$ is 1 if the condition x is true, and 0 otherwise. We define $\text{cr}(M) = \min \text{cr}(R)$, where the minimum is taken over all realizations R of M . We define $\text{pcr}(M)$ and $\text{ocr}(M)$ analogously.

Remark 3 For every map M , $\text{ocr}(M) \leq \text{pcr}(M) \leq \text{cr}(M)$.

Conjecture 1 For every map M , $\text{cr}(M) = \text{pcr}(M)$.

Lemma 2.1 If Conjecture 1 is true, then $\text{cr}(G) = \text{pcr}(G)$ for every graph G .

Proof Let D be a drawing of G with minimal pair crossing number. Drill small holes at the vertices. We obtain a drawing R of a weighted map M . If Conjecture 1 is true, there exists a drawing of M with the same crossing number. Collapse the holes to vertices to obtain a drawing D' of G with $\text{cr}(D') \leq \text{pcr}(G)$. \square

However, we show below that we can separate the odd crossing number from the crossing number for weighted maps, even in the annulus (a disk with a hole).

When analyzing crossing numbers of drawings on the annulus, we describe curves with respect to an initial drawing of the curve and a number of *Dehn twists*. Consider, for example, the four curves in the left part of Figure 1. Comparing them to the

³If we take a realization R of a map M , and contract each boundary component to a vertex, we obtain a drawing of a graph with a given rotation system [3]. For our purposes, maps are a more visual way to look at graphs with a rotation system.

corresponding curves in the right part, we see that the curves labeled c and d have not changed, but the curves labeled a and b have each undergone a single clockwise twist.

Two curves are *isotopic rel boundary* if they can be obtained from each other by a continuous deformation which does not move the boundary ∂M . Isotopy rel boundary is an equivalence relation, its equivalence classes are called *isotopy classes*. An *isotopy class* on the annulus is determined by a properly embedded arc connecting the endpoints, together with the number of twists performed.

Lemma 2.2 *Let $a \leq b \leq c \leq d$ be such that $a + c \geq d$. For the weighted map M in Figure 1 we have $\text{cr}(M) = \text{pcr}(M) = ac + bd$ and $\text{ocr}(M) = bc + ad$.*

Proof The upper bounds follow from the drawings in Figure 1, the left drawing for crossing and pair crossing number, the right drawing for odd crossing number.

Claim $\text{pcr}(M) \geq ac + bd$.

Proof of the Claim Let R be a drawing of M minimizing $\text{pcr}(R)$. We can apply twists so that the thick edge d is drawn as in the left part of Figure 1. Let α, β, γ be the number of clockwise twists applied to the ends of arcs a, b, c on the inner boundary to obtain the drawing R , where $\alpha = \beta = \gamma = 0$ corresponds to the drawing shown in the left part of Figure 1. Then,

$$\begin{aligned} \text{pcr}(R) = & cd[\gamma \neq 0] + bd[\beta \neq -1] + ad[\alpha \neq 0] + bc[\beta \neq \gamma] \\ & + ab[\alpha \neq \beta] + ac[\alpha \neq \gamma + 1]. \end{aligned} \quad (1)$$

If $\gamma \neq 0$, then $\text{pcr}(R) \geq cd + ab$ because at least one of the last five conditions in (1) must be true; the last five terms contribute at least ab (since $d \geq c \geq b \geq a$), and the first term contributes cd . Since $d(c - b) \geq a(c - b)$, $cd + ab \geq ac + bd$, and the claim is proved in the case that $\gamma \neq 0$.

Now assume that $\gamma = 0$. Equation (1) becomes

$$\text{pcr}(R) = bd[\beta \neq -1] + bc[\beta \neq 0] + ad[\alpha \neq 0] + ac[\alpha \neq 1] + ab[\alpha \neq \beta]. \quad (2)$$

If $\beta \neq -1$, then $\text{pcr}(R) \geq bd + ac$ because either $\alpha \neq 0$ or $\alpha \neq 1$. Since $bd + ac \geq bc + ad$, the claim is proved in the case that $\beta \neq -1$.

This leaves us with the case that $\beta = -1$. Equation (2) becomes

$$\text{pcr}(R) = bc + ad[\alpha \neq 0] + ac[\alpha \neq 1] + ab[\alpha \neq -1]. \quad (3)$$

The right-hand side of Equation (3) is minimized for $\alpha = 0$. In this case $\text{pcr}(R) = bc + ac + ab \geq ac + bd$ because we assume that $a + c \geq d$. \square

Claim $\text{ocr}(M) \geq bc + ad$.

Proof of the Claim Let R be a drawing of M minimizing $\text{ocr}(R)$. Let α, β, γ be as in the previous claim. We have

$$\text{ocr}(R) = cd[\gamma]_2 + bd[\beta + 1]_2 + ad[\alpha]_2 + bc[\beta + \gamma]_2 + ab[\alpha + \beta]_2 + ac[\alpha + \gamma + 1]_2, \quad (4)$$

where $[x]_2$ is 0 if $x \equiv 0 \pmod{2}$, and 1 otherwise.

If $\beta \not\equiv \gamma \pmod{2}$, then the claim clearly follows unless $\gamma = 0$, $\beta = 1$, and $\alpha = 0$ (all modulo 2). In that case $\text{ocr}(R) \geq bc + ab + ac \geq bc + ad$. Hence, the claim is proved if $\beta \not\equiv \gamma \pmod{2}$.

Assume then that $\beta \equiv \gamma \pmod{2}$. Equation (4) becomes

$$\text{ocr}(R) = cd[\beta]_2 + bd[\beta + 1]_2 + ad[\alpha]_2 + ab[\alpha + \beta]_2 + ac[\alpha + \beta + 1]_2. \quad (5)$$

If $\alpha \equiv 1 \pmod{2}$, then the claim clearly follows because either cd or bd contributes to the ocr . Thus we can assume $\alpha \equiv 0 \pmod{2}$. Equation (5) becomes

$$\text{ocr}(R) = (cd + ab)[\beta]_2 + (bd + ac)[\beta + 1]_2. \quad (6)$$

For both $\beta \equiv 0 \pmod{2}$ and $\beta \equiv 1 \pmod{2}$ we get $\text{ocr}(R) \geq bc + ad$. □

We get a separation of pcr and ocr for maps with small integral weights.

Corollary 2.3 *There is a weighted map M on the annulus with edges of weight $a = 1$, $b = c = 3$, and $d = 4$ for which $\text{cr}(M) = \text{pcr}(M) = 15$ and $\text{ocr}(M) = 13$.*

Optimizing the gap over the reals yields $b = c = 1$, $a = (\sqrt{3} - 1)/2$, and $d = 1 + a$, giving us the following separation of $\text{pcr}(M)$ and $\text{ocr}(M)$.

Corollary 2.4 *There exists a weighted map M on the annulus with $\text{ocr}(M) \leq \sqrt{3}/2 \text{pcr}(M)$.*

Conjecture 2 *For every weighted map M on the annulus, $\text{ocr}(M) \geq \frac{\sqrt{3}}{2} \text{pcr}(M)$.*

3 Upper Bounds on Crossing Numbers

In Section 5 we will transform the separation of ocr and pcr on maps into a separation on graphs. In particular, we will show that for every $\varepsilon > 0$ there is a graph G so that

$$\text{ocr}(G) < (\sqrt{3}/2 + \varepsilon) \text{cr}(G).$$

The gap cannot be arbitrarily large, as Pach and Tóth showed.

Theorem 3.1 (Pach and Tóth [7]) *Let G be a graph. Then $\text{cr}(G) \leq 2(\text{ocr}(G))^2$.*⁴

This result suggests the question whether the linear separation can be improved. We do not believe this to be possible:

⁴Better upper bounds on $\text{cr}(G)$ in terms of $\text{pcr}(G)$ are known [5, 11].




Fig. 2 Pulling an endpoint (*left*) and contracting the edge (*right*)

Conjecture 3 *There is a $c > 0$ so that $\text{cr}(G) < c \cdot \text{ocr}(G)$.*

In this section, we will show that our approach of comparing the different crossing numbers for maps with a fixed number of holes will not lead to a super-linear separation. Namely, for a (weighted) map M on a plane with n holes, we always have

$$\text{cr}(M) \leq \text{ocr}(M) \binom{n+4}{4} / 5, \quad (7)$$

with strict inequality if $n > 1$. It follows that for fixed n , there *is* only a constant factor separating $\text{cr}(M)$ and $\text{ocr}(M)$. And only fixed, small n are computationally feasible in analyzing potential counterexamples.

Observe that as a special case of Equation (7), if M is a (weighted) map on the annulus ($n = 2$) we get that $\text{cr}(M) < 3 \text{ocr}(M)$, which comes reasonably close to the $\sqrt{3}/2$ lower bound from the previous section.

Before proving Equation (7) in full generality, we first consider the case of unit weights.

For this section only, we will switch our point of view from maps as curves between holes on a plane to maps as graphs with a rotation system; that is, we contract each hole to a vertex, and record the order, in which the curves (edges) leave the vertex. Our basic operation will be the contraction of an edge by pulling one of its endpoints along the edge, until it coincides with the other endpoint (the rotations of the vertices merge). Figure 2 illustrates pulling v towards u along uv .⁵

Consider a drawing of G with the minimum number of odd pairs (edge pairs that cross an odd number of times), $\text{ocr}(G)$. We want to contract edges without creating too many new odd pairs. For each edge e , let o_e be the number of edges that cross e an odd number of times. Then $\sum_{e \in E(G)} o_e = 2 \text{ocr}(G)$, and since each edge is incident to exactly two vertices,

$$\sum_{v \in V(G)} \sum_{e \ni v} o_e = 4 \text{ocr}(G).$$

Applying the pigeonhole principle twice, there must be a vertex $v \in V(G)$ with $\sum_{e \ni v} o_e \leq 4 \text{ocr}(G)/n$, and there is a non-loop edge e incident to v with $o_e \leq 4 \text{ocr}(G)/(n \cdot d^*(v))$ (where $d^*(v)$ counts the number of non-loop edges incident to v). Contracting v to its neighbor along e creates at most $o_e(d^*(v) - 1) <$

⁵The illustration is taken from [8], where we investigate some other uses of this operation for graph drawings.

$4\text{ocr}(G)/n$ odd pairs (only edges intersecting e oddly will lead to odd intersections, and the parity of intersection along loops with endpoint v does not change; self-intersections can be removed). Repeating this operation $n - 1$ times, we transform G into a bouquet of loops at a single vertex with at most $\text{ocr}(G) \prod_{i=0}^{n-2} (1 + 4/(n - i))$ odd pairs (strictly less if $n > 1$). Without changing the rotation of the vertex, we can redraw all loops so that each odd pair intersects exactly once, and other pairs do not cross at all. We can then undo the contractions of the edges in reverse order without creating any new crossings. This yields a drawing of G with the original vertex rotations with at most $\text{ocr}(G) \prod_{i=0}^{n-2} (1 + 4/(n - i))$ crossings. Since the product term equals $\binom{n+4}{4}/5$, we have shown that $\text{cr}(G) \leq \text{ocr}(G) \binom{n+4}{4}/5$ (strict inequality for $n > 1$), as desired.

This argument proves Equation (7) for maps with unit weights. The next step is to extend this lemma to maps with arbitrary weights.

Consider two curves γ_1, γ_2 whose endpoints are adjacent and in the same order. In a drawing minimizing one of the crossing numbers we can always assume that the two curves are routed in parallel, following the curve that minimizes the total number of intersections with all curves other than γ_1 and γ_2 . The same argument holds for a block of curves with adjacent endpoints in the same order. This allows us to claim Equation (7) for maps with integer weights: a curve with integral weight w is replaced by w parallel duplicates of unit weight.

If we scale all the weights in a map M by a factor α , all the crossing numbers will change by a factor of α^2 . Hence, the case of rational weights can be reduced to integer weights. Finally, we observe that if we consider any of the crossing numbers as a function of the weights of M , this function is continuous: This is obvious for a fixed drawing of M , so it remains true if we minimize over a finite set of drawings of M . The maximum difference in the number of twists in an optimal drawing is bounded by a function of the crossing number; and thus it suffices to consider a finite set of drawings of M . We have shown:

Theorem 3.2 $\text{cr}(M) \leq \text{ocr}(M) \binom{n+4}{4}/5$ for weighted maps M on the plane with n holes.

4 Computing Crossing Numbers on the Annulus

Let M be a map on the annulus. We explained earlier that as far as crossing numbers are concerned, we can describe a curve in the realization of M by a properly embedded arc γ_{ab} connecting endpoints a and b on the inner and outer boundary of the annulus, and an integer $k \in \mathbb{Z}$, counting the number of twists applied to the curve γ_{ab} . Our goal is to compute the number of intersections between two arcs after applying a number of twists to each one of them. Since twists can be positive and negative and cancel each other out, we need to count crossings more carefully. Let us orient all arcs from the inner boundary to the outer boundary. Traveling along an arc α , a crossing with β counts as $+1$ if β crosses from right to left, and as -1 if it crosses from left to right. Summing up these numbers over all crossings for two arcs α and β yields

$\hat{\iota}(\alpha, \beta)$, the *algebraic crossing number* of α and β . Tutte [10] introduced the notion

$$\text{acr}(G) = \min_D \sum_{\{e, f\} \in \binom{E}{2}} |\hat{\iota}(\gamma_e, \gamma_f)|,$$

the *algebraic crossing number* of a graph, a notion that apparently has not drawn any attention since.

Let $D^k(\gamma)$ denote the result of adding k twists to the curve γ . For two curves α and β connecting the inner and outer boundary we have:

$$\hat{\iota}(D^k(\alpha), D^\ell(\beta)) = k - \ell + \hat{\iota}(\alpha, \beta). \quad (8)$$

Note that $\iota(\alpha, \beta) = |\hat{\iota}(\alpha, \beta)|$ for any two curves α, β on the annulus.

Let π be a permutation of $[n]$. A map M_π corresponding to π is constructed as follows. Choose $n + 1$ points on each of the two boundaries and number them $0, 1, \dots, n$ in the clockwise order. Let a_i be the vertex numbered i on the outer boundary and b_i be the vertex numbered π_i on the inner boundary, $i = 1, \dots, n$. We ask a_i to be connected to b_i in M_π .

We will encode a drawing R of M_π by a sequence of n integers x_1, \dots, x_n as follows. Fix a curve β connecting the a_0 and b_0 and choose γ_i so that $\iota(\beta, \gamma_i) = 0$ (for all i). We will connect a_i, b_i with the arc $D^{x_i}(\gamma_i)$ in R . Note that for $i < j$, $\hat{\iota}(\gamma_i, \gamma_j) = [\pi_i > \pi_j]$ and hence

$$\hat{\iota}(D^{x_i}(\gamma_i), D^{x_j}(\gamma_j)) = x_i - x_j + [\pi_i > \pi_j].$$

We have

$$\text{acr}(M_\pi) = \text{cr}(M_\pi) = \min \left\{ \sum_{i < j} |x_i - x_j + [\pi_i > \pi_j]| w_i w_j : x_i \in \mathbb{Z}, i \in [n] \right\}, \quad (9)$$

$$\text{pcr}(M_\pi) = \min \left\{ \sum_{i < j} [x_i - x_j + [\pi_i > \pi_j] \neq 0] w_i w_j : x_i \in \mathbb{Z}, i \in [n] \right\}, \quad (10)$$

and

$$\text{ocr}(M_\pi) = \min \left\{ \sum_{i < j} [x_i - x_j + [\pi_i > \pi_j] \not\equiv 0 \pmod{2}] w_i w_j : x_i \in \mathbb{Z}, i \in [n] \right\}. \quad (11)$$

Consider the relaxation of the integer program for $\text{cr}(M_\pi)$:

$$\text{cr}'(M_\pi) = \min \left\{ \sum_{i < j} |x_i - x_j + [\pi_i > \pi_j]| w_i w_j : x_i \in \mathbb{R}, i \in [n] \right\}. \quad (12)$$

Since (12) is a relaxation of (9), we have $\text{cr}'(M_\pi) \leq \text{cr}(M_\pi)$. The following lemma shows that $\text{cr}'(M_\pi) = \text{cr}(M_\pi)$.

Lemma 4.1 *Let n be a positive integer. Let $b_{ij} \in \mathbb{Z}$ and let $a_{ij} \in \mathbb{R}$ be non-negative, $1 \leq i < j \leq n$. Then*

$$\min \left\{ \sum_{i < j} a_{ij} |x_i - x_j + b_{ij}| : x_i \in \mathbb{R}, i \in [n] \right\}$$

has an optimal solution with $x_i \in \mathbb{Z}, i \in [n]$.

Proof Let \bar{x}^* be an optimal solution which satisfies the maximum number of $x_i - x_j + b_{ij} = 0, 1 \leq i < j \leq n$. Without loss of generality, we can assume $x_1^* = 0$. Let G be a graph on the vertex set $\{1, \dots, n\}$ with an edge between vertices i, j if $x_i^* - x_j^* + b_{ij} = 0$. Note that if i, j are connected by an edge and one of x_i^*, x_j^* is an integer, then both x_i^* and x_j^* are integers. It is then enough to show that G is connected.

Suppose that G is not connected. There exists a non-empty $A \subsetneq V(G)$ so that there are no edges between A and $V(G) - A$. Let χ_A be the characteristic vector of the set A , that is, $(\chi_A)_i = [i \in A]$. Let $f(\lambda)$ be the value of the objective function on $\bar{x} = \bar{x}^* + \lambda \cdot \chi_A$. Let I be the interval on which the signs of the $x_i - x_j + b_{ij}, 1 \leq i < j \leq n$ are the same as for \bar{x}^* . Then I is not the entire line (otherwise G would be connected). Since f is linear on I , f is optimal at $\lambda = 0$, and I contains a neighborhood of 0, it must be that f is constant on I . Choosing $x = x^* + \lambda \chi_A$ for λ an endpoint of I gives an optimal solution satisfying more $x_i - x_j + b_{ij} = 0, 1 \leq i < j \leq n$, a contradiction. \square

Theorem 4.2 *The crossing number of maps on the annulus can be computed in polynomial time.*

Proof Note that $cr'(M_\pi)$ is computed by the following linear program L_π :

$$\begin{aligned} \min \quad & \sum_{i < j} y_{ij} w_i w_j, \\ & y_{ij} \geq x_i - x_j + [\pi_i > \pi_j], \quad 1 \leq i < j \leq n, \\ & y_{ij} \geq -x_i + x_j - [\pi_i > \pi_j], \quad 1 \leq i < j \leq n. \end{aligned}$$

\square

Question 1 *Let M be a map on the annulus. Can $ocr(M)$ be computed in polynomial time?*


We conjectured earlier that crossing number and odd crossing number agree on maps. A more moderate goal would be to establish the following conjecture.

Conjecture 4 *For any map M on the annulus $cr(M) = pcr(M)$.*

5 Separating Crossing Numbers of Graphs

We modify the map from Lemma 2.2 to obtain a graph G separating $ocr(G)$ and $pcr(G)$. The graph G will have integral weights on edges. From G we can get an

Fig. 3 The inside flipped



unweighted graph G' with $\text{ocr}(G') = \text{ocr}(G)$ and $\text{pcr}(G') = \text{pcr}(G)$ by replacing an edge of weight w by w parallel edges of weight 1 (this does not change any of the crossing numbers). If needed we can get rid of parallel edges by subdividing edges, which does not change any of the crossing numbers.

We start with the map M from Lemma 2.2 with the following integral weights:

$$a = \left\lfloor \frac{\sqrt{3}-1}{2}m \right\rfloor, \quad b = c = m, \quad d = \left\lfloor \frac{\sqrt{3}+1}{2}m \right\rfloor,$$

where $m \in \mathbb{N}$ will be chosen later.

We replace each pair (a_i, b_i) of M by w_i pairs $(a_{i,1}, b_{i,1}), \dots, (a_{i,w_i}, b_{i,w_i})$ where the $a_{i,j}$ ($b_{i,j}$) occur on ∂S in clockwise order in a small interval around a_i (b_i). As before, we can argue that all the curves corresponding to (a_i, b_i) can be routed in parallel in an optimal drawing and, therefore, the resulting map N with unit weights will have the same crossing numbers as M .

We then replace the boundaries of the annulus by cycles (using one vertex for each $a_{i,j}$ and $b_{i,j}$), obtaining a graph G . We assign weight $W = 1 + \text{pcr}(N)$ to the edges in the cycles. This ensures that in a drawing of G minimizing any of the crossing numbers the boundary cycles are embedded without any intersections. Consequently, a drawing of G on the sphere that minimizes any one of the crossing numbers looks very much like the drawing of a map on the annulus. With one subtle difference: one of the boundaries may flip.

Given the map N on the annulus, the *flipped map* N' is obtained by flipping the order of the points on one of the boundaries. In other words, there are essentially two different ways of embedding the two boundary cycles of G on the sphere without intersections depending on the relative orientation of the boundaries. In one of the cases the drawing D of G gives a drawing of N , in the other case it gives a drawing of the flipped map N' . Fortunately, in the flipped case the group of edges corresponding to the weighted edge from a_i to b_i must intersect often with each other (as illustrated in Figure 3).

Now we know that

$$\begin{aligned} \text{ocr}(G) &\leq \text{ocr}(N) \quad (\text{since every drawing of } N \text{ is a drawing of } G) \\ &\leq w_1 w_3 + w_2 w_4 \quad (\text{by Lemma 2.2}) \\ &\leq \frac{3}{2}m^2 \quad (\text{by the choice of weights}). \end{aligned}$$

We will presently prove the following estimate on the flipped map.

Lemma 5.1 $\text{ocr}(N') \geq 2m^2 - 4m$.

With that estimate and our discussion of flipped maps, we have

$$\begin{aligned} \text{pcr}(G) &= \min\{\text{pcr}(N), \text{pcr}(N')\} \\ &\geq \min\{\text{pcr}(N), \text{ocr}(N')\} \quad (\text{since } \text{ocr} \leq \text{cr}) \\ &\geq \min\{\sqrt{3}m^2 - 2m, 2m^2 - 4m\} \quad (\text{choice of } w, \text{ and Lemma 5.1}). \end{aligned}$$

By making m sufficiently large, we can make the ratio of $\text{ocr}(G)$ and $\text{pcr}(G)$ arbitrarily close to $\sqrt{3}/2$.

Theorem 5.2 *For any $\varepsilon > 0$ there is a graph G such that*

$$\text{ocr}(G) < (\sqrt{3}/2 + \varepsilon) \text{pcr}(G).$$

The proof of Lemma 5.1 will require the following estimate.

Lemma 5.3 *Let $0 \leq a_1 \leq a_2 \leq \dots \leq a_n$ be such that $a_n \leq a_1 + \dots + a_{n-1}$. Then*

$$\max_{|y_i| \leq a_i} \left(\left(\sum_{i=1}^n y_i \right)^2 - 2 \sum_{i=1}^n y_i^2 \right) = \left(\sum_{i=1}^n a_i \right)^2 - 2 \sum_{i=1}^n a_i^2.$$

Proof of Lemma 5.1 Let $w_1 = a, w_2 = b, w_3 = d, w_4 = c$ (with a, b, c, d as in the definition of N). In any drawing of N' each group of the edges split into two classes, those with an even number of twists and those with an odd number of twists (two twists make the same contribution to $\text{ocr}(M')$ as no twists). Consequently, we can estimate $\text{ocr}(N')$ as follows.

$$\begin{aligned} \text{ocr}(N') &= \min_{k_i \in \{0, 1, \dots, w_i\}} \left(\sum_{i=1}^4 \binom{k_i}{2} + \sum_{i=1}^4 \binom{w_i - k_i}{2} + \sum_{i \neq j} k_i (w_j - k_j) \right) \\ &\geq -\frac{1}{2} \sum_{i=1}^4 w_i + \min_{0 \leq x_i \leq w_i} \left(\sum_{i=1}^4 \frac{x_i^2}{2} + \sum_{i=1}^4 \frac{(w_i - x_i)^2}{2} + \sum_{i \neq j} x_i (w_j - x_j) \right) \\ &= -\frac{1}{2} \sum_{i=1}^4 w_i + \frac{1}{4} \left(\sum_{i=1}^4 w_i \right)^2 + \min_{|y_i| \leq w_i/2} \left(2 \sum_{i=1}^4 y_i^2 - \left(\sum_{i=1}^4 y_i \right)^2 \right) \\ &\geq \frac{1}{2} \sum_{i=1}^4 w_i^2 - \frac{1}{2} \sum_{i=1}^4 w_i \quad (\text{using Lemma 5.3}) \\ &\geq \frac{1}{2} \left(\left(\frac{\sqrt{3}+1}{2} m - 1 \right)^2 + 2m^2 + \left(\frac{\sqrt{3}-1}{2} m - 1 \right)^2 - 4m \right) \\ &\geq 2m^2 - 4m. \end{aligned} \tag{13}$$

The equality between the second and third line can be verified by substituting $y_i = x_i - w_i/2$. \square

Proof of Lemma 5.3 Let y_1, \dots, y_n achieve the maximum value. Replacing the y_i by $|y_i|$ does not decrease the objective function. Without loss of generality, we can assume $0 \leq y_1 \leq y_2 \leq \dots \leq y_n$. Note that if $y_i < y_j$, then $y_i = a_i$ (otherwise increasing y_i by ε and decreasing y_j by ε increases the objective function for small ε).

Let k be the largest i such that $y_i = a_i$. Let $k = 0$ if no such i exists. We have $y_i = a_i$ for $i \leq k$ and $y_{k+1} = \dots = y_n$. If $k = n$ we are done.

We conclude the proof by showing that $k < n$ is not possible. Let t be the common value of $y_{k+1} = \dots = y_n$. Note that we have $t = y_{k+1} \leq a_{k+1}$.

Let

$$f(t) = \left(\left(\sum_{i=1}^k a_i \right) + (n-k)t \right)^2 - 2 \left(\left(\sum_{i=1}^k a_i^2 \right) + (n-k)t^2 \right).$$

We have

$$f'(t) = 2(n-k) \left(\left(\sum_{i=1}^k a_i \right) + (n-k-2)t \right).$$

Note that $f'(t) > 0$ for $t < a_{k+1}$. (This is easy to see when $k < n-1$; for $k = n-1$ we make use of the assumption that $a_n \leq \sum_{i=1}^{n-1} a_i$.) Therefore, $f(t)$ will be maximized by $t = a_{k+1}$ over values $t \leq a_{k+1}$. Hence, $y_{k+1} = a_{k+1}$, contradicting our choice of k . \square

6 Conclusion

The relationship between the different crossing numbers remains mysterious, and we have already mentioned several open questions and conjectures. Here we want to revive a question first asked by Tutte (in slightly different form). Recall the definition of the algebraic crossing number from Section 4:

$$\text{acr}(G) = \min_D \sum_{\{e,f\} \in \binom{E}{2}} |\hat{t}(\gamma_e, \gamma_f)|,$$

where γ_e is a curve representing edge e in a drawing D of G . It is clear that

$$\text{acr}(G) \leq \text{cr}(G).$$

Does equality hold?

Acknowledgement Thanks to the anonymous referee for helpful comments.

References

1. Archdeacon, D.: Problems in topological graph theory. <http://www.emba.uvm.edu/~archdeac/problems/altercross.html> (accessed April 7th, 2005)
2. Garey, M.R., Johnson, D.S.: Crossing number is NP-complete. *SIAM J. Algebr. Discrete Methods* **4**(3), 312–316 (1983)
3. Gross, J.L., Tucker, T.W.: *Topological Graph Theory*. Dover, Mineola (2001). Reprint of the 1987 original
4. Chojnacki, C. (Haim Hanani): Über wesentlich unplättbare Kurven im drei-dimensionalen Raume. *Fundam. Math.* **23**, 135–142 (1934)
5. Kolman, P., Matoušek, J.: Crossing number, pair-crossing number, and expansion. *J. Comb. Theory Ser. B* **92**(1), 99–113 (2004)
6. Pach, J.: Crossing numbers. In: *Discrete and Computational Geometry (Tokyo, 1998)*. Lecture Notes in Comput. Sci., vol. 1763, pp. 267–273. Springer, Berlin (2000)
7. Pach, J., Tóth, G.: Which crossing number is it anyway? *J. Comb. Theory Ser. B* **80**(2), 225–246 (2000)
8. Pelsmayer, M.J., Schaefer, M., Štefankovič, D.: Removing even crossings. In: *EuroComb*, April 2005
9. Székely, L.A.: A successful concept for measuring non-planarity of graphs: the crossing number. *Discrete Math.* **276**(1–3), 331–352 (2004). 6th International Conference on Graph Theory
10. Tutte, W.T.: Toward a theory of crossing numbers. *J. Comb. Theory* **8**, 45–53 (1970)
11. Valtr, P.: On the pair-crossing number. In: *Combinatorial and Computational Geometry*. MSRI Publications, vol. 52, pp. 545–551 (2005)
12. West, D.: Open problems—graph theory and combinatorics. <http://www.math.uiuc.edu/~west/openp/> (accessed April 7th, 2005)