

Received Date : 21-Jan-2016

Revised Date : 14-Jun-2016

Accepted Date : 14-Jun-2016

Article type : Review Article

Estrogens and progestagens: synthesis and action in the brain

Authors: Rossetti María F.^{1,2,*}; Cambiasso María J.^{3,4}; Holschbach M. Allie⁵; Cabrera Ricardo⁶.

Authors Affiliations:

¹ Departamento de Bioquímica Clínica, Facultad de Bioquímica y Ciencias Biológicas, Universidad Nacional del Litoral. Santa Fe, Argentina.

² Instituto de Salud y Ambiente del Litoral, CONICET-Universidad Nacional del Litoral. Santa Fe, Argentina.

³ Instituto de Investigación Médica Mercedes y Martín Ferreyra, INIMEC-CONICET-Universidad Nacional de Córdoba. Córdoba, Argentina.

⁴ Departamento de Biología Bucal, Facultad de Odontología -Universidad Nacional de Córdoba. Córdoba, Argentina.

⁵ Biomedical Sciences, Colorado State University. Fort Collins, Colorado, United States of America.

⁶ Instituto de Investigaciones Biomédicas. INBIOMED-IMBECU-CONICET. Universidad de Mendoza. Mendoza, Argentina.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi:

10.1111/jne.12402

This article is protected by copyright. All rights reserved.

Email addresses:

mfrossetti@fcb.unl.edu.ar

jcambiasso@immf.uncor.edu

maholsch@colostate.edu

ricabre@gmail.com

* Corresponding author. Departamento de Bioquímica Clínica, Facultad de Bioquímica y Ciencias Biológicas, Universidad Nacional del Litoral. Santa Fe, Argentina. E-mail: mfrossetti@fcb.unl.edu.ar.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Acknowledgement

We want to thank the International Workshop of Neuroendocrinology (IWNE) 2015.

Abbreviations

17 β -HSD: 17 β -hydroxysteroid dehydrogenase

3 α -HSD: 3 α -hydroxysteroid dehydrogenase

3 β -HSD: 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4-isomerase

5 α R: steroid 5 α -reductase

AD: Alzheimer's disease

Allop: Allopregnanolone

AP: activator protein

This article is protected by copyright. All rights reserved.

BST: Bed nucleus of the stria terminalis
DHEA: Dehydroepiandrosterone
DHEAS: Dehydroepiandrosterone sulfate
E2: Estradiol
ER: Estrogen receptor
ERE: Estrogen response element
GABA_A: Type-A γ -aminobutyric acid receptor
GPER1: G protein-coupled estrogen receptor 1
LH: Luteinizing hormone
MBP: Myelin basic protein
mPR: Membrane bound progesterone receptor
NF: Nuclear Factor
Ngn3: Neurogenin 3
NMDA: N-methyl-D-aspartate receptor
Oct-1: octamer-binding factor-1
OVX: Ovariectomized
P: Progesterone
P450arom: cytochrome P450arom
P450c17: 17 α -hydroxylase / C17-20 lyase
P450scc: cytochrome P450 side-chain cleavage
PGMRC1: Progesterone membrane receptor component 1
PR: Progesterone receptor
PRE: Progesterone response element
Preg: Pregnenolone
PregS: Pregnenolone sulfate
Sp: selective promoter factor
SREBP-: sterol regulatory element-binding protein
StAR: Steroidogenic acute regulatory protein

T: Testosterone

TBI: Traumatic brain injury

TFs: Transcription factors

TSPO: Translocator protein

Abstract

When steroids, such as pregnenolone, progesterone and estrogen, are synthesized *de novo* in neural tissues, they are more specifically referred to as neurosteroids. These neurosteroids bind specific receptors to promote essential brain functions. Pregnenolone supports cognition and protects mouse hippocampal cells against glutamate and amyloid peptide-induced cell death. Progesterone promotes myelination, spinogenesis, synaptogenesis, neuronal survival and dendritic growth. Allopregnanolone increases hippocampal neurogenesis, neuronal survival and cognitive functions. Estrogens, such as estradiol, regulate synaptic plasticity, reproductive behavior, aggressive behavior and learning. In addition, neurosteroids are neuroprotective in animal models of Alzheimer's disease, Parkinson's disease, brain injury, and aging. Using *in situ* hybridization and/or immunohistochemistry, steroidogenic enzymes, including cytochrome P450 side-chain cleavage, 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4 isomerase, cytochrome P450arom, steroid 5 α -reductase and 3 α -hydroxysteroid dehydrogenase, have been detected in numerous brain regions, including the hippocampus, hypothalamus and cerebral cortex. In this article, we summarize some of the studies related to synthesis and function of estrogens and progestagens in the central nervous system.

Keywords

Estrogen; progestagen; brain; neurosteroidogenic enzymes.

1. Introduction.

It is well known that steroid hormones regulate various functions associated with the brain including the development of the central and peripheral nervous system, the regulation of neurotransmitter systems, synaptic connectivity, dendritic branching, and myelination. They are also involved in cognition, emotion, mood, sexual behavior, and social behavior (1-12). Historically, the synthesis of these steroids was thought to be constrained to

This article is protected by copyright. All rights reserved.

steroidogenic glands such as gonads, adrenal cortex and placenta, and could then cross the blood-brain barrier to act in the brain (13). However, it is now well known that steroids are also synthesized *de novo* from cholesterol or steroidal precursors imported from peripheral sources in numerous brain regions, including the hippocampus, hypothalamus and cerebral cortex, by both neurons and glia. These locally synthesized hormones are called neurosteroids (13). Brain cholesterol is converted into pregnenolone (Preg) by the cholesterol side-chain cleavage enzyme (P450scc) in the mitochondria, as in peripheral steroidogenic organs. Subsequently, Preg is metabolized in the smooth endoplasmic reticulum into neurosteroids, such as progesterone (P), allopregnanolone (Allop), dehydroepiandrosterone (DHEA) or estradiol (E2) (14).

Neurosteroids regulate several central nervous system functions by binding their respective receptors and have key roles in neurodegenerative disease and aging. The present article focuses on two major groups of neurosteroids: estrogens (E2) and progestagens (P and Allop) and their common precursor Preg. Here, we summarize some of the studies related to the steroidogenic enzymes and classical receptors that are involved in their synthesis and action in the central nervous system.

2. Steroidogenic enzymes in the brain

The nervous system expresses all the enzymes necessary for the synthesis of neurosteroids (Fig. 1). Astrocytes express P450scc, cytochrome P450 17 α -hydroxylase/c17,20-lyase (P450c17), 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4-isomerase (3 β -HSD), 3 α -hydroxysteroid dehydrogenase (3 α -HSD), 17 β -hydroxysteroid dehydrogenase (17 β -HSD), steroid 5 α -reductase (5 α R) and cytochrome P450 aromatase (P450arom), producing Preg, P, DHEA, androstenedione, testosterone (T), Allop and E2 (15-19). Oligodendrocytes express P450scc, 5 α R and 3 β -HSD and produce Preg, P and Allop (18; 20). The neurons express P450scc, P450c17, 3 β -HSD, 5 α R and P450arom and produce Preg, DHEA, androstenedione, Allop and E2 (19; 21). Distinct patterns of expression of steroidogenic enzymes in neurons, oligodendrocytes, and astrocytes, suggest that some kind of cooperation between neurons and glial cells coordinates the metabolism of various sex steroids, particularly Allop, T and P (19).

2.1. Cytochrome P450 side chain cleavage

The first enzymatic reaction of steroidogenesis is the transformation of cholesterol into Preg, catalyzed by P450scc (Fig. 1). P450scc also known as CYP11A1 (Cytochrome P450 family 11 subfamily A member 1), is located in the

inner mitochondrial membrane, where it catalyzes the conversion of cholesterol to Preg via three reactions (19). The first two steps involve hydroxylation of cholesterol side chain, generating first, 22R-hydroxycholesterol and then 20 α , 22R-dihydroxycholesterol. Finally, P450_{scc} cleaves the bond between carbons 20 and 22, resulting in the production of Preg and isocaproic aldehyde. Each step of the monooxygenase reaction requires 2 electrons (reducing equivalents), which are transferred from NADPH to P450_{scc} through transfer proteins (22).

P450_{scc} is always active, but its activity is limited by the availability of cholesterol in the inner membrane. Cholesterol supply to this membrane is mainly mediated by steroidogenic acute regulatory protein (StAR) and the translocator protein (TSPO) of 18 kDa (23). Increasing TSPO-mediated translocation of cholesterol from the outer to the inner mitochondrial membrane by applying TSPO agonists, stimulates steroid production (24). TSPO likely functions as a channel, accommodating a cholesterol molecule in the space delineated by five transmembrane domains. The mechanisms by which the mitochondria-targeted protein StAR drives the transfer of cholesterol and increases steroidogenesis are less well understood (24).

The expression of P450_{scc} and the production of Preg by astrocytes, oligodendrocytes and neurons was previously described by Zwain and Yen (20). They noted that oligodendrocytes are the main source of Preg in the brain, as these cells produce Preg from cholesterol at a higher level than astrocytes or neurons, confirming previous suggestions that oligodendrocytes are primarily responsible for P450_{scc} activity in the brain (20). Interestingly, in rat embryos, P450_{scc} expression was mainly found in sensory structures of the peripheral nervous system, suggesting a possible role of this enzyme in the development and maturation of the brain (16). In the adult brain, P450_{scc} enzyme was found in the cortex, amygdala, hippocampus, and midbrain (11; 19). In the human brain, the presence of mRNA of P450_{scc} has been described in the olfactory bulb, corpus callosum, amygdala, hippocampus and cerebral cortex (19).

2.2. 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4-isomerase

Preg can be converted to P by the enzyme 3 β -HSD (Fig. 1). 3 β -HSD is a membrane-bound protein that has two distinct enzymatic activities: 3 β -dehydrogenation and isomerization of the double bond from C5,6 in the B ring (Δ 5-steroids) to C4,5 in the A ring (Δ 4-steroids) (11). 3 β -HSD is expressed in neurons, oligodendrocytes and astrocytes

and each of these types of cells can convert Preg to P (20). Two isoforms of 3 β -HSD have been described in humans, four in rats, and six in mice. These isoforms are expressed in a tissue- and developmentally specific manner and fall into two functionally distinct groups NAD⁺-dependent dehydrogenase/isomerases and NADPH-dependent 3-keto steroidreductases. In rats, 3 β -HSD mRNA type I has been detected in several regions of central nervous system, including the cerebellum, hippocampus, cortex and hypothalamus (19). In the human brain, type II 3 β -HSD mRNA has been detected by RT-PCR in the amygdala, hippocampus, cerebellum and spinal cord, among others. 3 β -HSD protein has also been detected in the brain of both rats and humans (11). Developmental changes in 3 β -HSD gene expression have been investigated in the brain of postnatal rodents. In the rat hippocampus, 3 β -HSD mRNA is 2–3-fold higher on postnatal day (PND) 7 and PND 14 than PND 70 (25); and is 9-fold higher in PND 90 than in PND 450 (26). In the cerebellum, 3 β -HSD is expressed transiently during PND 7 and PND 14 and disappears in the adult (25). These changes in enzyme expression could explain the decline in P levels during aging (27) that contribute to neurodegenerative diseases (9; 28-30). Gene regulation of 3 β -HSD is complex and involves several different factors, such as cyclic adenosine monophosphate and protein kinase-C (31), as well as DNA methylation (32). However, most of the studies have been conducted in rat and human adrenal glands and gonads; thus, the regulation of 3 β -HSD expression in the brain is still largely unknown.

2.3. Steroid 5 α -reductase

P can be converted to 5 α -dihydroprogesterone, by the enzyme 5 α R (Fig. 1) (1). Melcangi *et al.* (18) showed that the formation of 5 α -dihydroprogesterone takes place preferentially in neurons; however, type 2 astrocytes and oligodendrocytes also possess considerable 5 α R activity, while activity in type 1 astrocytes is much lower. Two isoforms of 5 α R, 5 α R type 1 (5 α R-1) and type 2 (5 α R-2), have been reported in rodents and humans. 5 α R-1 is the most abundant molecular form in the brain; it is present in many regions including the hypothalamus, hippocampus, cerebellum and cerebral cortex. In contrast, 5 α R-2 is only expressed exclusively in the late fetal and early postnatal period in the rat brain, and it is virtually undetectable in the hypothalamus, cerebellum, pons and medulla oblongata in humans. The fact that this pattern of expression correlates with T synthesis in the fetal testis suggests that 5 α R -2 could be involved in the control of brain sex differentiation (19).

The expression of 5 α R-1 is relatively constant over the period PND 1 – PND 84 but then decreases at least 2-fold from PND 90 to PND 450 in the rat hippocampus. The mRNA levels of 5 α R-2 also decreased between PND 1 to PND 84 (25). In addition, alterations of mRNA levels of 5 α R-1 in the brain are related to neurodegenerative diseases such as Niemann-Pick Disease Type C, Parkinson's disease and multiple sclerosis (28; 33). These changes may explain the decline in Allop levels during aging and neurodegenerative disease (4; 27); and therefore, the deterioration of neuronal and cognitive functions (9; 34). Moreover, some forms of enrichment, including sensory and social stimuli change the expression of 5 α R-1 in the brain. Specifically, rats housed in groups of eight animals in large cages and provided with an assortment of objects including large plastic tubes, rodent dwellings and toys of various shapes, sizes and colors during 105 days, increased the transcription of 5 α R-1 in the hippocampus (26). Since this brain structure is associated with learning and memory, these results suggested that 5 α R may play an important role in cognition.

Transcriptional regulation of 5 α R is not well understood. Some authors suggest that changes in the DNA methylation transcription factors (TFs) could be involved. The promoter regions of 5 α R-1 and 5 α R-2 genes were identified and several TFs such as selective promoter factor 1 (Sp1), Sp3, activator protein 2 (AP-2) and nuclear factor 1 (NF-1) can interact with those promoters (35; 36). In addition, Blanchard *et al.* (35) described an important CpG Island of about 1000 pb in the 5 α R-1 gene. CpG Islands are DNA strands of more than 200 bp where a cytosine (C) nucleotide is followed by a guanine (G) nucleotide at more than 50% above expected CG distribution. Cytosines in CG dinucleotides can be methylated to form 5-methylcytosine and thus the gene is silenced. Such methylation constitutes an important mechanism of transcriptional regulation. Accordingly, changes in the methylation patterns of specific sites located in 5 α R-1 gene CpG Island were correlated with alterations in the mRNA levels in the rat hippocampus (26; 37).

2.4. 3 α -hydroxysteroid dehydrogenase

The enzyme 3 α -HSD is involved in the synthesis of Allop (Fig. 1). Interestingly, the expression of 3 α -HSD appears to be mainly, if not exclusively, present in type 1 astrocytes. The compartmentalization of two strictly correlated enzymes (5 α R and 3 α -HSD) in separate central nervous system cell populations suggests the simultaneous participation of neurons and glia in the 5 alpha-reductive metabolism of hormonal steroids such as 5 α -

dihydroprogesterone (18). In humans, four functional isoforms of 3 α -HSD (1, 2, 3, 20a) were identified; types 2, 3 and 20a are widely expressed in the central nervous system (19). In contrast, rodents have a single 3 α -HSD isozyme that is expressed in the cortex, hippocampus, olfactory bulb, basal ganglia, hypothalamus, thalamus and cerebellum (11).

3 α -HSD expression is age-dependent. Higo *et al.* (25) showed that mRNA expression of this enzyme is 2.28-fold higher on PND 10 than PND 84 in the rat hippocampus. Recently, Rossetti *et al.* (26) showed that 3 α -HSD expression also decreased 2.3-fold from PND 90 to PND 450. In addition, the 3 α -HSD expression is reduced by the emergence of neurodegenerative diseases, such as Niemann-Pick Disease Type C (28), and increased by environmental stimuli (26), suggesting that 3 α -HSD enzyme is neuroprotective. The similar expression patterns of 3 α -HSD and 5 α R suggests that the synthesis of Allop is paramount for learning and memory and other hippocampal-dependent mechanisms.

The involvement of certain TFs in the regulation of 3 α -HSD gene expression has been examined by several authors. The 5'-flanking regions of the rat and human genes contain consensus sequences for AP-1, octamer-binding factor 1 (Oct-1) and steroid hormone response elements, which may comprise a steroid response unit (38). These Oct factors increase gene transcription whereas glucocorticoid response elements reduce transcription of this gene Penning (39). Hung and Penning (40) also suggest that NF-1 would upregulate the expression of 3 α -HSD enzyme in rat liver. Recently, Rossetti *et al.* (26) proposed that the transcriptional regulation of the 3 α -HSD gene in the rat brain would be mediated by differential methylation mechanisms (26; 37). Particularly, they found that the promoter was mostly methylated at a potential binding site for the sterol regulatory element-binding protein (SREBP-) and this change was correlated with alteration in the mRNA levels (26), suggesting that SREBP- also affects gene expression.

2.5. Cytochrome P450arom

P450arom catalyzes the last and obligatory step in the biosynthesis of estrogens (Fig. 1) and is necessary for sexual differentiation of the brain (41). Zwain and Yen (20) demonstrated that astrocytes and neurons, but not oligodendrocytes, express P450arom and produce E2 from T. Neurons appear to be more active than astrocytes in

aromatization of androgen to estrogen. Neurons cannot produce T, but astrocytes may provide T as a substrate for neurons to produce E2. Developmental expression of P450arom has been classified into three different groups: 1- fetal group (includes the anterior medial preoptic nucleus, the periventricular preoptic nucleus, neurons associated with the strial part of the preoptic area, and the rostral portion of the medial preoptic nucleus); 2- fetal/neonatal group (from the medial preoptic nucleus to the principal nucleus of the bed nucleus of the stria terminalis and the posterodorsal part of the medial amygdaloid nucleus) and 3- young/adult group (42). In the adult brain the pattern of P450arom distribution is restricted to interconnected nuclei, including the nucleus of the posteromedial amygdala, encapsulated region of the bed nucleus of the stria terminalis, ventrolateral portion of the ventromedial hypothalamic nucleus, and central component of the medial preoptic nucleus (43). These spatial variations of P450arom mRNA and protein provide evidence that estrogens play fundamental roles during brain development. Interestingly, P450arom is also expressed in the hippocampus, cerebral cortex, midbrain, spinal cord and cerebellum (43), suggesting that, in addition to reproductive functions, P450arom may play a role in modulation of mood, affective behaviors (e.g. depression) and/or learning and memory (44). It also plays an important role in neuroprotection after excitatory injury, experimental stroke, global ischaemia, reperfusion, and elevated intracranial pressure (45).

The regulation of P450arom in the brain is complex and not completely understood. Studies in several species have led to new perspectives on the control of this enzyme by both transcriptional and posttranscriptional mechanisms (37; 46; 47). In addition, there are regionally specific sex differences in P450arom expression during the critical period of sexual differentiation (48). However, some of the sex differences in P450arom expression could not be explained by organizational actions of gonadal hormones. Instead, genetic sex determines the expression of P450arom in specific brain areas during development, as demonstrated using the four core genotypes mouse model, in which the testis-determining gene Sry is moved from the Y chromosome onto an autosome to separate effects of gonadal sex from genetic sex (49). Individuals carrying the XY chromosome complement have higher expression levels of P450arom (mRNA and protein) in the stria terminalis and anterior amygdaloid area than individuals carrying XX chromosomes, irrespective of gonadal status (testes vs. ovary), indicating that brain P450arom at E16 is determined by sex chromosomes rather than gonadal hormones. The biological meaning of this effect is unknown; however, such differences *in vivo* could reflect differences in the local production of E2 by aromatization of T in these specific brain areas. According to these findings, amygdala neurons of genetic males would be exposed to

greater neurotogenic effects of E2, leading to larger dendritic trees and greater synaptic connectivity than neurons of genetic females .

3. Steroid hormones and neurotransmitter receptors in the brain

Neurosteroids exert several biological actions in the brain by both genomic actions mediated by nuclear /membrane steroid receptors and non-genomic actions mediated by neurotransmitter receptors. Thus, we focus this section on two main classical receptors; estrogen and progesterone receptors. It is important to consider the patterns of expression of these receptors within the brain, and understand how activating these receptors affects brain cell physiology and how these patterns of expression are controlled by hormones, age, sex, and experience. In addition, we briefly discuss the implication of the neurotransmitter receptors on neurosteroid actions.

3.1. Estrogen Receptors

There are two isoforms of the classical estrogen receptor (ER), ER α and ER β , which are transcribed from unique genes (50; 51). Activating ER α or ER β causes translocation of receptor-ligand dimers to the nucleus where they bind to estrogen response elements (EREs) on DNA (52) to control protein transcription (53) by recruiting various co-activators and co-repressors (51; 54).

In addition to this classical mode of ER expression and activation, ERs, including ER α and ER β , are also expressed on the membrane (55-59). The mechanism of this cell membrane association is unclear but involves posttranslational lipid modification (palmitoylation) and interaction with membrane/cytoplasmic scaffolding proteins (e.g. caveolins) (55-57). In addition to membrane-associated (m) ER α and mER β , G protein-coupled estrogen receptor 1 (GPER1) is another membrane-associated ER (58). Activating mERs alters membrane permeability (60) and activates second messenger cascades, including mitogen-activated protein kinases, extracellular-regulated kinases, and Src kinases, among others(59) and hyperpolarizes neurons in the preoptic area (61; 62). Interestingly, increasing data indicate that the potent androgen, dihydrotestosterone, can be metabolized to 3 β -Diol, a steroid that binds to ER β and may play a role in the estrogenic effects on pathological and physiological functions (63), such as anxiety (64; 65), cognition(66) and sexual differentiation of the brain (67).

ERs are found throughout the brain, but some areas with dense expression are highlighted below. Nuclear receptors are expressed in the pituitary, hypothalamus, hippocampus, amygdala, and prefrontal cortex (68; 69). Many cells with ER α are found in the bed nucleus of the stria terminalis (BST), medial amygdala, preoptic area, and various other hypothalamic nuclei (70-73). High levels are also seen in olfactory regions, the periaqueductal gray, area postrema, cerebellum, and parabrachial nucleus (72; 74-77). Like ER α , ER β is also found in the BST, lateral septum, the medial and basolateral amygdala, the trigeminal nuclei, the preoptic region, and other hypothalamic nuclei (69; 76; 78; 79). In addition, ER β is found in some regions with low or no ER α , such as the diagonal band of Broca, supraoptic area, and paraventricular nucleus (72; 76; 79; 80). Moderate levels are also seen in the hippocampus, substantia nigra and dorsal raphe (76; 81; 82).

ER α is expressed in cell nuclei as well as in dendrites and terminals in the hypothalamus (83). mER α is expressed on both the cytoplasmic surface and exterior surface of the cell membrane and is mostly found within presynaptic compartments of hippocampal neurons but is also seen in postsynaptic compartments and glia (84-86). mER β is also found on the cytoplasmic surface of the cell membrane, but is primarily expressed in postsynaptic dendrites with lower expression in presynaptic axons and glia (81). GPER1, like mER α and mER β , is found on the cytoplasmic surface of the cell membrane in pre and post-synaptic sites of hippocampal neurons (87-89) and is also found in clusters of vesicles in axon terminals (89). All three receptor types were also found in striatal neurons (mostly presynaptic sites) and glia (90). Similarly, the prefrontal cortex also has a preponderance of mERs at presynaptic sites and glia (91). GPER1 is highly expressed in the olfactory bulbs, hypothalamus, motor cortex, somatosensory piriform cortex, hippocampus, and the habenular nucleus of the epithalamus, nucleus of the solitary tract, and cerebellum (87; 89; 92-94).

Many factors influence expression of ERs. GPER1 expression is equivalent between the sexes but females in proestrus have higher GPER1 expression than estrus females, suggesting GPER expression can be modulated by even acute changes in circulating hormones (89). Estrogen replacement reduces ER α in preoptic and hypothalamic regions of female rodents (95). Extranuclear ER α levels in hippocampal neurons of mice are highest when estrogen is low, either during diestrus or after ovariectomy, but in rats they peak during proestrus when estrogens are high (84; 85; 96). Although aged and youthful monkeys have similar levels of ER α , GPER1, and P Receptor (PR)

This article is protected by copyright. All rights reserved.

protein, aged female rats have reduced nuclear ER expression in preoptic nuclei (97). Aged female rats also have fewer ER α -containing synapses in the hippocampus (98). Male rats retain ER α across adulthood but do have decreased ER α in response to circulating testosterone (99).

3.2. Progesterone Receptor

Like the ER, PRs are classically defined as ligand-activated transcription factors. There are two isoforms of the nuclear PR, but unlike ER α and ER β , PRA and PRB are transcribed from the same gene. PRA is an N-terminal truncated form of the full-length isomer, PRB. Unbound PR exists as a complex with chaperone proteins that are necessary for its subsequent binding (100). Bound PR dissociates from the chaperone proteins, undergoes conformational changes, dimerizes, and interacts directly with specific P response elements (PREs) in promoter regions of targeted genes by binding to steroid receptor coactivators (101). In addition to nuclear PRs, many membrane bound (m) PRs have been identified. mPRs activate G-proteins but are not GPCRs; they are members of the progestin and adipoQ receptor family (102; 103), which have 7 transmembrane domains. In addition, the b5-like heme/steroid-binding protein family includes P membrane receptor component one (PGMRC1) (104).

PR has been seen in many brain regions, including the hippocampus, frontal cortex, hypothalamus, and cerebellum (105). It is also densely expressed in the BST and the centromedial amygdala (106). mPRs are also widely distributed throughout the brain (103), but only at very low levels throughout most of the forebrain except for dense mPR β in the nucleus of the oculomotor cranial nerve (107). PGMRC1, but not mPRs, are abundant in forebrain structures that regulate neuroendocrine function (102). PGMRC1 is also in the hippocampus, cortex, and cerebellum (105).

Unlike ER, estrogens increase PR expression (108). Estrogen treatment increases PRA expression in male but not rat female cerebellum (105), but there is no sex difference in PR expression within the BST or centromedial amygdala (106). Estrogen also increases PRA expression in the hippocampus and olfactory bulb, whereas P has no effect (105). The effects of aging on PR expression are less clear. Whereas one group has found less cytosolic PR binding in the preoptic area of middle-aged ovariectomized (OVX) rats after 2 days of estrogen exposure compared to young OVX rats (109), other groups have reported no age differences on mRNA levels or PR binding (97; 110). It does

seem clear that whereas neonatal rodents have nuclear PR, expression of PR is largely extra nuclear in adult mice and rats (85).

3.3. Neurotransmitter Receptors

One of the best-documented examples of a non-genomic action of a steroid is the ability of several progesterone derivatives to activate type-A γ -aminobutyric acid (GABA_A) receptors, which are members of the ligand gated ion channel family and contain many distinct binding sites for GABA, benzodiazepines, barbiturates and convulsants. The GABA_A receptor is the principal inhibitory neurotransmitter receptor in the brain and can be made up of different subunits of the α , β , γ and δ subtypes, and their composition is region and developmental stage specific (10). Although there is no absolute specificity for neurosteroid modulation of GABA_A receptors, the α - and γ -subunits also affect GABA_A neuromodulation by either positive modulators, such as Preg and Allop, or negative modulators, such as Preg sulfate (PregS), DHEA and DHEA sulfate (DHEAS) (111). GABA_A receptors containing the δ -subunit can be less sensitive to neurosteroid modulation (10). Fluctuations in the concentration of neurosteroids and changes in GABAergic signaling have been implicated in a variety of physiological and pathophysiological conditions including stress, pregnancy, reproductive/sexual behaviors, depression, anxiety, seizure and epilepsy (7; 112-114), suggesting that GABA_A receptors are important mediators of the action of these compounds.

N-methyl-D-aspartate (NMDA) receptors are tetrameric ion channels containing two of four possible GluN2 subunits. These receptors have been implicated for decades in neurological diseases such as stroke, traumatic brain injury, dementia and schizophrenia. The GluN2 subunits substantially contribute to functional diversity of NMDA receptors and are distinctly expressed during development and among brain regions (115). Some neurosteroids such as E2 act as a negative modulator, whereas DHEA, Preg and their sulfate esters are thought to be positive allosteric modulators of NMDA receptors. Unlike GABA_A receptor interactions, the interaction of neurosteroids with the NMDA receptor is not well documented, and no specific interactions have been described (7; 10).

Although GABA_A and NMDA receptors appear to be primarily responsible for the action of neurosteroids; other kind of receptors have also been studied in the literature, such as sigma receptors, AMPA receptors and kainate receptors (10; 116). These receptors are also regulated allosterically by several neurosteroids such as PregS, DHEAS and P and have been implicated in different nervous system functions and pathologies.

4. Neurosteroids and their effects on nervous system functions.

4.1. Pregnenolone and sulfated neurosteroids

Preg and DHEA are precursors of estrogens, progestins and androgens, but they also influence neuronal functions and are likely to play particularly important roles in the aging nervous system (13). Preg is essential in maintaining cognitive functions and protects mouse hippocampal cells against glutamate and amyloid peptide-induced cell death (13). Preg also regulates neurotransmission, acting at both pre and postsynaptic sites to control synaptic release of neurotransmitters such as GABA, glutamate, noradrenaline, dopamine and serotonin (10).

In addition to these neurosteroids, their sulfated counterparts have distinct effects in the nervous system. DHEAS and PregS are the most abundant sulfated neurosteroids in the brain. Whereas Preg is a positive allosteric modulator of GABA receptors, PregS is a negative modulator of the same receptor. Sulfated neurosteroids are involved in a large number of biological functions in humans and other mammals (117-120). For example, PregS increases luteinizing hormone (LH) secretion by modulating GABA and glutamate receptors. This LH surge is inhibited by COUMATE (an irreversible inhibitor of the steroid sulfatase (STS) enzyme), showing that imbalances in neurosulfation can indirectly affect reproductive function (121-124). Furthermore, we observed that this hypothalamic sulfation prevents variations in steroid hormones by reducing STS gene expression and reduces receptive behaviors such as lordosis. This indicates that neurosulfation can fine-tune reproductive physiology and behavior by controlling expression and activity of enzymes involved (125). Thus, neurosteroid sulfation in the hypothalamus plays a key role in the reproductive function of the female rat.

4.2. Progestagens

4.2.1. Progesterone

In the central nervous system, P increases the number of oligodendrocytes expressing the myelin basic protein (MBP). MBP has many splice variants, which are developmentally regulated. In adult myelin, the role of the predominant 18.5 kDa isoform is to maintain the structural integrity and compaction of the myelin sheaths (126). This suggested that P promotes myelination by increasing the transcription of certain myelin genes (126). In the brain, P regulates spinogenesis, synaptogenesis, neuronal survival and dendritic growth (29; 30; 127; 128) and plays a neuroprotective role in numerous animal models of neurodegenerative diseases (24; 34). For example, after traumatic brain injury (TBI) P decreases cell death, gliosis, and cognitive deficits (9). Gonzalez Deniselle *et al.* (129) indicated that P restores motoneuron morphology and the expression of $\alpha 3$ subunit Na,K-ATPase mRNA, a neuronal enzyme controlling ion fluxes, neurotransmission, membrane potential, and nutrient uptake, and increased both muscle strength and survival time of Wobbler mice, a genetic model of spinal cord motor neuron disease. P prevents depression-like behavior in a model of Parkinson's disease induced by 6-hydroxydopamine in male rats (130). P also exerts marked neuroprotective effects after spinal cord injury, cerebral ischemia and stroke (34). Moreover, Liu *et al.* (131) showed that β -amyloid peptide ($A\beta_{25-35}$), a main etiological factor of Alzheimer's disease (AD), is exacerbated by low levels of P *in vivo* in the rat prefrontal cortex and hippocampus, suggesting that P is also essentially for learning and memory. In addition to its role in organization and protection of neural structures, P activity in the brain is a necessary component in female reproduction (132).

4.2.2. Allopregnanolone

P also acts on the nervous system through one of its most important metabolites, Allop. Allop increases neurogenesis and neuronal survival and reduces apoptosis in the hippocampus (7; 26). Allop also increases the density of dendritic spines as well as drebrin clusters in cultured hippocampal neurons, indicating that it increases excitatory synapse density (133). Allop also seems to be involved in learning and memory. For example, Frye (134) showed that young rats in proestrus and late pregnancy (reproductive states associated with higher cortical Allop levels) exhibit better performance on the object recognition task than diestrus rats or rats in early pregnancy. The infusion of E2 benzoate alone or with P into the hippocampus of ovariectomized rats has amnesic effects and Allop

can reverse this effect, suggesting that these effects are not mediated through the P receptor and beneficial effects may include the promotion of the cognitive performance of the hippocampus (135).

Interestingly, Allop has a key role in neurodegenerative disease. Reduced Allop levels were observed in the prefrontal cortex and in temporal cortex of patients with AD (26). Similarly, Allop content in the white matter (33) and in cerebrospinal fluid (136) is reduced in patients with multiple sclerosis and Parkinson's disease, respectively. In the triple transgenic mouse model of AD (3xTgAD), Allop prevents neurogenic and cognitive deficits (26). In addition, Allop restores hippocampal-dependent learning and memory and neural progenitor survival in aging wild type mice (26). Allop also increases Purkinje and granule cell survival in a mouse model of the human neurodegenerative disease Niemann–Pick disease type C (26). Allop is also neuroprotective in other experimental models, such as TBI, ischemia and spinal cord injury (34).

4.3. Estrogens

E2 has multiple important physiologic effects on several tissues and cellular phenotypes. E2 acts permanently on the developing brain to establish sex differences by regulating the growth, differentiation and survival of neurons and glia (137; 138). In the adult brain, E2 acts as an autocrine-paracrine factor that regulates synaptic plasticity, adult neurogenesis, reproductive behavior, aggressive behavior, pain processing and cognition (5; 45; 138-141). E2 also impacts cellular physiology by modulating calcium handling, immediate-early-gene expression (c-Fos) and kinase activity (cAMP response element-binding and extracellular-signal-regulated kinases) (142-144). The specific mechanisms of E2 action in the brain are region- and sex-specific and often involve neuronal/glial cross-talk (145). For instance, in primary cultures of dissociated neurons, the neuritogenic effect of E2 is differentially exerted depending on the genetic sex of the neurons (146), the region of the brain (2; 146-148), the maturity of the neurons (146; 149; 150) and the glial environment in which they develop (151-154). Moreover, at 14 days of embryonic age (E14), E2 only stimulates axogenesis in hypothalamic neurons from males, and this effect is contingent on the upregulation of neurogenin 3 (Ngn3), a proneural gene that is involved in neuritogenesis (142). The silencing of Ngn3 abolished sex differences in neuritogenesis, decreasing the differentiation of female neurons. Moreover, the sex difference in Ngn3 expression is determined by sex chromosomes, as demonstrated using the four core genotypes mouse model, in which the testis-determining gene Sry is moved from the Y chromosome onto an

autosomal to separate effects of gonadal sex from genetic sex (49). These findings indicate that the developmental actions of estrogen on neurite growth and differentiation can result from modulatory interactions with endogenous proneural factors (Ngn3) and the genetic sex rather than directly from estrogenic action alone (146).

5. Conclusions and future perspectives

Neurosteroids are synthesized within the nervous system. Steroidogenic enzymes and receptors are widely distributed in specific populations of neurons, neuronal precursors, and glia. Neurosteroids exert several neurotrophic and neuroprotective actions and therefore provide tremendous opportunities for developing therapeutic approaches. Although hormone replacement therapies have been studied in numerous laboratories, the results are somewhat contradictory. Hopefully, future studies will elucidate the mechanisms that regulate steroid synthesis and action in the brain and explore more alternatives to estrogen and progestagens (P and Allop) replacement.

6. References

- [1] Mellon SH, LD Griffin, NA Compagnone. Biosynthesis and action of neurosteroids. *Brain Res Brain Res Rev* 2001; 37: 3-12.
- [2] Haraguchi S, K Sasahara, H Shikimi, S Honda, N Harada, K Tsutsui. Estradiol promotes purkinje dendritic growth, spinogenesis, and synaptogenesis during neonatal life by inducing the expression of BDNF. *Cerebellum* 2012; 11: 416-7.
- [3] Hojo Y, G Murakami, H Mukai, S Higo, Y Hatanaka, M Ogiue-Ikeda, H Ishii, T Kimoto, S Kawato. Estrogen synthesis in the brain--role in synaptic plasticity and memory. *Mol Cell Endocrinol* 2008; 290: 31-43.
- [4] Marx CE, WT Trost, LJ Shampine, RD Stevens, CM Hulette, DC Steffens, JF Ervin, MI Butterfield, DG Blazer, MW Massing, JA Lieberman. The neurosteroid allopregnanolone is reduced in prefrontal cortex in Alzheimer's disease. *Biol Psychiatry* 2006; 60: 1287-94.
- [5] Rune GM, M Frotscher. Neurosteroid synthesis in the hippocampus: role in synaptic plasticity. *Neuroscience* 2005; 136: 833-42.
- [6] Charalampopoulos I, VI Alexaki, C Tsatsanis, V Minas, E Dermitzaki, I Lasaridis, L Vardouli, C Stournaras, AN Margioris, E Castanas, A Gravanis. Neurosteroids as endogenous inhibitors of neuronal cell apoptosis in aging. *Ann N Y Acad Sci* 2006; 1088: 139-52.
- [7] Charalampopoulos I, E Remboutsika, AN Margioris, A Gravanis. Neurosteroids as modulators of neurogenesis and neuronal survival. *Trends Endocrinol Metab* 2008; 19: 300-7.
- [8] Brinton RD. Neurosteroids as regenerative agents in the brain: therapeutic implications. *Nat Rev Endocrinol* 2013; 9: 241-50.

- [9] Djebaili M, Q Guo, EH Pettus, SW Hoffman, DG Stein. The neurosteroids progesterone and allopregnanolone reduce cell death, gliosis, and functional deficits after traumatic brain injury in rats. *J Neurotrauma* 2005; 22: 106-18.
- [10] Mellon SH, LD Griffin. Neurosteroids: biochemistry and clinical significance. *Trends Endocrinol Metab* 2002; 13: 35-43.
- [11] Compagnone NA, SH Mellon. Neurosteroids: biosynthesis and function of these novel neuromodulators. *Front Neuroendocrinol* 2000; 21: 1-56.
- [12] Asarian L, N Geary. Cyclic estradiol treatment normalizes body weight and restores physiological patterns of spontaneous feeding and sexual receptivity in ovariectomized rats. *Horm Behav* 2002; 42: 461-71.
- [13] Schumacher M, S Weill-Engerer, P Liere, F Robert, RJ Franklin, LM Garcia-Segura, JJ Lambert, W Mayo, RC Melcangi, A Parducz, U Suter, C Carelli, EE Baulieu, Y Akwa. Steroid hormones and neurosteroids in normal and pathological aging of the nervous system. *Prog Neurobiol* 2003; 71: 3-29.
- [14] Sierra A, E Lavaque, M Perez-Martin, I Azcoitia, DB Hales, LM Garcia-Segura. Steroidogenic acute regulatory protein in the rat brain: cellular distribution, developmental regulation and overexpression after injury. *Eur J Neurosci* 2003; 18: 1458-67.
- [15] Akwa Y, N Sananes, M Guezou, P Robel, EE Baulieu, C Le Goascogne. Astrocytes and neurosteroids: metabolism of pregnenolone and dehydroepiandrosterone. Regulation by cell density. *J Cell Biol* 1993; 121: 135-43.
- [16] Compagnone NA, A Bulfone, JL Rubenstein, SH Mellon. Expression of the steroidogenic enzyme P450scc in the central and peripheral nervous systems during rodent embryogenesis. *Endocrinology* 1995; 136: 2689-96.
- [17] Zwain IH, SS Yen, CY Cheng. Astrocytes cultured in vitro produce estradiol-17beta and express aromatase cytochrome P-450 (P-450 AROM) mRNA. *Biochim Biophys Acta* 1997; 1334: 338-48.
- [18] Melcangi RC, F Celotti, P Castano, L Martini. Differential localization of the 5 alpha-reductase and the 3 alpha-hydroxysteroid dehydrogenase in neuronal and glial cultures. *Endocrinology* 1993; 132: 1252-9.
- [19] Do Rego JL, JY Seong, D Burel, J Leprince, V Luu-The, K Tsutsui, MC Tonon, G Pelletier, H Vaudry. Neurosteroid biosynthesis: enzymatic pathways and neuroendocrine regulation by neurotransmitters and neuropeptides. *Front Neuroendocrinol* 2009; 30: 259-301.
- [20] Zwain IH, SS Yen. Neurosteroidogenesis in astrocytes, oligodendrocytes, and neurons of cerebral cortex of rat brain. *Endocrinology* 1999; 140: 3843-52.
- [21] Hojo Y, TA Hattori, T Enami, A Furukawa, K Suzuki, HT Ishii, H Mukai, JH Morrison, WG Janssen, S Kominami, N Harada, T Kimoto, S Kawato. Adult male rat hippocampus synthesizes estradiol from pregnenolone by cytochromes P45017alpha and P450 aromatase localized in neurons. *Proc Natl Acad Sci U S A* 2004; 101: 865-70.
- [22] Chung BC, KJ Matteson, R Voutilainen, TK Mohandas, WL Miller. Human cholesterol side-chain cleavage enzyme, P450scc: cDNA cloning, assignment of the gene to chromosome 15, and expression in the placenta. *Proc Natl Acad Sci U S A* 1986; 83: 8962-6.
- [23] Sierra A. Neurosteroids: the StAR protein in the brain. *J Neuroendocrinol* 2004; 16: 787-93.
- [24] Schumacher M, C Mattern, A Ghomari, JP Oudinet, P Liere, F Labombarda, R Sitruk-Ware, AF De Nicola, R Guennoun. Revisiting the roles of progesterone and allopregnanolone in the nervous system: resurgence of the progesterone receptors. *Prog Neurobiol* 2014; 113: 6-39.
- [25] Higo S, Y Hojo, H Ishii, T Kominami, K Nakajima, D Poirier, T Kimoto, S Kawato. Comparison of sex-steroid synthesis between neonatal and adult rat hippocampus. *Biochem Biophys Res Commun* 2009; 385: 62-6.

- [26] Rossetti MF, J Varayoud, GS Moreno-Piovan, EH Luque, JG Ramos. Environmental enrichment attenuates the age-related decline in the mRNA expression of steroidogenic enzymes and reduces the methylation state of the steroid 5 α -reductase type 1 gene in the rat hippocampus. *Mol Cell Endocrinol* 2015; 412: 330-8.
- [27] Caruso D, M Pesaresi, F Abbiati, D Calabrese, S Giatti, LM Garcia-Segura, RC Melcangi. Comparison of plasma and cerebrospinal fluid levels of neuroactive steroids with their brain, spinal cord and peripheral nerve levels in male and female rats. *Psychoneuroendocrinology* 2013; 38: 2278-90.
- [28] Mellon SH, W Gong, MD Schonemann. Endogenous and synthetic neurosteroids in treatment of Niemann-Pick Type C disease. *Brain Res Rev* 2008; 57: 410-20.
- [29] McEwen BS, CS Woolley. Estradiol and progesterone regulate neuronal structure and synaptic connectivity in adult as well as developing brain. *Exp Gerontol* 1994; 29: 431-6.
- [30] Zhang Z, R Yang, R Zhou, L Li, M Sokabe, L Chen. Progesterone promotes the survival of newborn neurons in the dentate gyrus of adult male mice. *Hippocampus* 2010; 20: 402-12.
- [31] Lavoie HA, SR King. Transcriptional regulation of steroidogenic genes: STARD1, CYP11A1 and HSD3B. *Exp Biol Med (Maywood)* 2009; 234: 880-907.
- [32] Vanselow J, M Spitschak, M Nimz, R Furbass. DNA methylation is not involved in preovulatory down-regulation of CYP11A1, HSD3B1, and CYP19A1 in bovine follicles but may have a role in permanent silencing of CYP19A1 in large granulosa lutein cells. *Biol Reprod* 2010; 82: 289-98.
- [33] Luchetti S, I Huitinga, DF Swaab. Neurosteroid and GABA-A receptor alterations in Alzheimer's disease, Parkinson's disease and multiple sclerosis. *Neuroscience* 2011; 191: 6-21.
- [34] Guennoun R, F Labombarda, MC Gonzalez Deniselle, P Liere, AF De Nicola, M Schumacher. Progesterone and allopregnanolone in the central nervous system: response to injury and implication for neuroprotection. *J Steroid Biochem Mol Biol* 2015; 146: 48-61.
- [35] Blanchard Y, S Seenundun, B Robaire. The promoter of the rat 5 α -reductase type 1 gene is bidirectional and Sp1-dependent. *Mol Cell Endocrinol* 2007; 264: 171-83.
- [36] Seenundun S, B Robaire. Cloning and characterization of the 5 α -reductase type 2 promoter in the rat epididymis. *Biol Reprod* 2005; 72: 851-61.
- [37] Rossetti MF, J Varayoud, GP Lazzarino, EH Luque, JG Ramos. Pregnancy and lactation differentially modify the transcriptional regulation of steroidogenic enzymes through DNA methylation mechanisms in the hippocampus of aged rats. *Mol Cell Endocrinol* 2016.
- [38] Lin HK, TM Penning. Cloning, sequencing, and functional analysis of the 5'-flanking region of the rat 3 α -hydroxysteroid/dihydrodiol dehydrogenase gene. *Cancer Res* 1995; 55: 4105-13.
- [39] Penning TM. 3 α -hydroxysteroid dehydrogenase: three dimensional structure and gene regulation. *J Endocrinol* 1996; 150 Suppl: S175-87.
- [40] Hung CF, TM Penning. Members of the nuclear factor 1 transcription factor family regulate rat 3 α -hydroxysteroid/dihydrodiol dehydrogenase (3 α -HSD/DD AKR1C9) gene expression: a member of the aldo-keto reductase superfamily. *Mol Endocrinol* 1999; 13: 1704-17.
- [41] MacLusky NJ, F Naftolin. Sexual differentiation of the central nervous system. *Science* 1981; 211: 1294-302.
- [42] Shinoda K, M Nagano, Y Osawa. Neuronal aromatase expression in preoptic, strial, and amygdaloid regions during late prenatal and early postnatal development in the rat. *J Comp Neurol* 1994; 343: 113-29.
- [43] Roselli CE, M Liu, PD Hurn. Brain aromatization: classic roles and new perspectives. *Semin Reprod Med* 2009; 27: 207-17.
- [44] Dalla C, K Antoniou, Z Papadopoulou-Daifoti, J Balthazart, J Bakker. Oestrogen-deficient female aromatase knockout (ArKO) mice exhibit depressive-like symptomatology. *Eur J Neurosci* 2004; 20: 217-28.

- [45] Arevalo MA, I Azcoitia, LM Garcia-Segura. The neuroprotective actions of oestradiol and oestrogen receptors. *Nat Rev Neurosci* 2015; 16: 17-29.
- [46] Balthazart J, GF Ball. New insights into the regulation and function of brain estrogen synthase (aromatase). *Trends Neurosci* 1998; 21: 243-9.
- [47] Tabatadze N, SM Sato, CS Woolley. Quantitative analysis of long-form aromatase mRNA in the male and female rat brain. *PLoS One* 2014; 9: e100628.
- [48] Lauber ME, A Sarasin, W Lichtensteiger. Transient sex differences of aromatase (CYP19) mRNA expression in the developing rat brain. *Neuroendocrinology* 1997; 66: 173-80.
- [49] Arnold AP, X Chen. What does the "four core genotypes" mouse model tell us about sex differences in the brain and other tissues? *Front Neuroendocrinol* 2009; 30: 1-9.
- [50] Toft D, J Gorski. A receptor molecule for estrogens: isolation from the rat uterus and preliminary characterization. *Proc Natl Acad Sci U S A* 1966; 55: 1574-81.
- [51] Kuiper GG, E Enmark, M Peltö-Huikko, S Nilsson, JA Gustafsson. Cloning of a novel receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci U S A* 1996; 93: 5925-30.
- [52] Kumar V, P Chambon. The estrogen receptor binds tightly to its responsive element as a ligand-induced homodimer. *Cell* 1988; 55: 145-56.
- [53] Nilsson S, S Makela, E Treuter, M Tujague, J Thomsen, G Andersson, E Enmark, K Pettersson, M Warner, JA Gustafsson. Mechanisms of estrogen action. *Physiol Rev* 2001; 81: 1535-65.
- [54] Rollerova E, M Urbancikova. Intracellular estrogen receptors, their characterization and function (Review). *Endocr Regul* 2000; 34: 203-18.
- [55] Boonyaratanakornkit V. Scaffolding proteins mediating membrane-initiated extra-nuclear actions of estrogen receptor. *Steroids* 2011; 76: 877-84.
- [56] Meitzen J, JI Luoma, MI Boulware, VL Hedges, BM Peterson, K Tuomela, KA Britson, PG Mermelstein. Palmitoylation of estrogen receptors is essential for neuronal membrane signaling. *Endocrinology* 2013; 154: 4293-304.
- [57] Pedram A, M Razandi, RC Sainson, JK Kim, CC Hughes, ER Levin. A conserved mechanism for steroid receptor translocation to the plasma membrane. *J Biol Chem* 2007; 282: 22278-88.
- [58] Carmeci C, DA Thompson, HZ Ring, U Francke, RJ Weigel. Identification of a gene (GPR30) with homology to the G-protein-coupled receptor superfamily associated with estrogen receptor expression in breast cancer. *Genomics* 1997; 45: 607-17.
- [59] Coleman KM, CL Smith. Intracellular signaling pathways: nongenomic actions of estrogens and ligand-independent activation of estrogen receptors. *Front Biosci* 2001; 6: D1379-91.
- [60] Fu XD, T Simoncini. Extra-nuclear signaling of estrogen receptors. *IUBMB Life* 2008; 60: 502-10.
- [61] Bologna CG, CM Revankar, SM Young, BS Edwards, JB Arterburn, AS Kiselyov, MA Parker, SE Tkachenko, NP Savchuck, LA Sklar, TI Oprea, ER Prossnitz. Virtual and biomolecular screening converge on a selective agonist for GPR30. *Nat Chem Biol* 2006; 2: 207-12.
- [62] Prossnitz ER, JB Arterburn, HO Smith, TI Oprea, LA Sklar, HJ Hathaway. Estrogen signaling through the transmembrane G protein-coupled receptor GPR30. *Annu Rev Physiol* 2008; 70: 165-90.
- [63] Handa RJ, TR Pak, AE Kudwa, TD Lund, L Hinds. An alternate pathway for androgen regulation of brain function: activation of estrogen receptor beta by the metabolite of dihydrotestosterone, 5alpha-androstane-3beta,17beta-diol. *Horm Behav* 2008; 53: 741-52.
- [64] Osterlund MK, MR Witt, JA Gustafsson. Estrogen action in mood and neurodegenerative disorders: estrogenic compounds with selective properties-the next generation of therapeutics. *Endocrine* 2005; 28: 235-42.
- [65] Krezel W, S Dupont, A Krust, P Chambon, PF Chapman. Increased anxiety and synaptic plasticity in estrogen receptor beta-deficient mice. *Proc Natl Acad Sci U S A* 2001; 98: 12278-82.
- [66] Fugger HN, TC Foster, J Gustafsson, EF Rissman. Novel effects of estradiol and estrogen receptor alpha and beta on cognitive function. *Brain Res* 2000; 883: 258-64.

- [67] Kudwa AE, V Michopoulos, JD Gatewood, EF Rissman. Roles of estrogen receptors alpha and beta in differentiation of mouse sexual behavior. *Neuroscience* 2006; 138: 921-8.
- [68] Montague D, CS Weickert, E Tomaskovic-Crook, DA Rothmond, JE Kleinman, DR Rubinow. Oestrogen receptor alpha localisation in the prefrontal cortex of three mammalian species. *J Neuroendocrinol* 2008; 20: 893-903.
- [69] Shughrue PJ, I Merchenthaler. Distribution of estrogen receptor beta immunoreactivity in the rat central nervous system. *J Comp Neurol* 2001; 436: 64-81.
- [70] Yokosuka M, H Okamura, S Hayashi. Postnatal development and sex difference in neurons containing estrogen receptor-alpha immunoreactivity in the preoptic brain, the diencephalon, and the amygdala in the rat. *J Comp Neurol* 1997; 389: 81-93.
- [71] Simonian SX, DP Spratt, AE Herbison. Identification and characterization of estrogen receptor alpha-containing neurons projecting to the vicinity of the gonadotropin-releasing hormone perikarya in the rostral preoptic area of the rat. *J Comp Neurol* 1999; 411: 346-58.
- [72] Laflamme N, RE Nappi, G Drolet, C Labrie, S Rivest. Expression and neuropeptidergic characterization of estrogen receptors (ERalpha and ERbeta) throughout the rat brain: anatomical evidence of distinct roles of each subtype. *J Neurobiol* 1998; 36: 357-78.
- [73] Chakraborty TR, PR Hof, L Ng, AC Gore. Stereologic analysis of estrogen receptor alpha (ER alpha) expression in rat hypothalamus and its regulation by aging and estrogen. *J Comp Neurol* 2003; 466: 409-21.
- [74] Simerly RB, C Chang, M Muramatsu, LW Swanson. Distribution of androgen and estrogen receptor mRNA-containing cells in the rat brain: an in situ hybridization study. *J Comp Neurol* 1990; 294: 76-95.
- [75] Belcher SM. Regulated expression of estrogen receptor alpha and beta mRNA in granule cells during development of the rat cerebellum. *Brain Res Dev Brain Res* 1999; 115: 57-69.
- [76] Mitra SW, E Hoskin, J Yudkovitz, L Pear, HA Wilkinson, S Hayashi, DW Pfaff, S Ogawa, SP Rohrer, JM Schaeffer, BS McEwen, SE Alves. Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* 2003; 144: 2055-67.
- [77] Shughrue PJ, PJ Scrimo, I Merchenthaler. Evidence for the colocalization of estrogen receptor-beta mRNA and estrogen receptor-alpha immunoreactivity in neurons of the rat forebrain. *Endocrinology* 1998; 139: 5267-70.
- [78] Creutz LM, MF Kritzer. Estrogen receptor-beta immunoreactivity in the midbrain of adult rats: regional, subregional, and cellular localization in the A10, A9, and A8 dopamine cell groups. *J Comp Neurol* 2002; 446: 288-300.
- [79] Shughrue PJ, MV Lane, I Merchenthaler. Biologically active estrogen receptor-beta: evidence from in vivo autoradiographic studies with estrogen receptor alpha-knockout mice. *Endocrinology* 1999; 140: 2613-20.
- [80] Platania P, F Laureanti, M Bellomo, R Giuffrida, AM Giuffrida-Stella, MV Catania, MA Sortino. Differential expression of estrogen receptors alpha and beta in the spinal cord during postnatal development: localization in glial cells. *Neuroendocrinology* 2003; 77: 334-40.
- [81] Milner TA, K Ayoola, CT Drake, SP Herrick, NE Tabori, BS McEwen, S Warriar, SE Alves. Ultrastructural localization of estrogen receptor beta immunoreactivity in the rat hippocampal formation. *J Comp Neurol* 2005; 491: 81-95.
- [82] Donner N, RJ Handa. Estrogen receptor beta regulates the expression of tryptophan-hydroxylase 2 mRNA within serotonergic neurons of the rat dorsal raphe nuclei. *Neuroscience* 2009; 163: 705-18.
- [83] Blaustein JD, MN Lehman, JC Turcotte, G Greene. Estrogen receptors in dendrites and axon terminals in the guinea pig hypothalamus. *Endocrinology* 1992; 131: 281-90.

- [84] Milner TA, BS McEwen, S Hayashi, CJ Li, LP Reagan, SE Alves. Ultrastructural evidence that hippocampal alpha estrogen receptors are located at extranuclear sites. *J Comp Neurol* 2001; 429: 355-71.
- [85] Mitterling KL, JL Spencer, N Dziedzic, S Shenoy, K McCarthy, EM Waters, BS McEwen, TA Milner. Cellular and subcellular localization of estrogen and progesterin receptor immunoreactivities in the mouse hippocampus. *J Comp Neurol* 2010; 518: 2729-43.
- [86] Gorosito SV, AG Lorenzo, MJ Cambiasso. Estrogen receptor alpha is expressed on the cell-surface of embryonic hypothalamic neurons. *Neuroscience* 2008; 154: 1173-7.
- [87] Funakoshi T, A Yanai, K Shinoda, MM Kawano, Y Mizukami. G protein-coupled receptor 30 is an estrogen receptor in the plasma membrane. *Biochem Biophys Res Commun* 2006; 346: 904-10.
- [88] Akama KT, LI Thompson, TA Milner, BS McEwen. Post-synaptic density-95 (PSD-95) binding capacity of G-protein-coupled receptor 30 (GPR30), an estrogen receptor that can be identified in hippocampal dendritic spines. *J Biol Chem* 2013; 288: 6438-50.
- [89] Waters EM, LI Thompson, P Patel, AD Gonzales, HZ Ye, EJ Filardo, DJ Clegg, J Gorecka, KT Akama, BS McEwen, TA Milner. G-protein-coupled estrogen receptor 1 is anatomically positioned to modulate synaptic plasticity in the mouse hippocampus. *J Neurosci* 2015; 35: 2384-97.
- [90] Almey A, EJ Filardo, TA Milner, WG Brake. Estrogen receptors are found in glia and at extranuclear neuronal sites in the dorsal striatum of female rats: evidence for cholinergic but not dopaminergic colocalization. *Endocrinology* 2012; 153: 5373-83.
- [91] Almey A, E Cannell, K Bertram, E Filardo, TA Milner, WG Brake. Medial prefrontal cortical estradiol rapidly alters memory system bias in female rats: ultrastructural analysis reveals membrane-associated estrogen receptors as potential mediators. *Endocrinology* 2014; 155: 4422-32.
- [92] Brailoiu E, SL Dun, GC Brailoiu, K Mizuo, LA Sklar, TI Oprea, ER Prossnitz, NJ Dun. Distribution and characterization of estrogen receptor G protein-coupled receptor 30 in the rat central nervous system. *J Endocrinol* 2007; 193: 311-21.
- [93] Hazell GG, ST Yao, JA Roper, ER Prossnitz, AM O'Carroll, SJ Lolait. Localisation of GPR30, a novel G protein-coupled oestrogen receptor, suggests multiple functions in rodent brain and peripheral tissues. *J Endocrinol* 2009; 202: 223-36.
- [94] Spary EJ, SE Chapman, JK Sinfield, A Maqbool, J Kaye, TF Batten. Novel G protein-coupled oestrogen receptor GPR30 shows changes in mRNA expression in the rat brain over the oestrous cycle. *Neurosignals* 2013; 21: 14-27.
- [95] Greco B, EA Allegretto, MJ Tetel, JD Blaustein. Coexpression of ER beta with ER alpha and progesterin receptor proteins in the female rat forebrain: effects of estradiol treatment. *Endocrinology* 2001; 142: 5172-81.
- [96] Romeo RD, JB McCarthy, A Wang, TA Milner, BS McEwen. Sex differences in hippocampal estradiol-induced N-methyl-D-aspartic acid binding and ultrastructural localization of estrogen receptor-alpha. *Neuroendocrinology* 2005; 81: 391-9.
- [97] Brown TJ, NJ MacLusky, M Shanabrough, F Naftolin. Comparison of age- and sex-related changes in cell nuclear estrogen-binding capacity and progesterin receptor induction in the rat brain. *Endocrinology* 1990; 126: 2965-72.
- [98] Adams MM, SE Fink, RA Shah, WG Janssen, S Hayashi, TA Milner, BS McEwen, JH Morrison. Estrogen and aging affect the subcellular distribution of estrogen receptor-alpha in the hippocampus of female rats. *J Neurosci* 2002; 22: 3608-14.
- [99] Wu D, AC Gore. Changes in androgen receptor, estrogen receptor alpha, and sexual behavior with aging and testosterone in male rats. *Horm Behav* 2010; 58: 306-16.
- [100] Pratt WB, DO Toft. Regulation of signaling protein function and trafficking by the hsp90/hsp70-based chaperone machinery. *Exp Biol Med (Maywood)* 2003; 228: 111-33.

- [101] Leonhardt SA, V Boonyaratanakornkit, DP Edwards. Progesterone receptor transcription and non-transcription signaling mechanisms. *Steroids* 2003; 68: 761-70.
- [102] Thomas P, Y Pang, J Dong, P Groenen, J Kelder, J de Vlieg, Y Zhu, C Tubbs. Steroid and G protein binding characteristics of the seatrout and human progestin membrane receptor alpha subtypes and their evolutionary origins. *Endocrinology* 2007; 148: 705-18.
- [103] Tang YT, T Hu, M Arterburn, B Boyle, JM Bright, PC Emtage, WD Funk. PAQR proteins: a novel membrane receptor family defined by an ancient 7-transmembrane pass motif. *J Mol Evol* 2005; 61: 372-80.
- [104] Losel R, S Breiter, M Seyfert, M Wehling, E Falkenstein. Classic and non-classic progesterone receptors are both expressed in human spermatozoa. *Horm Metab Res* 2005; 37: 10-4.
- [105] Guerra-Araiza C, O Villamar-Cruz, A Gonzalez-Arenas, R Chavira, I Camacho-Arroyo. Changes in progesterone receptor isoforms content in the rat brain during the oestrous cycle and after oestradiol and progesterone treatments. *J Neuroendocrinol* 2003; 15: 984-90.
- [106] Auger CJ, GJ De Vries. Progestin receptor immunoreactivity within steroid-responsive vasopressin-immunoreactive cells in the male and female rat brain. *J Neuroendocrinol* 2002; 14: 561-7.
- [107] Intlekofer KA, SL Petersen. Distribution of mRNAs encoding classical progestin receptor, progesterone membrane components 1 and 2, serpine mRNA binding protein 1, and progestin and ADIPOQ receptor family members 7 and 8 in rat forebrain. *Neuroscience* 2011; 172: 55-65.
- [108] Blaustein JD, JC Turcotte. Estradiol-induced progestin receptor immunoreactivity is found only in estrogen receptor-immunoreactive cells in guinea pig brain. *Neuroendocrinology* 1989; 49: 454-61.
- [109] Wise PM, B Parsons. Nuclear estradiol and cytosol progestin receptor concentrations in the brain and the pituitary gland and sexual behavior in ovariectomized estradiol-treated middle-aged rats. *Endocrinology* 1984; 115: 810-6.
- [110] Funabashi T, SP Kleopoulos, PJ Brooks, F Kimura, DW Pfaff, K Shinohara, CV Mobbs. Changes in estrogenic regulation of estrogen receptor alpha mRNA and progesterone receptor mRNA in the female rat hypothalamus during aging: an in situ hybridization study. *Neurosci Res* 2000; 38: 85-92.
- [111] Baulieu EE. Neurosteroids: a novel function of the brain. *Psychoneuroendocrinology* 1998; 23: 963-87.
- [112] MacKenzie G, J Maguire. Neurosteroids and GABAergic signaling in health and disease. *Biomol Concepts* 2013; 4: 29-42.
- [113] Murphy DD, NB Cole, V Greenberger, M Segal. Estradiol increases dendritic spine density by reducing GABA neurotransmission in hippocampal neurons. *J Neurosci* 1998; 18: 2550-9.
- [114] Pluchino N, P Drakopoulos, F Bianchi-Demicheli, JM Wenger, P Petignat, AR Genazzani. Neurobiology of DHEA and effects on sexuality, mood and cognition. *J Steroid Biochem Mol Biol* 2015; 145: 273-80.
- [115] Ogden KK, SF Traynelis. New advances in NMDA receptor pharmacology. *Trends Pharmacol Sci* 2011; 32: 726-33.
- [116] Niitsu T, M Iyo, K Hashimoto. Sigma-1 receptor agonists as therapeutic drugs for cognitive impairment in neuropsychiatric diseases. *Curr Pharm Des* 2012; 18: 875-83.
- [117] El-Etr M, Y Akwa, EE Baulieu, M Schumacher. The neuroactive steroid pregnenolone sulfate stimulates the release of gonadotropin-releasing hormone from GT1-7 hypothalamic neurons, through N-methyl-D-aspartate receptors. *Endocrinology* 2006; 147: 2737-43.
- [118] Gibbs TT, SJ Russek, DH Farb. Sulfated steroids as endogenous neuromodulators. *Pharmacol Biochem Behav* 2006; 84: 555-67.
- [119] Nanfaro F, R Cabrera, V Bazzocchini, M Laconi, R Yunes. Pregnenolone sulfate infused in lateral septum of male rats impairs novel object recognition memory. *Pharmacol Rep* 2010; 62: 265-72.

- [120] Schumacher M, P Liere, Y Akwa, K Rajkowski, W Griffiths, K Bodin, J Sjoval, EE Baulieu. Pregnenolone sulfate in the brain: a controversial neurosteroid. *Neurochem Int* 2008; 52: 522-40.
- [121] Cabrera RJ, CE Navarro. Progesterone in vitro increases NMDA-evoked [3H] dopamine release from striatal slices in proestrus rats. *Neuropharmacology* 1996; 35: 175-8.
- [122] Giuliani FA, R Yunes, CE Mohn, M Laconi, V Rettori, R Cabrera. Allopregnanolone induces LHRH and glutamate release through NMDA receptor modulation. *Endocrine* 2011; 40: 21-6.
- [123] Laconi MR, RJ Cabrera. Effect of centrally injected allopregnanolone on sexual receptivity, luteinizing hormone release, hypothalamic dopamine turnover, and release in female rats. *Endocrine* 2002; 17: 77-83.
- [124] Brann DW, VB Mahesh. Glutamate: a major neuroendocrine excitatory signal mediating steroid effects on gonadotropin secretion. *J Steroid Biochem Mol Biol* 1995; 53: 325-9.
- [125] Christensen A, GE Bentley, R Cabrera, HH Ortega, N Perfito, TJ Wu, P Micevych. Hormonal regulation of female reproduction. *Horm Metab Res* 2012; 44: 587-91.
- [126] Schumacher M, R Hussain, N Gago, JP Oudinet, C Mattern, AM Ghomari. Progesterone synthesis in the nervous system: implications for myelination and myelin repair. *Front Neurosci* 2012; 6: 10.
- [127] Mellon SH. Neurosteroid regulation of central nervous system development. *Pharmacol Ther* 2007; 116: 107-24.
- [128] Tsutsui K. Progesterone biosynthesis and action in the developing neuron. *Endocrinology* 2008; 149: 2757-61.
- [129] Gonzalez Deniselle MC, JJ Lopez-Costa, JP Saavedra, L Pietranera, SL Gonzalez, L Garay, R Guennoun, M Schumacher, AF De Nicola. Progesterone neuroprotection in the Wobbler mouse, a genetic model of spinal cord motor neuron disease. *Neurobiol Dis* 2002; 11: 457-68.
- [130] Casas S, S Garcia, R Cabrera, F Nanfaro, C Escudero, R Yunes. Progesterone prevents depression-like behavior in a model of Parkinson's disease induced by 6-hydroxydopamine in male rats. *Pharmacol Biochem Behav* 2011; 99: 614-8.
- [131] Liu S, H Wu, G Xue, X Ma, J Wu, Y Qin, Y Hou. Metabolic alteration of neuroactive steroids and protective effect of progesterone in Alzheimer's disease-like rats. *Neural Regen Res* 2013; 8: 2800-10.
- [132] Kuo J, P Micevych. Neurosteroids, trigger of the LH surge. *J Steroid Biochem Mol Biol* 2012; 131: 57-65.
- [133] Shimizu H, Y Ishizuka, H Yamazaki, T Shirao. Allopregnanolone increases mature excitatory synapses along dendrites via protein kinase A signaling. *Neuroscience* 2015; 305: 139-45.
- [134] Frye CA. Neurosteroids' effects and mechanisms for social, cognitive, emotional, and physical functions. *Psychoneuroendocrinology* 2009; 34 Suppl 1: S143-61.
- [135] Escudero C, S Casas, F Giuliani, V Bazzocchini, S Garcia, R Yunes, R Cabrera. Allopregnanolone prevents memory impairment: effect on mRNA expression and enzymatic activity of hippocampal 3-alpha hydroxysteroid oxidoreductase. *Brain Res Bull* 2012; 87: 280-5.
- [136] di Michele F, P Longone, E Romeo, S Lucchetti, L Brusa, M Pierantozzi, A Bassi, G Bernardi, P Stanzione. Decreased plasma and cerebrospinal fluid content of neuroactive steroids in Parkinson's disease. *Neurol Sci* 2003; 24: 172-3.
- [137] Lenz KM, MM McCarthy. Organized for sex - steroid hormones and the developing hypothalamus. *Eur J Neurosci* 2010; 32: 2096-104.
- [138] Campbell RE, AE Herbison. Gonadal steroid neuromodulation of developing and mature hypothalamic neuronal networks. *Curr Opin Neurobiol* 2014; 29: 96-102.

- [139] Remage-Healey L, SM Dong, A Chao, BA Schlinger. Sex-specific, rapid neuroestrogen fluctuations and neurophysiological actions in the songbird auditory forebrain. *J Neurophysiol* 2012; 107: 1621-31.
- [140] Zhang QG, R Wang, H Tang, Y Dong, A Chan, GR Sareddy, RK Vadlamudi, DW Brann. Brain-derived estrogen exerts anti-inflammatory and neuroprotective actions in the rat hippocampus. *Mol Cell Endocrinol* 2014; 389: 84-91.
- [141] Vierk R, N Brandt, GM Rune. Hippocampal estradiol synthesis and its significance for hippocampal synaptic stability in male and female animals. *Neuroscience* 2014; 274: 24-32.
- [142] Arevalo MA, I Ruiz-Palmero, MJ Scerbo, E Acaz-Fonseca, MJ Cambiasso, LM Garcia-Segura. Molecular mechanisms involved in the regulation of neuritogenesis by estradiol: Recent advances. *J Steroid Biochem Mol Biol* 2012; 131: 52-6.
- [143] Cattaneo E, A Maggi. c-fos induction by estrogen in specific rat brain areas. *Eur J Pharmacol* 1990; 188: 153-9.
- [144] Auger AP, JD Blaustein. Progesterone enhances an estradiol-induced increase in Fos immunoreactivity in localized regions of female rat forebrain. *J Neurosci* 1995; 15: 2272-9.
- [145] Arevalo MA, I Ruiz-Palmero, J Simon-Areces, E Acaz-Fonseca, I Azcoitia, LM Garcia-Segura. Estradiol meets notch signaling in developing neurons. *Front Endocrinol (Lausanne)* 2011; 2: 21.
- [146] Scerbo MJ, A Freire-Regatillo, CD Cisternas, M Brunotto, MA Arevalo, LM Garcia-Segura, MJ Cambiasso. Neurogenin 3 mediates sex chromosome effects on the generation of sex differences in hypothalamic neuronal development. *Front Cell Neurosci* 2014; 8: 188.
- [147] von Schassen C, L Fester, J Prange-Kiel, C Lohse, C Huber, M Bottner, GM Rune. Oestrogen synthesis in the hippocampus: role in axon outgrowth. *J Neuroendocrinol* 2006; 18: 847-56.
- [148] Ruiz-Palmero I, M Hernando, LM Garcia-Segura, MA Arevalo. G protein-coupled estrogen receptor is required for the neuritogenic mechanism of 17beta-estradiol in developing hippocampal neurons. *Mol Cell Endocrinol* 2013; 372: 105-15.
- [149] Diaz H, A Lorenzo, HF Carrer, A Caceres. Time lapse study of neurite growth in hypothalamic dissociated neurons in culture: sex differences and estrogen effects. *J Neurosci Res* 1992; 33: 266-81.
- [150] Cambiasso MJ, H Diaz, A Caceres, HF Carrer. Neuritogenic effect of estradiol on rat ventromedial hypothalamic neurons co-cultured with homotopic or heterotopic glia. *J Neurosci Res* 1995; 42: 700-9.
- [151] Cambiasso MJ, JA Colombo, HF Carrer. Differential effect of oestradiol and astroglia-conditioned media on the growth of hypothalamic neurons from male and female rat brains. *Eur J Neurosci* 2000; 12: 2291-8.
- [152] Hu R, WQ Cai, XG Wu, Z Yang. Astrocyte-derived estrogen enhances synapse formation and synaptic transmission between cultured neonatal rat cortical neurons. *Neuroscience* 2007; 144: 1229-40.
- [153] Lenz KM, BM Nugent, R Haliyur, MM McCarthy. Microglia are essential to masculinization of brain and behavior. *J Neurosci* 2013; 33: 2761-72.
- [154] Acaz-Fonseca E, R Sanchez-Gonzalez, I Azcoitia, MA Arevalo, LM Garcia-Segura. Role of astrocytes in the neuroprotective actions of 17beta-estradiol and selective estrogen receptor modulators. *Mol Cell Endocrinol* 2014; 389: 48-57.

Figure Legends

Figure 1. Pathway of neurosteroid synthesis. Steroidogenic acute regulatory protein (StAR); cytochrome P450 side chain cleavage (P450_{scc}); 3 β -hydroxysteroid dehydrogenase/ Δ 5- Δ 4-isomerase (3 β -HSD); cytochrome P450 17 α -hydroxylase/c17,20-lyase (P450_{c17}); steroid 5 α -reductase (5 α R); 3 α -hydroxysteroid dehydrogenase (3 α -HSD), 17 β -hydroxysteroid dehydrogenase (17 β -HSD) and cytochrome P450_{arom} (P450_{arom}). Astrocytes express P450_{scc}, P450_{c17}, 3 β -HSD, 3 α -HSD, 17 β -HSD, 5 α R and P450_{arom}, producing pregnenolone (Preg), progesterone (P), dehydroepiandrosterone (DHEA), androstenedione, testosterone, allopregnanolone (Allop) and estradiol (E2) (15-19). Oligodendrocytes express P450_{scc}, 5 α - reductase and 3 β -HSD and produce Preg, P and Allop (18; 20). The neurons express P450_{scc}, P450_{c17}, 3 β -HSD, 5 α -reductase and P450_{arom} and produce Preg, DHEA, androstenedione, AlloP and E2 (19).

