

## REVIEW

# Oestrogens and temperature-dependent sex determination in reptiles: all is in the gonads

C Pieau and M Dorizzi

Institut Jacques Monod, UMR 7592, CNRS et Universités Paris 6 et 7, 2 place Jussieu, 75251 Paris Cedex 05, France  
(Requests for offprints should be addressed to C Pieau; Email: pieau@ijm.jussieu.fr)

### Abstract

In many species of oviparous reptiles, the first steps of gonadal sex differentiation depend on the incubation temperature of the eggs. Feminization of gonads by exogenous oestrogens at a male-producing temperature and masculinization of gonads by antioestrogens and aromatase inhibitors at a female-producing temperature have irrefutably demonstrated the involvement of oestrogens in ovarian differentiation. Nevertheless, several studies performed on the entire gonad/adrenal/mesonephros complex failed to find differences between male- and female-producing temperatures in oestrogen content, aromatase activity and *aromatase* gene expression during the thermosensitive period for sex determination. Thus, the key role of aromatase and oestrogens in the first steps of ovarian differentiation has been questioned, and extragonadal organs or tissues, such as adrenal, mesonephros, brain or yolk, were considered as possible targets of temperature and sources of the oestrogens acting on gonadal sex differentiation.

In disagreement with this view, experiments and assays carried out on the gonads alone, i.e. separated from the adrenal/mesonephros, provide evidence that the gonads themselves respond to temperature shifts by modifying their sexual differentiation and are the site of aromatase activity and oestrogen synthesis during the thermosensitive period. Oestrogens act locally on both the cortical and the medullary part of the gonad to direct ovarian differentiation. We have concluded that there is no objective reason to search for the implication of other organs in the phenomenon of temperature-dependent sex determination in reptiles. From the comparison with data obtained in other vertebrates, we propose two main directions for future research: to examine how transcription of the *aromatase* gene is regulated and to identify molecular and cellular targets of oestrogens in gonads during sex differentiation, in species with strict genotypic sex determination and species with temperature-dependent sex determination.

*Journal of Endocrinology* (2004) **181**, 367–377

### Introduction

Many species of oviparous reptiles, including crocodylians, a majority of turtles, some lizards and the two closely related species of *Sphenodon* have been shown to display temperature-dependent sex determination (TSD). In these species, the differentiation of gonads into ovaries or testes depends on the incubation temperature of the eggs during a critical period of embryonic development designated the thermosensitive period (TSP). Various treatments prior to and/or during this period have demonstrated the involvement of oestrogens in gonadal sex differentiation. Treatments with exogenous oestrogens result in ovarian differentiation at a male-producing temperature, whereas treatments with antioestrogens or aromatase inhibitors result in testicular differentiation at a female-producing temperature (for review see Pieau *et al.* 1999).

Several studies have sought for oestrogen content, aromatase activity and *aromatase* gene expression in the gonad/adrenal/mesonephros (GAM) complexes during the embryonic development of turtles and crocodylians (White & Thomas 1992*a,b*, Smith & Joss 1994*a*, Smith *et al.* 1995, Jeyasuria & Place 1997, 1998, Willingham *et al.* 2000, Gabriel *et al.* 2001, Murdoch & Wibbels 2003). In only one of these studies were differences between male- and female-producing temperatures found during the TSP (Jeyasuria & Place 1998). In the other studies, differences were found only at the end of the TSP and/or after the TSP, or were not found during and after the TSP. Therefore, some authors inferred that oestrogens are not involved in the early steps of ovarian differentiation (Smith & Joss 1994*a*, Smith *et al.* 1995, Murdoch & Wibbels 2003). Other authors inferred that temperature acts on extragonadal tissues, in particular the adrenal and/or

mesonephros (White & Thomas 1992a), brain (Jeyasuria & Place 1998, Salame-Mendez *et al.* 1998, Gutierrez-Ospina *et al.* 1999, Willingham *et al.* 2000, Place *et al.* 2001) and yolk (Conley *et al.* 1997, Janzen *et al.* 1998, Bowden *et al.* 2000, 2002, Elf *et al.* 2002, for review see Elf 2003) since these tissues could be the sources of the oestrogens acting on gonadal sex differentiation.

We have analysed data obtained from the gonads alone, in particular in freshwater and marine turtles, that do not support the involvement of extragonadal tissues. This analysis has shown that (1) temperature has a direct effect on the gonads for their sexual differentiation, (2) during the TSP, aromatase activity in the gonads is dependent on the incubation temperature of the eggs, (3) oestrogens are synthesized by the gonads during the TSP and gonadal oestrogen synthesis depends on the incubation temperature of the eggs, and (4) oestrogens act on both the cortical and the medullary parts of the gonads.

Taking into account these data, we attempt to place the phenomenon of TSD in the larger context of gonadal sex differentiation in vertebrates and propose avenues for future research.

### Key morphological events in turtle gonads during embryonic development

Most data on the gonads alone were obtained in the freshwater turtle, *Emys orbicularis* and two marine turtles, *Lepidochelys olivacea* and *Dermochelys coriacea*. Figures 1 and 2 illustrate the main morphological events occurring in the gonads during embryonic development in *E. orbicularis*. In this species, 25 °C yields 100% males, 30 °C yields 100% females and 28.5 °C (pivotal temperature) yields males, females and intersexes at hatching (Pieau 1976, Raynaud & Pieau 1985). In intersexes, the gonads are ovotestes (Fig. 2F). After hatching, ovotestes generally evolve as typical testes. However, some oocytes may persist at the surface of gonads through to adulthood (Pieau *et al.* 1998).

Based on the developmental stages defined in another freshwater turtle, *Chelydra serpentina* (Yntema 1968), the TSP extends between stages 16 and 22 of embryonic development in *E. orbicularis* (Pieau & Dorizzi 1981). Prior to the TSP, the gonads are indifferent (or bipotential) (Fig. 1). They are composed of the germinal epithelium and, in the inner part, of thin cords of epithelial cells (the so-called 'sex-cords') within loose mesenchymal tissue (Fig. 2A). At 25 °C, the first histological signs of differentiation of testicular cords (future seminiferous cords) appear at stage 17 (Figs 1 and 2B). During the TSP, the germinal epithelium flattens, and germ cells leave it to migrate between the epithelial cells (future Sertoli cells) of the testicular cords. At hatching, testicular cords are wide and display high Sertolian epithelium, whereas the surface epithelium is generally very thin (Fig. 2D). However, some remnants of the germinal epithelium enclosing germ

cells may persist at the surface of testes. At 30 °C, in differentiating female gonads, during the TSP the initial sex-cords become thin, appear fragmented, or evolve as small lacunae bordered by a flat epithelium, whereas the germinal epithelium evolves as an ovarian cortex. Cortex development is mainly due to the *in situ* proliferation of germ cells. By the end of the TSP, some germ cells have entered into meiosis (Figs 1 and 2C). At hatching, primordial follicles with a growing oocyte are formed in the internal side of the cortex, in contact with the ovarian medulla (Figs 1 and 2E).


Gonadal development in other freshwater turtles is very similar to that in *E. orbicularis*. In marine turtles, it is roughly similar, although displaying some specificity. Thus, in *L. olivacea* and *D. coriacea*, germ cells have not entered into meiosis at the end of embryonic development (for review see Pieau *et al.* 1999). Developmental stages in embryos of marine turtles have been defined by Miller (1985). The correspondence of key stages with those defined in freshwater turtles is shown in Fig. 1.

### Temperature has a direct effect on the gonads for their sexual differentiation


The direct effect of temperature on the gonads was demonstrated by experiments carried out *in vitro* in the marine turtle *L. olivacea* (Moreno-Mendoza *et al.* 2001). In this species, eggs incubated at 26 °C yield males, whereas eggs incubated at 33 °C yield females. Gonadal masculinization was characterized by the expression of the SOX9 protein during normal (*in ovo*) embryonic development. From stages 21 to 24 of embryonic development (see Fig. 1 for stages in marine turtles), SOX9 is expressed at both temperatures, in gonads which are still histologically undifferentiated. Thereafter, SOX9 remains positive at 26 °C in the medullary cords of differentiating male gonads, whereas it is downregulated at 33 °C in differentiating female gonads (Moreno-Mendoza *et al.* 1999).

The effects of temperature shifts on the expression of SOX9 were compared in gonads cultured *in vitro* and in gonads of whole embryos maintained *in ovo*. The paired gonads of embryos were explanted at stage 24 of development at 33 °C (Fig. 3 series 1A) and at stage 23 of development at 26 °C (Fig. 3 series 1B). One gonad of each pair was cultured at 33 °C and the contralateral one at 26 °C. Immunological detection of SOX9 was performed after different times of culture. SOX9 expression was maintained in gonads cultured at 26 °C, whereas it was downregulated in gonads cultured at 33 °C, following the same time-course as that in gonads of whole embryos submitted to the same thermal treatments. Thirteen days after explantation (Fig. 3 series 1B) the gonads cultured at 26 °C presented well-developed medullary cords expressing SOX9, whereas the gonads cultured at 33 °C presented an ovarian cortex and fragmented medullary cords without SOX9-positive cells.

**FRESHWATER TURTLE (*Emys orbicularis*)**



**MARINE TURTLE (*Dermochelys coriacea*)**




**Figure 1** The main steps in the differentiation of gonads in relation to the age and developmental stage in embryos of the freshwater turtle *E. orbicularis*. Corresponding ages and stages in the marine turtle *D. coriacea* (according to Renous *et al.* (1989) and Desvages *et al.* (1993)).

In another series of experiments, gonads from embryos incubated at 33 °C were explanted at stage 24, cultured at 33 °C and shifted to 26 °C at different times; they were

then cultured at 26 °C for 9 or 10 days (Fig. 3 series 2). When the shift was performed at 0, 2 and 3 days, SOX9 expression was maintained, but at day 4 SOX9 expression





**Figure 3** Effects of temperature shifts on SOX9 protein expression in gonads cultured *in vitro* at stages 24 and 23 of embryonic development in the marine turtle *L. olivacea*. Series 1A, 1B and 2 correspond to three distinct experiments (adapted from Moreno-Mendoza *et al.* (2001)).

was no longer observed in cultured gonads, indicating the commitment of ovarian differentiation after 4 days of culture at the female-producing temperature. Again the

response was similar to that of control gonads of whole embryos shifted from 33 to 26 °C at the same ages (Moreno-Mendoza *et al.* 2001).



The similarity of response to temperature shifts in gonads cultured *in vitro* and in gonads developing in whole embryos clearly showed a direct effect of temperature on the gonads for their sexual differentiation.

### During the TSP aromatase activity in the gonads depends on the incubation temperature of the eggs

Aromatase is the enzyme complex that converts androgens to oestrogens. Aromatase activity was measured in the gonads separated from the adrenal/mesonephros complexes in the freshwater turtle *E. orbicularis* and the marine turtle *D. coriacea*.

In *E. orbicularis* the TSP extends between stages 16 and 22 (Fig. 1). In a first series of experiments, aromatase activity was measured in pools of gonads from embryos incubated at 25 °C (male-producing temperature) or at 30 °C (female-producing temperature). At the beginning of the TSP, the aromatase activity was very low at both temperatures but was somewhat higher at 30 °C than at 25 °C. It thereafter remained low in differentiating testes at 25 °C, whereas it increased exponentially during the TSP and formed a peak by the end of embryonic development in differentiating ovaries at 30 °C (Desvages & Pieau 1992a). The same profiles were obtained by measuring aromatase activity in the paired gonads of each individual. In addition, by this method it was shown that during the TSP aromatase activity in testes and ovotestes at 28.5 °C (pivotal temperature) was slightly higher than in testes at 25 °C, and aromatase activity in ovaries at 28.5 °C was slightly lower than in ovaries at 30 °C (Pieau *et al.* 1998).

In a second series of experiments, aromatase activity was measured in gonads of embryos of *E. orbicularis* shifted from a male- to a female-producing temperature and vice versa at different stages of embryonic development and for different times. Shifts performed during the TSP resulted in an increase or a decrease of aromatase activity, whereas shifts after the TSP were ineffective (Desvages & Pieau 1992a). In addition, these experiments showed that the temperature acts on the regulation of aromatase synthesis rather than on the protein activity itself (Desvages & Pieau 1992b).

In *D. coriacea* eggs incubated below 29 °C yield males, whereas eggs incubated above 29.5 °C yield females (Rimblot *et al.* 1985). As shown above, the equivalents of stages 16 and 22 (delimiting the TSP) in *E. orbicularis* are stages 23 and 27 in *D. coriacea* (Fig. 1). Eggs of *D. coriacea* were shifted from 27 °C (masculinizing) to 35 °C (highly feminizing) at different embryonic stages between stages 22 and 29, and then incubated for 6 days at 35 °C. Only shifts performed between stages 23 and 27 resulted in a significant increase of gonadal aromatase activity compared with that in embryos maintained at 27 °C (Desvages *et al.* 1993). It is noteworthy that in *L. olivacea*, changes in

expression of SOX9 protein resulting from temperature shifts were obtained during the same period (Moreno-Mendoza *et al.* 2001, see above). Therefore, the sensitive period to temperature shifts that induce changes of gonadal aromatase activity in *D. coriacea* and of SOX9 expression in *L. olivacea* coincides with the TSP for gonadal sex differentiation in *E. orbicularis*.

Differences between aromatase activities at male- and at female-producing temperatures were not seen during the TSP in the entire GAM complexes of *Crocodylus porosus* (Smith & Joss 1994a), *Alligator mississippiensis* (Smith *et al.* 1995) and *Trachemys scripta* (Willingham *et al.* 2000). Aromatase activity in the gonads was probably masked by that in the adrenal/mesonephros. Indeed, during the TSP, the gonad represents a very small portion of the GAM complex and, as shown in *E. orbicularis*, some aromatase activity is also present in the adrenal/mesonephros (Richard-Mercier *et al.* 1995). For the same reason, using RT-PCR, differences in *aromatase* mRNA levels between male and female GAMs were not seen during the TSP in *A. mississippiensis* (Gabriel *et al.* 2001) and *T. scripta* (Murdoch & Wibbels 2003). Therefore, extrapolating to the gonads data obtained for the entire GAMs may lead to misinterpretations.

### Oestrogens are synthesized by the gonads during the TSP and gonadal oestrogen synthesis depends on the incubation temperature of the eggs

Using a histochemical method,  $\Delta^5$ - $3\beta$ -hydroxysteroid dehydrogenase ( $3\beta$ -HSD) activity was found to be present in gonads of *E. orbicularis* already at stage 15, showing that steroidogenesis begins very early, prior to the TSP, in the indifferent gonads (Fig. 1). At this stage and at later stages,  $3\beta$ -HSD activity was higher at the male- than at the female-producing temperature (Pieau 1974).

A study of steroid metabolism performed during and after the TSP confirmed this preliminary result and, in addition, showed that in the gonads all active enzymes involved in complete steroidogenesis are present during the TSP (Desvages & Pieau 1991).

Sensitive methods were used to measure gonadal oestrogen content. Assays were performed in embryos of *E. orbicularis* (Dorizzi *et al.* 1991) and *D. coriacea* (Desvages *et al.* 1993). In both species, the gonads were separated from the adrenal/mesonephros complexes and pooled. In *E. orbicularis*, detection of oestrogens (oestrone and oestradiol-17 $\beta$ ) was carried out with a sensitive radioimmunoassay at stages 16, 17 and 18, i.e. at the beginning of the TSP. At stage 16, very low amounts of oestrogens were found at both male-producing temperature (25 °C) and female-producing temperature (30 °C). At stages 17 and 18, oestrogens remained detectable in very low amounts or were not detectable at the male-producing temperature, whereas higher levels of oestrogens were found at the female-producing temperature (Dorizzi *et al.* 1991).

In *D. coriacea*, detection of oestrogens was performed using an enzymatic method (Nicolas *et al.* 1979), at stages 26 (during the TSP) and 31 (hatching). At both stages, oestrogens were detected at the female-producing temperature (30.5 °C) but not at the male-producing temperature (27 °C). As expected, their levels at 30.5 °C, higher at stage 31 than at stage 26, were well correlated with aromatase activity (Desvages *et al.* 1993).

Given that the gonads synthesize oestrogens as early as at the beginning of the TSP and that gonadal oestrogen synthesis depends on the temperature during this period, it can be assumed that gonadal endogenous oestrogens but not extragonadal oestrogens – synthesized in the brain or contained in the egg yolk – are involved in TSD.

### Oestrogens act on both the cortical and the medullary parts of the gonads

In reptiles at the end of embryonic development, the surface epithelium of the testes is generally very thin, but remnants of the initial germinal epithelium enclosing germ cells may persist locally. Previous early studies reported that oestrogenic treatment of immature alligators (*A. mississippiensis*, Forbes 1938) and newly hatched or juvenile turtles (*Malaclemys terrapin*, Risley 1941, *Mauremys leprosa*, Vivien & Stéfan 1958) induced proliferation and entry into meiosis of germ cells in remnants of the germinal epithelium. These results indicated that the development of an ovarian cortex from the germinal epithelium is controlled by oestrogens. However, after such late treatments, testicular cords were maintained in the medullary part of the gonads.

At the time of the discovery of TSD in turtles, oestradiol was injected into eggs incubated at a male-producing temperature in two species, *Testudo graeca* and *E. orbicularis*. In *T. graeca*, treatments performed early – when embryonic gonads still appeared histologically undifferentiated – inhibited development of testicular cords (they became thin or lacunar), but not later treatments (Pieau 1970). In *E. orbicularis*, treatments were performed at stages 19–20 (i.e. during the TSP) with different doses of oestradiol. In general, the higher the doses of hormone, the stronger the inhibition of testicular cord development. In all cases, an ovarian cortex was formed. Thus, the gonads of treated embryos were typical ovaries, hypo-ovaries (with a very reduced medullary part) or ovotestes (Pieau 1974).


Since then, ovary differentiation at a male-producing temperature under the effects of oestrogenic treatment has been obtained in all other reptilian species studied, including crocodylian, lizard and turtle species (Bull *et al.* 1988, Pieau *et al.* 1994a). The time-period during which the treatment is efficient coincides with the TSP (Gutzke & Chymiy 1988). During this period, binding of tritiated oestradiol by the gonads was shown to occur at both male- and female-producing temperatures (Smith & Joss 1994b).

Moreover, *oestrogen receptor* mRNAs were detected in putative male and female gonads as early as at the beginning of the TSP. Transcripts were distributed throughout the cortex and medulla in differentiating ovaries, whereas they were found in medullary testicular cords but not in the flattened parts of the surface epithelium in differentiating testes (Bergeron *et al.* 1998). However, as shown above, the development of an ovarian-like cortex under the effects of exogenous oestradiol remains possible after the TSP in the remnants of the germinal epithelium, indicating the persistence of oestrogen receptors in these regions.

In the freshwater turtle *T. scripta*, shifts from a male- to a female-producing temperature as well as oestrogenic treatments at the same embryonic stages during the TSP had similar chronological effects on gonadal structure, i.e. inhibition of testicular cord development preceding the proliferation of germ cells in the cortex (Wibbels *et al.* 1993). It can therefore be assumed that, at a female-producing temperature, oestrogens synthesized by the gonads themselves (see above) direct their differentiation into ovaries.

The implication of oestrogens in ovary differentiation has been irrefutably confirmed by the results of treatments with antioestrogens and aromatase inhibitors prior to and/or during the TSP. It suffices to recall here that treatments with the antioestrogen tamoxifen in *E. orbicularis* and the aromatase inhibitors fadrozole or letrozole in *E. orbicularis*, *T. scripta* and *C. serpentina* resulted in differentiation of testicular cords in gonads at a female-producing temperature: at hatching, the gonads were testes or ovotestes (for review see Pieau *et al.* 1999). In *E. orbicularis*, treatment with letrozole somewhat after the TSP also resulted in masculinization of gonads but was considerably less efficient than during the TSP; not only were the masculinized individuals less numerous, but all these individuals displayed ovotestes (Dorizzi *et al.* 1996, Belaïd *et al.* 2001).

A characteristic of masculinization of gonads under the effects of aromatase inhibitors is the presence of lacunar tubes, or lacunae, bordered by a flat epithelium and a higher Sertolian-like epithelium. These structures were observed in *E. orbicularis* after treatment during the TSP (Dorizzi *et al.* 1994, Richard-Mercier *et al.* 1995) and also after the TSP (Dorizzi *et al.* 1996, Belaïd *et al.* 2001). They were also observed in the sex-reversed gonads of genetic female chickens treated with an aromatase inhibitor prior to sexual differentiation (Vaillant *et al.* 2001a). In these gonads, the thickened part of the epithelium of the lacunae was shown to express the anti-Müllerian hormone (AMH), *SOX9* and *SF1* mRNAs as does the Sertolian epithelium of control testicular cords, whereas the flattened part did not. Lacunae bordered by a flat epithelium are normally formed in the ovarian medulla. Thus, during sex reversal, ovarian lacunae evolve as testicular cords, by transdifferentiation of their epithelium into a Sertolian epithelium (Vaillant *et al.* 2001b). In *E. orbicularis*,



**Figure 4** Molecular and physiological events in the gonads consistent with the involvement of endogenous oestrogens in the first steps of ovarian differentiation in freshwater and marine turtles (ER, oestrogen receptor).

testicular cords of ovotestes, resulting from treatment with letrozole, express *SOX9* mRNAs as testicular cords of control testes (Belaid *et al.* 2004).

### Conclusions and perspectives

Taken together, the data obtained from experiments and assays carried out on the gonads alone in turtle embryos provide evidence that, in reptiles exhibiting TSD, the gonads are the targets of temperature for their sexual differentiation and are the site of aromatase activity and oestrogen synthesis during the TSP. Gonadal oestrogen synthesis depends on the incubation temperature. At female-producing temperatures, as early as at the beginning of the TSP, endogenous oestrogens act locally on both the cortical and the medullary parts of the gonad to direct ovarian sex differentiation. The molecular and physiological events that support this major role of oestrogens are shown in Fig. 4.

In our opinion, contrary to the suggestion of Murdoch & Wibbels (2003), there cannot exist on one hand reptilian species such as *E. orbicularis*, *D. coriacea* and *M. Terrapin*, in which oestrogens are involved in ovarian sex differentiation as early as at the beginning of the TSP, and on the other hand reptilian species such as *C. porosus*, *A. mississippiensis* and *T. scripta*, in which oestrogens are involved in later stages of ovarian differentiation, i.e. only at the end or after the TSP. Using appropriate methods, aromatase activity, *aromatase* mRNA levels and oestrogen content in the gonads should be higher at the female- than at the male-producing temperature during the TSP in all reptilian species exhibiting TSD. Should this be otherwise, the results of treatments with exogenous estrogens and treatments with antioestrogens or aromatase inhibitors would be very difficult to interpret, as recognized by Murdoch & Wibbels (2003) themselves.

The thermosensitivity of gonadal sex differentiation has been demonstrated not only in many reptile species but also in several fish and some amphibian species. In



thermosensitive species, mitotic sex chromosomes are not distinguishable by classical cytological studies. However, a mechanism of genotypic sex determination (GSD) – male heterogamety, female heterogamety or polygenic – has been shown in these species by various methods, such as breeding tests after hormonal or thermal treatments, examination of lampbrush chromosomes or the electrophoretic pattern of sex-linked enzymes (fish: for review see Baroiller & Guiguen 2001, amphibians: for review see Chardard *et al.* 2004). Therefore, the effects of temperature are superimposed on the GSD mechanism.

As in reptiles, in fish and amphibians oestrogens are involved in ovarian differentiation, and aromatase plays a key role in this process. In the tilapia *Oreochromis niloticus*, the differentiating ovary displays a stronger expression of the *aromatase* gene and higher levels of oestradiol-17 $\beta$  than the differentiating testis (D’Cotta *et al.* 2001). In the newt *Pleurodeles walil*, transcripts of the *aromatase* gene have been detected as early as at the beginning of the TSP in gonad/mesonephros complexes and their levels are higher in female gonads than in male gonads by the end of this period (Kuntz *et al.* 2003). It is therefore reasonable to assume that the same mechanism of action of temperature, resulting in the activation or the repression of transcription of the *aromatase* gene in gonads, exists in fish, amphibians and reptiles. Some possible mechanisms have been considered (Pieau 1996), but to date the molecular target of temperature remains unknown.

Oestrogens are also involved in ovarian differentiation of birds that all display a male ZZ/ female ZW mechanism of GSD (Elbrecht & Smith 1992). In chickens, *aromatase* transcripts have been detected in the two female gonads but not in the male gonads at day 6–6.5 of embryonic development, i.e. just prior to or at the very beginning of the first histological signs of gonadal sex differentiation (for review see Vaillant *et al.* 2001a). During the first steps of gonadal sex differentiation (days 7 and 8), *aromatase* gene expression is upregulated in female gonads while it remains undetectable in male gonads (Nishikimi *et al.* 2000, Oréal *et al.* 2002).

In marsupials, the implication of oestrogens in the first steps of ovarian differentiation can be expected since treatment of male pouch young with oestrogens leads to sex reversal of the testes: the gonads resemble ovotestes or ovaries with a clearly defined cortex containing germ cells having entered into meiosis (Burns 1961, Coveney *et al.* 2001). In eutherian mammals, *in vivo* treatment of gravid females with exogenous oestrogens does not lead to sex reversal of XY genetic male embryos (Greene *et al.* 1940). However, definitive evidence that oestrogens play a critical role in maintaining ovarian structure has been found recently. In XX genetic female mice with targeted disruption of the genes encoding oestrogen receptors  $\alpha$  and  $\beta$  (ER $\alpha\beta$ KO mice) or aromatase (ArKO mice), Leydig cells differentiate from interstitial cells, and seminiferous tubule-like structures presenting Sertoli cells expressing

SOX9 differentiate from pre-existing granulosa cells after birth (Couse *et al.* 1999, Britt *et al.* 2002, Dupont *et al.* 2003). If ArKO female mice are fed a diet containing phyto-oestrogens, the appearance of Leydig and Sertoli cells is postponed and reduced. Likewise, if oestradiol-17 $\beta$  is administered to such females, the number of Leydig and Sertoli cells decreases in the ovaries (Britt *et al.* 2002, for review see Britt & Findlay 2003). From these findings, it may be asked whether, in eutherian mammals, oestrogens are not also involved in the first steps of ovarian differentiation.

A very simple scheme for the involvement of the *aromatase* gene and oestrogens in gonadal sex differentiation of vertebrates including mammals has been proposed previously (Pieau *et al.* 1994b). In accordance with this view and considering the evolutionary aspects, Crews (1994) estimates that ‘vestigial remnants of a sex hormone mediation of sex determination might be present in mammals’.

Given the converging data obtained in non-mammalian vertebrates that provide evidence of a key role of aromatase and oestrogens in the first steps of ovarian differentiation and their possible implication in mammals, it seems to us that future research should be carried out in two main directions. The first direction should be to examine how transcription of the *aromatase* gene is regulated in gonads at the time of sex differentiation in species with strict GSD and species with TSD. The second direction should be to identify molecular and cellular targets of oestrogens in the cortical and the medullary parts of the gonads during sex differentiation. *A priori*, the same targets should be found in all vertebrates, whatever the mechanism of sex determination.

## Acknowledgements

We thank Michael Richardson (Leiden University) for providing photographs of *E. orbicularis* embryos and Myriam Barre for help with the figures.

## References

- Baroiller J-F & Guiguen Y 2001 Endocrine and environmental aspects of sex differentiation in gonochoristic fish. In *Genes and Mechanisms in Vertebrate Sex Determination*, pp 178–201. Eds G Scherer & M Schmid. Basel: Birkhäuser.
- Belaid B, Richard-Mercier N, Pieau C & Dorizzi M 2001 Sex reversal and aromatase in the European pond turtle: treatment with letrozole after the thermosensitive period for sex determination. *Journal of Experimental Zoology* **290** 490–497.
- Belaid B, NGuyen Cong H, Devilliers G, Richard-Mercier N, Pieau C & Dorizzi M 2004 Lymphoid aggregates in gonads of embryos, hatchlings and young of turtles with temperature-dependent sex determination. *Journal of Experimental Zoology* **301A** 160–168.
- Bergeron JM, Gahr M, Horan K, Wibbels T & Crews D 1998 Cloning and *in situ* hybridization analysis of estrogen receptor in the

- developing gonad of the red-eared slider turtle, a species with temperature-dependent sex determination. *Development, Growth and Differentiation* **40** 243–254.
- Bowden RM, Ewert MA & Nelson CE 2000 Environmental sex determination in a reptile varies seasonally and with yolk hormones. *Proceedings of the Royal Society* **267** 1745–1749.
- Bowden RM, Ewert MA & Nelson CE 2002 Hormone levels in yolk decline throughout development in the red-eared slider turtle (*Trachemys scripta elegans*). *General and Comparative Endocrinology* **129** 171–177.
- Britt KL & Findlay JK 2003 Regulation of the phenotype of ovarian somatic cells by estrogen. *Molecular and Cellular Endocrinology* **202** 11–17.
- Britt KL, Kerr J, O'Donnell L, Jones MEE, Drummond AE, Davis SR, Simpson ER & Findlay JK 2002 Estrogen regulates development of the somatic cell phenotype in the eutherian ovary. *FASEB Journal* **16** 1389–1397.
- Bull JJ, Gutzke WHN & Crews D 1988 Sex reversal by estradiol in three reptilian orders. *General and Comparative Endocrinology* **70** 425–428.
- Burns RK 1961 Role of hormones in the differentiation of sex. In *Sex and the Internal Secretions*, vol. 1, edn 3, pp 76–158. Ed. WC Young. Baltimore: Williams & Wilkins Co.
- Chardard D, Penrad-Mobayed M, Chesnel A, Pieau C & Dourmon C 2004 Thermal sex reversals in amphibians. In *Temperature-dependent Sex Determination in Vertebrates*. Eds N Valenzuela & V Lance. Washington DC: Smithsonian Books (In Press).
- Conley AJ, Elf P, Corbin CJ, Dubowsky S, Fivizzani A & Lang J W 1997 Yolk steroids decline during sexual differentiation in the alligator. *General and Comparative Endocrinology* **107** 191–200.
- Couse JF, Hewitt SC, Bunch DO, Sar M, Walker VR, Davis BJ & Korach KS 1999 Postnatal sex reversal of the ovaries in mice lacking estrogen receptors  $\alpha$  and  $\beta$ . *Science* **286** 2328–2331.
- Coveney D, Shaw G & Renfree MB 2001 Estrogen-induced gonadal sex reversal in the tammar wallaby. *Biology of Reproduction* **65** 613–621.
- Crews D 1994 Temperature, steroids and sex determination. *Journal of Endocrinology* **142** 1–8.
- D'Cotta H, Fostier A, Guiguen Y, Govoroun M & Baroiller J-F 2001 Aromatase plays a key role during normal and temperature-induced sex differentiation of tilapia *Oreochromis niloticus*. *Molecular Reproduction and Development* **59** 265–276.
- Desvages G & Pieau C 1991 Steroid metabolism in gonads of turtle embryos as a function of the incubation temperature of eggs. *Journal of Steroid Biochemistry and Molecular Biology* **39** 203–213.
- Desvages G & Pieau C 1992a Aromatase activity in gonads of turtle embryos as a function of the incubation temperature of eggs. *Journal of Steroid Biochemistry and Molecular Biology* **41** 851–853.
- Desvages G & Pieau C 1992b Time required for temperature-induced changes in gonadal aromatase activity and related gonadal structure in turtle embryos. *Differentiation* **52** 13–18.
- Desvages G, Girondot M & Pieau C 1993 Sensitive stages for the effects of temperature on gonadal aromatase activity in embryos of the marine turtle *Dermochelys coriacea*. *General and Comparative Endocrinology* **92** 54–61.
- Dorizzi M, Mignot T-M, Guichard A, Desvages G & Pieau C 1991 Involvement of oestrogens in sexual differentiation of gonads as a function of temperature in turtles. *Differentiation* **47** 9–17.
- Dorizzi M, Richard-Mercier N, Desvages G, Girondot M & Pieau C 1994 Masculinization of gonads by aromatase inhibitors in a turtle with temperature-dependent sex determination. *Differentiation* **58** 1–8.
- Dorizzi M, Richard-Mercier N & Pieau C 1996 The ovary retains male potential after the thermosensitive period for sex determination in the turtle *Emys orbicularis*. *Differentiation* **60** 193–201.
- Dupont S, Dennefeld C, Krust A, Chambon P & Mark M 2003 Expression of *Sox9* in granulosa cells lacking the estrogen receptors, ER $\alpha$  and ER $\beta$ . *Developmental Dynamics* **226** 103–106.
- Elbrecht A & Smith RG 1992 Aromatase enzyme activity and sex determination in chickens. *Science* **255** 467–470.
- Elf PK 2003 Yolk steroid hormones and sex determination in reptiles with TSD. *General and Comparative Endocrinology* **132** 349–355.
- Elf PK, Lang JW & Fivizzani AJ 2002 Dynamics of yolk steroid hormones during development in a reptile with temperature-dependent sex determination. *General and Comparative Endocrinology* **127** 34–39.
- Forbes TR 1938 Studies on the reproductive system of the alligator. II. The effects of prolonged injections of estrone in the immature alligator. *Journal of Experimental Zoology* **78** 335–367.
- Gabriel WN, Blumberg B, Sutton S, Place AR & Lance VA 2001 Alligator aromatase cDNA sequence and its expression in embryos at male and female incubation temperatures. *Journal of Experimental Zoology* **290** 439–448.
- Greene RR, Burrill MW & Ivy AC 1940 Experimental intersexuality. The effects of oestrogens on the antenatal development of the rat. *American Journal of Anatomy* **67** 305–345.
- Gutierrez-Ospina G, Jiménez-Trejo FJ, Favila R, Moreno-Mendoza NA, Granados Rojas L, Barrios FA, Diaz-Cintra S & Merchant-Larios H 1999 Acetylcholinesterase-positive innervation is present at undifferentiated stages of the sea turtle *Lepidochelys olivacea* embryo gonads: implications for temperature-dependent sex determination. *Journal of Comparative Neurology* **410** 90–98.
- Gutzke WHN & Chymiy DB 1988 Sensitive periods during embryogeny for hormonally induced sex determination in turtles. *General and Comparative Endocrinology* **71** 265–267.
- Janzen FJ, Wilson ME, Tucker JK & Ford SP 1998 Endogenous yolk steroid hormones in turtles with different sex-determining mechanisms. *General and Comparative Endocrinology* **111** 306–317.
- Jeyasuria P & Place AR 1997 Temperature-dependent aromatase expression in developing diamondback terrapin (*Malaclemys terrapin*) embryos. *Journal of Steroid Biochemistry and Molecular Biology* **61** 415–425.
- Jeyasuria P & Place AR 1998 Embryonic brain–gonadal axis in temperature-dependent sex determination of reptiles: a role for P450 aromatase (CYP 19). *Journal of Experimental Zoology* **281** 428–449.
- Kuntz S, Chesnel A, Duterque-Coquillaud M, Grillier-Vuissoz I, Callier M, Dourmon C, Flament S & Chardard D 2003 Differential expression of P450 aromatase during gonadal sex differentiation and sex reversal of the newt *Pleurodeles waltl*. *Journal of Steroid Biochemistry and Molecular Biology* **84** 89–100.
- Miller JD 1985 Embryology of marine turtles. In *Biology of the Reptilia*, vol. 14, pp 270–328. Eds C Gans, F Billett & PFA Maderson. New York: Wiley.
- Moreno-Mendoza N, Harley VR & Merchant-Larios H 1999 Differential expression of SOX9 in gonads of the sea turtle *Lepidochelys olivacea* at male- or female-promoting temperatures. *Journal of Experimental Zoology* **284** 705–710.
- Moreno-Mendoza N, Harley VR & Merchant-Larios H 2001 Temperature regulates SOX9 expression in cultured gonads of *Lepidochelys olivacea*, a species with temperature sex determination. *Developmental Biology* **229** 319–326.
- Murdock C & Wibbels T 2003 Cloning and expression of aromatase in a turtle with temperature-dependent sex determination. *General and Comparative Endocrinology* **130** 109–119.
- Nicolas JC, Boussioux AM, Descomps B & Crastes de Paulet A 1979 Enzymatic determination of estradiol and estrone in plasma and urine. *Clinica Chimica Acta* **92** 1–9.
- Nishikimi H, Kansaku N, Saito N, Usami M, Ohno Y & Shimada K 2000 Sex differentiation and mRNA expression of P450c17, P450 arom and AMH in gonads of the chicken. *Molecular Reproduction and Development* **55** 20–30.

- Oréal E, Mazaud S, Picard J-Y, Magre S & Carré-Eusèbe D 2002 Different patterns of anti-Müllerian hormone expression, as related to DMRT1, SF-1, WT1, GATA-4, Wnt-4, and Lhx9 expression, in the chick differentiating gonads. *Developmental Dynamics* **225** 221–232.
- Pieau C 1970 Effets de l'œstradiol sur l'appareil génital de l'embryon de tortue mauresque (*Testudo graeca* L.). *Archives d'Anatomie Microscopique et de Morphologie Expérimentale* **59** 295–318.
- Pieau C 1974 Différenciation du sexe en fonction de la température chez les embryons d'*Emys orbicularis* L. (Chélonien); effets des hormones sexuelles. *Annales d'Embryologie et de Morphogénèse* **7** 365–394.
- Pieau C 1976 Données récentes sur la différenciation sexuelle en fonction de la température chez les embryons d'*Emys orbicularis* L. (Chélonien). *Bulletin de la Société Zoologique de France* **101** (Suppl. 4) 46–53.
- Pieau C 1996 Temperature variation and sex determination in reptiles. *BioEssays* **18** 19–26.
- Pieau C & Dorizzi M 1981 Determination of temperature sensitive stages for sexual differentiation of the gonads in embryos of the turtle, *Emys orbicularis*. *Journal of Morphology* **170** 373–382.
- Pieau C, Girondot M, Richard-Mercier N, Desvages G, Dorizzi M & Zaborski P 1994a Temperature sensitivity of sexual differentiation of gonads in the European pond turtle: hormonal involvement. *Journal of Experimental Zoology* **270** 86–94.
- Pieau C, Girondot M, Desvages G, Dorizzi M, Richard-Mercier N & Zaborski P 1994b Environmental control of gonadal differentiation. In *The Differences Between The Sexes*, pp 433–448. Eds RV Short & E Balaban. Cambridge: Cambridge University Press.
- Pieau C, Dorizzi M, Richard-Mercier N & Desvages G 1998 Sexual differentiation of gonads as a function of temperature in the turtle *Emys orbicularis*: endocrine function, intersexuality and growth. *Journal of Experimental Zoology* **281** 400–408.
- Pieau C, Dorizzi M & Richard-Mercier N 1999 Temperature-dependent sex determination and gonadal differentiation in reptiles. *Cellular and Molecular Life Sciences* **55** 887–900.
- Place AR, Lang J, Gavasso S & Jeyasuria P 2001 Expression of P450 arom in *Malacemys terrapin* and *Chelydra serpentina*: a tale of two sites. *Journal of Experimental Zoology* **290** 673–690.
- Raynaud A & Pieau C 1985 Embryonic development of the genital system. In *Biology of the Reptilia*, vol. 15, pp 149–300. Eds C Gans & F Billett. New York: Wiley.
- Renous S, Rimblot-Baly F, Fretey J & Pieau C 1989 Caractéristiques du développement embryonnaire de la Tortue Luth, *Dermochelys coriacea* (Vandelli, 1761). *Annales des Sciences Naturelles, Zoologie, Paris* **10** 197–229.
- Richard-Mercier N, Dorizzi M, Desvages G, Girondot M & Pieau C 1995 Endocrine sex reversal of gonads by the aromatase inhibitor letrozole (CGS 20267) in *Emys orbicularis*, a turtle with temperature-dependent sex determination. *General and Comparative Endocrinology* **100** 314–326.
- Rimblot F, Fretey J, Mrosovsky N, Lescure J & Pieau C 1985 Sexual differentiation as a function of the incubation temperature of eggs in the sea-turtle *Dermochelys coriacea* (Vandelli, 1761). *Amphibia-Reptilia* **6** 83–92.
- Risley PL 1941 A comparison of effects of gonadotropic and sex hormones on the urogenital systems of juvenile terrapins. *Journal of Experimental Zoology* **87** 477–515.
- Salame-Mendez A, Herrera-Munoz J, Moreno-Mendoza N & Merchante-Larios H 1998 Response of diencephalon but not the gonad to female-promoting temperature with elevated estradiol levels in the sea turtle *Lepidochelys olivacea*. *Journal of Experimental Zoology* **280** 304–313.
- Smith CA & Joss JMP 1994a Steroidogenic enzyme activity and ovarian differentiation in the saltwater crocodile, *Crocodylus porosus*. *General and Comparative Endocrinology* **93** 232–245.
- Smith CA & Joss JMP 1994b Uptake of <sup>3</sup>H-estradiol by embryonic crocodile gonads during the period of sexual differentiation. *Journal of Experimental Zoology* **270** 219–224.
- Smith CA, Elf PK, Lang JW & Joss JMP 1995 Aromatase enzyme activity during gonadal sex differentiation in alligator embryos. *Differentiation* **58** 281–290.
- Vaillant S, Dorizzi M, Pieau C & Richard-Mercier N 2001a Sex reversal and aromatase in chicken. *Journal of Experimental Zoology* **290** 727–740.
- Vaillant S, Magre S, Dorizzi M, Pieau C & Richard-Mercier N 2001b Expression of *AMH*, *SF1* and *SOX9* in gonads of genetic female chickens during sex reversal induced by an aromatase inhibitor. *Developmental Dynamics* **22** 228–237.
- Vivien JH & Stéfan Y 1958 Féminisation des gonades de jeunes tortues d'eau *Emys leprosa* mâles par action du diéthylstilboestrol. *Comptes Rendus des Séances de la Société de Biologie* **152** 649–652.
- White RB & Thomas P 1992a Adrenal-kidney and gonadal steroidogenesis during sexual differentiation in a reptile with temperature-dependent sex determination. *General and Comparative Endocrinology* **88** 10–19.
- White RB & Thomas P 1992b Stimulation of *in vitro* steroidogenesis by pituitary hormones in a turtle (*Trachemys scripta*) within the temperature-sensitive period for sex determination. *Biology of Reproduction* **47** 952–959.
- Wibbels T, Gideon P, Bull JJ & Crews D 1993 Estrogen- and temperature-induced medullary cord regression during gonadal differentiation in a turtle. *Differentiation* **53** 149–154.
- Willingham E, Baldwin R, Skipper J & Crews D 2000 Aromatase activity during embryogenesis in the brain and adrenal-kidney-gonad of the red-eared slider turtle, a species with temperature-dependent sex determination. *General and Comparative Endocrinology* **119** 202–207.
- Yntema CL 1968 A series of stages in the embryonic development of *Chelydra serpentina*. *Journal of Morphology* **125** 219–252.

Received 27 January 2004

Accepted 18 March 2004

Made available online as an

Accepted Preprint 25 March 2004