

# OGLE-TR-211 – a new transiting inflated hot Jupiter from the OGLE survey and ESO LP666 spectroscopic follow-up program\*

A. Udalski<sup>1,2</sup>, F. Pont<sup>3</sup>, D. Naef<sup>4</sup>, C. Melo<sup>4</sup>, F. Bouchy<sup>5</sup>, N. C. Santos<sup>6</sup>, C. Moutou<sup>7</sup>, R. F. Díaz<sup>8</sup>, W. Gieren<sup>9</sup>, M. Gillon<sup>3</sup>, S. Hoyer<sup>10</sup>, M. Mayor<sup>3</sup>, T. Mazeh<sup>11</sup>, D. Minniti<sup>12</sup>, G. Pietrzyński<sup>1,2,9</sup>, D. Queloz<sup>3</sup>, S. Ramirez<sup>5</sup>, M. T. Ruiz<sup>10</sup>, A. Shporer<sup>11</sup>, O. Tamuz<sup>11</sup>, S. Udry<sup>3</sup>, M. Zoccali<sup>12</sup>, M. Kubiak<sup>1,2</sup>, M. K. Szymański<sup>1,2</sup>, I. Soszyński<sup>1,2</sup>, O. Szewczyk<sup>1,2,9</sup>, K. Ulaczyk<sup>1,2</sup>, and Ł. Wyrzykowski<sup>2,13</sup>

<sup>1</sup> Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warsaw, Poland  
e-mail: udalski@astrouw.edu.pl

<sup>2</sup> The OGLE Team

<sup>3</sup> Observatoire de Genève, 51 Chemin des Maillettes, 1290 Sauverny, Switzerland

<sup>4</sup> European Southern Observatory, Casilla 19001, Santiago 19, Chile

<sup>5</sup> Institut d'Astrophysique de Paris, 98bis Bd Arago, 75014 Paris, France

<sup>6</sup> Centro de Astrofísica, Universidade do Porto, Rua das Estrelas, 4150-762 Porto, Portugal

<sup>7</sup> Laboratoire d'Astrophysique de Marseille, Traverse du Siphon, BP 8, Les Trois Lucs, 13376 Marseille Cedex 12, France

<sup>8</sup> Instituto de Astronomía y Física del Espacio (CONICET-UBA), Buenos Aires, Argentina

<sup>9</sup> Departamento de Física, Astronomy Group, Universidad de Concepción, Casilla 160-C, Concepción, Chile

<sup>10</sup> Department of Astronomy, Universidad de Chile, Santiago, Chile

<sup>11</sup> School of Physics and Astronomy, R. and B. Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel

<sup>12</sup> Departamento de Astronomía y Astrofísica, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile

<sup>13</sup> Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, UK

Received 26 November 2007 / Accepted 4 February 2008

## ABSTRACT

We present results of the photometric campaign for planetary and low-luminosity object transits conducted by the OGLE survey in the 2005 season (Campaign #5). About twenty of the most promising candidates discovered in these data were subsequently verified spectroscopically with the VLT/FLAMES spectrograph.

One of the candidates, OGLE-TR-211, reveals clear changes of radial velocity with a small amplitude of 82 m/s, varying in phase with photometric transit ephemeris. Further analysis confirms the planetary nature of this system. Follow-up precise photometry of OGLE-TR-211 with VLT/FORS, together with radial velocity spectroscopy, supplemented with high-resolution, high S/N VLT/UVES spectra allowed us to derive parameters of the planet and host star. OGLE-TR-211b is a hot Jupiter orbiting an F7-8 spectral type dwarf star with a period of 3.68 days. The mass of the planet is equal to  $1.03 \pm 0.20 M_{\text{Jup}}$ , while its radius  $1.36^{+0.18}_{-0.09} R_{\text{Jup}}$ . The radius is about 20% larger than the typical radius of hot Jupiters of similar mass. OGLE-TR-211b is, then, another example of inflated hot Jupiters – a small group of seven exoplanets with large radii and unusually low densities – objects that are a challenge to the current models of exoplanets.

**Key words.** planetary systems – stars: individual: OGLE-TR-211

## 1. Introduction

The discovery of the first transit of an extrasolar planet around HD 209458 (Charbonneau et al. 2000; Henry et al. 2000) opened a new era in the extrasolar planet field – the epoch of searches for extrasolar planets with large-scale photometric surveys. While this channel of finding exoplanets developed relatively slowly in the first couple of years after that discovery, the past two years brought fast acceleration due to the search methods maturing. At present about 30 transiting planets are known (cf. <http://www.inscience.ch/transits/>) and they constitute about 10% of all known exoplanets.

Transiting planets play an especially important role among the known exoplanets. These are the only objects for which the

most important parameters (e.g., radius, mass, density) can be precisely derived from observations, thereby providing direct comparison with models and thus allowing a better understanding of the exoplanet structure and evolution. Also, a wide variety of follow-up observations can be performed on transiting system to allow, for example, studies of planetary atmospheres and search for other planetary companions, etc.

Transiting planets are discovered via two channels: photometric follow-up of spectroscopically found exoplanets and the classical transit method approach, i.e., large scale photometric surveys providing transiting candidates that have to then be verified and confirmed spectroscopically. Both approaches have been successful in providing examples of very wide diversity among transiting planets – regular hot Jupiters, very hot Jupiters on extremely short 1–2 day period orbits, inflated objects with radii much larger than expected from modeling, small Neptune-sized objects, or very massive planets on eccentric orbits. Large numbers of new extrasolar transiting planets are expected to be

\* Based on observations made with the FORS1 camera and the FLAMES/UVES spectrograph at the VLT, ESO, Chile (program 07.C-0706, 076.C-0122, and 177.C-0666) and 1.3-m Warsaw Telescope at Las Campanas Observatory, Chile.

discovered in the next couple of years from space missions like Corot or Kepler, in particular small-size planets undetectable in ground-based searches.

The Optical Gravitational Lensing Experiment (OGLE) was the first successful photometric survey to discover a large number of transiting candidates that were subsequently verified spectroscopically (Udalski et al. 2002a,b,c, 2003). Five of these candidates, OGLE-TR-56, OGLE-TR-113, OGLE-TR-132, OGLE-TR-111, and OGLE-TR-10, turned out to be extrasolar planetary systems (Konacki et al. 2003; Bouchy et al. 2004; Konacki et al. 2004; Pont et al. 2004; Bouchy et al. 2005; Konacki et al. 2005), the first extrasolar planets discovered with the transit method. Moreover, two planet-sized stars with the lowest known masses were also found (Pont et al. 2005b, 2006a). In addition, the OGLE photometric transit campaigns provided huge amounts of observational material allowing better understanding of the problems of photometric transit searches, data systematics, etc. (Gould et al. 2006; Pont et al. 2006b).

The OGLE transit campaigns have been conducted every year since 2001. Results of the spectroscopic follow-up of candidates discovered during Campaigns #1 (2001) and #2 (2002) were published by Bouchy et al. (2005) and Pont et al. (2005a), respectively. Recently, analysis of the near threshold candidates from Campaign #2 led to the discovery of the sixth OGLE transiting exoplanet, OGLE-TR-182 (Pont et al. 2007).

In this paper we present results of OGLE Campaign #5 conducted in 2005 and spectroscopic follow-up of the best transiting candidates detected in these data. A new transiting extrasolar planetary system OGLE-TR-211 was found among them. In the following sections we provide details of the photometric and spectroscopic observations of OGLE new candidates and derive parameters of a planetary companion in OGLE-TR-211 system.

## 2. OGLE planetary transit campaign #5

OGLE planetary transit campaigns became a standard part of the OGLE observing schedule. About 75% of observing time was devoted to this sub-project of the OGLE survey every southern fall – from February to April. The OGLE planetary campaign #5 was conducted by OGLE in the 2005 observing season from February 2, 2005 to June 23, 2005. Observations were carried out with the 1.3-m Warsaw Telescope at Las Campanas Observatory, equipped with the  $8192 \times 8192$  pixel CCD mosaic camera (Udalski 2003). The observing strategy was similar to previous campaigns, in particular #3 and #4 (Udalski et al. 2004). Because the experience gathered during the spectroscopic observations of the first OGLE candidates indicated some trouble with reasonable spectroscopic follow-up of the faintest objects from typical OGLE candidate lists, the exposure time during campaign #5 was shortened to 120 s. This allowed one more field to be added thereby increasing the number of observed fields to four and covering about 1.4 square degrees of the Galactic disk regions in the Carina constellation. The cadence time for each of these four fields was about 16 min, and their equatorial coordinates are listed in Table 1. All observations were collected through the *I*-band filter. The median seeing of the all observations was about 1.1 arcsec. Altogether, about 1320 images of each of the fields were secured during the campaign. Also a few *V*-band observations were taken for color information and CMD construction.

Collected data were reduced in a similar way to those of Campaigns #3 and #4 (Udalski et al. 2004). The selection of transit candidates was also performed similarly: all non-variable objects, that is, those with the rms of the average magnitude of

**Table 1.** Equatorial coordinates of the fields observed during the OGLE Campaign #5.

Field	RA(J2000)	Dec(J2000)
CAR107	10:47:15	−62:00:25
CAR108	10:47:15	−61:24:35
CAR109	10:42:10	−62:10:25
CAR110	10:42:15	−61:34:35

all observations smaller than 0.015 mag, were subject to the de-trending algorithm of Kruszcwski & Semeniuk (2003) and then to a transit search procedure using the BLS algorithm of Kovács et al. (2002). Altogether, light curves of about 50 000 stars were analyzed. The list of transiting candidates was prepared after careful visual inspection of objects that passed the BLS criteria. Clear false cases triggered, for example, by noisy light curves were removed. We also excluded objects with any depth of transits greater than 50 mmag, with a clear signature of ellipsoidal effect indicating massive secondary or those with clear V-shape of transits indicating grazing eclipses of a binary star. The minimum number of individual transits collected during the campaign was set to three. For all the candidates, the limits on the size of a transiting companion were calculated as in Udalski et al. (2004). All candidates with a companion whose lower limit of radius was larger than  $0.2 R_{\odot}$  were removed from the final list.

Table 2 lists the transit candidates from the OGLE Campaign #5 that passed the photometric search criteria. The naming convention follows the standard OGLE convention, i.e., OGLE-TR-NNN. About twenty candidates for transiting planetary systems were found. These objects were selected as targets for spectroscopic follow-up observations conducted under the LP666 program. The photometry of all selected candidates is available from the OGLE Internet archive at <ftp://ftp.astrouw.edu.pl/ogle/ogle3/transits/tr201-219/>

## 3. Spectroscopic follow-up

Spectroscopic follow-up observations of the OGLE Campaign #5 transit candidates were carried out during three observing slots allocated to the LP666 program in April/May 2006, February 2007, and April 2007 on VLT with the FLAMES spectrograph. Also, part of the Geneva group observing time under program 07.C-0706 on the same instrument in February 2006 was used. Unfortunately the weather conditions during all these observing runs were exceptionally unfavorable with many cloudy or large seeing nights ( $>2$  arcsec), considerably limiting the results. In particular after the initial screening of all candidates, only the most promising objects were observed further. The remaining objects, in contrast to the previous follow-up observations (Bouchy et al. 2004; Pont et al. 2005a), were left without any full characterization. The strategy of the spectroscopic follow-up was identical to the one described by Pont et al. (2007).

Table 2 includes the column with spectroscopic status resulting from the collected spectra. Only one candidate turned out to be a very promising planetary candidate revealing low-amplitude radial-velocity variation in phase with photometric ephemeris, namely OGLE-TR-211. It was then extensively observed during all following spectroscopic runs, allowing confirmation of its planetary nature. Three more candidates do not show significant radial velocity variations. Table 3 lists these objects and provides the upper limits on radial velocity semi-amplitudes,  $K$ , at the  $2\text{-}\sigma$  level, the corresponding maximum

**Table 2.** Planetary and low-mass object transit candidates from the OGLE Campaign #5 conducted in 2005.

Name	RA [2000]	Dec [2000]	Period [days]	$T_c$ [Hel.JD]	$I$ [mag]	Depth [mag]	Status
OGLE-TR-201	10:48:31.61	-62:01:01.8	2.3680	2453404.860	15.6	0.016	fast rotator
OGLE-TR-202	10:46:06.06	-61:52:11.3	1.6545	2453404.438	13.6	0.017	not observed
OGLE-TR-203	10:49:32.04	-61:35:38.2	3.3456	2453406.178	15.6	0.014	not observed
OGLE-TR-204	10:47:39.44	-61:19:00.8	3.1097	2453405.515	14.8	0.026	SB2
OGLE-TR-205	10:45:53.40	-61:09:22.1	1.7501	2453404.601	16.0	0.015	not observed
OGLE-TR-206	10:45:22.00	-61:28:28.8	3.2658	2453403.246	13.8	0.006	no variation
OGLE-TR-207	10:40:10.41	-61:53:55.5	4.8170	2453406.978	14.3	0.021	SB2
OGLE-TR-208	10:39:54.18	-61:58:07.7	4.5025	2453406.567	15.3	0.022	SB2
OGLE-TR-209	10:40:56.27	-62:14:20.2	2.2056	2453403.657	15.0	0.022	no variation
OGLE-TR-210	10:40:44.21	-62:13:21.6	2.2427	2453403.012	15.2	0.032	fast rotator
OGLE-TR-211	10:40:14.51	-62:27:19.8	3.6772	2453406.271	14.3	0.008	planet
OGLE-TR-212	10:43:37.95	-61:45:01.5	2.2234	2453404.170	16.3	0.016	blend?
OGLE-TR-213	10:43:00.85	-61:51:10.2	6.5746	2453403.301	15.3	0.036	SB2
OGLE-TR-214	10:44:27.96	-61:35:35.7	3.6010	2453403.090	16.5	0.023	SB1
OGLE-TR-215	10:43:48.69	-61:40:00.3	4.9237	2453407.616	14.8	0.016	no variation
OGLE-TR-216	10:42:02.43	-61:20:16.0	1.9763	2453403.064	14.6	0.011	blend?
OGLE-TR-217	10:40:59.73	-61:30:38.4	5.7208	2453404.104	16.1	0.037	no ccf
OGLE-TR-218	10:41:12.74	-61:28:20.8	2.2488	2453404.697	14.5	0.020	fast rotator
OGLE-TR-219	10:40:53.40	-61:43:15.1	9.7466	2453405.683	15.1	0.032	SB2

Notes: SB1 – single line spectroscopic binary; SB2 – double line spectroscopic binary; no ccf – lack of the cross-correlation function.

**Table 3.** OGLE transit candidates with no significant radial velocity variation.

Object	$K$ limit [m/s]	Mass limit [ $M_{\text{Jup}}$ ]	Radius limit [ $R_{\text{Jup}}$ ]
OGLE-TR-206	<59	<0.42	>0.6
OGLE-TR-209	<16	<0.16	>0.8
OGLE-TR-215	<63	<0.52	>1.5

mass of a potential planet for a  $1 M_{\odot}$  host star, and the lower limit on the planetary radius (i.e. central transit) for such a primary.


#### 4. OGLE-TR-211 candidate

OGLE-TR-211 turned out to be the most promising candidate for a transiting planetary system from the OGLE Campaign #5 sample. After the initial spectroscopic follow-up observations confirming this status, it was decided to allocate a considerable amount of observing time to a precise characterization of this object. OGLE-TR-211 is relatively bright ( $I \approx 14.3$  mag) in the OGLE sample. Its equatorial coordinates are listed in Table 2, while Fig. 1 shows the finding chart –  $2' \times 2'$  OGLE  $I$ -band image. OGLE-TR-211 is listed in the 2MASS and DENIS catalogs as 2MASS10401438-6227201 and J104014.3-622720, respectively.

##### 4.1. Photometry

The original OGLE 2005 season photometric data covered eight partial transits. When folded with the period of 3.677 days, they revealed a clear transit about 8 mmag deep. The estimated lower limit of the transiting companion radius was  $0.12 R_{\odot}$  making OGLE-TR-211 a very good candidate for a transiting planet.


Due to the limited ( $\approx 3$  months long) duration of the OGLE campaign, the photometric period has a limited accuracy of  $\approx 10^{-4}P$  or worse in the case of low depth transits. In order to

**Fig. 1.** Finding chart for OGLE-TR-211. Size of the  $I$ -band OGLE sub-frame is  $120'' \times 120''$ , North is up and East to the left.

confirm the presence of transits in OGLE-TR-211 and refine the ephemeris, OGLE observed the star in the next observing seasons: 2006 and 2007. The first additional transit data points were collected in January 2006, providing a necessary confirmation and update of the photometric orbit for the coming spectroscopic follow-up observations. Up to June 2007 eight additional transits of OGLE-TR-211 were covered. Thanks to the over two year-long baseline, these observations allowed us to derive precise photometric ephemeris of OGLE-TR-211, as listed in Table 5 (see Sect. 4.4).

The very small depth of transits combined with  $\approx 5$  mmag accuracy of individual OGLE measurements makes determining the precise shape of transit difficult. Therefore, similarly to other cases of OGLE transiting planets, it was decided to conduct a





**Fig. 2.** *Upper panel:* OGLE light curve of OGLE-TR-211, *lower panel:* VLT observations. The continuous line represents the best model of the transit.

photometric follow-up of this object on a much larger telescope. Three photometric runs on VLT with the FORS camera were performed in the period of March/June 2006.

The first-run VLT images of OGLE-TR-211 were obtained on March 16, 2006 through a *V*-band filter with the exposure time of 8 s. They cover the pre-ingress, ingress, and bottom of the transit. Unfortunately the egress of the transit was missed. Two additional runs were carried out on May 25, 2006 and June 30, 2006. In these cases observations were obtained through *V* and *R* filters and the exposure time varied from 25 to 80 s depending on seeing. Unfortunately, both of these runs cover a practically non-variable part of the transit light curve. The May observations show only the flat bottom of the transit with only a small trace of starting egress at the very end. The June observations cover only the post transit light curve with the very end of egress at the beginning. Thus, the full reconstruction of the precise VLT light curve of OGLE-TR-211 became difficult.

The VLT data were reduced using the OGLE data pipeline (Udalski 2003), based on the difference image analysis (DIA) method. Because the March 2006 observations that defined a considerable part of the transit shape were obtained in the *V*-band, only *V*-band images from the remaining observing runs were used. To have similar time resolution and to lower the scatter of the March observations, these data were binned to have an effective resolution of 3 min. The March light curve indicates that the transit depth is about 8 mmag – practically identical with what resulted from OGLE observations, which is reassuring. Therefore the May data were phased with the photometric period resulting from OGLE observations and adjusted in magnitude scale in such a way that the observed bottom of the transit corresponds to the level of the bottom observed in March 2006. Similarly, the June photometry was shifted so the post-egress part of transit corresponds to the pre-ingress part observed in March. The lower panel of Fig. 2 shows the reconstructed VLT light curve of OGLE-TR-211, while the OGLE light curve is presented in the upper panel.

The VLT data were also reduced using the DecPhot PSF fitting software (Gillon et al. 2006). While the results were generally similar, this photometry was not used for further analysis because of difficulties with the normalization between runs.

**Table 4.** Radial velocity measurements for OGLE-TR-211.

Date	Phase	$VR$	$VR_{\text{rectified}}$	$\sigma_{VR}$	$BS$	$\sigma_{BS}$
[JD-2 450 000]		[ $\text{km s}^{-1}$ ]	[ $\text{km s}^{-1}$ ]	[ $\text{km s}^{-1}$ ]	[ $\text{km s}^{-1}$ ]	[ $\text{km s}^{-1}$ ]
3793.81192	0.3889	18.784	18.784	0.055	0.132	0.071
3794.75446	0.6452	18.866	18.866	0.053	0.073	0.069
3852.50001	0.3487	18.720	18.720	0.053	0.007	0.033
3853.51724	0.6254	18.875	18.875	0.053	-0.055	0.016
3854.54151	0.9039	18.841	18.841	0.051	0.038	0.028
3855.50251	0.1652	18.711	18.711	0.055	-0.020	0.088
3855.59958	0.1916	18.816	18.816	0.060	0.025	0.034
3858.49787	0.9798	18.906	18.906	0.054	0.027	0.058
3881.57542	0.2556	18.766	18.766	0.053	–	–
4143.83440	0.5753	18.860	18.860	0.046	-0.004	0.018
4144.71247	0.8141	18.911	18.911	0.046	-0.043	0.032
4145.72709	0.0900	18.762	18.762	0.048	0.249	0.100
4149.81577	0.2019	18.681	18.681	0.047	-0.146	0.031
4150.77874	0.4638	18.891	18.891	0.046	-0.070	0.122
4203.55370	0.8156	18.745	18.925	0.046	0.038	0.052
4204.51070	0.0759	18.623	18.803	0.046	-0.036	0.058
4205.59031	0.3695	18.657	18.837	0.046	-0.043	0.047
4207.70041	0.9433	18.658	18.838	0.046	0.073	0.061
4208.68020	0.2097	18.578	18.758	0.046	0.074	0.051
4209.61470	0.4639	18.517	18.697	0.049	-0.118	0.074


#### 4.2. Radial velocity follow-up

Twenty high-resolution spectroscopic observations of OGLE-TR-211 were obtained with the FLAMES/UVES spectrograph on VLT during four spectroscopic follow-up runs. The strategy of observations and reduction procedures were identical to the description in Pont et al. (2007). The resulting radial velocities of OGLE-TR-211 with their errors are listed in Table 4. The errors include additional 45 m/s term derived from the estimation of stability of the instrument and added in quadrature to the photon noise radial velocity errors to account for possible systematic errors.

After two 2006 runs, it was clearly found that the radial velocities of OGLE-TR-211 follow the photometric period and reveal  $\approx 100$  m/s semi-amplitude variation in appropriate phase with planetary interpretation. However, during the last 2007 April run, while this variation was clearly seen, the  $\gamma$  zero point of radial velocities was shifted by about 180 m/s, which is more than  $3\text{-}\sigma$  of the typical accuracy of observations. This is a considerably larger shift than the shifts noted by Pont et al. (2007) during the analysis of the recently discovered transiting planet OGLE-TR-182b.

Although it cannot be fully ruled out that the shift has some instrumental origin, it is more likely that it corresponds to a real change in the average radial velocity. Unfortunately, the collected dataset is too limited to draw any sound conclusions about its origin. We can only speculate that it may be caused by the presence of an additional companion in the OGLE-TR-211 system having a wide orbit with a period of several years. Additional spectroscopic observations are necessary to verify this interpretation.

For the purpose of characterizing the planetary companion of OGLE-TR-211, the  $\gamma$  velocity was fitted separately for each observing run for a preliminary fit of the spectroscopic orbit. Only the shift of  $\gamma$  velocity of the 2007 April run turned out to be statistically significant. It was subtracted from original radial velocities for the final fit presented in the upper panel of Fig. 3. Rectified radial velocities of OGLE-TR-211 are also listed in Table 4.



**Fig. 3.** Upper panel: radial velocity observations for OGLE-TR-211, phased with the photometric transit ephemeris. Solid line represents the best fit assuming no eccentricity. Two periods are presented for clarity. Lower panel: bisector span values plotted as a function of phase for testing blending scenarios.

Because blending of an eclipsing star with a bright third light source can cause radial velocity variations comparable to a planetary signal, line profiles of the spectra of OGLE-TR-211 were examined for the correlation of the average line bisector spans (BS – see the last columns of Table 4) with phase and observed radial velocities (e.g., Santos et al. 2002; Torres et al. 2004). No such correlation was found, allowing us to rule out blending scenario (lower panel of Fig. 3).

#### 4.3. Host star spectroscopy

To derive parameters of the host star of OGLE-TR-211, high-resolution spectra were obtained with the UVES spectrograph on the VLT working in the slit mode on May 7 and 8, 2007. The spectra were summed so the coadded spectrum reached an S/N of about 100 in the  $\lambda \approx 6707 \text{ \AA}$  lithium line region. Measurements followed the method described in Santos et al. (2006), and the results are listed in Table 5. Temperature and gravity indicates that the host star is a dwarf of F7-8 spectral type and metallicity somewhat larger than solar.

#### 4.4. OGLE-TR-211 parameters

The parameters of OGLE-TR-211 were derived with the standard method used in the case of the previous OGLE planets (Bouchy et al. 2004; Pont et al. 2007). The OGLE and VLT photometry with a red noise value equal to the photon noise was fitted for  $P$ ,  $T$ ,  $b$ ,  $R/R_*$ , and  $a/R_*$ . The eccentricity of the orbit in the final fitting was assumed to be zero, as practically all the observed orbits of short period transiting planets are circularized by tidal interactions. Also our preliminary fit of the radial velocity curve with the orbital period fixed at the photometric value yielded eccentricity  $e = 0.16 \pm 0.22$ , consistent with circular orbit at better than  $1-\sigma$  level. Model parameters with their  $1-\sigma$  errors from chi-square analysis including the effect of correlated noise are listed in Table 5.

**Table 5.** Parameters of the OGLE-TR-211 system.

Period [days]	$3.67724 \pm 0.00003$
Transit epoch [Hel. JD]	$2453428.334 \pm 0.003$
Transit duration [days]	$0.1838 \pm 0.0055$
$V/R$ semi-amplitude [m/s]	$82 \pm 16$
Semi-major axis [AU]	$0.051 \pm 0.001$
Orbit-radius ratio [ $a/R_*$ ]	$6.75^{+0.23}_{-2.28}$
Radius ratio [ $R/R_*$ ]	$0.085 \pm 0.004$
$b$	$0.68^{+0.07}_{-0.68}$
Orbital angle [ $^\circ$ ]	$>82.7 (1-\sigma)$
Light Curve Model:	
OGLE rms [mmag]	4.1
VLT rms [mmag]	1.6
Radial Velocity Model:	
rms [m/s]	48
$T_{\text{eff}}$ [K]	$6325 \pm 91$
$\log g$	$4.22 \pm 0.17$
$\eta$ [ $\text{km s}^{-1}$ ]	$1.63 \pm 0.21$
[Fe/H]	$0.11 \pm 0.10$
Stellar radius [ $R_\odot$ ]	$1.64^{+0.21}_{-0.07}$
Stellar mass [ $M_\odot$ ]	$1.33 \pm 0.05$
Planet radius [ $R_{\text{Jup}}$ ]	$1.36^{+0.18}_{-0.09}$
Planet mass [ $M_{\text{Jup}}$ ]	$1.03 \pm 0.20$

Then the radius and mass of the host star were estimated with the maximum-likelihood method combining the observed spectroscopic parameters of the host star, transit parameters obtained from the light curve fit and Girardi et al. (2002) stellar evolution models. Table 5 lists the parameters of the host star and transiting planet in the OGLE-TR-211 system.


The transit shape is not precise enough to determine the inclination angle very precisely, mainly because of the effect of red noise with only partial transits measured with the VLT. Nevertheless, the size of the planet is relatively well-constrained thanks to the accuracy of the stellar spectroscopic parameters and the weak dependence of the planetary radius on the impact parameter for central transits.

The O–C weighted rms scatter around the final photometric model of the OGLE-TR-211 system is equal to  $\sigma = 1.6 \text{ mmag}$  and  $\sigma = 4.1 \text{ mmag}$  for the VLT and OGLE dataset, respectively. The weighted rms of radial velocity observations relative to the best fit is  $\sigma_{\text{RV}} = 48 \text{ m/s}$ .

## 5. Discussion

Parameters of the transiting companion of the OGLE-TR-211 host star indicate that it is a planetary object belonging to the hot Jupiter class of extrasolar planets. It brings the total number of extrasolar planets discovered by the OGLE survey to seven – significantly lowering the discrepancy between the number of typical hot Jupiters and shortest period, very hot Jupiters in the OGLE sample of extrasolar transiting planets (Pont et al. 2005a; Gould et al. 2006).

OGLE-TR-211b orbits an F7-8 main sequence star. The system parameters closely resemble those of the recently discovered system HAT-P-6 (Noyes et al. 2007) with the only difference that the host star in OGLE-TR-211 is more metal abundant by  $\approx 0.2 \text{ dex}$ . The position of the planet on the mass-orbital period diagram is similar to the position of other transiting exoplanets



**Fig. 4.** Mass-radius diagram for transiting hot Jupiters ( $0.5\text{--}1.5 M_J$ ). Closed symbols indicate host stars with  $T_{\text{eq}} > 1500$  K. OGLE-TR-211 is indicated by the circled dot. The lines show the models of Guillot (2005) in two extreme cases: cold gas planet with a  $15 M_{\oplus}$  core and  $T_{\text{eq}} = 2000$  K planet without a core. The six large planets with radius error bars above the upper models are all strongly irradiated.

with similar mass and period, supporting the relation noted first by Mazeh et al. (2005). Also, the surface gravity of OGLE-TR-211b of about  $10.3 \text{ m/s}^2$  is consistent with the correlation between period and surface gravity noted by Southworth et al. (2007).

Recently Hansen & Barman (2007) have proposed the division of hot Jupiters into two classes based on their location on the (zero albedo) equilibrium temperature – Safronov number,  $\Theta$ , diagram. The position of OGLE-TR-211b in this diagram ( $T_{\text{eq}} = 1720$  K,  $\Theta = 0.059$ ) implies that this planet belongs to the class I of hot Jupiters according to their classification.

While the mass of a new hot Jupiter OGLE-TR-211b is very similar to the mass of another hot Jupiter, OGLE-TR-182b, recently found by Pont et al. (2007), its radius is about 20% larger. The mean density of OGLE-TR-211b is equal to  $0.54^{+0.15}_{-0.24} \text{ g/cm}^3$ , placing it among a group of seven objects with the lowest densities: Trés-4, HD 209458, HAT-P-1, WASP-1, HAT-P-4, and HAT-P-6 (Kovács et al. 2007). All these objects have inflated radii compared to other transiting hot Jupiters of similar mass. Such planets pose a problem for the planetary models and are crucial objects for testing the proposed scenarios explaining “bloated” hot Jupiters (Burrows et al. 2007; Guillot et al. 2006). It is worth noting that the metallicity of the OGLE-TR-211 host star falls in the middle of the range of metallicities of “bloated” Jupiters host stars (recently increased significantly by the HAT-P-6 system) suggesting that metallicity cannot be the only parameter responsible for inflating the radii of hot Jupiters.

On the other hand, the characteristics of the OGLE-TR-211 system reinforce the association of “bloated” close-in giant planets with strong irradiation from the host star. Figure 4 shows the mass-radius diagram for the presently known (January 2008) transiting planets, classified according to their equilibrium temperature<sup>1</sup>.

All anomalously large planets have equilibrium temperatures hotter than 1500 K. This would require an increasingly unlikely coincidence if the anomalously large radii were explained by dynamical effects such as obliquity tide or eccentricity pumping by an unseen companion (Winn & Holman 2005). It instead tends to favor explanations directly linking the excess of radius with the incident flux, such as proposed by Guillot et al. (2006) (that

a fraction of the incident flux energy is converted to mechanical energy by a yet undetermined physical process) – over explanations with a more indirect causal relation, such as put forward by Burrows et al. (2007) and Chabrier & Baraffe (2007) invoking increased opacity or internal density gradients.

Further additional high-accuracy photometric follow-up observations of the OGLE-TR-211 transits are necessary for better constraining the inclination and radius of the planet. Additional high-resolution spectroscopy planned in the 2008 observing season should lower any uncertainty about the planet mass. Although OGLE-TR-211, similar to other OGLE transiting planets, is too faint for the currently available facilities to successfully carry out some follow-up observations like IR photometry, the precise long-term transit timing is feasible allowing search for another planetary objects in the OGLE-TR-211 system.

*Acknowledgements.* We thank the anonymous referee for constructive remarks and comments that improved the paper. The OGLE project is partially supported by the Polish MNiSW grant N20303032/4275. W.G., D.M., G.P., M.T.R., and M.Z. gratefully acknowledge support for this work from the Chilean FONDAF Center of Astrophysics 15010003. N.C.S. acknowledges the support from Fundação para a Ciência e a Tecnologia, Portugal, in the form of a grant (reference POCI/CTE-AST/56453/2004), and support by the EC’s FP6 and by FCT (with POCI2010 and FEDER funds), within the HELAS international collaboration.

## References

- Bouchy, F., Pont, F., Santos, N. C., et al. 2004, *A&A*, 421, L13  
 Bouchy, F., Pont, F., Melo, C., et al. 2005, *A&A*, 431, 1105  
 Burrows, A., Hubeny, I., Budaj, J., & Hubbard, W. B. 2007, *ApJ*, 661, 502  
 Chabrier, G., & Baraffe, I. 2007, *ApJ*, 661, L81  
 Charbonneau, D., Brown, T. M., Latham, D. W., & Mayor, M. 2000, *ApJ*, 529, L45  
 Gillon, M., Pont, F., Moutou, C., et al. 2006, *A&A*, 459, 249  
 Girardi, L., Bertelli, G., Bressan, A., et al. 2002, *A&A*, 391, 195  
 Gould, A., Dorsher, S., Gaudi, B. S., & Udalski, A. 2006, *Acta Astron.*, 56, 1  
 Guillot, T. 2005, *Ann. Rev. of Earth and Planetary Sciences*, 33, 493  
 Guillot, T., Santos, N. C., Pont, F., et al. 2006, *A&A*, 453, L21  
 Hansen, B. M. S., & Barman, T. 2007, *ApJ*, 671, 861  
 Henry, G. W., Marcy, G. W., Butler, R. P., & Vogt, S. S. 2000, *ApJ*, 529, L41  
 Konacki, M., Torres, G., Jha, S., & Sasselov, D. D. 2003, *Nature*, 421, 507  
 Konacki, M., Torres, G., Sasselov, D. D., et al. 2004, *ApJ*, 609, L37  
 Konacki, M., Torres, G., Sasselov, D. D., & Jha, S. 2005, *ApJ*, 624, 372  
 Kovács, G., Zucker, S., & Mazeh, T. 2002, *A&A*, 391, 369  
 Kovács, G., Bakos, G. Á., Torres, G., et al. 2007, *ApJ*, 670, L41  
 Kruszcowski, A., & Semeniuk, I. 2003, *Acta Astron.*, 53, 241  
 Mazeh, T., Zucker, S., & Pont, F. 2005, *MNRAS*, 356, 955  
 Noyes, R. W., Bakos, G. Á., Torres, G., et al. 2007, *ApJ*, 673, L79  
 Pont, F., Bouchy, F., Queloz, D., et al. 2004, *A&A*, 426, L15  
 Pont, F., Bouchy, F., Melo, C., et al. 2005a, *A&A*, 438, 1123  
 Pont, F., Melo, C., Bouchy, F., et al. 2005b, *A&A*, 433, L21  
 Pont, F., Moutou, C., Bouchy, F., et al. 2006a, *A&A*, 447, 1035  
 Pont, F., Zucker, S., & Queloz, D. 2006b, *MNRAS*, 373, 231  
 Pont, F., Tamuz, O., Udalski, A., et al. 2007, *A&A*, in press [arXiv:astro-ph/0710.5278]  
 Santos, N. C., Mayor, M., Naef, D., et al. 2002, *A&A*, 392, 215  
 Santos, N. C., Pont, F., Melo, C., et al. 2006, *A&A*, 450, 825  
 Southworth, J., Wheatley, P. J., & Sams, G. 2007, *MNRAS*, 379, L11  
 Sozzetti, A., Torres, G., Charbonneau, D., et al. 2007, *ApJ*, 664, 1190  
 Torres, G., Konacki, M., Sasselov, D. D., & Jha, S. 2004, *ApJ*, 614, 979  
 Udalski, A. 2003, *Acta Astron.*, 53, 291  
 Udalski, A., Paczynski, B., Żebruń, K., et al. 2002a, *Acta Astron.*, 52, 1  
 Udalski, A., Żebruń, K., Szymański, M., et al. 2002b, *Acta Astron.*, 52, 115  
 Udalski, A., Szweczyk, O., Żebruń, K., et al. 2002c, *Acta Astron.*, 52, 317  
 Udalski, A., Pietrzyński, G., Szymański, M., et al. 2003, *Acta Astron.*, 53, 133  
 Udalski, A., Szymański, M. K., Kubiak, M., et al. 2004, *Acta Astron.*, 54, 313  
 Winn, J. N., & Holman, M. J. 2005, *ApJ*, 628, L159

<sup>1</sup>  $T_{\text{eq}} = T_*(1 - A)/4F)^{1/4}(R_*/a)^{1/2}$  using for all planets a Bond albedo  $A = 0.3$  and a heat redistribution factor  $F = 1/2$ .