

## ON A CLASS OF EXACT LOCALLY CONFORMAL COSYMPLECTIC MANIFOLDS

I. MIHAI and L. VERSTRAELEN

Department Wiskunde, Katholieke Universiteit Leuven  
Celestijnenlaan 200 B, B - 3000 Leuven, Belgium

R. ROSCA

59 Avenue Emile Zola  
75015 Paris, France

(Received May 6, 1993 and in revised form June 1994)

**ABSTRACT.** An almost cosymplectic manifold  $M$  is a  $(2m + 1)$ -dimensional oriented Riemannian manifold endowed with a 2-form  $\Omega$  of rank  $2m$ , a 1-form  $\eta$  such that  $\Omega^m \wedge \eta \neq 0$  and a vector field  $\xi$  satisfying  $i_\xi \Omega = 0$  and  $\eta(\xi) = 1$ . Particular cases were considered in [3] and [6].

Let  $(M, g)$  be an odd dimensional oriented Riemannian manifold carrying a globally defined vector field  $T$  such that the Riemannian connection is parallel with respect to  $T$ . It is shown that in this case  $M$  is a hyperbolic space form endowed with an exact locally conformal cosymplectic structure. Moreover  $T$  defines an infinitesimal homothety of the connection forms and a relative infinitesimal conformal transformation of the curvature forms.

The existence of a structure conformal vector field  $C$  on  $M$  is proved and their properties are investigated. In the last section, we study the geometry of the tangent bundle of an exact locally conformal cosymplectic manifold.

**KEY WORDS AND PHRASES:** Locally conformal cosymplectic manifold,  $T$ -parallel connection, infinitesimal homothety, infinitesimal conformal transformation, Hamiltonian vector field, tangent bundle, Liouville vector field, complete lift, mechanical system.

**1991 AMS SUBJECT CLASSIFICATION CODES:** 53C15, 53C05, 53C20, 58A15, 58A30.

### 1. INTRODUCTION

In the last decade a series of papers have been devoted to almost cosymplectic manifolds  $M(\Omega, \eta, \xi, g)$ . As is well known, an almost cosymplectic manifold  $M$  is an odd dimensional (say  $2m + 1$ ) oriented manifold, where the triple  $(\Omega, \eta, \xi)$  of tensor fields is

- i) a 2-form  $\Omega$  of rank  $2m$
- ii) a 1-form  $\eta$  such that  $\Omega^m \wedge \eta \neq 0$
- iii) a vector field (called the Reeb vector field) such that  $i_\xi \Omega = 0$  and  $\eta(\xi) = 1$ .

One has the following more studied cases:

1°  $\Omega$  and  $\eta$  are both closed forms. Then  $M$  is called a cosymplectic manifold.

2°  $d\eta = 0$ ,  $d\Omega = 2\eta \wedge \Omega$ . Then  $M$  is called a Kenmotsu manifold.

3°  $d\eta = \omega \wedge \eta$ ,  $d\Omega = 2\omega \wedge \Omega$ . Then  $M$  is called a locally conformal cosymplectic manifold (see [3],[16]). In this case  $\omega$  and its dual vector  $T = b^{-1}(\omega)$  with respect to  $g$  is called the Lee form (or characteristic form) and Lee vector field respectively.

In the present paper we consider an almost cosymplectic manifold  $M(\Omega, \eta, \xi, g)$  carrying a globally defined vector field  $T$  whose dual form  $b(T)$  is denoted by  $\omega$ .

Next denote by  $0 = \text{vect}\{e_A; A = 0, 1, \dots, 2m\}$  an orthonormal vector basis on  $M$  and by  $\left\{ \begin{smallmatrix} A \\ \theta_B \end{smallmatrix} \right\}$  the associated connection forms. If the connection forms satisfy

$$\theta_B^A = \langle T, e_B \wedge e_A \rangle; \quad \wedge \text{ is the wedge product,}$$

then one has

$$\nabla_{T'} e_A = 0$$

Therefore we agree to say that  $M$  is structured by a  $T$ -parallel connection. In this condition the following significative fact emerges: the almost cosymplectic structure  $1 \times Sp(2m, \mathbf{R})$  of  $M$  moves to an exact locally conformal cosymplectic structure  $1 \times Sp(2m, \mathbf{R})$  (abbreviated exact L.C.C.), having  $T$  (resp.  $\omega = -df/f$ ) as Lee vector field (resp. Lee form).

Moreover any such a manifold  $M$  is a space form of curvature  $-2c$  and  $f$  is the energy function corresponding to a Hamiltonian vector field associated with  $T$  (in the sense of [3]). If  $\theta$  (resp.  $\Theta$ ) represents the indexless (or generic) connection forms (resp. curvature forms) of  $M$ , then  $T$  defines an infinitesimal homothety of  $\theta$ , i.e.  $L_T \theta = 2c\theta$ , and a relative infinitesimal  $T$  conformal transformation of  $\Theta$  and  $\Omega$ , i.e.

$$d(L_T \Theta) = 2c\omega \wedge \Theta, \quad d(L_T \Omega) = 2c\omega \wedge \Omega.$$

In Section 3 the existence of a structure conformal vector field  $C$  on  $M$  is proved, i.e.

$$\nabla_Z C = \lambda Z + g(Z, T)C - g(Z, C)T; \quad \lambda \in C^\infty M, \quad Z \in \Gamma(TM).$$

Moreover  $C$  is a divergence conformal vector field, i.e.  $\text{grad}(\text{div } C)$  is a concurrent vector field and it defines an infinitesimal conformal transformation of:

- i) the conformal cosymplectic form  $\Omega$ , i.e.  $L_C \Omega = \rho\Omega, \rho = 2\lambda$ ;
- ii) the dual forms  $\omega^A$ , i.e.  $L_C \omega^A = \frac{\rho}{2}\omega^A$ ;
- iii) the curvature forms  $\Theta_b^A$ , i.e.  $L_C \Theta_b^A = \rho\Theta_b^A$ ;
- iv) all the  $(2q + 1)$ -forms  $\alpha_q = b(C) \wedge \Omega^q$ , i.e.  $L_C \alpha_q = (1 + q)\rho\alpha_q$ ;
- v) all the functions  $g(C, Z)$ , i.e.  $L_C g(C, Z) = \rho g(C, Z), Z \in \Gamma(TM)$ .

In the last section, we discuss some properties of the tangent bundle manifold  $TM$  having as basis the exact (L.C.C.)-manifold  $M$ . Denote by  $V, \gamma$  and  $\nu$  the Liouville vector field ([13]), the Liouville 1-form and the Liouville function respectively, on  $TM$ .

The following properties are proved:

- i) the complete lift  $\Omega^c$  of  $\Omega$  is a  $d^w$ -exact 2-form ( $d^w$  is the cohomological operator [11]) and is homogeneous of class 1, i.e.

$$L_\nu \Omega^c = \Omega^c;$$

- ii)  $\gamma$  satisfies  $d^{-w}\gamma = \psi$  and  $\psi$  is a Finslerian form, i.e.

$$L_\nu \psi = \psi, \quad i_\nu \psi = 0$$

( $i_\nu$  denotes the vertical differentiation operator [11]);

- iii) the vertical lift  $T^\nu$  of  $T$  defines an infinitesimal automorphism of  $\psi$ , i.e.  $LT^\nu \psi = 0$ ;
- iv) the function  $r = f\nu$  and the 2-form  $f\psi$  define a regular mechanical system  $\mathcal{M}$  ([13]) having  $r$  as kinetic energy and  $f\psi$  as canonical symplectic (exact) form.

### 1. PRELIMINARIES

Let  $(M, g)$  be a Riemannian  $C^\infty$ -manifold and let  $\nabla$  be the covariant differential operator with respect to the metric tensor  $g$ . Assume that  $M$  is oriented and  $\nabla$  is a Levi-Civita connection. Let  $\Gamma(TM) = \chi(M)$  and  $b: TM \rightarrow T^*M$  be the set of sections of the tangent bundle  $TM$  and the musical isomorphism ([18]) defined by  $g$ , respectively. Following [18] we set

$$A^q(M, TM) = \Gamma \text{Hom}(\Lambda^q TM, TM)$$

and notice that elements of  $A^q(M, TM)$  are vector valued  $q$ -forms ( $q \leq \dim M$ ).

Denote by  $d^\nabla: A^q(M, TM) \rightarrow A^{q+1}(M, TM)$  the exterior covariant derivative operator with respect to  $\nabla$ . It should be noticed that generally  $d^\nabla = d^\nabla \circ d^\nabla \neq 0$  unlike  $d^2 = d \circ d = 0$ . If  $p \in M$ , then the vector valued 1-form  $dp \in A^1(M, TM)$  is the canonical vector valued 1-form of  $M$  ([5]) and since  $\nabla$  is symmetric one has  $d^\nabla(dp) = 0$ . The operator

$$d^\omega = d + e(\omega) \tag{1.1}$$

acting on  $\Lambda M$ , where  $e(\omega)$  means the exterior product by the closed 1-form  $\omega$ , is called the cohomological operator ([11]). One has

$$d^\omega \circ d^\omega = 0. \tag{1.2}$$

Any form  $u \in \Lambda M$  such that  $d^\omega u = 0$  is said to be  $d^\omega$ -closed and if  $\omega$  is an exact form, then  $u$  is said to be a  $d^\omega$ -exact form. Any vector field  $Z \in \Gamma(TM)$  such that

$$d^\nabla(\nabla Z) = \nabla^2 Z = \pi \wedge dp \in A^2(M, TM) \tag{1.3}$$

for some 1-form  $\pi$ , is said to be an exterior concurrent vector field ([17]). The form  $\pi$  which is called the concurrence form is given by

$$\pi = \lambda b(Z); \quad \lambda \in C^\infty M. \tag{1.4}$$

A non flat manifold of dimension  $m > 2$  is an elliptic or hyperbolic space-form if and only if every vector field on  $M$  is an exterior concurrent one ([17]). On the tangent bundle manifold  $TM$ ,  $d_\nu$  and  $i_\nu$  define the vertical differentiation and the vertical derivation operators respectively ([7]).  $d_\nu$  is an anti-derivation of degree 1 on  $\Lambda(TM)$  and  $i_\nu$  is a derivation of degree 0 on  $\nabla(TM)$ .

In an  $n$ -dimensional Riemannian manifold  $M$ , denote by

$$O = \text{vect}\{e_A; A = 1, \dots, n\}$$

a local field of orthonormal frames and let

$$O^* = \text{covect}\{\omega^A; A = 1, \dots, n\}$$

be its associated coframe.

The soldering form  $dp$  is expressed by

$$dp = \omega^A \otimes e_A \tag{1.5}$$

and E. Cartan's structure equations written indexless manner are

$$\nabla e = \theta \otimes e \tag{1.6}$$

$$d\omega = -\theta \wedge \omega \tag{1.7}$$

$$d\theta = -\theta \wedge \theta + \Theta \tag{1.8}$$

Any vector field  $T$  such that

$$\nabla T = s dp + u \otimes T, \quad u \in \Lambda^1 M \tag{1.9}$$

is called a torse forming (K. Yano [20]). If  $du = 0$ , then  $T$  is a closed torse forming, which implies that  $T$  is an exterior concurrent vector field, and if  $u = 0$ , then  $T$  is a concurrent vector field ([22]).

Let now  $W$  be any conformal vector field on  $M$  (i.e. the conformal version of Killing's equations). As is well known,  $W$  satisfies

$$L_w g = \rho g \quad \text{or} \quad g(\nabla_z W, Z') + g(\nabla_{z'} W, Z) = \rho g(Z, Z') \tag{1.10}$$

where the conformal scalar  $\rho$  is defined by

$$\rho = \frac{2}{n}(\text{div} W). \tag{1.11}$$

We recall some basic formulas which we shall use in the following sections.

$$L_w b(Z) = \rho b(Z) + b[W, Z] \quad (\text{Orsted lemma}) \quad (1.12)$$

$$L_w K = (n - 1)\Delta\rho - K\rho \quad (1.13)$$

$$2L_w S(Z, Z') = (\Delta)\rho g(Z, Z') - (n - 2) (\text{Hess}\nabla^p)(Z, Z'). \quad (1.14)$$

In the above equations  $L_w, K, \Delta$  and  $S$  denote the *Lie* derivative with respect to  $W$ , the scalar curvature of  $M$ , the Laplacian and the Ricci tensor field of  $\nabla$ , respectively. One has

$$(\text{Hess}_{\nabla}\rho)(Z, Z') = g(Z, H_\rho Z'), \quad H_\rho Z' = \nabla_{Z'}(\text{grad } \rho)$$

(see also [2]).

**2. EXACT LOCALLY CONFORMAL COSYMPLECTIC MANIFOLDS**

Let  $(M, g)$  be a  $(2m + 1)$ -dimensional oriented Riemannian  $C^\infty$ -manifold and let  $T = \sum_{A=0}^{2m} t^A e_A$  and  $\omega = b(T)$  be a globally defined vector field on  $M$  and its dual form respectively.

Denote by  $\mathbf{O} = \text{vect}\{e_A; A = 0, 1, \dots, 2m\}$  (resp.  $\theta_B^A$ ) a local field of orthonormal frames on  $M$  (resp. the associated connection forms). Recall that the vectorial wedge product  $\wedge$  is defined by

$$(X \wedge Y)Z = g(Y, Z)X - g(X, Z)Y; \quad Z \in \Gamma(TM)$$

i.e.

$$X \wedge Y = b(Y) \otimes X - b(X) \otimes Y.$$

Assume now that all the connection forms  $\theta$  satisfy

$$\theta_B^A = \langle T, e_B \wedge e_A \rangle. \quad (2.1)$$

Then by the structure equations (1.6), it follows at once

$$\theta_B^A = t^B \omega^A - t^A \omega^B. \quad (2.2)$$

It should be noticed that if  $\theta$  satisfy (2.2) one has  $\theta(T) = 0$  and the above equation shows that all the connection forms  $\theta$  are relations of integral invariance for the vector field  $T$  (in the sense of A. Lichnerowicz [14]).

Next by the structure equations (1.6) and by (2.2) one obtains

$$\nabla e_A = t^A dp - \omega^A \otimes T \quad (2.3)$$

and the above equation implies

$$\nabla_T e_A = 0. \quad (2.4)$$

From (2.4) the following significant fact emerges: all the vectors of the  $\mathbf{O}$ -basis are  $T$ -parallel. Therefore we agree to say that the Riemannian manifold under consideration is structured by a  $T$ -parallel connection (abr. T.P.).

Further again by (2.2) one derives by the structure equations (1.7)

$$d\omega^A = \omega \wedge \omega^A; \quad \omega = b(T) = t^A \omega^A \quad (2.5)$$

which by a simple argument implies that the dual form  $\omega$  of  $T$  is closed, i.e.

$$d\omega = 0. \quad (2.6)$$

Thus in terms of  $d^\omega$ -cohomology, (2.5) may be written as

$$d^{-\omega}\omega^A = 0 \quad (2.7)$$

and  $\mathbf{O}^* = \{\omega^A\}$  is defined as a  $d^{-\omega}$ -closed covector basis.

Now for reasons which will soon appear, we set

$$\omega^0 = \eta, \quad e_0 = \xi \quad (2.8)$$

and consider on  $M$  the globally defined 2-form  $\Omega$  of rank  $2m$  given by

$$\Omega = \sum \omega^a \wedge \omega^{a^*}; \quad a = 1, \dots, m; \quad a^* = a + m. \tag{2.9}$$

Then since  $\Omega^m \wedge \eta \neq 0, i_\xi \Omega = 0$ , one may say that the triple  $(\Omega, \eta, \xi)$  defines an almost cosymplectic structure  $1 \times Sp(2m, \mathbf{R})$  having  $\xi$  as Reeb's vector field.

Next taking the exterior differential of  $\Omega$  a short calculation gives with the help of (2.5)

$$d\Omega = 2\omega \wedge \Omega \Leftrightarrow d^{-2\omega}\Omega = 0 \tag{2.10}$$

and by (2.5) we may write

$$d\eta = \omega \wedge \eta \Leftrightarrow d^{-\omega}\eta = 0. \tag{2.11}$$

We conclude that any odd dimensional Riemannian manifold  $M$  structured by a  $T$ -parallel connection is endowed with a locally conformal cosymplectic structure  $1 \times CSP(2n, \mathbf{R})$  (abr. L.C.C.). We notice that the vector field  $T$  (resp. the 1-form  $\omega = b(T)$ ) is the Lee vector field (resp. the Lee form) of this structure.

Moreover since  $\omega = t^A \omega^A$ , then by a simple argument it follows on behalf of (2.5) that one may set

$$dt^A = f\omega^A; \quad f \in C^\infty M \tag{2.12}$$

which by exterior differentiation gives instantly

$$\omega = -df/f. \tag{2.13}$$

Therefore since  $\omega$  is an exact form, it follows on behalf of a known terminology, that the manifold  $M$  under consideration is an exact (L.C.C.)-manifold. We agree to call  $f$  the distinguished scalar field associated with the exact (L.C.C.)-structure.

Now taking the covariant differential of  $T$  one finds by (2.3) and (2.12)

$$\nabla T = (f + 2l)dp - \omega \otimes T \tag{2.14}$$

where we have set

$$g(T, T) = 2l. \tag{2.15}$$

Using (2.12) and (2.15), we have

$$dl = f\omega \Rightarrow l + f = c = \text{const} \neq 0 \tag{2.16}$$

and (2.14) becomes

$$\nabla T = (l + c)dp - \omega \otimes T. \tag{2.17}$$

Hence, by (1.9) and (2.6)  $T$  is a closed torse forming and consequently an exterior concurrent (abr. E.C.)-vector field.

Operating now on  $\nabla e_A$  and  $\nabla T$  by the exterior covariant derivative operator  $d^\nabla$ , one gets by (2.12) and (2.16)

$$d^\nabla(\nabla e_A) = \nabla^2 e_A = 2c\omega^A \wedge dp \tag{2.18}$$

$$d^\nabla(\nabla T) = \nabla^2 T = 2c\omega \wedge dp. \tag{2.19}$$

From the above equations it is seen that any vector field  $Z$  on  $M$  is E.C. with constant conformal scalar  $2c$ . Therefore on behalf of the general properties of E.C.-vector fields ([17]), we may state the following striking property: the exact L.C.C.-manifold  $M(\Omega, \eta, \xi)$  under discussion is a space-form of curvature  $-2c$ .

As a consequence, it follows that the curvature forms  $\Theta$  are expressed by

$$\Theta_B^A = -2c\omega^A \wedge \omega^B \tag{2.20}$$

Next taking the exterior differential of the forms  $\Theta$ , one quickly finds by

$$d\Theta_B^A = 2\omega \wedge \Theta_B^A \Leftrightarrow d^{-2\omega}\Theta_B^A = 0 \tag{2.21}$$

which shows that all the curvature forms  $\Theta$  are  $d^{-2\omega}$ -exact.

On the other hand taking the Lie derivatives of the covectors  $\omega^A$  of  $\mathbf{O}^*$  one derives by (2.12) and (2.16)

$$L_I \omega^A = (l + c)\omega^A - t^A \omega. \tag{2.22}$$

Therefore since  $L_I$  satisfies Leibniz rule one deduces by (2.20)

$$L_I \Theta_B^A = 2(l + c)\Theta_B^A + 2c\theta_B^A \wedge \omega \tag{2.23}$$

Similarly, we obtain

$$d\Theta_B^A = 2f\omega^B \wedge \omega^A + \omega \wedge \theta_B^A \tag{2.24}$$

Clearly by (2.12) one has  $L_I t^A = f t^A$  and with the help of (2.22) we deduce

$$L_I \theta_B^A = 2c\theta_B^A. \tag{2.25}$$

Accordingly by the above equations we may say that the Lie vector field  $T$  defines on infinitesimal homothety of all the connection forms  $\theta$ .

Taking now the exterior differential of the equations (2.23), a standard calculation gives

$$d(L_I \Theta_B^A) = 8c\omega \wedge \Theta_B^A \tag{2.26}$$

which proves that  $T$  defines a relative infinitesimal conformal transformation ([19]) of the curvature forms.

let  $\mu : TM \rightarrow T^*M$ ,  $\mu(Z) = i_Z \Omega$  be the bundle isomorphism defined by  $\Omega$  and set  $\bar{\omega} = \mu(T)$ , i.e.

$$\bar{\omega} = i_T \Omega = \sum_{a=1}^m (t^a \omega^{a*} - t^{a*} \omega^a) \tag{2.27}$$

for the dual form of  $T$  with respect to  $\Omega$ . By (2.5) and (2.12) an easy calculation gives

$$d\bar{\omega} = 2f\Omega + \omega \wedge \bar{\omega} \tag{2.28}$$

and by (2.10) and (2.13) one gets

$$L_T \Omega = 2(l + c)\Omega + \bar{\omega} \wedge \omega \tag{2.29}$$

and consequently by (2.28) it follows

$$d(L_T \Omega) = 2c\omega \wedge \Omega. \tag{2.30}$$

Hence as for the curvature forms  $\Theta$ ,  $T$  defines a relative conformal transformation of the structure 2-form  $\Omega$ .

Consider now the vector valued 1-form

$$F = \omega^a \otimes e_{a*} - \omega^{a*} \otimes e_a \in A^1(M, TM). \tag{2.31}$$

If  $Z$  is any vector field, a simple calculation gives

$$\langle F, Z \rangle = Z^a e_{a*} - Z^{a*} e_a = \bar{Z} \tag{2.32}$$

which implies

$$g(Z, Z') + g(Z, \bar{Z}') = 0, \quad Z, Z' \in \Gamma(TM) \tag{2.33}$$

and  $\langle F, d\rho \rangle = 2\Omega$ .

On the other hand since  $\bar{\omega}(T) = 0$  one gets by (2.27)

$$L_T \bar{\omega} = 2c\bar{\omega} \tag{2.34}$$

that is  $T$  defines an infinitesimal homothety of  $\bar{\omega} = (\mu \circ b)T$ .

Next by (2.12) and (2.13) one easily gets

$$i_T \Omega = \frac{df}{f} - \frac{1}{f} \xi(f)\eta . \tag{2.35}$$

Therefore by reference to [3] one may call  $\bar{T}$  the cosymplectic Hamiltonian vector field of  $M$  and the distinguished scalar  $f$  turns out to be the energy function corresponding to  $\bar{T}$ .

Moreover by (2.35) one derives

$$L_T \Omega = \eta(\bar{T})\eta \wedge \omega \Rightarrow d(L_T \Omega) = 0 \tag{2.36}$$

which shows that  $\bar{T}$  defines a relative infinitesimal automorphism (R. Abraham [1]) of  $\Omega$ .

Summing up, we state the following

**THEOREM.** Let  $M$  be a  $(2m + 1)$ -dimensional Riemannian manifold and let  $T$  be a globally defined vector field on  $M$ . If  $M$  is structured by a  $T$ -parallel connection, then  $M$  is endowed with an exact locally conformal cosymplectic structure  $1 \times CSp(2m, \mathbf{R})$ , having  $T$  (resp.  $\omega = b(T)$ ) as Lee vector (resp. Lee form) and any such an  $M$  is a space-form of curvature  $-2c$ .

Moreover one has the following properties:

- i)  $T$  defines an infinitesimal homothety of the connection forms  $\theta$  and of the 1-form  $\mu(T)$ , i.e.

$$L_T \theta = 2c \theta , \quad L_T \mu(T) = 2c \mu(T)$$

- ii)  $T$  defines a relative infinitesimal conformal transformation of the curvature forms  $\Theta$  and of the structure 2-form  $\Omega$ , i.e.

$$d(L_T \Theta) = 8c \omega \wedge \Theta , \quad d(L_T \Omega) = 2c \omega \wedge \Omega$$

- iii) the vector field  $\bar{T} = (b^{-1} \circ \mu) T$  (resp.  $f$ ) is the cosymplectic Hamiltonian associated with the  $1 \times CSp(2m, \mathbf{R})$ -structure of  $M$  (resp. its corresponding energy function) and  $\bar{T}$  defines a relative infinitesimal automorphism of  $\Omega$ .

Let now  $\Phi : M \rightarrow \bar{M}$  be a conformal diffeomorphism (abr. C.D.) that is

$$\Phi : g \rightarrow e^{2\sigma} g = \bar{g} ; \quad \sigma \in C^\infty M .$$

One also say that  $g$  and  $\bar{g}$  are conformally equivalent metrics and setting  $e^{2\sigma} = v^2$ , we agree to call the function  $v$  the argument of the C.D.

As is shown one has for  $Z, Z' \in \Gamma(TM)$

$$\hat{\nabla} Z = \nabla Z + b(\text{grad } \sigma) \otimes Z - b(Z) \otimes \text{grad } \sigma + g(Z, \text{grad } \sigma) dp \tag{2.37}$$

or equivalently

$$\hat{\nabla}_Z Z' = \nabla_Z Z' + Z'(\sigma)Z + Z(\sigma)Z' - g(Z, Z') \text{grad } \sigma \tag{2.38}$$

and if  $K$  and  $\bar{K}$  denote the scalar curvature of  $M$  and  $\bar{M}$  respectively then one has ([8])

$$\bar{K} = e^{-2\sigma} \{ K + 2(n - 1)(n - 2) \| \text{grad } \sigma \|^2 \} \tag{2.39}$$

( $n = \dim M$ ).

If  $M$  is an exact (L.C.C.)-manifold, its Ricci tensor field  $S$  satisfies

$$S(Z, Z') = -4mc g(Z, Z') ; \quad Z, Z' \in \Gamma(TM) \tag{2.40}$$

and the scalar curvature  $K$  is given by

$$K = -4m(2m + 1)c . \tag{2.41}$$

Perform now a conformal transformation of  $M$  having as argument  $e^\sigma$  the energy function  $f$ . It is obvious that

$$(2.42) d \sigma = df/f = -\omega . \tag{2.42}$$

Then we have  $\text{grad } \sigma = -T$ , which implies

$$\Delta\sigma = \operatorname{div} T = (2m + 1)c + (2m - 1)l. \tag{2.43}$$

Hence by (2.41) and (2.43) we derive at once from (2.39),  $\vec{K} = 0$ , that is  $\vec{M}$  is a flat manifold. We notice that this fact is in accordance with the known

**PROPOSITION.** A Riemannian manifold of constant curvature is conformally flat, provided  $n \geq 3$ .

Using (2.37) one may prove that all vectors  $\bar{e}_A$  are parallel (the connection forms  $\bar{\theta}_B^A$  vanish, i.e.  $\bar{\nabla}$  is a flat connection). Thus we have

**PROPOSITION.** If  $M$  is an exact (L.C.C.)-manifold with metric tensor  $g$  and energy function  $f$ , then the metric  $f^2g$  is flat.

### 3. STRUCTURE CONFORMAL VECTOR FIELDS ON AN EXACT (L.C.C.)-MANIFOLD

In consequence of some conformal properties induced by the  $T$ -parallel connection which structures  $M(\Omega, \eta, \xi, g)$  we are naturally led to see if the manifold  $M$  under consideration carries a structure conformal vector field  $C$  in the sense of [6], [15]. Therefore the covariant differential of  $C$  is expressed by

$$\nabla C = \lambda dp + C \wedge T = \lambda dp + \omega \otimes C - \alpha \otimes T; \quad \lambda \in C^\infty M, \quad \alpha = b(C). \tag{3.1}$$

Put

$$C = C^A e_A \Rightarrow b(C) = \alpha = C^A \omega^A \tag{3.2}$$

and  $s = g(C, T)$ . Then by (2.3) and (3.1) one quickly gets

$$dC^A = (\lambda - s)\omega^A + C^A \omega \tag{3.3}$$

$$d\alpha = 2\omega \wedge \alpha \Rightarrow d^{-2\omega}\alpha = 0. \tag{3.4}$$

Next since  $ds = \langle \nabla C, T \rangle + \langle \nabla T, C \rangle$ , a short calculation gives

$$ds = \lambda \omega - (l - c)\alpha \tag{3.5}$$

$$ds = d\lambda. \tag{3.6}$$

By (3.4), (3.5) and (3.6) it is seen that the existence of  $C$  is assured by an exterior differential system  $\Sigma$  whose characteristic numbers are

$$r = 3, \quad s_0 = 2, \quad s_1 = 1.$$

Then  $\Sigma$  is in involution in the sense of E. Cartan (i.e.  $r = s_c + s_l$ ). Accordingly one may say that the existence of  $C$  depends on 2 arbitrary functions of one argument (E. Cartan's test). The conformal scalar  $\rho$  associated with  $C(L_c g = \rho g)$  is given by

$$\rho = 2\lambda. \tag{3.7}$$

By a short calculation one has

$$[C, T] = -\lambda T - (l - c)C; \quad [ \ ]: \text{ Lie bracket} \tag{3.8}$$

and from (3.5) it follows

$$L_c \omega = ds = \lambda \omega - (l - c)\alpha. \tag{3.9}$$

This equation matches by Orsted's lemma (1.12) the expression of  $[C, T]$ .

On the other hand since  $C$  is necessarily an E. C. vector field ( $M$  is a space-form), then operating (3.1) by  $d^\nabla$  and taking account of (3.4) and (3.5), one derives

$$d^\nabla(\nabla C) = \nabla^2 C = 2c\alpha \wedge dp. \tag{3.10}$$

The above equation is coherent with the properties obtained in Section 2.

Setting now

$$\bar{\alpha} = i_c \Omega = \Sigma(C^a \omega^{a*} - C^{a*} \omega^a) \tag{3.11}$$



one gets by (3.4) and (2.5)

$$d\bar{\alpha} = 2(\lambda - s)\Omega + 2\omega \wedge \bar{\alpha} \tag{3.12}$$

and one follows

$$L_C \Omega = \rho \Omega . \tag{3.13}$$

Hence (3.13) reveals that  $C$  defines an infinitesimal conformal transformation (abr. I.C.T.) of the conformal cosymplectic form  $\Omega$ .

By similar methods, one gets by (2.5), (2.24), (2.20) and (2.21)

$$L_C \omega^A = \frac{\rho}{2} \omega^A, \quad L_C \theta_B^A = \frac{\rho}{2} \theta_B^A, \quad L_C \Theta_B^A = \rho \Theta_B^A . \tag{3.14}$$

Therefore one may say that  $C$  defines an I.C.T. of the exact (L.C.C.)-structure of  $M$ .

Moreover let  $L$  be the operator of type (1.1) on forms defined by  $S$ . Goldberg ([8]), that is  $Lu = u \wedge \Omega; u \in \Lambda^1 M$ , and consider on  $M$  the  $(2q + 1)$ -forms

$$L^q \alpha = \alpha_q = \alpha \wedge \Omega^q . \tag{3.15}$$

Since by Orsted's lemma one has

$$L_C \alpha = \rho \alpha \tag{3.16}$$

then by (3.13) and a standard calculation one derives

$$L_C \alpha_q = (q + 1)\rho \alpha_q . \tag{3.17}$$

Hence  $C$  defines an (I.C.T.) of all the  $(2q + 1)$ -forms  $\alpha_q$ .

Next since  $C$  is a conformal vector field, then as is known (see (1.11)) one has

$$\operatorname{div} C = (\rho/2)(2m + 1) \tag{3.18}$$

and since  $\rho = 2\lambda$  it follows by (3.5) and (3.6) that

$$\operatorname{grad} \rho = \rho T + 2(c - l)C . \tag{3.19}$$

Further by (2.16) and taking account of (2.14) and (3.1) it is easily deduced

$$\nabla \operatorname{grad} \rho = 2c \rho dp . \tag{3.20}$$

Thus one may state the following relevant property: the gradient of the associated scalar  $\rho$  of  $C$  is a concurrent vector field (K. Yano and B. Y. Chen [22]). We agree to call a conformal vector field such that the gradient of its conformal scalar  $\rho$  is a concurrent vector field, a divergence conformal vector field. Such a situation occurs also when studying conformal vector fields on Lorentzian P.S. manifolds (see I. Mihai and R. Rosca [15]).

On the other hand from (2.14) one derives

$$\operatorname{div} T = (2m - 1)l + (2m + 1)c \tag{3.21}$$

and since  $\operatorname{div} C = (2m + 1)\lambda$ , one gets on behalf of (3.20)

$$\Delta \rho = -\operatorname{div}(\operatorname{grad} \rho) = -2(2m + 1)c \rho \tag{3.22}$$

which shows that  $\rho$  is an eigenfunction of  $\Delta$ .

$C$  being an E.C. vector field satisfying (3.10), one has ([17])

$$S(C, Z) = -4mc g(C, Z), \quad Z \in \Gamma(TM) \tag{3.23}$$

where  $S$  denotes the Ricci tensor field of  $\nabla$ .

Now making use of (1.14) and carrying out the calculations, one finds by (3.19) and (3.22)

$$L_C g(C, Z) = \rho g(C, Z) . \tag{3.24}$$

Hence the vector field  $C$  defines an I.C.T. of all the functions  $g(C, Z)$ , where  $Z \in \Gamma(TM)$ .

Concluding, we have proved the following

**THEOREM.** Let  $M$  be the exact (L.C.C.) manifold defined in Section 2 and  $C$  a structure conformal vector field on  $M$  (which existence is proved), i.e.

$$\nabla C = \frac{\rho}{2} dp + C \wedge T; \quad L_C g = \rho g$$

Then  $C$  is a divergence conformal vector field (i.e.  $\text{grad}(\text{div } C)$  is a concurrent vector field) and it defines the following infinitesimal conformal transformations

$$L_C \Omega = \rho \Omega, \quad L_C \omega^i = \frac{\rho}{2} \omega^i, \quad L_C \theta_B^i = \frac{\rho}{2} \theta_B^i$$

$$L_C \Theta_B^i = \rho \Theta_B^i, \quad L_C \alpha_q = (1 + q)\rho \alpha_q, \quad L_C g(C, Z) = \rho g(C, Z) (Z \in \Gamma(TM))$$

where  $\Omega, \omega^i, \theta_B^i, \Theta_B^i$  and  $\alpha_q = b(C) \wedge \Omega^q$  are the conformal symplectic 2-form, the dual forms, the connection forms, the curvature forms and the  $(2q + 1)$ -forms defined by the  $(1,1)$ -operator  $L$ , respectively on  $M$ .

**4. GEOMETRY OF THE TANGENT BUNDLE OF AN EXACT (L.C.C.)-MANIFOLD**

Let now  $TM$  be the tangent bundle manifold having the exact (L.C.C.)-manifold  $M$  discussed in Section 2 as a basis.

Denote by  $V(v^A) (A = 0, 1, \dots, 2m)$  the Liouville vector field (or the canonical vector field [7]). Accordingly we may consider the set  $B^* = \{\omega^A, dv^A\}$  as an adapted cobasis in  $TM$ . Following Godbillon ([7]) we denote by  $d_v$  and  $i_v$  the vertical differentiation and the vertical derivative operators with respect to  $B^*$ , respectively ( $d_v$  is an antiderivation of degree 1 on  $\Lambda(TM)$  and  $i_v$  is a derivation of degree 0 on  $\Lambda(TM)$ ). Let  $T_s^r M$  be the set of all tensor fields of type  $(r, s)$  on  $M$ .

In general as is known ([23]) the vertical and complete lifts are linear mappings of  $T_s^r M$  into  $T_s^r(TM)$  and one has

$$(T_1 \otimes T_2)^c = T_1^v \otimes T_2^c + T_1^c \otimes T_2^v. \tag{4.1}$$

In the case under discussion we may define the complete lift  $\Omega^c$  of the structure 2-form  $\Omega$  of  $M$  by the 2-form of rank  $4m$  on  $TM$

$$\Omega^c = \Sigma(dv^a \wedge \omega^{a^*} + \omega^a \wedge dv^{a^*}), \quad a = 1, \dots, m; \quad a^* = a + m. \tag{4.2}$$

On the other hand since the Liouville vector field  $V$  is expressed by

$$V = \Sigma v^A \frac{\partial}{\partial v^A} \tag{4.3}$$

then as is known the basic 1-form

$$\gamma = \Sigma v^A \omega^A \tag{4.4}$$

is called the Liouville form (see also [13]).

Taking now the exterior differential of  $\Omega^c$  one finds by (2.5)

$$d\Omega^c = \omega \wedge \Omega^c \Leftrightarrow d^{-\omega} \Omega^c = 0 \tag{4.5}$$

which shows that  $\Omega^c$  is similarly as  $\Omega$  a  $d^{-\omega}$ -exact form. We recall that in general conformal properties are not preserved by complete lifts ([23]).

One has

$$i_v \Omega^c = \Sigma(v^a \omega^{a^*} - v^{a^*} \omega^a) \tag{4.6}$$

which implies  $\omega(V) = 0$  and so by (4.5) and (4.6) one gets

$$L_v \Omega^c = \Omega^c. \tag{4.7}$$

Accordingly on behalf of a known definition ([13]), the above equation shows that  $\Omega^c$  is of class 1, a homogeneous form on  $TM$ . Taking now the exterior differential of the Liouville form  $\gamma$  defined by (4.4), one gets at once by (2.5)

$$d\gamma = \omega \wedge \gamma + \psi \Leftrightarrow d^{-\omega} \gamma = \psi \tag{4.8}$$

where we have set

$$\psi = \sum d v^A \wedge \omega^A . \tag{4.9}$$

From (4.8) and (1.2) one obtains instantly

$$d^{m+1} \psi = 0 \Leftrightarrow d \psi = \omega \wedge \psi . \tag{4.10}$$

Since clearly the 2-form  $\psi$  is of maximal rank, we agree to call  $\psi$  the canonical conformal symplectic form of  $M$ . Noticing that one has

$$i_V \psi = \gamma , \quad \omega(V) = 0 \tag{4.11}$$

which implies

$$L_V \psi = \psi . \tag{4.12}$$

Hence  $\psi$  is as  $\Omega^2$  a homogeneous of class 1, 2-form.

Next making use of the vertical operator  $i_v$  defined by  $i_v \lambda = 0, i_v d v^A = \omega^A, i_v \omega^A = 0 (\lambda \in C^\infty M)$  one quickly finds by (4.9)

$$i_v \psi = 0 \tag{4.13}$$

and the above equation together with (4.12) proves that  $\psi$  is a Finslerian form ([7]).

We recall that the vertical lift  $Z^v$  ([23]) of a vector field  $Z \in \Gamma(TM)$  with components  $Z^A$  in  $M$ , has as components

$$Z^v = \begin{pmatrix} 0 \\ Z^A \end{pmatrix} = Z^A \frac{\partial}{\partial v^A} .$$

Hence in the case under consideration one has

$$T^v = \sum t^A \frac{\partial}{\partial v^A} ; \quad A = 0, 1, \dots, 2m \tag{4.14}$$

and by (4.9) one gets

$$i_{T^v} \psi = \omega . \tag{4.15}$$

Therefore by (4.10) one derives

$$L_{T^v} \psi = 0 \tag{4.16}$$

and one may say that  $T^v$  defines an infinitesimal automorphism of  $\psi$ .

Finally we set

$$r = f v \tag{4.17}$$

where

$$v = \frac{1}{2} \sum (v^A)^2 \tag{4.18}$$

denotes the Liouville function on  $M$  ([9]).

Operating on  $r$  by the vertical differentiation operator  $d_v$  ([7]) one gets

$$d_v r = f \sum_A v^A \omega^A = f \mu \tag{4.19}$$

and taking the exterior differential of (4.19) we obtain by (2.13) and (4.9)

$$d(d_v r) = f \sum d v^A \wedge \omega^A = f \psi . \tag{4.20}$$

Next putting  $H = f \psi$  it follows by (2.13)

$$dH = 0 . \tag{2.21}$$

Therefore the exact symplectic form  $H$  can be viewed as the canonical symplectic form of the  $(4m + 2)$ -dimensional manifold  $TM$  ([13]).





Finally by reference to [13] one may consider that the pair  $(r, H)$  defines a regular mechanical system  $\mathcal{M}$  (in the sense of Klein [13]) having the scalar  $r$  as kinetic energy.

**THEOREM.** Let  $TM$  be the tangent bundle manifold having as basis the exact (L.C.C.)-manifold  $M(\Omega, T, \omega)$  discussed in Section 2. Let  $V, \gamma$  and  $v$  be the Liouville vector field, the Liouville form and the Liouville function of  $TM$ , respectively. One has the following properties:

- i) the complete lift  $\Omega^c$  on  $TM$  of the conformal cosymplectic form  $\Omega$  of  $M$  is a homogeneous of class 1, 2-form, i.e.  $L_V \Omega^c = \Omega^c$ , and it is  $d^{\text{con}}$ -exact, i.e.  $d^{\text{con}} \Omega^c = 0$ ;
- ii)  $\gamma$  satisfies  $d^{\text{con}} \gamma = \psi \Rightarrow d^{\text{con}} \psi = 0$  and  $\psi$  is the canonical conformal symplectic form of  $TM$  and  $\psi$  enjoys also the property to be a Finslerian form;
- iii) the vertical lift  $T^c$  of  $T$  defines an infinitesimal automorphism of  $\psi$ , i.e.  $L_{T^c} \psi = 0$ ;
- iv)  $r = fV$  and  $f\psi$  define a regular mechanical system on  $TM$  having  $r$  as kinetic energy and  $f\psi$  as canonical symplectic form (where  $f$  is the energy function of  $M$ ).

### REFERENCES

- [1] ABRAHAM, R. Foundations of Mechanics, W. A. Benjamin Inc., New York (1967).
- [2] BRANSON, T. Conformally covariant equations of differential forms, Comm. Partial Diff. Equations, 7 (1982), 393-431.
- [3] CHINEA, D., DE LEON, M. and MORRERO, J. C. Locally conformal cosymplectic manifolds and time-dependent Hamiltonian systems, Comm. Math. Univ. Carolinae, 32 (1991), 383-387.
- [4] DATTA, D. K. Exterior recurrent forms in a manifold, Tensor N.S., 36 (1982), 115-120.
- [5] DIEUDONNÉ, J. Treaties on Analysis, Vol. 4, Academic Press, New York (1974).
- [6] DONATO, S. and ROSCA, R. Structure conformal vector fields on almost paracontact manifolds with parallel structure vector, Osterreichische Akademie des Wissenschaften, Wien, 198 (1989), 201-209.
- [7] GODBILLON, C. P. Géométrie Différentielle et Mécanique Analytique, Hermann, Paris (1969).
- [8] GOLDBERG, S. Curvature and Homology, Academic Press, New York (1962).
- [9] GOLDBERG, V. V. and ROSCA, R. Pseudo-Sasakian manifolds endowed with a contact conformal connection, Internat. J. Math. and Math. Sci., 9 (1986), 733-747.
- [10] GOLDBERG, V. V. and ROSCA, R. Foliate conformal Kählerian manifolds, Rend. Sem. Mat. Messina, Serie II, Vol. I (1991), 105-122.
- [11] GUEDIRA, F. and LICHNEROWICZ, A. Géométrie des algèbres de Lie locales de Kirilov, J. Math. Pures Appl., 63 (1984), 407-494.
- [12] KERMOTSU, K. A class of almost contact Riemannian manifolds, Tohoku Math. J., 24 (1972), 93-103.
- [13] KLEIN, I. Espaces variationnels et mécanique, Ann. Inst. Fourier, 12 (1962), 1-124.
- [14] LICHNEROWICZ, A. Les relations intégrales d'invariance et leurs applications à la dynamique, Bull. Sci. Math., 70 (1946), 82-95.
- [15] MIHAI, I. and ROSCA, R. On Lorentzian P-Sasakian manifolds, Classical Analysis, World Scientific Publ., Singapore (1992), 155-169.
- [16] OLCSAK, Z. and ROSCA, R. Normal locally conformal almost cosymplectic manifolds, Publicationes Math. (Debrecen), 39 (1991), 315-323.
- [17] PETROVIC, M., ROSCA, R. and VERSTRAELEN, L. On exterior concurrent vector fields I. Some general results, Socehow J. Math., 15 (1989), 179-187.
- [18] POOR, W. A. Differential Geometric Structures, McGraw Hill Book Co., New York (1981).
- [19] ROSCA, R. On some infinitesimal transformations in Riemannian and pseudo-Riemannian manifolds (Preprint).
- [20] YANO, K. On the torse-forming directions in Riemannian spaces, Proc. Imp. Acad., Tokyo, 20 (1944), 340-345.
- [21] YANO, K. Integral Formulas in Riemannian Geometry, M. Dekker, New York (1970).
- [22] YANO, K. and CHEN, B. Y. On the concurrent vector fields of immersed manifolds, Kodai Math. Sem. Rep., 23 (1971), 343-350.
- [23] YANO, K. and ISHIHARA, S. Differential Geometry of Tangent and Cotangent Bundles, M. Dekker, New York (1973).

**Hindawi**

Submit your manuscripts at  
<http://www.hindawi.com>

