

ON A COEFFICIENT PROBLEM FOR BI-UNIVALENT FUNCTIONS

M. LEWIN

1. **Introduction.** Let S be the family of analytic functions:

$$(1) \quad f(z) = z + a_2 z^2 + \dots,$$

which are regular and univalent in $|z| < 1$. Let $f^{-1}(z)$ be the inverse function of $f(z)$. We have:

$$(2) \quad f^{-1}(z) = z - a_2 z^2 + (2a_2^2 - a_3) z^3 + \dots$$

The function $f(z)$ will be called bi-univalent if both $f(z)$ and $f^{-1}(z)$ are univalent in $|z| < 1$; $f(z)$ will be said to belong to σ iff (i) $f(z) \in S$ and (ii) there exists a function $g(z) \in S$ such that $f(g(z)) = g(f(z)) = z$ in some neighborhood of the origin.

Z. Nehari remarked¹ that if $\phi(z) = \phi_1 z + \phi_2 z^2 + \dots$ and $\psi(z) = \psi_1 z + \psi_2 z^2 + \dots$, with $\psi_1 = \phi_1$, are two functions mapping the open unit circle onto a schlicht domain containing the open unit circle, then the function

$$(3) \quad f(z) = \phi[\psi^{-1}(z)]$$

is bi-univalent. It will belong to σ if both $f(z)$ and $f^{-1}(z)$ are regular in $|z| < 1$.

It is not clear whether all functions of σ are representable in the form (3). If they are, the representation may not be unique. The function $f(z)$ in (3) will belong to σ even though $\phi(z)$ or $\psi(z)$ or both have poles in $|z| < 1$, on condition that if $z = p_\phi$ and $z = p_\psi$ are the respective poles, the following two inequalities be satisfied:

$$|\phi(p_\psi)| \geq 1 \quad \text{and} \quad |\psi(p_\phi)| \geq 1.$$

It is also not clear whether all functions of σ , even if they are of the form (3), can be represented by functions $\phi(z)$ and $\psi(z)$ which are both regular in $|z| < 1$.

It is the purpose of this paper to obtain the following theorem: For $f(z) \in \sigma$, $|a_2| < 1.51$.

2. **Application of the area-theorem.** The area-theorem applied to the function $f(z) \in S$ in (1) yields:

Received by the editors February 20, 1966 and, in revised form, July 16, 1966.

¹ Oral communication.

$$(4) \quad \sum_{p=1}^{\infty} p |c_p|^2 \leq 1,$$

where

$$\frac{1}{f(z)} = z^{-1} + \sum_{p=0}^{\infty} c_p z^p.$$

Let $f_n(z) = [f(z^n)]^{(1/n)}$ for integral $n \geq 1$. If $f(z) \in S$, then $f_n(z) \in S$. It follows that if $f(z) \in \sigma$ then $f_n(z) \in \sigma$. Applying inequality (3) to $f_n(z)$ we obtain

$$(n-1) |a_2|^2 + (2n-1) |a_3 - ((n+1)/2n)a_2^2|^2 \leq n^2$$

and hence:

$$(5) \quad \left| a_3 - \frac{n+1}{2n} a_2^2 \right| \leq \left(\frac{n^2 - (n-1) |a_2|^2}{2n-1} \right)^{1/2}.$$

Applying (4) to the function $f_n^{-1}(z)$ we get:

$$(6) \quad \left| a_3 - \frac{3n-1}{2n} a_2^2 \right| \leq \left(\frac{n^2 - (n-1) |a_2|^2}{2n-1} \right)^{1/2}.$$

Inequalities (5) and (6) imply:

$$\frac{(n-1)}{2n} |a_2|^2 \leq \left(\frac{n^2 - (n-1) |a_2|^2}{2n-1} \right)^{1/2}.$$

By squaring both sides a quadratic inequality for $|a_2|^2$ is obtained, whence:

$$(7) \quad |a_2|^2 \leq \frac{2n^2}{(n-1)((2n)^{1/2} + 1)}.$$

For $n=2, 3$ and 4 the right-hand side of (7) takes the values 2.67, 2.61 and 2.78 respectively. We thus get the best result for $n=3$, so that:

$$|a_2|^2 < 2.61,$$

or:

$$(8) \quad |a_2| < 1.62,$$

This bound can be improved by the use of Grunsky's inequalities.

3. The l -sequence $\{l_k\}$. We use the sequence $\{l_k\}$ introduced by E. Jabotinsky [3]. Let $f(s, z)$ be the s -iterate of

$$f(z) = z + a_2 z^2 + \dots$$

That is, let $f(s, z)$ be defined for integer s by:

$$f(1, z) = f(z) \quad \text{and} \quad f(s, z) = f[(s-1), f(z)].$$

Then:

$$f(s, z) = z + \sum_{p=2}^{\infty} a_p(s)z^p,$$

where $a_p(s)$ is a polynomial of degree $(p-1)$ in s . We define:

$$(9) \quad l_k = \frac{d}{ds} a_{k-1}(s) \Big|_{s=0}.$$

The sequences $\{a_i\}$ and $\{l_k\}$ determine each other uniquely. In particular:

$$(10) \quad a_2 = l_1, \quad a_3 = l_2 + l_1^2, \quad a_4 = l_3 + \frac{5}{2} l_1 l_2 + l_1^3;$$

also:

$$(11) \quad l_1 = a_2, \quad l_2 = a_3 - a_2^2, \quad l_3 = a_4 - \frac{5}{2} a_2 a_3 + \frac{3}{2} a_2^3.$$

The sequence $\{l_k\}$ has the following two properties:

(A) Let $\{\lambda_k\}$ be the l -sequence of the function $f^{-1}(z) = f(-1, z)$, then:

$$(12) \quad \lambda_k = -l_k.$$

(B) Let $\{l_k^{(n)}\}$ be the l -sequence of the function $f_n(z) = [f(z^n)]^{1/n}$.

We have:

$$(13) \quad l_k^{(n)} = \begin{cases} (1/n)l_{k/n} & \text{if } n \mid k; \\ 0 & \text{if } n \nmid k. \end{cases}$$

PROOF OF A. Let $L(z) = \sum_{k=1}^N l_k z^{k+1}$ (where N is a large but otherwise arbitrary integer). $L(z)$ then defines an analytic function $f^*(z) = z + a_2 z^2 + \dots$ and $L(z)$ is the coefficient of s in the Taylor expansion of $f^*(s, z)$ about $s=0$. The existence of such an expansion is proved in [2]. On the other hand the first $N+1$ coefficients of the respective power-series expansions of $f^*(z)$ and $f(z)$ coincide. (If $f(s, z)$ admits a power-series expansion in s , we may define $L(z) = \sum_{k=1}^{\infty} l_k z^{k+1}$ which has then a positive radius of convergence [3], so that $f^*(z) \equiv f(z)$.) Let now $g^*(z)$ be the inverse of $f^*(z)$ so that

$$(14) \quad g^*(s, z) = f^*(-s, z).$$

It follows immediately that

$$(15) \quad L^{(g^*)}(z) \equiv -L^{(f^*)}(z)$$

whence (12) holds for every $k \leq N$.

Since N may be taken arbitrarily large, (12) holds for every k .

PROOF OF B. Define $F_n(z) = [f^*(z^n)]^{1/n}$.

It is easily seen that

$$(16) \quad F_n(s, z) = [f^*(s, z^n)]^{1/n}.$$

Differentiating both sides with respect to s gives:

$$(17) \quad \frac{\partial F_n(s, z)}{\partial s} = \frac{1}{n} f^*(s, z^n)^{1/n-1} \frac{\partial f^*(s, z^n)}{\partial s}.$$

Putting now $s=0$ and noting that $f^*(0, z) \equiv z$, we obtain

$$(18) \quad L^{(n)}(z) = \frac{1}{n} z^{1-n} L(z^n),$$

or

$$(19) \quad \sum_{\nu=1}^{\infty} l_{\nu}^{(n)} z^{\nu+1} = \frac{1}{n} z^{1-n} \sum_{k=1}^N l_k(z^n)^{k+1} = \sum_{k=1}^N \frac{1}{n} l_k z^{nk+1},$$

whence, by equating coefficients, B is obtained immediately.

4. Grunsky's inequalities. Grunsky [1] established a necessary and sufficient condition for the function $f(z) = z + a_2 z^2 + \dots$ to be univalent in $|z| < 1$. Let the numbers g_{mn} be defined by:

$$\log \frac{f(z) - f(w)}{z - w} = \log \frac{f(z)}{z} + \log \frac{f(w)}{w} + \sum_{m,n}^{\infty} g_{mn} z^m w^n.$$

Then Grunsky's condition is:

$$(20) \quad \left| \sum_{m,n=1}^N g_{mn} x_m x_n \right| \leq \sum_{p=1}^N \frac{|x_p|^2}{p}$$

for every choice of the complex numbers x_k and all N .

In [3] the g_{mn} corresponding to the function $f(z)$ are given explicitly in terms of the l_k of that function. In particular:

$$g_{11} = l_2,$$

$$g_{12} = g_{21} = l_3 + \frac{1}{2} l_1 l_2,$$

$$g_{13} = g_{31} = l_4 + l_1 l_3 + \frac{1}{2} l_2^2 + \frac{1}{3} l_1^2 l_2,$$

$$g_{22} = l_4 + l_1 l_3 + \frac{1}{3} l_1^2 l_2,$$

$$g_{33} = l_6 + 2l_1 l_5 + l_2 l_4 + \frac{1}{2} l_1^2 l_2^2 + 2l_1^2 l_4 + \frac{4}{3} l_1 l_2 l_3 + \frac{1}{3} l_2^3 + l_1^3 l_3 + \frac{1}{5} l_1^4 l_2.$$

For bi-univalent functions, Grunsky's inequalities (20) will come in pairs for every N , obtained from each other by replacing, according

to (12), every l_k by $-l_k$. Hence, in these inequalities, odd powers of the l 's can be separated from the even powers, yielding separate inequalities, by using the fact that if $|a+b| \leq k$ and $|a-b| \leq k$ then $|a| \leq k$ and $|b| \leq k$.

5. **A bound for $|l_1 l_3|$.** Grunsky's inequality (20) for $N=2$ becomes:

$$(21) \quad |l_2 x_1^2 + 2(l_3 + \frac{1}{2}l_1 l_2)x_1 x_2 + (l_4 + l_1 l_3 + \frac{1}{3}l_1^2 l_2)x_2^2| \leq |x_1|^2 + \frac{1}{2}|x_2|^2.$$

Taking $x_1=0$, $x_2=1$ and omitting all odd powers of the l 's, we get, for bi-univalent functions:

$$(22) \quad |l_1 l_3| \leq \frac{1}{2}.$$

This result can be improved for functions in σ by the use of (8) (valid for functions of σ only). Indeed, choosing $x_1=l_1$, $x_2=\beta>0$ and omitting all odd powers of the l 's, (21) yields, for functions in σ :

$$|l_1 l_3| (2\beta + \beta^2) \leq |l_1|^2 + \frac{1}{2}\beta^2.$$

Here the best result for $|l_1 l_3|$, given $|l_1|^2$, is obtained by taking:

$$\beta = |l_1|^2 + (|l_1|^4 + 2|l_1|^2)^{1/2},$$

whence:

$$(23) \quad |l_1 l_3| \leq \frac{1}{(1+2/|l_1|^2)^{1/2}+1}.$$

Using the fact that $l_1=a_2$, and the bound (8) obtained by taking $n=3$ in (7):

$$(24) \quad |l_1 l_3| \leq \frac{3}{3 + (11 + 2.6)^{1/2}} < 0.43.$$

We shall use this bound to improve (8).

6. **An improved bound for $|a_2| = |l_1|$.** Grunsky's inequality (20) for $N=3$, written for $f_2(z) = (f(z^2))^{1/2}$, using (13) and taking $x_2=0$, is:

$$(25) \quad \left| \frac{1}{2}l_1 x_1^2 + (l_2 + \frac{1}{4}l_1^2)x_1 x_3 + (\frac{1}{2}l_3 + \frac{1}{4}l_1 l_2 + \frac{1}{24}l_1^3)x_3^2 \right| \leq |x_1|^2 + \frac{1}{3}|x_3|^2,$$

Choosing $x_1=l_1$, $x_3=\beta>0$ and omitting all even powers of the l 's, (25) becomes:

$$|l_1^3(\frac{1}{2} + \frac{1}{4}\beta + \frac{1}{24}\beta^2) + \frac{1}{2}l_3\beta^2| \leq |l_1|^2 + \frac{1}{3}\beta^2,$$

whence:

$$(26) \quad |l_1|^3(\frac{1}{2} + \frac{1}{4}\beta + \frac{1}{24}\beta^2) \leq |l_1|^2 + \frac{1}{3}\beta^2 + \frac{1}{2}|l_3|\beta^2.$$

Multiplying by $|l_1|$ and substituting for $|l_1 l_3|$ from (24) this becomes

$$|l_1|^4 \left(\frac{1}{2} + \frac{1}{4}\beta + \frac{1}{24}\beta^2 \right) - |l_1|^3 - \frac{1}{3}\beta |l_1| < \frac{0.43}{2} \beta^2.$$

Choosing $\beta = 1.5$ this yields, for functions in σ :

$$(25) \quad |a_2| < 1.51$$

Further improvement of both bounds (8) and (24) by the interplay of inequalities (23) and (26) yields very little. We have not succeeded in bringing down the bound in (25) to $\frac{3}{2}$.

BIBLIOGRAPHY

1. H. Grunsky, *Koeffizientenbedingungen für schlicht abbildende meromorphe Funktionen*, Math. Z. **45** (1939), 29–61.
4. P. Erdős and E. Jabotinsky, *On analytic iteration*, J. Analyse Math. **8** (1960–1961), 361–376.
3. E. Jabotinsky, *Analytic iteration*, Trans. Amer. Math. Soc. **108** (1963), 457–477.

TECHNION, HAIFA, ISRAEL