

## ON A FIXED POINT THEOREM OF GREGUŠ

**BRIAN FISHER**

Department of Mathematics  
University of Leicester  
Leicester LE1 7RH, England

and

**SALVATORE SESSA**

Istituto Matematico  
Facoltà Di Architettura  
Università Di Napoli  
Via Monteoliveto 3 80134 Naples, Italy

(Received May 18, 1984)

ABSTRACT. We consider two selfmaps  $T$  and  $I$  of a closed convex subset  $C$  of a Banach space  $X$  which are weakly commuting in  $X$ , i.e.

$$\|T I x - I T x\| \leq \|I x - T x\| \text{ for any } x \text{ in } X,$$

and satisfy the inequality

$$\|T x - T y\| \leq a \|I x - I y\| + (1 - a) \max \{ \|T x - I x\|, \|T y - I y\| \}$$

for all  $x, y$  in  $C$ , where  $0 < a < 1$ . It is proved that if  $I$  is linear and non-expansive in  $C$  and such that  $IC$  contains  $TC$ , then  $T$  and  $I$  have a unique common fixed point in  $C$ .

KEY WORDS AND PHRASES. *Common fixed point, Banach space.*

1980 MATHEMATICS SUBJECT CLASSIFICATION CODE. 54H25, 47H10.

### 1. INTRODUCTION.

The second author [1], generalizing a result of Das and Naik [2], defined two mappings  $T$  and  $I$  of a metric space  $(X, d)$  into itself to be weakly commuting if

$$d(TIx, ITx) \leq d(Ix, Tx) \tag{1.1}$$

for all  $x$  in  $X$ . Two commuting mappings clearly satisfy (1.1) but the converse is not generally true as is shown with the following example:

EXAMPLE 1. Let  $X = [0, 1]$  with the Euclidean metric and define  $T$  and  $I$  by

$$Tx = x/(x+4), \quad Ix = x/2$$

for all  $x$  in  $X$ . Then

$$\begin{aligned} d(TIx, ITx) &= \frac{x}{x+8} - \frac{x}{2x+8} = \frac{x^2}{2(x+8)(x+4)} \\ &\leq \frac{x^2 + 2x}{2(x+4)} = \frac{x}{2} - \frac{x}{x+4} = d(Ix, Tx) \end{aligned}$$

for all  $x$  in  $X$  but for any  $x \neq 0$ :

$$TIx = x/(x+8) > x/(2x+8) = ITx.$$

From now on,  $C$  denotes a closed convex subset of a Banach space  $X$ . In a recent paper Gregus [3] proved the following theorem:

**THEOREM 1.** Let  $T$  be a mapping of  $C$  into itself satisfying the inequality

$$\|Tx - Ty\| \leq a\|x - y\| + b\|Tx - x\| + c\|Ty - y\| \quad (1.2)$$

for all  $x, y$  in  $C$ , where  $0 < a < 1$ ,  $b \geq 0$ ,  $c \geq 0$  and  $a + b + c = 1$ . Then  $T$  has a unique fixed point.

Mappings satisfying inequality (1.2) with  $a = 1$  and  $b = c = 0$  are called nonexpansive and were considered by Kirk [4].

Wong [5] studied mappings satisfying inequality (1.2) with  $a = 0$  and  $b = c = \frac{1}{2}$ .

## 2. MAIN RESULTS.

We now prove the following generalization of Theorem 1:

**THEOREM 2.** Let  $T$  and  $I$  be two weakly commuting mappings of  $C$  into itself satisfying the inequality

$$\|Tx - Ty\| \leq a\|Ix - Iy\| + (1-a) \max \{ \|Tx - Ix\|, \|Ty - Iy\| \} \quad (2.1)$$

for all  $x, y$  in  $C$ , where  $0 < a < 1$ . If  $I$  is linear, nonexpansive in  $C$  and such that  $IC$  contains  $TC$ , then  $T$  and  $I$  have a unique common fixed point in  $C$ .

**PROOF.** Let  $x = x_0$  be an arbitrary point in  $C$  and choose points  $x_1, x_2, x_3$  in  $C$  such that

$$Ix_1 = Tx, Ix_2 = Tx_1, Ix_3 = Tx_2.$$

This can be done since  $IC$  contains  $TC$ . Then for  $r = 1, 2, 3$  we have on using inequality (2.1)

$$\begin{aligned} \|Tx_r - Ix_r\| &= \|Tx_r - Tx_{r-1}\| \\ &\leq a\|Ix_r - Ix_{r-1}\| + (1-a) \max \{ \|Tx_r - Ix_r\|, \|Tx_{r-1} - Ix_{r-1}\| \} \\ &= a\|Tx_{r-1} - Ix_{r-1}\| + (1-a) \max \{ \|Tx_r - Ix_r\|, \|Tx_{r-1} - Ix_{r-1}\| \} \end{aligned}$$

and so

$$\|Tx_r - Ix_r\| \leq \|Tx_{r-1} - Ix_{r-1}\|.$$

It follows that

$$\|Tx_r - Ix_r\| \leq \|Tx - Ix\| \quad (2.2)$$

for  $r = 1, 2, 3$ .

Further

$$\begin{aligned} \|Tx_2 - Tx\| &\leq a\|Ix_2 - Ix\| + (1-a) \max \{ \|Tx_2 - Ix_2\|, \|Tx - Ix\| \} \\ &\leq a(\|Tx_1 - Ix_1\| + \|Tx - Ix\|) + (1-a) \|Tx - Ix\| \\ &\leq (1+a) \|Tx - Ix\| \end{aligned}$$

on using inequality (2.2). Thus

$$\|Tx_2 - Ix_1\| \leq (1+a) \|Tx - Ix\|. \quad (2.3)$$

We will now define a point  $z$  by

$$z = \frac{1}{2} x_2 + \frac{1}{2} x_3.$$

Since  $C$  is convex the point  $z$  is in  $C$  and being  $I$  linear, we have

$$Iz = \frac{1}{2} Ix_2 + \frac{1}{2} Ix_3 = \frac{1}{2} Tx_1 + \frac{1}{2} Tx_2.$$

It follows that

$$\begin{aligned} \|Tz - Iz\| &\leq \frac{1}{2} \|Tz - Tx_1\| + \frac{1}{2} \|Tz - Tx_2\| \\ &\leq \frac{1}{2} [a\|Iz - Ix_1\| + (1-a) \max \{\|Tz - Iz\|, \|Tx_1 - Ix_1\|\}] \\ &\quad + \frac{1}{2} [a\|Iz - Ix_2\| + (1-a) \max \{\|Tz - Iz\|, \|Tx_2 - Ix_2\|\}] \\ &= \frac{1}{2} a(\|Iz - Ix_1\| + \|Iz - Ix_2\|) + (1-a) \max \{\|Tz - Iz\|, \|Tx - Ix\|\} \end{aligned}$$

on using inequalities (2.1) and (2.2). Now

$$\begin{aligned} \|Iz - Ix_1\| &\leq \frac{1}{2} \|Ix_2 - Ix_1\| + \frac{1}{2} \|Ix_3 - Ix_1\| \\ &= \frac{1}{2} \|Tx_1 - Ix_1\| + \frac{1}{2} \|Tx_2 - Ix_1\| \\ &\leq (1 + \frac{1}{2} a) \|Tx - Ix\| \end{aligned}$$

from inequalities (2.2) and (2.3) and

$$\|Iz - Ix_2\| = \frac{1}{2} \|Ix_3 - Ix_2\| = \frac{1}{2} \|Tx_2 - Ix_2\| \leq \frac{1}{2} \|Tx - Ix\|.$$

It follows that

$$\|Tz - Iz\| \leq \frac{1}{4} a(3+a) \|Tx - Ix\| + (1-a) \max \{\|Tz - Iz\|, \|Tx - Ix\|\}$$

and so

$$\|Tz - Iz\| \leq \lambda \|Tx - Ix\|$$

where

$$\lambda = (4 - a + a^2)/4 < 1.$$

We therefore have

$$\inf \{\|Tz - Iz\| : z = \frac{1}{2} x_2 + \frac{1}{2} x_3\} \leq \lambda \cdot \inf \{\|Tx - Ix\| : x \in C\}$$

and since we obviously have

$$\inf \{\|Tz - Iz\| : z = \frac{1}{2} x_2 + \frac{1}{2} x_3\} \geq \inf \{\|Tx - Ix\| : x \in C\},$$

it follows that

$$\inf \{ \|Tx - Ix\| : x \in C \} = 0.$$

Each of the sets

$$K_n = \{x \in C : \|Tx - Ix\| \leq 1/n\}, H_n = \{x \in C : \|Tx - Ix\| \leq (a+1)/an\}$$

(for  $n = 1, 2, \dots$ ) must therefore be non-empty and obviously

$$K_1 \supseteq K_2 \supseteq \dots \supseteq K_n \supseteq \dots$$

Thus each of the sets  $\overline{TK}_n$ , where  $\overline{TK}_n$  denotes the closure of  $TK_n$ , must be non-empty for  $n = 1, 2, \dots$  and

$$\overline{TK}_1 \supseteq \overline{TK}_2 \supseteq \dots \supseteq \overline{TK}_n \supseteq \dots$$

Further, for arbitrary  $x, y$  in  $K_n$ ,

$$\|Tx - Ty\| \leq a \|Ix - Iy\| + (1-a) \max \{ \|Tx - Ix\|, \|Ty - Iy\| \}$$

$$\leq a(\|Tx - Ix\| + \|Tx - Ty\| + \|Ty - Iy\|) + (1-a)/n$$

$$\leq (a+1)/n + a\|Tx - Ty\|$$

and so

$$\|Tx - Ty\| \leq \frac{a+1}{(1-a)n}.$$

Thus

$$\lim_{n \rightarrow \infty} \text{diam}(TK_n) = \lim_{n \rightarrow \infty} \text{diam}(\overline{TK}_n) = 0.$$

It follows, by a well known result of Cantor (see, for example [6], p. 156), that the intersection  $\bigcap_{n=1}^{\infty} \overline{TK}_n$  contains exactly one point  $w$ .

Now let  $y$  be an arbitrary point in  $\overline{TK}_n$ . Then for arbitrary  $\epsilon > 0$  there exists a point  $y'$  in  $K_r$  such that

$$\|Ty' - y\| < \epsilon \tag{2.4}$$

and so, using the weak commutativity of  $T$  and  $I$  and the nonexpansiveness of  $I$ , we have from (2.1) and (2.4):

$$\begin{aligned} \|Ty - Iy\| &\leq \|Ty - TIy'\| + \|TIy' - ITy'\| + \|ITy' - Iy\| \\ &\leq a\|Iy - I^2y'\| + (1-a) \max \{ \|Ty - Iy\|, \|TIy' - I^2y'\| \} \\ &\quad + \|Iy' - Ty'\| + \|Ty' - y\| \\ &\leq a\|y - Iy'\| + (1-a) \max \{ \|Ty - Iy\|, \|TIy' - ITy'\| + \|Ty' - Iy'\| \} \\ &\quad + 1/n + \epsilon \\ &\leq a(\|y - Ty'\| + 1/n) + (1-a) \max \{ \|Ty - Iy\|, 2/n \} + 1/n + \epsilon \\ &\leq (1+a)\epsilon + (a+1)/n + (1-a) \max \{ \|Ty - Iy\|, 2/n \}. \end{aligned}$$

Since  $\epsilon$  is arbitrary it follows that

$$\|Ty - Iy\| \leq (a+1)/n + (1-a) \max \{ \|Ty - Iy\|, 2/n \}. \tag{2.5}$$

If  $\|Ty - Iy\| \leq 2/n$ , then we have

$$\|Ty - Iy\| \leq 2/n < (a+1)/an.$$

If  $||Ty - Iy|| > 2/n$ , (2.5) implies

$$||Ty - Iy|| \leq (a+1).n + (1-a).||Ty - Iy||.$$

So in both cases  $y$  lies in  $H_n$ . Thus  $\overline{TK_n} \subseteq H_n$  and so the point  $w$  must be in  $H_n$  for  $n = 1, 2, \dots$ . It follows that

$$||Tw - Iw|| \leq (a+1)/an$$

for  $n = 1, 2, \dots$  and so  $Tw = Iw$ .

Since (1.1) holds, we also have  $ITw = TIw$ . Thus

$$\begin{aligned} ||T^2w - Tw|| &\leq a||ITw - Iw|| + (1-a) \max \{ ||T^2w - ITw||, ||Tw - Iw|| \} \\ &= a||T^2w - Tw|| \end{aligned}$$

and it follows that  $Tw = w'$  is a fixed point of  $T$  since  $a < 1$ . Further  $Iw' = ITw + TIw = ITw = Tw' = w'$  and so  $w'$  is also a fixed point of  $I$ . Now suppose that  $T$  and  $I$  have a second common fixed point  $w''$ . Then

$$\begin{aligned} ||w' - w''|| &= ||Tw' - Tw''|| \\ &\leq a||Iw' - Iw''|| + (1-a) \max \{ ||Tw' - Iw'||, ||Tw'' - Iw''|| \} \\ &\leq a||w' - w''|| \end{aligned}$$

and the uniqueness of the common fixed point follows since  $a < 1$ . This completes the proof of the theorem.

EXAMPLE 2. Let  $X = \mathbb{R}$  and  $C = [0, 1]$  with the usual norm. Let  $T$  and  $I$  be as in example 1.  $I$  is clearly linear and nonexpansive and further

$$TC = [0, 1/5] \subset [0, 1/2] = IC.$$

Thus

$$||Tx - Ty|| = \frac{4||x - y||}{(x+4)(y+4)} \leq \frac{1}{2} \cdot \frac{||x - y||}{2} = \frac{1}{2} ||Ix - Iy||$$

for all  $x, y$  in  $C$  and inequality (2.1) is satisfied for  $a = 1/2$ .

So all the assumptions of Theorem 2 hold and  $w = 0$  is the unique common fixed point of  $T$  and  $I$ .

Letting  $I$  be the identity mapping in Theorem 2, we have the following corollary which extends Theorem 1:

COROLLARY. Let  $T$  be a mapping of  $C$  into itself satisfying the inequality

$$||Tx - Ty|| \leq a||x - y|| + (1-a) \max \{ ||Tx - x||, ||Ty - y|| \}$$

for all  $x, y$  in  $C$ , where  $0 < a < 1$ . Then  $T$  has a unique fixed point.

The result of this corollary was given in [7].

We note that the weak commutativity in Theorem 2 is a necessary condition. It suffices to consider the following example:

EXAMPLE 3. Let  $X = \mathbb{R}$  and let  $C = [0, 1]$  with the usual norm.

Define  $T$  and  $I$  by  $Tx = 1/3$ ,  $Ix = x/2$  for any  $x$  in  $C$ .

It is easily seen that all the conditions of Theorem 2 are satisfied except that of weak commutativity since with  $x = 1/2$

$$\|T(1/2) - IT(1/2)\| = 1/6 > 1/12 = \|T(1/2) - I(1/2)\| .$$

However  $T$  and  $I$  do not have a common fixed point.

We conclude that although the mappings  $T$  and  $I$  in Theorem 2 have a unique common fixed point in  $C$ , it is possible for them to have other fixed points, as proved in the next example:

Example 4. Let  $X = C = \mathbb{R}^2$  with norm

$$\|(x,y)\| = \max \{|x|, |y|\}$$

for all  $(x,y)$  in  $\mathbb{R}^2$ . Define mappings  $T$  and  $I$  on  $\mathbb{R}^2$  by

$$T(x,y) = (0,y), \quad I(x,y) = (x,-y)$$

for all  $(x,y)$  in  $\mathbb{R}^2$ . Then for all  $(x,y) \in \mathbb{R}^2, (x',y') \in \mathbb{R}^2$

$$\|T(x,y) - T(x',y')\| = |y - y'|$$

and

$$\begin{aligned} a\|I(x,y) - I(x',y')\| &+ (1-a) \max \{\|T(x,y) - I(x,y)\|, \|T(x',y') - I(x',y')\|\} \\ &= a \max \{|x-x'|, |y-y'|\} + (1-a) \max \{|x|, 2|y|, |x'|, 2|y'|\} \\ &\geq a|y - y'| + 2(1-a) \max \{|y|, |y'|\} \\ &\geq a|y - y'| + (1-a) (|y| + |y'|) \\ &\geq |y-y'| \end{aligned}$$

if  $0 < a < 1$ . Since  $T$  commutes with  $I$  and  $I$  is a linear isometry, it follows that all the conditions of Theorem 2 are satisfied but  $T$  and  $I$  each have an infinite number of fixed points.

#### REFERENCES

1. SESSA, S. On a Weak Commutativity Condition of Mappings in Fixed Point Considerations, Publ. Inst. Math., 32 (46) (1982), 149-153.
2. DAS, K.M. and NAIK, K.V. Common Fixed Point Theorems for Commuting Maps on a Metric Space, Proc. Amer. Math. Soc., 77 (1979), 369-373.
3. GREGUŠ, Jr., M. A Fixed Point Theorem in Banach Space, Boll. Un. Mat. Ital., (5) 17-A (1980), 193-198.
4. KIRK, W.A. A Fixed Point Theorem for Mappings Which do not Increase Distances, Amer. Math Monthly, 72(1965), 1004-1006.
5. WONG, Ch. S. On Kannan Maps, Proc. Amer. Math. Soc., 47 (1975), 105-111.
6. DUGUNDIJ, J. and GRANAS, A. Fixed Point Theory I, Polish Scientific Publishers, Warsaw (1982).
7. FISHER, B. Common Fixed Points on a Banach Space, Chung Juan J., XI (1982), 12-15.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

