

On a free group of transformations defined by an automaton

Mariya Vorobets* and Yaroslav Vorobets*[†]

Abstract

We prove that three automorphisms of the rooted binary tree defined by a certain 3-state automaton generate a free non-Abelian group of rank 3.

1 Introduction

A Mealy automaton A over a finite alphabet X is determined by the set of internal states Q , the state transition function $\phi : Q \times X \rightarrow Q$, and the output function $\psi : Q \times X \rightarrow X$. The automaton starts its work at some state $q \in Q$ and a sequence of letters $x_1, x_2, \dots, x_n \in X$ is input into A . The automaton uses the function ψ to produce an output sequence y_1, y_2, \dots, y_n ($y_i \in X$) while changing its states according to the function ϕ . This gives rise to a transformation $A_q : X^* \rightarrow X^*$ of the set X^* of finite words over alphabet X . A detailed account of the theory of Mealy automata is given in the survey paper [GNS].

An automaton is called finite if it has only finitely many states. An automaton is called initial if it has a fixed initial state. An initial automaton over an alphabet X defines a transformation of the set X^* . A finite non-initial automaton defines a finite number of transformations, each being assigned to an internal state. Assuming all of them are invertible, the finite automaton defines a finitely generated transformation group. This group is an example of a self-similar group (see [Nek]).

*Partially supported by the NSF grants DMS-0308985 and DMS-0456185.

[†]The second author is supported by a Clay Research Scholarship.




Figure 1: Aleshin's automata.

All invertible transformations defined by finite initial automata over a fixed alphabet X form a countable group $\mathcal{G}(X)$ (see [GNS]). The set X^* is endowed with the structure of a rooted regular tree so that elements of $\mathcal{G}(X)$ are automorphisms of the tree. The groups of finite automaton transformations contain many finitely generated subgroups with remarkable properties (e.g., infinite torsion groups, groups of intermediate growth) but their complete structure is yet to be understood.

The problem of embedding the free non-Abelian group into a group $\mathcal{G}(X)$ of finite automaton transformations turned out to be hard. The first attempt to solve it was made by Aleshin [A]. He introduced two finite initial automata over alphabet $\{0, 1\}$ and claimed that two transformations of the rooted binary tree $\{0, 1\}^*$ defined by these automata generate a free group. However the paper [A] does not contain a proof. There is only a sketch that raises reasonable doubts as to whether Aleshin's argument was complete.

Aleshin's automata are depicted in Figure 1 by means of Moore diagrams. The Moore diagram of an automaton is a directed graph with labeled edges. The vertices are the states of the automaton and edges are state transition routes. Each label consists of two letters from the alphabet. The left one is the input field, it is used to choose a transition route. The right one is the output generated by the automaton. Aleshin considered these automata as initial, with initial state b .

The first reliable result was obtained years later by Brunner and Sidki [BS]. They showed that the group $\text{GL}(n, \mathbb{Z})$ can be embedded into the group

of finite automaton transformations over the alphabet of cardinality 2^n . Since $\text{SL}(2, \mathbb{Z})$ contains a free group, this yields two finite initial automata over a 4-letter alphabet generating the free group. An example of a free group generated by finite automaton transformations over a 2-letter alphabet was given by Olijnyk and Sushchanskij [OS].

A harder problem is to present the free group as the group defined by a single finite non-initial automaton. Recently it was solved by Glasner and Mozes [GM]. They constructed infinitely many finite automata of algebraic origin that define transformation groups with various properties, in particular, free groups. The simplest example from [GM] is a pair of automata A_1 and A_2 . The automaton A_1 is a 14-state automaton over a 6-letter alphabet while A_2 is a 6-state automaton over a 14-letter alphabet. The automata define free groups on 7 and 3 generators, respectively. Note that the number of free generators is half of the number of states. The 14 transformations defined by the automaton A_1 form a symmetric set of free generators. That is, they split into two sets of free generators such that the elements in each set are inverses of elements in the other set. The 6 transformations defined by A_2 also form a symmetric set of free generators. The automata A_1 and A_2 are dual in that either of them can be obtained from the other by interchanging the alphabet with the set of internal states and the state transition function with the output function.

In this paper, we consider the problem of finding a finite automaton that defines a free group such that the rank of the group (i.e., the number of free generators) is equal to the number of states of the automaton. A candidate for a solution has been available for some time. Namely, Brunner and Sidki conjectured (see [S]) that the first of two Aleshin's automata shown in Figure 1 is the required one. Here we prove that this is indeed the case.

Theorem 1.1 *Three automorphisms of the rooted binary tree defined by the first Aleshin automaton generate a free transformation group on three generators.*

The proof of the theorem is based on the dual automaton approach (already used in [GM]). The main idea of this approach is that properties of the group defined by a finite automaton are determined by orbits of the transformation group defined by the dual automaton.

The paper is organized as follows. Section 2 addresses some general constructions concerning automata and their properties. In Section 3 we consider the Aleshin automaton along with a number of related automata and

establish some properties of these automata. In Section 4 we use results of Sections 2 and 3 to prove Theorem 1.1.

2 Automata

An *automaton* A is a quadruple (Q, X, ϕ, ψ) consisting of two nonempty sets Q, X along with two maps $\phi : Q \times X \rightarrow Q, \psi : Q \times X \rightarrow X$. The set X is to be finite, it is called the *input/output alphabet* of the automaton. We say that A is an automaton over the alphabet X . Q is called the set of *internal states* of A . The automaton A is called *finite* if the set Q is finite. ϕ and ψ are called the *state transition function* and the *output function*, respectively. One may regard these functions as a single map $(\phi, \psi) : Q \times X \rightarrow Q \times X$.

The automaton A functions as follows. The active automaton is always supposed to be in some state $q \in Q$. The automaton reads an input letter $x \in X$. Then it performs two independent tasks: passes to the state $\phi(q, x)$ and sends the letter $\psi(q, x)$ to the output. After that the automaton is ready to accept another input letter. Usually the automaton job consists of transducing the whole sequence of input letters. Suppose that A started its work in a state $q \in Q$ (the *initial state*) and a word $w = x_1x_2 \dots x_n$ over the alphabet X was input into A . As the result of automaton's job, we obtain two sequences: a sequence of states $q_0 = q, q_1, \dots, q_n$, which describes the internal work of the automaton, and the output word $v = y_1y_2 \dots y_n$. Here $q_i = \phi(q_{i-1}, x_i)$ and $y_i = \psi(q_{i-1}, x_i)$ for $1 \leq i \leq n$.

Let X^* denote the set of words over the alphabet X . A *word* $w \in X^*$ is merely a finite sequence whose elements belong to X . However the elements of w are called *letters* and w is usually written so that its elements are not separated by delimiters. The number of letters of a word w is called its *length* and denoted by $|w|$. It is assumed that X^* contains the empty word \emptyset . The set X is naturally embedded in X^* . If $w_1 = x_1 \dots x_n$ and $w_2 = y_1 \dots y_m$ are words over the alphabet X then w_1w_2 denotes their concatenation $x_1 \dots x_ny_1 \dots y_m$. X^* is the free monoid generated by all elements of X relative to the operation $(w_1, w_2) \mapsto w_1w_2$. Another structure on X^* is that of a rooted k -regular tree, where k is the cardinality of X . Namely, we consider a graph with the set of vertices X^* where two vertices $w_1, w_2 \in X^*$ are joined by an edge if $w_1 = w_2x$ or $w_2 = w_1x$ for some $x \in X$. The root of the tree is the empty word.

In the above description of the automaton's functions, it is shown how

the output function ψ can be extended to a map $\psi^* : Q \times X^* \rightarrow X^*$. By definition, ψ^* satisfies the recursive relations $\psi^*(q, \emptyset) = \emptyset$, $\psi^*(q, xw) = \psi(q, x)\psi^*(\phi(q, x), w)$ for all $x \in X$, $w \in X^*$, $q \in Q$. Moreover, ψ^* is uniquely determined by these relations. Now for any $q \in Q$ we define a transformation $A_q : X^* \rightarrow X^*$, $A_q(w) = \psi^*(q, w)$ for all $w \in X^*$. We say that the transformation A_q is defined by the automaton A with the initial state q . Clearly, A_q preserves length of words. Besides, A_q transforms words from the left to the right, that is, the first n letters of a word $A_q(w)$ depend only on the first n letters of w . This implies that A_q is an endomorphism of X^* as a rooted regular tree. If A_q is invertible then it belongs to the group $\text{Aut}(X^*)$ of automorphisms of the tree.

The semigroup of transformations of X^* generated by A_q , $q \in Q$ is denoted by $S(A)$. The automaton A is called *invertible* if A_q is invertible for all $q \in Q$. If A is invertible then A_q , $q \in Q$ generate a transformation group $G(A)$, which is a subgroup of $\text{Aut}(X^*)$. We say that $S(A)$ (resp. $G(A)$) is the semigroup (resp. group) defined by the automaton A .

Lemma 2.1 *Suppose the automaton A is invertible. Then the actions of the semigroup $S(A)$ and the group $G(A)$ on X^* have the same orbits.*

Proof. The action of the group $G(A)$ on X^* preserves length of words. Hence an arbitrary word $w \in X^*$ belongs to some finite set $W \subset X^*$ that is invariant under the $G(A)$ action. Since the automaton A is invertible, every transformation $g \in S(A)$ is invertible. The restriction of g to W is a bijection, which is of finite order since the set W is finite. It follows that the semigroup $S(A)$ of transformations of X^* becomes a group when restricted to W . In particular, the restrictions to W of the semigroup $S(A)$ and of the group $G(A)$ coincide. Then the orbits $\{g(w) \mid g \in S(A)\}$ and $\{g(w) \mid g \in G(A)\}$ coincide as well. ■

One way to picture an automaton, which we use in this paper, is the *Moore diagram*. The Moore diagram of an automaton $A = (Q, X, \phi, \psi)$ is a directed graph with labeled edges defined as follows. The vertices of the graph are states of the automaton A . Every edge carries a label of the form $x|y$, where $x, y \in X$. The left field x of the label is referred to as the *input field* while the right field y is referred to as the *output field*. The set of edges of the graph is in a one-to-one correspondence with the set $Q \times X$. Namely, for any $q \in Q$ and $x \in X$ there is an edge that goes from the vertex q to $\phi(q, x)$ and carries the label $x|\psi(q, x)$. The Moore diagram of an automaton

can have loops (edges joining a vertex to itself) and multiple edges. To simplify pictures, we do not draw multiple edges in this paper. Instead, we use multiple labels.

The transformations A_q , $q \in Q$ can be defined in terms of the Moore diagram of the automaton A . For any $q \in Q$ and $w \in X^*$ we find a path δ in the Moore diagram such that δ starts at the vertex q and the word w can be obtained by reading the input fields of labels along δ . Such a path exists and is unique. Then the word $A_q(w)$ is obtained by reading the output fields of labels along the path δ .

Let Γ denote the Moore diagram of the automaton A . We associate to Γ two directed graphs Γ_1 and Γ_2 with labeled edges. Γ_1 is obtained from Γ by interchanging the input and output fields of all labels. That is, a label $x|y$ is replaced by $y|x$. Γ_2 is obtained from Γ by reversing all edges. The *inverse automaton* of A is the automaton whose Moore diagram is Γ_1 . The *reverse automaton* of A is the automaton whose Moore diagram is Γ_2 . If one of the graphs Γ_1 and Γ_2 is the Moore diagram of an automaton then the automaton shares with A the alphabet and internal states. Moreover, its state transition and output functions are uniquely determined by the graph. However neither Γ_1 nor Γ_2 must be the Moore diagram of an automaton. The necessary and sufficient condition for such a labelling to be the Moore diagram of an automaton is that for any $q \in Q$ and $x \in X$ there is exactly one edge of the graph that goes out of the vertex q and has x as the input field of its label.

Proposition 2.2 *An automaton $A = (Q, X, \phi, \psi)$ is invertible if and only if for any $q \in Q$ the map $\psi(q, \cdot) : X \rightarrow X$ is bijective. The inverse automaton I is well defined if and only if A is invertible. If this is the case, then $I_q = A_q^{-1}$ for all $q \in Q$.*

Proof. Let Γ be the Moore diagram of A and Γ_1 be the graph obtained from Γ by interchanging the input and output fields of all labels. For any $x \in X$ and $q \in Q$ the number of edges of Γ_1 that go out of the vertex q and have x as the input field of their labels is equal to the number of $y \in X$ such that $\psi(y, q) = x$. The inverse automaton of A is well defined if this number is always equal to 1, that is, if and only if the map $\psi(\cdot, q) : X \rightarrow X$ is bijective for any $q \in Q$.

Suppose there exists $q \in Q$ such that the map $\psi(\cdot, q)$ is not bijective. Since X is a finite set, $\psi(x_1, q) = \psi(x_2, q)$ for some $x_1, x_2 \in X$, $x_1 \neq x_2$. Then $A_q(x_1) = A_q(x_2)$; in particular, A_q is not invertible.

Now assume that the map $\psi(\cdot, q)$ is bijective for any $q \in Q$. By the above the inverse automaton I of A is well defined. The graphs Γ and Γ_1 , which are the Moore diagrams of automata A and I , differ only in labels. In particular, they share all paths. Given such a path δ , a word $w \in X^*$ can be obtained by reading the input fields of labels along δ in one of the graphs if and only if the same word w is obtained by reading the output fields of labels along δ in the other graph. It follows that for any $q \in Q$ and $w_1, w_2 \in X^*$ we have $w_2 = A_q(w_1)$ if and only if $w_1 = I_q(w_2)$. This means that A_q is invertible for any $q \in Q$ and $I_q = A_q^{-1}$. ■

For any word $w = x_1x_2 \dots x_n$ over an alphabet X we denote by \overleftarrow{w} the reversed word $x_n \dots x_2x_1$.

Proposition 2.3 *Given an automaton $A = (Q, X, \phi, \psi)$, the reverse automaton R is well defined if and only if for any $x \in X$ the map $\phi(\cdot, x) : Q \rightarrow Q$ is bijective. Assume this is the case and let $w_1, w_2 \in X^*$. Then $w_2 = g(w_1)$ for some $g \in S(A)$ if and only if $\overleftarrow{w_2} = h(\overleftarrow{w_1})$ for some $h \in S(R)$.*

Proof. Let Γ be the Moore diagram of A and Γ_2 be the graph obtained from Γ by reversing all edges. For any $q \in Q$ and $x \in X$ the number of edges of Γ_2 that go out of the vertex q and have x as the input field of their labels is equal to the number of $p \in Q$ such that $\phi(p, x) = q$. The reverse automaton R of A is well defined if this number is always equal to 1, that is, if and only if $\phi(\cdot, x) : Q \rightarrow Q$ is bijective for any $x \in X$.

Assume the reverse automaton R is well defined. The graph Γ_2 is the Moore diagram of R . Any path δ in the graph Γ is assigned a “reversed” path δ' in the graph Γ_2 such that a word $w \in X^*$ can be obtained by reading the input (resp. output) fields of labels along δ if and only if the reversed word \overleftarrow{w} is obtained by reading the input (resp. output) fields of labels along δ' . It follows that, given $w_1, w_2 \in X^*$, we have $w_2 = A_q(w_1)$ for some $q \in Q$ if and only if $\overleftarrow{w_2} = R_p(\overleftarrow{w_1})$ for some $p \in Q$.

Let $w_1, w_2 \in X^*$. If $w_2 = g(w_1)$ for some $g \in S(A)$ then there is a sequence of words $u_1 = w_1, u_2, \dots, u_n = w_2$ ($n > 1$) such that $u_{i+1} = A_{q_i}(u_i)$ for some $q_i \in Q$, $1 \leq i \leq n-1$. By the above $u_{i+1} = R_{p_i}(\overleftarrow{u_i})$, where $p_i \in Q$, $1 \leq i \leq n-1$. Hence $\overleftarrow{w_2} = h(\overleftarrow{w_1})$ for some $h \in S(R)$. Similarly, $\overleftarrow{w_2} = h(\overleftarrow{w_1})$ for some $h \in S(R)$ implies that $w_2 = g(w_1)$ for some $g \in S(A)$. ■

Let $A = (Q, X, \phi, \psi)$ be an automaton. For any nonempty word $\xi = q_1q_2 \dots q_n \in Q^*$ we let $A_\xi = A_{q_n} \dots A_{q_2}A_{q_1}$. Also, we let $A_\emptyset = 1$ (here 1 stands

for the unit element of the group $\text{Aut}(X^*)$, i.e., the identity mapping on X^*). Obviously, any element of the semigroup $S(A)$ is represented as A_ξ for a nonempty word $\xi \in Q^*$. The map $X^* \times Q^* \rightarrow X^*$ given by $(w, \xi) \mapsto A_\xi(w)$ defines a right action of the monoid Q^* on the rooted regular tree X^* . That is, $A_{\xi_1\xi_2}(w) = A_{\xi_2}(A_{\xi_1}(w))$ for all $\xi_1, \xi_2 \in Q^*$ and $w \in X^*$.

To each finite automaton $A = (Q, X, \phi, \psi)$ we associate a *dual automaton* D , which is obtained from A by interchanging the alphabet with the set of internal states and the state transition function with the output function. To be precise, $D = (X, Q, \tilde{\phi}, \tilde{\psi})$, where $\tilde{\phi}(x, q) = \psi(q, x)$ and $\tilde{\psi}(x, q) = \phi(q, x)$ for all $x \in X$ and $q \in Q$. Unlike the inverse and reverse automata, the dual automaton is always well defined. It is easy to see that A is the dual automaton of D .

The dual automaton D defines a right action of the monoid X^* on Q^* given by $(\xi, w) \mapsto D_w(\xi)$. This action and the action of Q^* on X^* defined by the automaton A are related in the following way.

Proposition 2.4 *For any $w, u \in X^*$ and $\xi, \eta \in Q^*$,*

$$A_\xi(wu) = A_\xi(w)A_{D_w(\xi)}(u),$$

$$D_w(\xi\eta) = D_w(\xi)D_{A_\xi(w)}(\eta).$$

Proof. First consider the case when one of words w and ξ is empty. If $w = \emptyset$ then $A_\xi(w) = \emptyset$ and $D_w = 1$. If $\xi = \emptyset$ then $A_\xi = 1$ and $D_w(\xi) = \emptyset$.

Further, consider the case when both w and ξ are one-letter words. In this case, $A_\xi(wu) = xA_q(u)$, where $x = \psi(\xi, w)$, $q = \phi(\xi, w)$, while $D_w(\xi\eta) = pD_y(\eta)$, where $p = \tilde{\psi}(w, \xi)$, $y = \tilde{\phi}(w, \xi)$. It remains to observe that $\tilde{\phi}(w, \xi) = \psi(\xi, w) = A_\xi(w)$ and $\phi(\xi, w) = \tilde{\psi}(w, \xi) = D_w(\xi)$.

In the general case, we prove the proposition by induction on the sum of lengths of words w and ξ . By the above the proposition holds when $|w| + |\xi| \leq 2$. Now let $n > 2$ and assume the proposition holds if $|w| + |\xi| < n$. Take arbitrary words $w, u \in X^*$ and $\xi, \eta \in Q^*$ such that $|w| + |\xi| = n$. Since $n > 2$, at least one of the words w, ξ has more than one letter. If $|w| > 1$ then $w = w_1w_2$ for some words $w_1, w_2 \in X^*$ that are shorter than w . Repeatedly using the inductive assumption, we obtain

$$\begin{aligned} A_\xi(wu) &= A_\xi(w_1w_2u) = A_\xi(w_1)A_{D_{w_1}(\xi)}(w_2u) = \\ &A_\xi(w_1)A_{D_{w_1}(\xi)}(w_2)A_{D_{w_2}(D_{w_1}(\xi))}(u) = A_\xi(w_1w_2)A_{D_{w_1w_2}(\xi)}(u). \end{aligned}$$

If $|\xi| > 1$ then $\xi = \xi_1\xi_2$ for some words $\xi_1, \xi_2 \in Q^*$ that are shorter than ξ . By the inductive assumption,

$$\begin{aligned} A_\xi(wu) &= A_{\xi_1\xi_2}(wu) = A_{\xi_2}(A_{\xi_1}(wu)) = A_{\xi_2}(A_{\xi_1}(w)A_{D_w(\xi_1)}(u)) = \\ &A_{\xi_2}(A_{\xi_1}(w))A_\zeta(A_{D_w(\xi_1)}(u)) = A_{\xi_1\xi_2}(w)A_{D_w(\xi_1)\zeta}(u), \end{aligned}$$

where $\zeta = D_{A_{\xi_1}(w)}(\xi_2)$. Since $|w| + |\xi_1| < n$, we have $D_w(\xi_1)\zeta = D_w(\xi_1\xi_2)$.

The equality $D_w(\xi\eta) = D_w(\xi)D_{A_\xi(w)}(\eta)$ is verified in the same way. ■

Corollary 2.5 *Suppose $A_\xi = 1$ for some $\xi \in Q^*$. Then $A_{g(\xi)} = 1$ for every $g \in S(D)$.*

Proof. $g \in S(D)$ means that $g = D_w$ for some word $w \in X^*$. By Proposition 2.4, $A_\xi(wu) = A_\xi(w)A_{g(\xi)}(u)$ for any $u \in X^*$. Since $A_\xi = 1$, we have $wu = wA_{g(\xi)}(u)$, which implies that $A_{g(\xi)}(u) = u$. ■

Let $A = (Q_1, X, \phi_1, \psi_1)$ and $B = (Q_2, X, \phi_2, \psi_2)$ be two automata over the same alphabet X . Assume $Q_1 \cap Q_2 = \emptyset$. The *disjoint union* of automata A and B is the automaton $U = (Q_1 \cup Q_2, X, \phi, \psi)$, where functions ϕ, ψ are such that $\phi = \phi_1$ and $\psi = \psi_1$ on $Q_1 \times X$ while $\phi = \phi_2$ and $\psi = \psi_2$ on $Q_2 \times X$. Obviously, $U_q = A_q$ for $q \in Q_1$ and $U_q = B_q$ for $q \in Q_2$. The Moore diagram of U is the disjoint union of the Moore diagrams of A and B . For example, Figure 2 may be regarded either as the Moore diagrams of two 3-state automata or as the Moore diagram of a single 6-state automaton, their disjoint union. If $Q_1 \cap Q_2 \neq \emptyset$, then the disjoint union of automata A and B is not defined (this is the case for two automata whose Moore diagrams are depicted in Figure 1). However we can rename some states of the automata so that they share no states anymore and then consider the disjoint union of the modified automata.

3 The Aleshin automaton

In this section we turn to the main object of study of this paper, which is the first of the two automata of Aleshin whose Moore diagrams are depicted in Figure 1. This is an automaton A over the alphabet $X = \{0, 1\}$ with the set of internal states $Q = \{a, b, c\}$. The state transition function ϕ and the output function ψ of A are defined as follows: $\phi(a, 0) = \phi(b, 1) = c$,




Figure 2: The Aleshin automaton and its inverse.

$\phi(a, 1) = \phi(b, 0) = b$, $\phi(c, 0) = \phi(c, 1) = a$; $\psi(a, 0) = \psi(b, 0) = \psi(c, 1) = 1$, $\psi(a, 1) = \psi(b, 1) = \psi(c, 0) = 0$.

Using Proposition 2.2, it is easy to verify that A is an invertible automaton. In particular, A_a , A_b , and A_c are automorphisms of the rooted binary tree X^* . Let I denote the inverse automaton of A . By I' denote the automaton obtained from I by renaming its states a , b , c to a^{-1} , b^{-1} , c^{-1} , respectively. Here, a^{-1} , b^{-1} , and c^{-1} are regarded as elements of the free group on generators a , b , c . Further, let B denote the disjoint union of automata A and I' . B is an automaton over the alphabet $X = \{0, 1\}$ with the set of internal states $Q_{\pm} = \{a, b, c, a^{-1}, b^{-1}, c^{-1}\}$. We have $B_a = A_a$, $B_b = A_b$, $B_c = A_c$, $B_{a^{-1}} = A_a^{-1}$, $B_{b^{-1}} = A_b^{-1}$, $B_{c^{-1}} = A_c^{-1}$. In particular, $S(B) = G(A)$. Figure 2 depicts the Moore diagram of the automaton B .

We are going to consider two automata over the alphabet Q_{\pm} . The first




Figure 3: The dual automaton D .




Figure 4: Automaton E .

one is the dual automaton D of the automaton B . It has two states 0 and 1; its Moore diagram is shown in Figure 3. The other one is an auxiliary automaton E . The automaton E has 3 internal states α , β , and γ . Its transition function ϕ_E is defined as follows. If $q \in \{a, b, a^{-1}, b^{-1}\}$ then $\phi_E(\alpha, q) = \beta$ and $\phi_E(\beta, q) = \alpha$. If $q \in \{c, c^{-1}\}$ then $\phi_E(\alpha, q) = \alpha$ and $\phi_E(\beta, q) = \beta$. Also, $\phi_E(\gamma, q) = \gamma$ for all $q \in Q_{\pm}$. The output function ψ_E of E is defined so that $\psi_E(\alpha, q) = \sigma_{\alpha}(q)$, $\psi_E(\beta, q) = \sigma_{\beta}(q)$, and $\psi_E(\gamma, q) = \sigma_{\gamma}(q)$ for all $q \in Q_{\pm}$, where $\sigma_{\alpha} = (a^{-1}b^{-1})$, $\sigma_{\beta} = (ab)$, and $\sigma_{\gamma} = (bc)(b^{-1}c^{-1})$ are permutations on the set Q_{\pm} . The Moore diagram of the automaton E is shown in Figure 4. Using Proposition 2.2, one can verify that D and E are invertible automata.

To each permutation τ on the set $Q = \{a, b, c\}$ we assign a mapping $\pi_{\tau} : Q_{\pm}^* \rightarrow Q_{\pm}^*$ as follows. First we define a permutation $\tilde{\tau}$ on the set Q_{\pm} by $\tilde{\tau}(q) = \tau(q)$, $\tilde{\tau}(q^{-1}) = (\tau(q))^{-1}$ for all $q \in Q$. Then for any nonempty word $\xi = q_1q_2 \dots q_n$ over the alphabet Q_{\pm} we let $\pi_{\tau}(\xi) = \tilde{\tau}(q_1)\tilde{\tau}(q_2) \dots \tilde{\tau}(q_n)$. Besides, we let $\pi_{\tau}(\emptyset) = \emptyset$. Clearly, the mapping π_{τ} is an automorphism of the monoid Q_{\pm}^* .

Lemma 3.1 $\pi_{\tau_1\tau_2} = \pi_{\tau_1}\pi_{\tau_2}$ and $\pi_{\tau^{-1}} = \pi_{\tau}^{-1}$ for any permutations τ, τ_1, τ_2 on the set $\{a, b, c\}$.

Proof. Let τ_1, τ_2 be permutations on $\{a, b, c\}$. Since $\pi_{\tau_1}, \pi_{\tau_2}$, and $\pi_{\tau_1\tau_2}$ are automorphisms of the monoid Q_{\pm}^* , it follows that $\pi_{\tau_1\tau_2} = \pi_{\tau_1}\pi_{\tau_2}$ whenever $\pi_{\tau_1\tau_2}(q) = \pi_{\tau_1}\pi_{\tau_2}(q)$ for all $q \in Q_{\pm}$. Given $q \in \{a, b, c\}$, we have $\pi_{\tau_1}\pi_{\tau_2}(q) = \pi_{\tau_1}(\tau_2(q)) = \tau_1\tau_2(q) = \pi_{\tau_1\tau_2}(q)$, $\pi_{\tau_1}\pi_{\tau_2}(q^{-1}) = \pi_{\tau_1}((\tau_2(q))^{-1}) = (\tau_1\tau_2(q))^{-1} = \pi_{\tau_1\tau_2}(q^{-1})$.

Let τ be a permutation on $\{a, b, c\}$. By the above $\pi_{\tau}\pi_{\tau^{-1}} = \pi_{\tau^{-1}}\pi_{\tau} = \pi_{\text{id}}$, where id denotes the identity map on $\{a, b, c\}$. Obviously, π_{id} is the identity map on Q_{\pm}^* . Hence $\pi_{\tau^{-1}} = \pi_{\tau}^{-1}$. \blacksquare

Lemma 3.2 $E_\alpha^2 = E_\beta^2 = E_\gamma^2 = 1$, $E_\alpha E_\beta = E_\beta E_\alpha = \pi_{(ab)}$, $E_\gamma = \pi_{(bc)}$.
The group $G(E)$ contains π_τ for any permutation τ on $\{a, b, c\}$.

Proof. It is easy to see that the inverse automaton of E coincides with E . Proposition 2.2 implies that $E_\alpha^2 = E_\beta^2 = E_\gamma^2 = 1$. The equality $E_\gamma = \pi_{(bc)}$ follows from the definition of the automaton E .

Consider the following permutations on the set Q_\pm : $\sigma_\alpha = (a^{-1}b^{-1})$, $\sigma_\beta = (ab)$, and $\sigma = (ab)(a^{-1}b^{-1})$. Given an arbitrary word $\xi = q_1q_2 \dots q_k \in Q_\pm^*$, the words $E_\alpha(\xi) = a_1a_2 \dots a_k$, $E_\beta(\xi) = b_1b_2 \dots b_k$, and $\pi_{(ab)}(\xi) = c_1c_2 \dots c_k$ are computed as follows. For any i , $1 \leq i \leq k$ we count the number of times when letters a, b, a^{-1}, b^{-1} occur in the sequence q_1, \dots, q_{i-1} . If this number is even then $a_i = \sigma_\alpha(q_i)$, $b_i = \sigma_\beta(q_i)$; otherwise $a_i = \sigma_\beta(q_i)$, $b_i = \sigma_\alpha(q_i)$. In any case, $c_i = \sigma(q_i)$. Since the set $\{a, b, a^{-1}, b^{-1}\}$ is invariant under permutations σ_α and σ_β , the equalities $\sigma_\alpha\sigma_\beta = \sigma_\beta\sigma_\alpha = \sigma$ imply that $E_\alpha E_\beta = E_\beta E_\alpha = \pi_{(ab)}$.

Since $\pi_{(ab)}, \pi_{(bc)} \in G(E)$ and the group of permutations on $\{a, b, c\}$ is generated by permutations (ab) and (bc) , it follows from Lemma 3.1 that $G(E)$ contains all transformations of the form π_τ . ■

Lemma 3.3 $D_0 = \pi_{(ac)}E_\alpha = \pi_{(abc)}E_\beta$, $D_1 = \pi_{(abc)}E_\alpha = \pi_{(ac)}E_\beta$.

Proof. Let $\xi = q_1q_2 \dots q_k$ be a word in the alphabet Q_\pm . According to the definitions of automata D and E , the words $E_\alpha(\xi) = a_1a_2 \dots a_k$, $E_\beta(\xi) = b_1b_2 \dots b_k$, $D_0(\xi) = c_1c_2 \dots c_k$, and $D_1(\xi) = d_1d_2 \dots d_k$ can be computed as follows. First we define 4 permutations on the set Q_\pm : $\sigma_\alpha = (a^{-1}b^{-1})$, $\sigma_\beta = (ab)$, $\sigma_0 = (ac)(a^{-1}b^{-1}c^{-1})$, $\sigma_1 = (abc)(a^{-1}c^{-1})$. Now for any i , $1 \leq i \leq k$ we count the number of times when letters a, b, a^{-1}, b^{-1} occur in the sequence q_1, \dots, q_{i-1} . If this number is even then $a_i = \sigma_\alpha(q_i)$, $b_i = \sigma_\beta(q_i)$, $c_i = \sigma_0(q_i)$, $d_i = \sigma_1(q_i)$. Otherwise $a_i = \sigma_\beta(q_i)$, $b_i = \sigma_\alpha(q_i)$, $c_i = \sigma_1(q_i)$, $d_i = \sigma_0(q_i)$.

Let us consider 2 more permutations on the set Q_\pm : $\tau_0 = (ac)(a^{-1}c^{-1})$ and $\tau_1 = (abc)(a^{-1}b^{-1}c^{-1})$. It is easy to verify that $\sigma_0 = \tau_0\sigma_\alpha = \tau_1\sigma_\beta$ and $\sigma_1 = \tau_1\sigma_\alpha = \tau_0\sigma_\beta$. Therefore for any i , $1 \leq i \leq k$ we have $c_i = \tau_0(a_i) = \tau_1(b_i)$ and $d_i = \tau_1(a_i) = \tau_0(b_i)$. This means that $D_0(\xi) = \pi_{(ac)}E_\alpha(\xi) = \pi_{(abc)}E_\beta(\xi)$ and $D_1(\xi) = \pi_{(abc)}E_\alpha(\xi) = \pi_{(ac)}E_\beta(\xi)$. ■

Proposition 3.4 $G(D) = G(E)$.

Proof. Lemmas 3.2 and 3.3 imply that $D_0, D_1 \in G(E)$; therefore $G(D) \subset G(E)$.

By Lemma 3.3, $D_0D_1^{-1} = \pi_{(ac)}E_\alpha(\pi_{(abc)}E_\alpha)^{-1} = \pi_{(ac)}\pi_{(abc)}^{-1}$. By Lemma 3.1, $\pi_{(ac)}\pi_{(abc)}^{-1} = \pi_{(ac)}\pi_{(acb)} = \pi_{(bc)} = E_\gamma$. Similarly,

$$D_0^{-1}D_1 = (\pi_{(ac)}E_\alpha)^{-1}\pi_{(abc)}E_\alpha = E_\alpha^{-1}\pi_{(ac)}^{-1}\pi_{(abc)}E_\alpha = E_\alpha^{-1}\pi_{(ab)}E_\alpha.$$

Lemma 3.2 implies that E_α and $\pi_{(ab)}$ commute, hence $D_0^{-1}D_1 = \pi_{(ab)}$. The group of all permutations on the set $\{a, b, c\}$ is generated by permutations (ab) and (bc) . Since $\pi_{(ab)}, \pi_{(bc)} \in G(D)$, it follows from Lemma 3.1 that $G(D)$ contains all transformations of the form π_τ . Then $E_\alpha, E_\beta \in G(D)$ by Lemma 3.3. Thus $G(E) \subset G(D)$. \blacksquare

4 Patterns and orbits

This section is devoted to the proof of Theorem 1.1. We use the notation of the previous section.

As shown in Section 2, the automaton B defines a right action $X^* \times Q_\pm^* \rightarrow X^*$ of the monoid Q_\pm^* on the rooted binary tree X^* given by $(w, \xi) \mapsto B_\xi(w)$. Since Q_\pm^* is the free monoid generated by $a, b, c, a^{-1}, b^{-1}, c^{-1}$, there exists a unique homomorphism $\chi : Q_\pm^* \rightarrow \{-1, 1\}$ such that $\chi(a) = \chi(b) = \chi(a^{-1}) = \chi(b^{-1}) = -1$, $\chi(c) = \chi(c^{-1}) = 1$.

Lemma 4.1 *Given $\xi \in Q_\pm^*$, the automorphism B_ξ of the rooted binary tree $\{0, 1\}^*$ acts trivially on the first level of the tree (i.e., on one-letter words) if and only if $\chi(\xi) = 1$.*

Proof. For any $\xi \in Q_\pm^*$ let $\tilde{\chi}(\xi) = 1$ if B_ξ acts trivially on the first level of the binary tree X^* and $\tilde{\chi}(\xi) = -1$ otherwise. If $\tilde{\chi}(\xi) = -1$ then $B_\xi(0) = 1$, $B_\xi(1) = 0$. Since $B_{\xi_1\xi_2} = B_{\xi_2}B_{\xi_1}$ for all $\xi_1, \xi_2 \in Q_\pm^*$, it follows that the map $\tilde{\chi} : Q_\pm^* \rightarrow \{-1, 1\}$ is a homomorphism of the monoid Q_\pm^* . By definition, $\tilde{\chi}(a) = \tilde{\chi}(b) = -1$, $\tilde{\chi}(c) = 1$, and $\tilde{\chi}(q^{-1}) = \tilde{\chi}(q)$ for any $q \in \{a, b, c\}$. Since $\tilde{\chi}$ is a homomorphism, it follows that $\tilde{\chi} = \chi$. \blacksquare

Now we introduce an alphabet consisting of two symbols $*$ and $*^{-1}$. A word over the alphabet $\{*, *^{-1}\}$ is called a *pattern*. Every word ξ over the alphabet Q_\pm is assigned a pattern v that is obtained from ξ by substituting $*$ for each occurrence of letters a, b, c and substituting $*^{-1}$ for each occurrence of letters a^{-1}, b^{-1}, c^{-1} . We say that v is the pattern of ξ or that ξ follows the pattern v .

A word $\xi = q_1q_2\dots q_n \in Q_{\pm}^*$ is called *freely irreducible* if none of its two-letter subwords $q_1q_2, q_2q_3, \dots, q_{n-1}q_n$ coincides with one of the following words: $aa^{-1}, bb^{-1}, cc^{-1}, a^{-1}a, b^{-1}b, c^{-1}c$. Otherwise ξ is called *freely reducible*.

Lemma 4.2 *For any nonempty pattern v there exist words $\xi_1, \xi_2 \in Q_{\pm}^*$ such that ξ_1 and ξ_2 are freely irreducible, follow the pattern v , and $\chi(\xi_2) = -\chi(\xi_1)$.*

Proof. Given a nonempty pattern v , let us substitute a for each occurrence of $*$ in v and b^{-1} for each occurrence of $*^{-1}$. We get a word $\xi_1 \in Q_{\pm}^*$ that follows the pattern v . Now let us modify ξ_1 by changing its first letter. If this letter is a , we change it to c . If the first letter of ξ_1 is b^{-1} , we change it to c^{-1} . This yields another word $\xi_2 \in Q_{\pm}^*$ that follows the pattern v . By construction, ξ_1 and ξ_2 are freely irreducible. Since $\chi(a) = \chi(b^{-1}) = -1$ and $\chi(c) = \chi(c^{-1}) = 1$, it follows that $\chi(\xi_2) = -\chi(\xi_1)$. ■

The following proposition is the main step in the proof of Theorem 1.1.

Proposition 4.3 *Suppose $\xi \in Q_{\pm}^*$ is a freely irreducible word. Then the orbit of ξ under the action of the group $G(E)$ on Q_{\pm}^* consists of all freely irreducible words following the same pattern as ξ .*

To prove Proposition 4.3, we need several lemmas.

Lemma 4.4 *Two words $\xi_1, \xi_2 \in Q_{\pm}^*$ are in the same orbit of the $G(E)$ action if and only if the reversed words $\overleftarrow{\xi_1}$ and $\overleftarrow{\xi_2}$ are in the same orbit of this action.*

Proof. Using Proposition 2.3, we verify that the reverse automaton R of E is well defined. The Moore diagram Γ_R of R is obtained from the Moore diagram Γ_E of E by reversing all edges. It is easy to observe that Γ_R can also be obtained by renaming vertices α and β of the graph Γ_E to β and α , respectively (see Figure 4). Hence $R_{\alpha} = E_{\beta}$, $R_{\beta} = E_{\alpha}$, $R_{\gamma} = E_{\gamma}$. In particular, $S(R) = S(E)$. By Lemma 3.2, E_{α} , E_{β} , and E_{γ} are involutions; therefore $S(E) = G(E)$. Now it follows from Proposition 2.3 that, given $\xi_1, \xi_2 \in Q_{\pm}^*$, we have $\xi_2 = g(\xi_1)$ for some $g \in G(E)$ if and only if $\overleftarrow{\xi_2} = h(\overleftarrow{\xi_1})$ for some $h \in G(E)$. ■

Let V^0 denote the set of patterns without double letters. That is, a pattern $v \in V^0$ does not contain subwords $**$ and $*^{-1}*^{-1}$. We consider 4 subsets of V^0 . By V_{++}^0 denote the set of $v \in V^0$ such that $*$ is the first and

the last letter of v . By V_{+-}^0 denote the set of $v \in V^0$ such that either $v = \emptyset$ or the first letter of v is $*$ while the last letter is $*^{-1}$. By V_{-+}^0 denote the set of $v \in V^0$ such that either $v = \emptyset$ or the first letter of v is $*^{-1}$ while the last letter is $*$. By V_{--}^0 denote the set of $v \in V^0$ such that $*^{-1}$ is the first and the last letter of v . Clearly, $V^0 = V_{++}^0 \cup V_{+-}^0 \cup V_{-+}^0 \cup V_{--}^0$.

Now we define W_{++}^0 (resp. $W_{+-}^0, W_{-+}^0, W_{--}^0$) as the set of all words over the alphabet $\{a, b, a^{-1}, b^{-1}\}$ that follow patterns from the set V_{++}^0 (resp. $V_{+-}^0, V_{-+}^0, V_{--}^0$). In particular, W_{++}^0 contains the words $a, ab^{-1}a, ab^{-1}ab^{-1}a, \dots$, W_{+-}^0 contains $\emptyset, ab^{-1}, ab^{-1}ab^{-1}, \dots$, W_{-+}^0 contains $\emptyset, b^{-1}a, b^{-1}ab^{-1}a, \dots$, and W_{--}^0 contains $b^{-1}, b^{-1}ab^{-1}, b^{-1}ab^{-1}ab^{-1}, \dots$.

Consider an endomorphism r of the free monoid Q_{\pm}^* defined by $r(a) = a, r(a^{-1}) = a^{-1}, r(b) = b, r(b^{-1}) = b^{-1}, r(c) = r(c^{-1}) = \emptyset$. For any $\xi \in Q_{\pm}^*$ the word $r(\xi)$ is obtained by deleting all letters c and c^{-1} in ξ . The restriction of r to the set $W_{++}^0 \cup W_{+-}^0 \cup W_{-+}^0 \cup W_{--}^0$ is the identity map. Let $W_{++} = r^{-1}(W_{++}^0), W_{+-} = r^{-1}(W_{+-}^0), W_{-+} = r^{-1}(W_{-+}^0),$ and $W_{--} = r^{-1}(W_{--}^0)$.

Lemma 4.5 (i) $E_{\alpha}(\xi) = \xi$ for all $\xi \in W_{++} \cup W_{+-}$ while $E_{\beta}(\xi) = \xi$ for all $\xi \in W_{-+} \cup W_{--}$.

(ii) If $\xi \in W_{++}$ then $E_{\alpha}(\xi a) = \xi b$ and $E_{\alpha}(\xi b) = \xi a$. If $\xi \in W_{+-}$ then $E_{\alpha}(\xi a^{-1}) = \xi b^{-1}$ and $E_{\alpha}(\xi b^{-1}) = \xi a^{-1}$. If $\xi \in W_{-+}$ then $E_{\beta}(\xi a) = \xi b$ and $E_{\beta}(\xi b) = \xi a$. If $\xi \in W_{--}$ then $E_{\beta}(\xi a^{-1}) = \xi b^{-1}$ and $E_{\beta}(\xi b^{-1}) = \xi a^{-1}$.

Proof. Let $\xi \in Q_{\pm}^*$. If $x \in \{c, c^{-1}\}$ then $E_{\alpha}(x\xi) = xE_{\alpha}(\xi)$ and $E_{\beta}(x\xi) = xE_{\beta}(\xi)$. Besides, ξ belongs to one of the sets $W_{++}, W_{+-}, W_{-+}, W_{--}$ if and only if $x\xi$ belongs to the same set. Further, if $x \in \{a, b\}$ then $E_{\alpha}(x\xi) = xE_{\beta}(\xi)$. In this case, $x\xi \in W_{++} \cup W_{+-}$ if and only if $\xi \in W_{-+} \cup W_{--}$. Finally, if $x \in \{a^{-1}, b^{-1}\}$ then $E_{\beta}(x\xi) = xE_{\alpha}(\xi)$. In this case, $x\xi \in W_{-+} \cup W_{--}$ if and only if $\xi \in W_{++} \cup W_{+-}$. The statement (i) of the lemma follows from the above by induction on the length of ξ .

Any word $\xi \in W_{++}$ contains an odd number of letters a, b, a^{-1}, b^{-1} . Therefore $E_{\alpha}(\xi\eta) = E_{\alpha}(\xi)E_{\beta}(\eta) = \xi E_{\beta}(\eta)$ for all $\eta \in Q_{\pm}^*$. In particular, $E_{\alpha}(\xi a) = \xi b$ and $E_{\alpha}(\xi b) = \xi a$. The rest of the statement (ii) is obtained in a similar way. We omit the details. \blacksquare

Given a freely irreducible word $\xi \in Q_{\pm}^*$, let $Z(\xi)$ denote the set of freely irreducible words in Q_{\pm}^* that follow the same pattern as ξ and match ξ completely or except for the last letter. Obviously, $\xi \in Z(\xi)$, and $\eta \in Z(\xi)$ if and only if $\xi \in Z(\eta)$. If $\xi \neq \emptyset$ then $Z(\xi)$ consists of 2 or 3 words. Namely, there are exactly 3 words in Q_{\pm}^* that follow the same pattern as ξ and match

ξ completely or except for the last letter. However if the last two letters in the pattern of ξ are distinct then one of these 3 words is freely reducible.

Lemma 4.6 *Suppose v is a pattern of length at least 2 such that the last two letters of v are distinct. Then there exist two freely irreducible words $\xi_a, \xi_b \in Q_{\pm}^*$ such that ξ_a and ξ_b follow the pattern v , $Z(\xi_a) = \{\xi_a, \xi_b\}$, and $\xi_b = g(\xi_a)$, where $g \in \{E_{\alpha}, E_{\beta}\}$.*

Proof. Let v_0 denote the pattern obtained by deleting the last letter of v . We replace each letter $*$ in the word v_0 by a if the next letter is $*^{-1}$ and by c otherwise. Each letter $*^{-1}$ is replaced by b^{-1} if the next letter is $*$ and by c^{-1} otherwise. We get a word $\eta \in Q_{\pm}^*$ that follows the pattern v_0 . For example, if $v_0 = ****^{-1}*^{-1}*^{-1}****^{-1}*^{-1}*^{-1}$ then $\eta = ccac^{-1}c^{-1}b^{-1}ccac^{-1}c^{-1}c^{-1}$. Now let $\xi_a = \eta a$, $\xi_b = \eta b$ if $v = v_0*$ and let $\xi_a = \eta a^{-1}$, $\xi_b = \eta b^{-1}$ if $v = v_0*^{-1}$. The words ξ_a and ξ_b follow the pattern v . By construction, η is freely irreducible and its last letter is c or c^{-1} . Since the last letters of patterns v and v_0 are different, it follows that ξ_a, ξ_b are freely irreducible and $Z(\xi_a) = \{\xi_a, \xi_b\}$.

It is easy to observe that η belongs to one of the sets $W_{++}, W_{+-}, W_{-+}, W_{--}$. Lemma 4.5 implies that $g(\xi_a) = \xi_b$ and $g(\xi_b) = \xi_a$, where $g \in \{E_{\alpha}, E_{\beta}\}$. Namely, $g = E_{\alpha}$ if the first letter of v is $*$ and $g = E_{\beta}$ otherwise. \blacksquare

Lemma 4.7 *Suppose v is a pattern of length at least 2 such that the first two letters of v coincide as well as the last two letters of v . Then there exist 9 freely irreducible words $\xi_{q_1 q_2} \in Q_{\pm}^*$, $q_1, q_2 \in \{a, b, c\}$ such that each $\xi_{q_1 q_2}$ follow the pattern v and for any $q \in \{a, b, c\}$ we have $Z(\xi_{qc}) = \{\xi_{qa}, \xi_{qb}, \xi_{qc}\}$, $Z(\xi_{cq}) = \{\xi_{aq}, \xi_{bq}, \xi_{cq}\}$, and $\xi_{qb} = g_q(\xi_{qa})$, where $g_q \in \{E_{\alpha}, E_{\beta}\}$.*

Proof. Let v_0 denote the pattern obtained by deleting the first and the last letters of v . We replace each letter $*$ in the word v_0 by a if the previous letter is $*^{-1}$ and by c otherwise. Each letter $*^{-1}$ is replaced by b^{-1} if the previous letter is $*$ and by c^{-1} otherwise. We get a word $\eta \in Q_{\pm}^*$ that follows the pattern v_0 . For example, if $v_0 = *** *^{-1} *^{-1} *^{-1} * * * *^{-1} *^{-1} *^{-1}$ then $\eta = cccb^{-1}c^{-1}c^{-1}accb^{-1}c^{-1}c^{-1}$. Now for any $q \in \{a, b, c\}$ let $\xi_q = q\eta$ if the first letter of v is $*$ and let $\xi_q = q^{-1}\eta$ otherwise. Further, for any $q_1, q_2 \in \{a, b, c\}$ let $\xi_{q_1 q_2} = \xi_{q_1} q_2$ if the last letter of v is $*$ and let $\xi_{q_1 q_2} = \xi_{q_1} q_2^{-1}$ otherwise. Each $\xi_{q_1 q_2}$ follows the pattern v . By construction, η is freely irreducible. Since the first two letters of v coincide and so do the last two letters, it

follows that each $\xi_{q_1 q_2}$ is freely irreducible. Then $Z(\xi_{qc}) = \{\xi_{qa}, \xi_{qb}, \xi_{qc}\}$ and $Z(\xi_{cq}) = \{\overset{\leftarrow}{\xi}_{aq}, \overset{\leftarrow}{\xi}_{bq}, \overset{\leftarrow}{\xi}_{cq}\}$ for any $q \in \{a, b, c\}$.

It is easy to observe that the words ξ_a and ξ_b belong to one of the sets $W_{++} \cup W_{+-}$ and $W_{-+} \cup W_{--}$ while η and ξ_c belong to the other. Lemma 4.5 implies that $g(\xi_{aa}) = \xi_{ab}$, $g(\xi_{ab}) = \xi_{aa}$, $g(\xi_{ba}) = \xi_{bb}$, and $g(\xi_{bb}) = \xi_{ba}$, where $g = E_\alpha$ if the first letter of v is $*$ and $g = E_\beta$ otherwise. Moreover, $h(\xi_{ca}) = \xi_{cb}$ and $h(\xi_{cb}) = \xi_{ca}$, where h is the element of $\{E_\alpha, E_\beta\}$ different from g . \blacksquare

Proof of Proposition 4.3. First we shall show that the $G(E)$ action on Q_\pm^* preserves patterns and free irreducibility of words. Suppose $q_1|q_2$ is the label assigned to an edge of the Moore diagram of the automaton E . Then either $q_1, q_2 \in \{a, b, c\}$ or $q_1, q_2 \in \{a^{-1}, b^{-1}, c^{-1}\}$ (see Figure 4). It follows that transformations E_α , E_β , and E_γ preserve patterns of words. So does any $g \in G(E)$. Further, it is easy to verify that sets $P_1 = \{aa^{-1}, bb^{-1}, cc^{-1}\}$ and $P_2 = \{a^{-1}a, b^{-1}b, c^{-1}c\}$ are invariant under transformations E_α , E_β , and E_γ . Therefore P_1 and P_2 are invariant under the $G(E)$ action on Q_\pm^* . Any freely reducible word $\xi \in Q_\pm^*$ is represented as $\xi_1 \xi_0 \xi_2$, where $\xi_0 \in P_1 \cup P_2$ and $\xi_1, \xi_2 \in Q_\pm^*$. Proposition 2.4 implies that for any $g \in S(E) = G(E)$ we have $g(\xi) = g(\xi_1)g_0(\xi_0)g_1(\xi_2)$, where $g_0, g_1 \in G(E)$. By the above $g(\xi)$ is freely reducible. Thus the $G(E)$ action preserves free reducibility of words. Since $G(E)$ is a group, its action on Q_\pm^* preserves free irreducibility as well.

Now we are going to prove that for any freely irreducible words $\xi_1, \xi_2 \in Q_\pm^*$ following the same pattern v there exists $g \in G(E)$ such that $\xi_2 = g(\xi_1)$. The claim is proved by induction on the length of the pattern v . The empty pattern is followed only by the empty word. As for one-letter patterns, it is enough to notice that $\pi_{(abc)}(a) = b$, $\pi_{(abc)}(b) = c$, $\pi_{(abc)}(c) = a$, $\pi_{(abc)}(a^{-1}) = b^{-1}$, $\pi_{(abc)}(b^{-1}) = c^{-1}$, $\pi_{(abc)}(c^{-1}) = a^{-1}$, and $\pi_{(abc)} \in G(E)$ by Lemma 3.2. Now let $n \geq 2$ and assume that the claim holds for all patterns of length less than n . Take any pattern v of length n . First consider the case when the last two letters of v are distinct. By Lemma 4.6, there exists a freely irreducible word $\xi \in Q_\pm^*$ such that ξ follows the pattern v and the set $Z(\xi)$ is contained in an orbit of the $G(E)$ action. Secondly, consider the case when the first two letters of v are distinct. Here, the last two letters of the reversed pattern \overleftarrow{v} are distinct. Then there is a freely irreducible word ξ_- such that ξ_- follows the pattern \overleftarrow{v} and $Z(\xi_-)$ is contained in an orbit of the $G(E)$ action. Clearly, $\xi_- = \overset{\leftarrow}{\xi}_+$, where ξ_+ is a freely irreducible word following the pattern v .

Now consider the case when the last two letters of v coincide as well as the first two letters. Let $\xi_{q_1q_2}$, $q_1, q_2 \in \{a, b, c\}$ be the words provided by Lemma 4.7. By ξ_a, ξ_b, ξ_c denote the words obtained by deleting the last letter of $\xi_{aa}, \xi_{ba}, \xi_{ca}$, respectively. Obviously, ξ_a, ξ_b , and ξ_c are freely irreducible and follow the same pattern of length $n - 1$. By the inductive assumption there exist $h_a, h_b \in G(E)$ such that $h_a(\xi_c) = \xi_a$, $h_b(\xi_c) = \xi_b$. Since the $G(E)$ action preserves patterns, it follows that $h_a(\xi_{cc}) \in Z(\xi_{ac})$, $h_b(\xi_{cc}) \in Z(\xi_{bc})$. Recall that there are involutions $g_a, g_b, g_c \in G(E)$ such that $g_q(\xi_{qa}) = \xi_{qb}$, $q \in \{a, b, c\}$. If $h_a(\xi_{cc}) = \xi_{aa}$ or $h_a(\xi_{cc}) = \xi_{ab}$, then the word $h_a^{-1}g_a h_a(\xi_{cc})$ matches ξ_{cc} except for the last letter. Hence $h_a^{-1}g_a h_a(\xi_{cc}) \in \{\xi_{ca}, \xi_{cb}\}$. Since $g_c(\xi_{ca}) = \xi_{cb}$, it follows that $Z(\xi_{cc}) = \{\xi_{ca}, \xi_{cb}, \xi_{cc}\}$ is contained in an orbit of the $G(E)$ action. If $h_b(\xi_{cc}) = \xi_{ba}$ or $h_b(\xi_{cc}) = \xi_{bb}$, we reach the same conclusion. On the other hand, if $h_a(\xi_{cc}) = \xi_{ac}$ and $h_b(\xi_{cc}) = \xi_{bc}$ then the set $\{\xi_{ac}, \xi_{bc}, \xi_{cc}\}$ is contained in an orbit. By Lemma 4.4, $Z(\xi_{cc}) = \{\overleftarrow{\xi_{ac}}, \overleftarrow{\xi_{bc}}, \overleftarrow{\xi_{cc}}\}$ is also contained in an orbit of the $G(E)$ action.

In any of the considered cases, there exists a freely irreducible word ξ such that ξ follows the pattern v and one of the sets $Z(\xi)$, $Z(\overleftarrow{\xi})$ is contained in an orbit of the $G(E)$ action. Take any freely irreducible words ξ_1, ξ_2 following the pattern v . First suppose that $Z(\xi)$ is contained in an orbit. Let η, η_1, η_2 be words obtained by deleting the last letter of ξ, ξ_1, ξ_2 , respectively. Then η, η_1, η_2 are freely irreducible and follow the same pattern obtained by deleting the last letter of v . By the inductive assumption there are $g_1, g_2 \in G(E)$ such that $\eta = g_1(\eta_1) = g_2(\eta_2)$. Since the $G(E)$ action preserves patterns and free irreducibility, it follows that $g_1(\xi_1), g_2(\xi_2) \in Z(\xi)$. As $Z(\xi)$ is contained in an orbit, there exists $g \in G(E)$ such that $g(g_1(\xi_1)) = g_2(\xi_2)$. Then $\xi_2 = g_2^{-1}g g_1(\xi_1)$. Now suppose that the set $Z(\overleftarrow{\xi})$ is contained in an orbit. Let ζ, ζ_1, ζ_2 be words obtained by deleting the last letter of $\overleftarrow{\xi}, \overleftarrow{\xi_1}, \overleftarrow{\xi_2}$, respectively. Clearly, ζ, ζ_1, ζ_2 are freely irreducible and follow the same pattern obtained by deleting the last letter of \overleftarrow{v} . By the inductive assumption there exist $h_1, h_2 \in G(E)$ such that $\zeta = h_1(\zeta_1) = h_2(\zeta_2)$. Since the $G(E)$ action preserves patterns and free irreducibility, we conclude that $h_1(\overleftarrow{\xi_1}), h_2(\overleftarrow{\xi_2}) \in Z(\overleftarrow{\xi})$. It follows that $\overleftarrow{\xi_1}$ and $\overleftarrow{\xi_2}$ are in the same orbit of the $G(E)$ action. By Lemma 4.4, ξ_1 and ξ_2 are also in the same orbit. The induction step is complete. \blacksquare

Proof of Theorem 1.1. The group $G(A)$ is the free non-Abelian group on generators A_a, A_b, A_c if and only if $A_{q_1}^{m_1} A_{q_2}^{m_2} \dots A_{q_n}^{m_n} \neq 1$ for any pair

of sequences q_1, \dots, q_n and m_1, \dots, m_n such that $n > 0$, $q_i \in \{a, b, c\}$ and $m_i \in \mathbb{Z} \setminus \{0\}$ for $1 \leq i \leq n$, and $q_i \neq q_{i+1}$ for $1 \leq i \leq n-1$. Since $B_a = A_a$, $B_b = A_b$, $B_c = A_c$, $B_{a^{-1}} = A_a^{-1}$, $B_{b^{-1}} = A_b^{-1}$, $B_{c^{-1}} = A_c^{-1}$, an equivalent condition is that $B_\xi \neq 1$ for any nonempty freely irreducible word $\xi \in Q_\pm^*$.

Suppose $B_\xi = 1$ for some freely irreducible word $\xi \in Q_\pm^*$. By Corollary 2.5, $B_{g(\xi)} = 1$ for all $g \in S(D)$. Then Proposition 2.1 imply that $B_{g(\xi)} = 1$ for all $g \in G(D)$. By Proposition 3.4, $G(D) = G(E)$. Now it follows from Proposition 4.3 that $B_\eta = 1$ for any freely irreducible word $\eta \in Q_\pm^*$ following the same pattern as ξ . In particular, B_η acts trivially on the first level of the binary tree $\{0, 1\}^*$. Finally, Lemmas 4.1 and 4.2 imply that ξ follows the empty pattern. Then ξ itself is the empty word. \blacksquare

References

- [A] S. V. Aleshin. A free group of finite automata. *Mosc. Univ. Math. Bull.* **38** (1983), no. 4, 10–13.
- [BS] A. M. Brunner, S. Sidki. The generation of $GL(n, \mathbb{Z})$ by finite state automata. *Int. J. Algebra Comput.* **8** (1998), no. 1, 127–139.
- [GM] Y. Glasner, S. Mozes. Automata and square complexes. *Geom. Dedicata* **111** (2005), 43–64.
- [GNS] R. I. Grigorchuk, V. V. Nekrashevich, V. I. Sushchanskii. Automata, dynamical systems, and groups. *Grigorchuk R. I. (ed.), Dynamical systems, automata, and infinite groups*. Proc. Steklov Inst. Math. **231** (2000), 128–203.
- [Nek] V. Nekrashevych. *Self-similar groups*. Math. Surveys and Monographs **117**. Amer. Math. Soc., Providence, RI, 2005.
- [OS] A. S. Olijnyk, V. I. Sushchanskij. Free group of infinite unitriangular matrices. *Math. Notes* **67** (2000), no. 3, 320–324.
- [S] S. Sidki. Automorphisms of one-rooted trees: growth, circuit structure, and acyclicity. *J. Math. Sci., NY* **100** (2000), no. 1, 1925–1943.

DEPARTMENT OF MATHEMATICS
 TEXAS A&M UNIVERSITY
 COLLEGE STATION, TX 77843–3368