

ON A HYPERGRAPH TURÁN PROBLEM OF FRANKL

PETER KEEVASH, BENNY SUDAKOV*

Received February 16, 2003

Let $\mathcal{C}_r^{(2k)}$ be the $2k$ -uniform hypergraph obtained by letting P_1, \dots, P_r be pairwise disjoint sets of size k and taking as edges all sets $P_i \cup P_j$ with $i \neq j$. This can be thought of as the ‘ k -expansion’ of the complete graph K_r : each vertex has been replaced with a set of size k . An example of a hypergraph with vertex set V that does not contain $\mathcal{C}_3^{(2k)}$ can be obtained by partitioning $V = V_1 \cup V_2$ and taking as edges all sets of size $2k$ that intersect each of V_1 and V_2 in an odd number of elements. Let $\mathcal{B}_n^{(2k)}$ denote a hypergraph on n vertices obtained by this construction that has as many edges as possible. For n sufficiently large we prove a conjecture of Frankl, which states that any hypergraph on n vertices that contains no $\mathcal{C}_3^{(2k)}$ has at most as many edges as $\mathcal{B}_n^{(2k)}$.

Sidorenko has given an upper bound of $\frac{r-2}{r-1}$ for the Turán density of $\mathcal{C}_r^{(2k)}$ for any r , and a construction establishing a matching lower bound when r is of the form $2^p + 1$. In this paper we also show that when $r = 2^p + 1$, any $\mathcal{C}_r^{(4)}$ -free hypergraph of density $\frac{r-2}{r-1} - o(1)$ looks approximately like Sidorenko’s construction. On the other hand, when r is not of this form, we show that corresponding constructions do not exist and improve the upper bound on the Turán density of $\mathcal{C}_r^{(4)}$ to $\frac{r-2}{r-1} - c(r)$, where $c(r)$ is a constant depending only on r .

The backbone of our arguments is a strategy of first proving approximate structure theorems, and then showing that any imperfections in the structure must lead to a suboptimal configuration. The tools for its realisation draw on extremal graph theory, linear algebra, the Kruskal–Katona theorem and properties of Krawtchouk polynomials.

Mathematics Subject Classification (2000): 05C35, 05C65, 05D05, 05E35

* Research supported in part by NSF grants DMS-0355497, DMS-0106589, and by an Alfred P. Sloan fellowship.

1. Introduction

Given an r -uniform hypergraph \mathcal{F} , the Turán number $ex(n, \mathcal{F})$ of \mathcal{F} is the maximum number of edges in an r -uniform hypergraph on n vertices that does not contain a copy of \mathcal{F} . Determining these numbers is one of the main challenges in Extremal Combinatorics. For ordinary graphs (the case $r=2$) a rich theory has been developed, initiated by Turán in 1941, who solved the problem for complete graphs. He also posed the question of finding $ex(n, \mathcal{K}_s^{(r)})$ for complete hypergraphs with $s > r > 2$, but to this day not one single instance of this problem has been solved. It seems hard even to determine the *Turán density*, which for general \mathcal{F} is defined as $\pi(\mathcal{F}) = \lim_{n \rightarrow \infty} ex(n, \mathcal{F}) / \binom{n}{r}$. The problem of finding the numbers $ex(n, \mathcal{F})$ when $r > 2$ is notoriously difficult, and exact results on hypergraph Turán numbers are very rare (see [3, 9] for surveys). In this paper we obtain such a result for a sequence of hypergraphs introduced by Frankl.

Let $\mathcal{C}_r^{(2k)}$ be the $2k$ -uniform hypergraph obtained by letting P_1, \dots, P_r be pairwise disjoint sets of size k and taking as edges all sets $P_i \cup P_j$ with $i \neq j$. This can be thought of as the ‘ k -expansion’ of the complete graph K_r : each vertex has been replaced with a set of size k . The Turán problem for $\mathcal{C}_3^{(2k)}$ was first considered by Frankl [2], who determined the density $\pi(\mathcal{C}_3^{(2k)}) = 1/2$.

Frankl obtained a large $\mathcal{C}_3^{(2k)}$ -free hypergraph on n vertices by partitioning an n -element set V into 2 parts V_1, V_2 and taking those edges which intersect each part V_i in an odd number of elements. When the parts have sizes $\frac{n}{2} \pm t$ we denote this hypergraph by $\mathcal{B}^{(2k)}(n, t)$. To see that it is $\mathcal{C}_3^{(2k)}$ -free, consider any P_1, P_2, P_3 that are pairwise disjoint sets of k vertices. Then $|V_1 \cap P_i|$ and $|V_1 \cap P_j|$ have the same parity for some pair ij , so $P_i \cup P_j$ is not an edge. Let t^* be chosen to maximise the number of edges in $\mathcal{B}^{(2k)}(n, t)$, and denote any hypergraph obtained in this manner by $\mathcal{B}_n^{(2k)}$. Write $b_{2k}(n)$ for the number of edges in $\mathcal{B}_n^{(2k)}$. Frankl [2] conjectured that the maximum number of edges in a $\mathcal{C}_3^{(2k)}$ -free hypergraph is always achieved by some $\mathcal{B}_n^{(2k)}$. Our first theorem proves this conjecture for n sufficiently large.

Theorem 1.1. *Let H be a $2k$ -uniform hypergraph on n vertices that does not contain a copy of $\mathcal{C}_3^{(2k)}$ and let n be sufficiently large. Then the number of edges in H is at most $b_{2k}(n)$, with equality only when H is a hypergraph of the form $\mathcal{B}_n^{(2k)}$.*

The proof of this theorem falls naturally into two parts. The first stage is to prove a ‘stability’ version, which is that any hypergraph with close to the

maximum number of edges looks approximately like some $\mathcal{B}^{(2k)}(n, t)$. Armed with this, we can analyse any imperfections in the structure and show that they must lead to a suboptimal configuration, so that the optimum is indeed achieved by the construction. This strategy was also used recently in [4] to prove the conjecture of Sós on the Turán number of the Fano plane, so this seems to be a useful tool for developing the Turán theory of hypergraphs.

For general r , Sidorenko [8] showed that the Turán density of $\mathcal{C}_r^{(2k)}$ is at most $\frac{r-2}{r-1}$. This is a consequence of Turán’s theorem applied to an auxiliary graph G constructed from a $2k$ -uniform hypergraph H ; the vertices of G are the k -tuples of vertices of H , and two k -tuples P_1, P_2 are adjacent if $P_1 \cup P_2$ is an edge of H . He also gave a construction for a matching lower bound when r is of the form $2^p + 1$, which we now describe. Let W be a vector space of dimension p over the field $GF(2)$, i.e. the finite field with 2 elements $\{0, 1\}$. Partition a set of vertices V as $\bigcup_{w \in W} V_w$. Given t and a t -tuple of vertices $X = x_1 \cdots x_t$ with $x_i \in V_{w_i}$ let $\Sigma X = \sum_1^t w_i$. Define a $2k$ -uniform hypergraph H , where a $2k$ -tuple X is an edge iff $\Sigma X \neq 0$. Observe that this doesn’t contain a copy of $\mathcal{C}_r^{(2k)}$. Indeed, if P_1, \dots, P_r are disjoint k -tuples then there is some $i \neq j$ with $\Sigma P_i = \Sigma P_j$ (by the pigeonhole principle). Then $\Sigma(P_i \cup P_j) = \Sigma P_i + \Sigma P_j = 0$, so $P_i \cup P_j$ is not an edge. To see that this construction can achieve the stated Turán density, choose the partition so that $|V_w| = |V|/(r-1)$. Then a random (average) $2k$ -tuple is an edge with probability $\frac{r-2}{r-1} + o(1)$, as can be seen by conditioning on the positions of all but one element.

This construction depends essentially on an algebraic structure, which only exists for certain values of r . We will show that this is an intrinsic feature of the problem, by proving a stronger upper bound on the Turán density of $\mathcal{C}_r^{(4)}$ when r is not of the form $2^p + 1$.

Theorem 1.2. *Suppose $r \geq 3$, and let H be a 4-uniform hypergraph on n vertices with at least $(\frac{r-2}{r-1} - 10^{-33}r^{-70})\binom{n}{4}$ edges. If H is $\mathcal{C}_r^{(4)}$ -free, then $r = 2^p + 1$ for some integer p .*

In contrast to Theorem 1.1 this is a result showing that certain constructions do not exist, so it is perhaps surprising that its proof also uses a stability argument. We study the properties of a $\mathcal{C}_r^{(4)}$ -free hypergraph with density close to $\frac{r-2}{r-1}$ and show that it gives rise to an edge colouring of the complete graph K_{r-1} with special properties. Next we prove that for such an edge-colouring there is a natural $GF(2)$ vector space structure on the colours. Of course, such a space has cardinality 2^p , for some p , so we get a contradiction unless $r = 2^p + 1$.

A complication arising in [Theorem 1.1](#) is that the optimum construction is not achieved by a partition into two equal parts. Finding t to maximise the number of edges in $\mathcal{B}^{(2k)}(n, t)$ is an interesting problem in enumerative combinatorics, equivalent to finding the minima of binary Krawtchouk polynomials. This is a family of polynomials orthogonal with respect to the uniform measure on a n -dimensional cube that play an important rôle in the analysis of binary Hamming association schemes (see, e.g., [5]). Despite some uncertainty in the location of their minima, the known bounds are sufficient for us to show that some $\mathcal{B}^{(2k)}(n, t)$ must be optimal.

In the case $k = 2$ one can compute the size of $\mathcal{B}^{(2k)}(n)$ precisely, and there are considerable simplifications of the argument, so in the [next section](#) for illustrative purposes we start by giving a separate proof for this case. [Section 3](#) contains a stability theorem for $\mathcal{C}_3^{(2k)}$ and the general case of [Theorem 1.1](#). Then in [Section 4](#) we prove a stability result for $\mathcal{C}_r^{(4)}$ for all r , and use it to establish [Theorem 1.2](#). The [final section](#) of the paper contains some concluding remarks.

We will assume throughout this paper that n is sufficiently large.

2. The Turán number of $\mathcal{C}_3^{(4)}$

We start by proving Frankl's conjecture for 4-uniform hypergraphs. This will serve to illustrate our method, as it has fewer complications than the general case. In addition, in this case it is easy to compute the Turán numbers of $\mathcal{C}_3^{(4)}$ precisely.

We recall that $\mathcal{C}_3^{(4)}$ is the 4-uniform hypergraph with three edges $\{abcd, abef, cdef\}$. We can obtain a large $\mathcal{C}_3^{(4)}$ -free graph on n vertices by partitioning an n -element set into 2 parts and taking those edges which have 1 point in either class and 3 points in the other. To see this, think of an edge as being the union of 2 different *types* of pairs of vertices: one type consisting of pairs with both vertices in one class, the other consisting of pairs that have one point of each class. Given any 3 pairs there are 2 of the same type, and these do not form an edge in the construction.

To maximise the number of edges in this bipartite construction, it is *not* the case that the two parts have sizes as equal as possible, but we will see that the difference in the sizes should be at most of order \sqrt{n} . Let $\mathcal{B}(n, t)$ denote the 4-uniform hypergraph obtained by partitioning an n -element set into 2 parts with sizes $\frac{n}{2} + t$ and $\frac{n}{2} - t$, and taking those edges which have 1 point in either class and 3 points in the other. Let $b(n, t)$ be the number of edges in $\mathcal{B}(n, t)$ and let $d(n, t)$ be the degree of any vertex belonging to

the side with size $\frac{n}{2} + t$. Then the vertices on the side with size $\frac{n}{2} - t$ have degree $d(n, -t)$. We will start with some estimates on these parameters. By definition,

$$\begin{aligned}
 b(n, t) &= \binom{\frac{n}{2} + t}{3} + \binom{\frac{n}{2} - t}{3} \\
 &= \frac{n^4 - 6n^3 + 8n^2 - 16t^4 - 32t^2 + 24t^2n}{48} \\
 (1) \quad &= \frac{1}{48} \left((n^2 - 3n + 4)^2 - (4t^2 - 3n + 4)^2 \right).
 \end{aligned}$$

Thus to maximise $b(n, t)$ we should pick a value of t that minimises $4t^2 - 3n + 4$, subject to the restriction that when n is even t has to be an integer, and when n is odd $t + \frac{1}{2}$ has to be an integer. Let \mathcal{B}_n denote a hypergraph $\mathcal{B}(n, t^*)$, where t^* is such a value of t . By symmetry we can take $t^* > 0$. There is usually a unique best choice of t^* , but for some n there are 2 equal choices of t^* . Note that for any best choice we certainly have $|t^* - \sqrt{3n/4 - 1}| \leq 1/2$.

Let $b(n)$ be the number of edges in \mathcal{B}_n . Then

$$|48b(n) - (n^2 - 3n + 4)^2| = |4(t^*)^2 - 3n + 4|^2 < 50n.$$

It will be useful later to consider the following estimate which follows immediately from the last inequality for sufficiently large n

$$(2) \quad b(n) - b(n - 1) > \frac{1}{12}n^3 - \frac{1}{2}n^2.$$

Next we give an explicit formula for the degrees in $\mathcal{B}(n, t)$

$$\begin{aligned}
 d(n, t) &= \binom{\frac{n}{2} - t}{2} \binom{\frac{n}{2} + t - 1}{2} + \binom{\frac{n}{2} - t}{3} \\
 (3) \quad &= \frac{n^3 - 6n^2 + 8n + 12t^2}{12} + \frac{6tn - 8t^3 - 16t}{12}.
 \end{aligned}$$

We finish these calculations with an upper bound on the maximum degree of \mathcal{B}_n

$$\begin{aligned}
 \Delta(n) &= \frac{1}{12} (n^3 - 6n^2 + 8n + 12(t^*)^2) + \frac{1}{12} |6t^*n - 8(t^*)^3 - 16t^*| \\
 (4) \quad &< \frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2}.
 \end{aligned}$$

The first step in the proof is to show that any $\mathcal{C}_3^{(4)}$ -free 4-uniform hypergraph H with density close to $1/2$ has the correct approximate structure. To do so we need a few definitions. If we have a partition of the vertex set of H as $V(H) = V_1 \cup V_2$ we call a 4-tuple of vertices *good* if it has either 1 point in V_1 and 3 points in V_2 or 1 point in V_2 and 3 points in V_1 ; otherwise we call it *bad*. With respect to H , we call a 4-tuple *correct* if it is either a good edge or a bad non-edge; otherwise we call it *incorrect*. We obtain the following stability result.

Theorem 2.1. *For every $\epsilon > 0$ there is $\eta > 0$ so that if H is a $\mathcal{C}_3^{(4)}$ -free 4-uniform hypergraph with $e(H) > b(n) - \eta n^4$ then there is a partition of the vertex set as $V(H) = V_1 \cup V_2$ such that all but ϵn^4 4-tuples are correct.*

In the proof of this result we need a special case of the Simonovits stability theorem [10] for graphs, which we recall. It states that for every $\epsilon' > 0$ there is $\eta' > 0$ such that if G is a triangle free graph on N vertices with at least $(1 - \eta') \binom{N}{2} / 2$ edges then there is a partition of the vertex set as $V(G) = U_1 \cup U_2$ with $e_G(U_1) + e_G(U_2) < \epsilon' N^2$.

Proof of Theorem 2.1. Define an auxiliary graph G whose vertices are all pairs of vertices of H , and where the pairs ab and cd are adjacent exactly when $abcd$ is an edge of H . Since H is $\mathcal{C}_3^{(4)}$ -free we see that G is triangle-free. Also, each edge of H creates exactly 3 edges in G (corresponding to the 3 ways of breaking a 4-tuple into pairs) so

$$e(G) > 3(b(n) - \eta n^4) > (1 - 50\eta) \frac{1}{2} \binom{\binom{n}{2}}{2}.$$

Choose η so that Simonovits stability applies with $\eta' = 50\eta$, $N = \binom{n}{2}$ and $\epsilon' = \epsilon^2/500$. We can also require that $\eta < \epsilon^2/500$. We get a partition of the pairs of vertices of H as $U_1 \cup U_2$, where all but $\epsilon' N^2 < \epsilon^2 n^4 / 2000$ edges of H are formed by taking a pair from U_1 and a pair from U_2 .

We will call the pairs in U_1 *red*, and the pairs in U_2 *blue*. A 4-tuple $abcd$ will be called *properly coloured* if either

- (i) $abcd$ is an edge of H and each of the 3 sets $\{ab, cd\}, \{ac, bd\}, \{ad, bc\}$ has one red pair and one blue pair, or
- (ii) $abcd$ is not an edge and each of the 3 sets $\{ab, cd\}, \{ac, bd\}, \{ad, bc\}$ consists of two pairs with the same colour.

An improperly coloured 4-tuple is either an edge that is the union of two pairs of the same colour or a non-edge which is the union of two pairs with different colours. There are at most $\epsilon^2 n^4 / 2000$ of the former 4-tuples, and

the number of latter is at most

$$|U_1||U_2| - (e(G) - \epsilon'N^2) \leq \frac{50\eta}{2} \frac{N^2}{2} + \epsilon'N^2 \leq \left(\frac{50\eta}{16} + \epsilon'/4\right) n^4 < \epsilon^2 n^4 / 140.$$

Therefore all but $(\epsilon^2/140 + \epsilon^2/2000)n^4 < \epsilon^2 n^4 / 130$ 4-tuples are properly coloured.

A simple counting argument shows that there is a pair ab so that for all but $\binom{4}{2}(\epsilon^2 n^4 / 130) / \binom{n}{2} < \epsilon^2 n^2 / 10$ other pairs cd the 4-tuple $abcd$ is properly coloured. Without loss of generality ab is red. Partition the vertices of $V - ab$ into 4 sets according to the colour of the edges they send to $\{a, b\}$. We label these sets RR, BB, RB, BR , where R means ‘red’, B means ‘blue’ and a vertex c belongs to the set that labels the colours of the edges ca, cb in this order. Note that if c is in RR and d is in RB then ca and db are coloured red and blue, whereas cb and da are both red, so $abcd$ is improperly coloured. We deduce that one of RR and RB has size at most $\epsilon n / 3$, since otherwise we would have at least $\epsilon^2 n^2 / 9$ improperly coloured 4-tuples containing ab . The same argument applies when take one point from each of BB and RB , or RR and BR , or BB and BR . Therefore, either RB and BR each have size at most $\epsilon n / 3$, or RR and BB each have size at most $\epsilon n / 3$.

In the case when RB and BR each have size at most $\epsilon n / 3$ we look at the pairs in $RR \cup BB$. If c and d are both in RR then both of the opposite pairs $\{ac, bd\}$ and $\{ad, bc\}$ are coloured red. If cd is coloured blue then $abcd$ is improperly coloured, so all but at most $\epsilon^2 n^2 / 10$ pairs in RR are coloured red. Similarly all but at most $\epsilon^2 n^2 / 10$ pairs in BB are coloured red, and all but at most $\epsilon^2 n^2 / 10$ pairs with one vertex in RR and one in BB are coloured blue. Define a partition $V = V_1 \cup V_2$, where V_1 contains RR , V_2 contains BB and the remaining vertices are distributed arbitrarily. Note that all the incorrect 4-tuples with respect to this partition belong to the one of the following three groups.

- (i) Improperly coloured 4-tuples. There are at most $\epsilon^2 n^4 / 130$ of those.
- (ii) Properly coloured 4-tuples which use at least one vertex in $RB \cup BR$. There are at most $(2\epsilon n / 3) \binom{n}{3}$ such 4-tuples.
- (iii) Properly coloured 4-tuples which contain either a red pair of vertices with one vertex in RR and one in BB , or contain a blue pair of vertices from RR or from BB . There at most $(3\epsilon^2 n^2 / 10) \binom{n}{2}$ such 4-tuples.

Therefore all but at most $\frac{\epsilon^2 n^4}{130} + 2\frac{\epsilon n}{3} \binom{n}{3} + 3\frac{\epsilon^2 n^2}{10} \binom{n}{2} < \epsilon n^4$ 4-tuples are correct with respect to this partition.

The case when RR and BB each have size at most $\epsilon n / 3$ can be treated similarly. Here the conclusion is that all but at most $\epsilon^2 n^2 / 5$ pairs within RB or BR are coloured blue, and all but at most $\epsilon^2 n^2 / 10$ pairs with one vertex

in RB and one in BR are coloured red. Then, similarly as above one can show that with respect to a partition where V_1 contains RB , V_2 contains BR and the remaining vertices are distributed arbitrarily, all but at most en^4 4-tuples are correct. ■

Using the stability theorem we can now prove the following exact Turán result.

Theorem 2.2. *Let H be a 4-uniform hypergraph on n vertices that does not contain a copy of $\mathcal{C}_3^{(4)}$ and let n be sufficiently large. Then the number of edges in H is at most $b(n)$, with equality only when H is one of at most 2 hypergraphs \mathcal{B}_n .*

Proof. Let H be a 4-uniform hypergraph on n vertices, which has $e(H) \geq b(n)$ and contains no $\mathcal{C}_3^{(4)}$. First we claim that we can assume that H has minimum degree at least $b(n) - b(n - 1)$. Indeed, suppose that we have proved the result under this assumption for all $n \geq n_0$. Construct a sequence of hypergraphs $H = H_n, H_{n-1}, \dots$ where H_{m-1} is obtained from H_m by deleting a vertex of degree less than $b(m) - b(m - 1)$. By setting $f(m) = e(H_m) - b(m)$ we have $f(n) \geq 0$ and $f(m) \geq f(m + 1) + 1$. If we can continue this process to obtain a hypergraph H_{n_0} then $n - n_0 \leq \sum_{m=n_0}^{n-1} (f(m) - f(m + 1)) \leq f(n_0) \leq \binom{n_0}{4}$, which is a contradiction for n sufficiently large. Otherwise we obtain a hypergraph $H_{n'}$ with $n > n' > n_0$ having minimal degree at least $b(n') - b(n' - 1)$ and without a $\mathcal{C}_3^{(4)}$. Then by the above assumption $e(H_{n'}) \leq b(n')$ and again we obtain a contradiction, since

$$e(H) = e(H_n) \leq b(n') + \sum_{n' < m \leq n} (b(m) - b(m - 1) - 1) < b(n).$$

Substituting from equation (2) we can assume H has minimum degree

$$(5) \quad \delta(H) \geq b(n) - b(n - 1) > \frac{1}{12}n^3 - \frac{1}{2}n^2.$$

Given a partition of $V(H) = V_1 \cup V_2$, we call an edge $abcd$ of H *good* if $abcd$ is a good 4-tuple (as defined before) with respect to this partition; otherwise we call it *bad*. By Theorem 2.1 there is a partition with all but at most $10^{-25}n^4$ edges of H being good. Let $V(H) = V_1 \cup V_2$ be the partition which minimises the number of bad edges. With respect to this partition, every vertex belongs to at least as many good edges as bad edges, or we can move it to the other class of the partition. Also, by definition, there are at most $b(n)$ good 4-tuples with respect to any partition. We must have

$||V_1| - n/2| < 10^{-6}n$ and $||V_2| - n/2| < 10^{-6}n$. Otherwise by equation (1) we get

$$e(H) < \frac{1}{48} \left((n^2 - 3n + 4)^2 - (4 \cdot 10^{-12}n^2 - 3n + 4)^2 \right) + 10^{-25}n^4 < b(n),$$

which is a contradiction.

Note that there is no pair of vertices ab for which there are both $10^{-10}n^2$ pairs cd such that $abcd$ is a good edge and $10^{-10}n^2$ pairs ef such that $abef$ is an bad edge. Indeed, each such cd and ef which are disjoint give a 4-tuple $cdef$ which is good, but cannot be an edge as it would create a $\mathcal{C}_3^{(4)}$. Moreover, every 4-tuple can be obtained at most 3 times in this way, and every cd is disjoint from all but at most $2n$ pairs ef . Thus at least $10^{-10}n^2(10^{-10}n^2 - 2n)/3 > 10^{-21}n^4$ good 4-tuples are not edges of H , and therefore $e(H) < b(n) - 10^{-21}n^4 + 10^{-25}n^4 < b(n)$, which is a contradiction.

The next step of the proof is the following claim.

Claim 2.3. *Any vertex of H is contained in at most $10^{-5}n^3$ bad edges.*

Proof. Suppose some vertex a belongs to $10^{-5}n^3$ bad edges. Call another vertex b *good* if there are at most $10^{-10}n^2$ pairs cd such that $abcd$ is a bad edge, otherwise call b *bad*. By the above discussion, for every bad vertex b there are at most $10^{-10}n^2$ pairs ef such that $abef$ is a good edge. Note that there are at least $10^{-5}n$ bad vertices, otherwise we would only have at most $10^{-5}n \cdot \binom{n}{2} + (1 - 10^{-5})n \cdot 10^{-10}n^2 < 10^{-5}n^3$ bad edges through a , which is contrary to our assumption. By choice of partition there are at least as many good edges containing a as bad. We know that a has degree at least $\frac{1}{12}n^3 - \frac{1}{2}n^2$, at least half of which is good, so there are at least $n/24$ good vertices.

Suppose that the number of good vertices is αn , and so there are $(1-\alpha)n-1$ bad vertices. We can count the edges containing a as follows. By definition there are at most $10^{-10}n^3$ such good edges containing a bad vertex, and at most $10^{-10}n^3$ such bad edges containing a good vertex. Now we bound the number of remaining good edges. Note that these edges only contain good vertices. Looking at the vertices of such an edge in some order, we can select the first 2 vertices in $\alpha n(\alpha n - 1)$ ways. Since the edge is good, the choice of 2 vertices together with a restricts the fourth vertex to lie in some particular class V_i , so it can be chosen in at most $(\frac{1}{2} + 10^{-6})n$ ways. Note that we have counted each edge 6 times, so we get at most $\alpha n(\alpha n - 1)(\frac{1}{2} + 10^{-6})n/6 < (\alpha^2 + \frac{1}{2} \cdot 10^{-5})\frac{1}{2}\binom{n}{3}$ edges. Similarly there are at most $((1-\alpha)^2 + \frac{1}{2} \cdot 10^{-5})\frac{1}{2}\binom{n}{3}$ remaining bad edges through a . Since $1/24 \leq \alpha \leq 1 - 10^{-5}$, in total the

number of edges containing a is bounded by

$$\begin{aligned} & \left(\alpha^2 + \frac{1}{2} \cdot 10^{-5}\right) \frac{1}{2} \binom{n}{3} + \left((1 - \alpha)^2 + \frac{1}{2} \cdot 10^{-5}\right) \frac{1}{2} \binom{n}{3} + 2 \cdot 10^{-10} n^3 \\ & < \frac{1}{12} n^3 - \frac{1}{2} n^2 < \delta(H). \end{aligned}$$

This contradiction proves the claim. ■

Now write $|V_1| = n/2 + t$, $|V_2| = n/2 - t$ with $-10^{-6}n < t < 10^{-6}n$. By possibly renaming the classes (i.e. replacing t with $-t$) we can assume that $d(n, t) \leq d(n, -t)$. Then any vertex of V_1 belongs to $d(n, t)$ good 4-tuples. Now $d(n, t)$ is the minimum degree of $\mathcal{B}(n, t)$, which is certainly at most the maximum degree of \mathcal{B}_n . Comparing with equation (4) we see that any vertex of V_1 belongs to at most $\frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2}$ good 4-tuples. From now on this will be the only property of V_1 we use that might possibly not be a property of V_2 . We will eventually end up showing the same bound on the number of good 4-tuples containing a vertex of V_2 . Then the whole argument will apply verbatim switching V_1 for V_2 .

We will use this property in the following manner. Suppose a is a vertex of V_1 for which K of the good 4-tuples containing a are not edges of H . Then there are at most $\frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2} - K$ good edges containing a , so by (5) there must be at least

$$\begin{aligned} & \delta(H) - \left(\frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2} - K\right) \\ & \geq \left(\frac{1}{12}n^3 - \frac{1}{2}n^2\right) - \left(\frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2} - K\right) = K - n^{3/2} \end{aligned}$$

bad edges containing a . Similarly, if a' is a vertex in V_2 then it belongs to at most

$$d(n, -t) = \frac{1}{12}(n^3 - 6n^2 + 8n + 12t^2) + \frac{1}{12}|6tn - 8t^3 - 16t| < \frac{1}{12}n^3 - \frac{1}{2}n^2 + 10^{-6}n^3$$

good edges. Thus, if it belongs to L good 4-tuples which are not edges of H then it must belong to at least $L - 10^{-6}n^3$ bad edges.

Suppose for the sake of contradiction that there is some bad edge incident with V_1 . Denote the set of bad edges containing some vertex v by $\mathcal{Z}(v)$. Let a be a vertex in V_1 belonging to the maximum number of bad edges and let $Z = |\mathcal{Z}(a)|$. Note that $Z > 0$. For every bad edge $abcd$ containing a , consider a partition of its vertices into pairs, say ac and bd . Recall that there are 2 types of pairs, one type consisting of pairs with both vertices in one

class, the other consisting of pairs that have one point of each class. By definition of a bad edge, ac and bd are pairs of the same type. If ef is any pair of the other type which is disjoint from both of them, then $acef$ and $bdef$ are good 4-tuples. One of them is not an edge of H , or we get a $C_3^{(4)}$. The number of such pairs ef is clearly at least

$$\begin{aligned} \min \left\{ (|V_1| - 4)(|V_2| - 4), \binom{|V_1| - 4}{2} + \binom{|V_2| - 4}{2} \right\} \\ \geq \left(\frac{1}{4} - 10^{-12} \right) n^2 - O(n) > n^2/5. \end{aligned}$$

Let $Z_1(a)$ be those bad edges for which there is some partition into pairs ac and bd , so that for at least $n^2/10$ of the pairs ef defined above, the good 4-tuple $acef$ is not an edge. Let $Z_2(a) = Z(a) - Z_1(a)$, and write $Z_i = |Z_i(a)|$ for $i=1, 2$. Then one of Z_1, Z_2 is at least $Z/2$.

Case 1. Suppose $Z_1 \geq Z/2$. Let C be the (non-empty) set of vertices c such that there is some edge $abcd$ in $Z_1(a)$, and $acef$ is a good non-edge for at least $n^2/10$ pairs ef . Then we have at least $|C|n^2/30$ good non-edges containing a , as we count each $acef$ at most 3 times. This implies that there are at least $|C|n^2/30 - n^{3/2} \geq |C|n^2/31$ bad edges containing a and therefore $n^2/31 \leq Z/|C|$. Since every edge in $Z_1(a)$ contains at most 3 vertices of C there exists $c \in C$ which is contained in at least $|Z_1(a)|/(3|C|) = Z_1/(3|C|) \geq Z/(6|C|) > n^2/200$ bad edges. Fix one such c .

Note that a graph with n vertices and m edges contains a matching of size at least $m/2n$, since otherwise there is a set of fewer than m/n vertices that cover all the edges of the graph, which is impossible by direct counting. Consider the set of pairs bd such that $abcd$ is a bad edge. Then there exists a matching M of size at least $n/400$ so that for each bd in M we have that $abcd$ is a bad edge of H . Partition such an edge into pairs ab and cd . Then, as we explained above, there are at least $n^2/5$ pairs ef such that one of the 4-tuples $abef$ and $cdef$ is a good non-edge. Since M is a matching we count each such 4-tuple at most 3 times, so one of a or c belongs to at least $\frac{1}{3} \cdot \frac{1}{2} \cdot \frac{n^2}{5} \cdot \frac{n}{400} = n^3/12000$ good non-edges. Therefore it belongs to at least $n^3/12000 - 10^{-6}n^3 > 10^{-5}n^3$ bad edges, which contradicts [Claim 2.3](#).

Case 2. Now suppose $Z_2 \geq Z/2$. Note that every bad edge containing a contains at least one other point of V_1 , so there is some $b \in V_1$ belonging to at least Z_2/n edges of $Z_2(a)$. Fix one such b . Suppose cd is a pair such that $abcd$ is in $Z_2(a)$, and consider any partition of $abcd$ into pairs p_1, p_2 with a in p_1 and b in p_2 . Then, by definition of $Z_2(a)$, there are at least $n^2/10$ pairs ef such that $p_2 \cup ef$ is a good non-edge. Let C be the set of vertices c for

which there exists a vertex d such that $abcd$ is an edge of $\mathcal{Z}_2(a)$. Then there are at least $|C|n^2/30$ good non-edges containing b , as we count each $bcef$ at most 3 times. Thus, there are at least $|C|n^2/30 - n^{3/2} > |C|n^2/50$ bad edges containing b . By maximality of Z we have $|C|n^2/50 \leq |\mathcal{Z}(b)| \leq Z$. Note that each edge in $\mathcal{Z}_2(a)$ that contains b is obtained by picking a pair of vertices in C , so $Z/(2n) \leq Z_2/n \leq \binom{|C|}{2} < 1250Z^2/n^4$. Therefore $Z \geq n^3/2500$, which again contradicts [Claim 2.3](#).

We conclude that there are no bad edges incident to the vertices of V_1 , i.e. all bad edges have all 4 vertices in V_2 . We can use this information to give more precise bounds on the sizes of V_1 and V_2 . We recall that $|V_1| = n/2 + t$, $|V_2| = n/2 - t$ and $d(n, t) \leq d(n, -t)$. Suppose that $|t| \geq \sqrt{n}$, so that $6|t|n - 8|t|^3 - 16|t| < -2n^{3/2}$ and by (3)

$$\begin{aligned} d(n, t) &= \frac{1}{12}(n^3 - 6n^2 + 8n + 12t^2) + \frac{1}{12}(6|t|n - 8|t|^3 - 16|t|) \\ &< \frac{1}{12}n^3 - \frac{1}{2}n^2 - \frac{n^{3/2}}{12} < \delta(H). \end{aligned}$$

This is a contradiction, since the vertices of V_1 only belong to good edges, of which there are at most $d(n, t) < \delta(H)$. Therefore $|t| < \sqrt{n}$. Now we can bound the number of good 4-tuples containing a vertex of V_2 . By (3), this number is at most

$$\begin{aligned} d(n, -t) &= \frac{1}{12}(n^3 - 6n^2 + 8n + 12t^2) + \frac{1}{12}(8|t|^3 - 6|t|n + 16|t|) \\ &< \frac{1}{12}n^3 - \frac{1}{2}n^2 + n^{3/2}. \end{aligned}$$

Now the same argument as we used to show that no bad edges are incident with the vertices of V_1 shows that none are incident with V_2 either. We conclude that all edges are good. Then by definition of $b(n)$ we have $e(H) \leq b(n)$, with equality only when H is a \mathcal{B}_n , so the theorem is proved. ■

3. Proof of Frankl’s conjecture

In this section we will prove the general case of the Frankl conjecture. We recall that $\mathcal{C}_3^{(2k)}$ is the $2k$ -uniform hypergraph with three edges $\{P_1 \cup P_2, P_2 \cup P_3, P_3 \cup P_1\}$, where P_1, P_2, P_3 are pairwise disjoint sets of k vertices. We can obtain a large $\mathcal{C}_3^{(2k)}$ -free graph on n vertices by partitioning an n -element set V into 2 parts V_1, V_2 and taking those edges which intersect each part V_i in an odd number of elements. To see this, consider any P_1, P_2, P_3 that

are pairwise disjoint sets of k vertices. Then $|V_1 \cap P_i|$ and $|V_1 \cap P_j|$ have the same parity for some pair ij , so $P_i \cup P_j$ is not an edge.

Note that this construction is the same as the one we described for $\mathcal{C}_3^{(4)}$ when $k=2$. In the 4-uniform case we were able to calculate the sizes of the parts that maximise the number of edges. For general k this is an interesting problem in enumerative combinatorics, that is equivalent to finding the minima of binary Krawtchouk polynomials. These polynomials play an important rôle in the analysis of binary Hamming association schemes and so many of their properties are well-known in this context (see, e.g., [5]). In particular, the location of their roots is an important problem, but we will need here only a crude estimate that follows easily from known results. In the [first subsection](#) of this section we will state this estimate and apply it to various parameters of our construction. The rest of the proof follows the same broad outline as that of the 4-uniform case, in that it falls naturally into two parts. We will prove the stability part in the [second subsection](#), and the full result we defer to the [final subsection](#).

3.1. Binary Krawtchouk polynomials

Let $\mathcal{B}^{(2k)}(n, t)$ denote the $2k$ -uniform hypergraph obtained by partitioning an n -element set into two parts with sizes $\frac{n}{2} + t$ and $\frac{n}{2} - t$, and taking as edges all $2k$ -tuples with odd intersection with each part. Let $b_{2k}(n, t)$ be the number of edges in $\mathcal{B}^{(2k)}(n, t)$ and let $d_{2k}(n, t)$ be the degree of any vertex belonging to the side with size $\frac{n}{2} + t$. Then the vertices on the side with size $\frac{n}{2} - t$ have degree $d_{2k}(n, -t)$.

The *binary Krawtchouk polynomials* $K_m^n(x)$ can be defined by the generating function

$$\sum_{m=0}^n K_m^n(x) z^m = (1 - z)^x (1 + z)^{n-x}.$$

From here we get the explicit expression $K_m^n(x) = \sum_{i=0}^m (-1)^i \binom{x}{i} \binom{n-x}{m-i}$. Recall that $b_{2k}(n, t)$ was the number of $2k$ -tuples with odd intersection with both parts in the above partition of an n -element set and so $\binom{n}{2k} - b_{2k}(n, t)$ is the number of $2k$ -tuples with even intersection with these parts. This implies that $(\binom{n}{2k} - b_{2k}(n, t)) - b_{2k}(n, t) = \sum_{i=0}^{2k} (-1)^i \binom{n/2+t}{i} \binom{n/2-t}{2k-i} = K_{2k}^n(n/2 + t)$, which gives

$$(6) \quad b_{2k}(n, t) = \frac{1}{2} \left(\binom{n}{2k} - K_{2k}^n(n/2 + t) \right),$$

so maximising $b_{2k}(n, t)$ is equivalent to finding the minimum of $K_{2k}^n(x)$. Similarly, we can also express the degrees of $\mathcal{B}^{(2k)}(n, t)$ in terms of Krawtchouk polynomials. Indeed, by definition, $d_{2k}(n, t)$ is the number of $(2k - 1)$ -tuples with even intersection with the first part in the partition of an $(n - 1)$ -element set in two parts with sizes $n/2 + t - 1$ and $n/2 - t$, and therefore $\binom{n-1}{2k-1} - d_{2k}(n, t)$ is the number of $(2k - 1)$ -tuples with odd intersection with this part. Then, $d_{2k}(n, t) - (\binom{n-1}{2k-1} - d_{2k}(n, t)) = \sum_{i=0}^{2k-1} (-1)^i \binom{n/2+t-1}{i} \binom{n/2-t}{2k-1-i} = K_{2k-1}^{n-1}(n/2 + t - 1)$, i.e.

$$(7) \quad d_{2k}(n, t) = \frac{1}{2} \left(\binom{n-1}{2k-1} + K_{2k-1}^{n-1}(n/2 + t - 1) \right).$$

Note that $K_m^n(x)$ is a polynomial of degree m . It is known that it has m simple roots, symmetric with respect to $n/2$. The smallest root is given by the following formula obtained by Levenshtein [6]:

$$r = n/2 - \max \left(\sum_{i=0}^{m-2} x_i x_{i+1} \sqrt{(i+1)(n-i)} \right),$$

where the maximum is taken over x_i with $\sum_{i=0}^{m-1} x_i^2 = 1$. From the Cauchy-Schwartz inequality we see that $n/2 - r < \sqrt{mn}$. Note that $K_{2k}^n(0) = K_{2k}^n(n) = \binom{n}{2k} > 0$, so the minimum of $K_{2k}^n(x)$ occurs in the range $n/2 \pm \sqrt{2kn}$.

Let t^* be chosen to maximise the number of edges in $\mathcal{B}^{(2k)}(n, t)$, and denote any hypergraph obtained in this manner by $\mathcal{B}_n^{(2k)}$. Note that t^* may not be unique, but must satisfy $|t^*| < \sqrt{2kn}$. Also, by symmetry we can assume that $t^* > 0$. Write $b_{2k}(n)$ for the number of edges in $\mathcal{B}_n^{(2k)}$.

Lemma 3.1. (i) $K_m^n(n/2 + t) = \sum_{i=0}^{m/2} (-1)^{i+m} \binom{n/2-t}{i} \binom{2t}{m-2i}$.

(ii) If $c > 1$ and $0 \leq s \leq c\sqrt{n}$ then $\left| d_{2k}(n, \pm s) - \frac{1}{2} \binom{n-1}{2k-1} \right| < (10c^2)^k n^{k-1/2}$.

(iii) $\left| b_{2k}(n) - \frac{1}{2} \binom{n}{2k} \right| < (20kn)^k$, $\left| d_{2k}(n, \pm t^*) - \frac{1}{2} \binom{n-1}{2k-1} \right| < (20k)^k n^{k-1/2}$.

(iv) If $C > 20^k$ then $d_{2k}(n, C\sqrt{n}) < \frac{1}{2} \binom{n-1}{2k-1} - 20^k n^{k-1/2}$.

(v) $\left| b_{2k}(n, \epsilon n) - \left(\frac{1}{2} \binom{n}{2k} - \frac{1}{2} \binom{2\epsilon n}{2k} \right) \right| < (10\epsilon)^k n^{2k-1}$, $\left| d_{2k}(n, \epsilon n) - \left(\frac{1}{2} \binom{n-1}{2k-1} - \frac{1}{2} \binom{2\epsilon n-1}{2k-1} \right) \right| < (10\epsilon)^k n^{2k-2}$.

Proof. (i) Rewrite the generating function as $\sum_{m=0}^n K_m^n(n/2 + t) z^m = (1 - z^2)^{n/2-t} (1 - z)^{2t}$ and expand.

(ii) Using part (i) with $t = s - 1/2$, and applying (7), we get

$$\begin{aligned} \left| d_{2k}(n, s) - \frac{1}{2} \binom{n-1}{2k-1} \right| &= \frac{1}{2} |K_{2k-1}^{n-1}(n/2 + s - 1)| \\ &= \frac{1}{2} \left| \sum_{i=0}^{k-1} (-1)^{i+1} \binom{n/2 - s}{i} \binom{2s - 1}{2k - 1 - 2i} \right| \\ &< k \cdot (2c\sqrt{n})^{2k-1} < (10c^2)^k n^{k-1/2}. \end{aligned}$$

The corresponding inequality for $d_{2k}(n, -s)$ can be obtained similarly.

(iii) The second statement follows from part (ii) with $c = \sqrt{2k} \geq t^*/\sqrt{n}$. To prove the first statement, we use (6), part (i) and again the fact that $0 < t^* < \sqrt{2kn}$. Altogether they imply

$$\begin{aligned} \left| b_{2k}(n) - \frac{1}{2} \binom{n}{2k} \right| &= \frac{1}{2} |K_{2k}^n(n/2 + t^*)| < \sum_{i=0}^k \binom{n/2}{i} \binom{2\sqrt{2kn}}{2k - 2i} \\ &< (k + 1)(2\sqrt{2kn})^{2k} < (20kn)^k. \end{aligned}$$

(iv) By (7) we have

$$\begin{aligned} d_{2k}(n, C\sqrt{n}) - \frac{1}{2} \binom{n-1}{2k-1} &= \frac{1}{2} K_{2k-1}^{n-1}(n/2 + C\sqrt{n} - 1) \\ &= \frac{1}{2} \sum_{i=0}^{k-1} (-1)^{i+1} \binom{n/2 - C\sqrt{n}}{i} \binom{2C\sqrt{n} - 1}{2k - 1 - 2i} \\ &< -\frac{1}{2} \binom{2C\sqrt{n} - 1}{2k - 1} + \frac{1}{2}(k - 1)(n/2) \binom{2C\sqrt{n} - 1}{2k - 3} \\ &= (1 + o(1)) \left(-\frac{(2C\sqrt{n})^{2k-1}}{2(2k - 1)!} + \frac{(k - 1)n(2C\sqrt{n})^{2k-3}}{4(2k - 3)!} \right) \\ &< -\left(\frac{C^2}{(2k - 1)^2} - \frac{k - 1}{2} \right) \frac{(2C)^{2k-3}}{(2k - 3)!} n^{k-1/2} < -20^k n^{k-1/2}. \end{aligned}$$

(v) Using the formula for $K_{2k}^n(n/2 + t)$ from part (i) together with (6) we obtain that

$$\begin{aligned}
 \left| b_{2k}(n, \epsilon n) - \left(\frac{1}{2} \binom{n}{2k} - \frac{1}{2} \binom{2\epsilon n}{2k} \right) \right| &= \left| \frac{1}{2} K_{2k}^n(n/2 + \epsilon n) - \frac{1}{2} \binom{2\epsilon n}{2k} \right| \\
 &= \frac{1}{2} \left| \sum_{i=1}^k (-1)^i \binom{n/2 - \epsilon n}{i} \binom{2\epsilon n}{2k - 2i} \right| \\
 &< \frac{n}{2} \cdot (2\epsilon n)^{2k-2} + O(n^{2k-2}) \\
 &< (10\epsilon)^k n^{2k-1}.
 \end{aligned}$$

The proof of the inequality for $d_{2k}(n, \epsilon n)$ can be obtained similarly and we omit it here. ■

We remark that these simple estimates are sufficient for our purposes, but the location of the roots and asymptotic values for Krawtchouk polynomials in the oscillatory region are known with more precision (see, e.g., [5]). With this information one could find better estimates for $b_{2k}(n)$, and possibly how many different choices of t give the maximum number of edges.

We conclude this section with an estimate on the difference of successive values of $b_{2k}(n)$.

Lemma 3.2. $b_{2k}(n) - b_{2k}(n - 1) \geq \frac{1}{2} \binom{n-1}{2k-1}$.

Proof. Suppose that $\mathcal{H} = \mathcal{B}^{(2k)}(n - 1)$ has $b_{2k}(n - 1)$ edges and has parts $V(\mathcal{H}) = A \cup B$. Let \mathcal{H}_1 be obtained from \mathcal{H} by adding a vertex v_1 to A , together with all the $2k$ -tuples containing v_1 and having odd intersections with $A \cup v_1$ and B . Let \mathcal{H}_2 be similarly obtained by adding a vertex v_2 to B , together with corresponding edges. By definition each \mathcal{H}_i has at most $b_{2k}(n)$ edges, so the degree of each v_i is a lower bound for $b_{2k}(n) - b_{2k}(n - 1)$. On the other hand, for each $(2k - 1)$ -tuple X of vertices in \mathcal{H} there is exactly one i such that $X \cup v_i$ is an edge of \mathcal{H}_i , so one of the v_i has degree at least $\frac{1}{2} \binom{n-1}{2k-1}$. ■

3.2. A stability result for $\mathcal{C}_3^{(2k)}$

In this subsection we prove a stability result for $\mathcal{C}_3^{(2k)}$. We start by recalling a version of the Kruskal–Katona theorem due to Lovász. Write $[m] = \{1, \dots, m\}$, let $[m]^{(k)}$ denote the subsets of $[m]$ of size k , and suppose $\mathcal{A} \subset [m]^{(k)}$. The *shadow* of \mathcal{A} is $\partial\mathcal{A} \subset [m]^{(k-1)}$ consisting of all sets of size $k - 1$ that are contained in some element of \mathcal{A} . For any real x write $\binom{x}{k} = x(x - 1) \cdots (x - k + 1)/k!$. The following result appears in [7] (Exercise 13.31).

Proposition 3.3. *If $\mathcal{A} \subset [m]^{(k)}$ and $|\mathcal{A}| = \binom{x}{k}$ then $|\partial\mathcal{A}| \geq \binom{x}{k-1}$. ■*

Suppose we have a $2k$ -uniform hypergraph H and a partition of the vertex set $V(H) = V_1 \cup V_2$. Our terminology for $2k$ -tuples matches that of the 4 -uniform case. We call a $2k$ -tuple of vertices *good* if it intersects each V_i in an odd number of elements; otherwise we call it *bad*. We call a $2k$ -tuple *correct* if it is either a good edge or a bad non-edge; otherwise we call it *incorrect*.

Theorem 3.4. *For every $\epsilon > 0$ there is $\eta > 0$ so that if H is a $\mathcal{C}_3^{(2k)}$ -free $2k$ -uniform hypergraph with $e(H) > \frac{1}{2} \binom{n}{2k} - \eta n^{2k}$ then there is a partition of the vertex set as $V(H) = V_1 \cup V_2$ such that all but ϵn^{2k} $2k$ -tuples are correct.*

Proof. Define an auxiliary graph G whose vertices are all k -tuples of vertices of H , and where the k -tuples P_1 and P_2 are adjacent exactly when $P_1 \cup P_2$ is an edge of H . Since H is $\mathcal{C}_3^{(2k)}$ -free we see that G is triangle-free. Also, each edge of H creates exactly $\frac{1}{2} \binom{2k}{k}$ edges in G (corresponding to the ways of breaking a $2k$ -tuple into two k -tuples) so

$$e(G) > \frac{1}{2} \binom{2k}{k} \left(\frac{1}{2} \binom{n}{2k} - \eta n^{2k} \right) > (1 - (k!)^2 2^{2k} \eta) \frac{1}{2} \binom{n}{2k}.$$

Choose η so that the Simonovits stability theorem (see Section 2) applies with $\eta' = (k!)^2 2^{2k} \eta$, $N = \binom{n}{k}$ and $\epsilon' = 10^{-6k^2} \epsilon^k$. We can also require that $\eta < 10^{-6k^2} \epsilon^k$. We get a partition of the k -tuples of vertices of H as $U_0 \cup U_1$, where all but $\epsilon' N^2 = \epsilon' \binom{n}{k}^2 < 10^{-6k^2} \epsilon^k n^{2k}$ edges of H are formed by taking a k -tuple from U_0 and a k -tuple from U_1 .

We will think of the sets U_i as determining a 2-colouring of all k -tuples, and say that the k -tuples in U_i have colour i . A $2k$ -tuple I will be called *properly coloured* if, either it is an edge of H and however we partition I into k -tuples P_1 and P_2 they have different colours, or it is not an edge of H and for any partition of I into two k -tuples they have the same colour.

An improperly coloured $2k$ -tuple is either an edge that is the union of two k -tuples of the same colour or a non-edge which is the union of two k -tuples with different colours. There are at most $10^{-6k^2} \epsilon^k n^{2k}$ of the former $2k$ -tuples, and the number of the latter is at most

$$\begin{aligned} |U_0||U_1| - (e(G) - \epsilon' N^2) &\leq \frac{(k!)^2 2^{2k} \eta}{2} \frac{N^2}{2} + \epsilon' N^2 \leq \left(\frac{(k!)^2 2^{2k} \eta}{4(k!)^2} + \frac{\epsilon'}{(k!)^2} \right) n^{2k} \\ &\leq 10^{-5k^2-1} \epsilon^k n^{2k}. \end{aligned}$$

Therefore all but $(10^{-6k^2} \epsilon^k + 10^{-5k^2-1}) \epsilon^k n^{2k} < 10^{-5k^2} \epsilon^k n^{2k}$ $2k$ -tuples are properly coloured.

A simple counting argument shows that there is a k -tuple P so that for all but $\binom{2k}{k} 10^{-5k^2} \epsilon^k n^{2k} / \binom{n}{k} < 10^{-4k^2} \epsilon^k n^k$ other k -tuples Q the $2k$ -tuple $P \cup Q$ is properly coloured. Without loss of generality P has colour 0. We will call a k -tuple Q *proper* if $P \cup Q$ is properly coloured; otherwise it is *improper*. Then by definition there are at most $10^{-4k^2} \epsilon^k n^k$ improper k -tuples. Call a $(k - 1)$ -tuple $X \subset V - P$ *abnormal* if there are at least $2^{-3k} \epsilon n$ vertices $x \in V - (P \cup X)$ for which $X \cup x$ is improper; otherwise call it *normal*. It is easy to see that there are at most $k \cdot 10^{-4k^2} \epsilon^k n^k / (2^{-3k} \epsilon n) < 10^{-3k^2} \epsilon^{k-1} n^{k-1}$ abnormal $(k - 1)$ -tuples.

We partition the vertices of $V - P$ according to the colour of the k -tuples that they form when they replace an element of P . To be precise, we fix an order p_1, \dots, p_k of P and partition into 2^k parts $V - P = \bigcup V_s$, where $s = (s_1, \dots, s_k) \in \{0, 1\}^k$ and a vertex x belongs to V_s iff $(P - p_i) \cup x$ has colour s_i for every $1 \leq i \leq k$.

Consider a $(k - 1)$ -tuple $X = x_1 \cdots x_{k-1}$ and suppose a is a vertex such that $X \cup a$ is proper. Fix $1 \leq i \leq k$ and consider the partitions $P \cup X \cup a = (P) \cup (X \cup a) = ((P - p_i) \cup a) \cup (X \cup p_i)$. Let V_s be the class containing a , so that $(P - p_i) \cup a$ has colour s_i . We recall that P has colour 0, so if also $s_i = 0$ then to be properly coloured $X \cup a$ must have the same colour as $X \cup p_i$. On the other hand, if $s_i = 1$ then $X \cup a$ and $X \cup p_i$ must have different colours. If we write $c_X(v)$ for the colour of $X \cup v$ for any vertex v , then this can be summarised as

$$(8) \quad \text{If } a \in V_s \text{ and } X \cup a \text{ is proper, then } c_X(a) + s_i = c_X(p_i) \pmod{2}.$$

Suppose there are 2 classes V_s and $V_{s'}$ both of size at least $2^{-2k} \epsilon n$. Since $\binom{2^{-2k} \epsilon n}{k-1} > 10^{-3k^2} \epsilon^{k-1} n^{k-1}$ some $(k - 1)$ -tuple $X \subset V_s$ is normal. This means that there are at most $2^{-3k} \epsilon n$ vertices $x \in V - (P \cup X)$ for which $X \cup x$ is improper, so there is $a \in V_s$ and $b \in V_{s'}$ such that $X \cup a$ and $X \cup b$ are proper. For any pair of indices i, j we have $c_X(a) + s_i = c_X(p_i)$, $c_X(a) + s_j = c_X(p_j)$, $c_X(b) + s'_i = c_X(p_i)$ and $c_X(b) + s'_j = c_X(p_j)$. Adding these equations gives $s_i + s_j + s'_i + s'_j = 0$. If s and s' differ in some co-ordinate i then this equation shows that they must also differ in any other co-ordinate j . In other words, if $s' \neq s$ we must have $s' = \bar{s}$, where \bar{s} denotes the sequence whose i th entry is $1 - s_i$.

Let $V_{\bar{s}}$ be the largest class, and write $m = |V_{\bar{s}}|$. Clearly $m \geq 2^{-k} (n - k)$. Then all other classes, except possibly $V_{\bar{s}}$, have size at most $2^{-2k} \epsilon n$. Let \mathcal{A}_i be the set of proper k -tuples contained in $V_{\bar{s}}$ that have colour i . Then $|\mathcal{A}_0| + |\mathcal{A}_1| > \binom{m}{k} - 10^{-4k^2} \epsilon^k n^k > (1 - 10^{-3k^2} \epsilon) \binom{m}{k}$. Write $|\mathcal{A}_i| = \alpha_i \binom{m}{k}$, so that $\alpha_0 + \alpha_1 > 1 - 10^{-3k^2} \epsilon$. Suppose both α_i are at least $10^{-2k^2} \epsilon$. Observe that

$|\mathcal{A}_i| = \binom{\alpha_i^{1/k} m}{k} + O(m^{k-1})$, so by Proposition 3.3 we have

$$|\partial\mathcal{A}_i| \geq \binom{\alpha_i^{1/k} m}{k-1} + O(m^{k-2}) = \alpha_i^{(k-1)/k} \binom{m}{k-1} + O(m^{k-2}).$$

Note that if $z \leq 2^{-k}$ we have that $z^{-1/k} \geq 2$ and therefore

$$\begin{aligned} z^{(k-1)/k} + (1 - 10^{-3k^2} \epsilon - z)^{(k-1)/k} &\geq z^{(k-1)/k} + (1 - 10^{-3k^2} \epsilon - z) \\ &\geq 2z + 1 - 10^{-3k^2} \epsilon - z = 1 + z - 10^{-3k^2} \epsilon. \end{aligned}$$

Since $z^{(k-1)/k}$ is concave, $\alpha_0 \geq 10^{-2k^2} \epsilon$ and $10^{-2k^2} \epsilon < 2^{-k}$ we have

$$\begin{aligned} \alpha_0^{(k-1)/k} + \alpha_1^{(k-1)/k} &> \alpha_0^{(k-1)/k} + (1 - 10^{-3k^2} \epsilon - \alpha_0)^{(k-1)/k} \\ &\geq (10^{-2k^2} \epsilon)^{(k-1)/k} + (1 - 10^{-3k^2} \epsilon - 10^{-2k^2} \epsilon)^{(k-1)/k} \\ &\geq 1 + 10^{-2k^2} \epsilon - 10^{-3k^2} \epsilon \geq 1 + 10^{-3k^2} \epsilon. \end{aligned}$$

We deduce that $|\partial\mathcal{A}_0 \cap \partial\mathcal{A}_1| > 0$, i.e. there is a $(k-1)$ -tuple X and points a_0, a_1 such that $X \cup a_i$ is proper, with $c_X(a_i) = i$. But equation (8) gives $i + s_1 = c_X(a_i) + s_1 = c_X(p_1)$, for $i = 0, 1$, which is a contradiction. We conclude that there is $t \in \{0, 1\}$ for which $\alpha_{1-t} < 10^{-2k^2} \epsilon$, and so all but at most $10^{-2k^2} \epsilon \binom{m}{k} + 10^{-4k^2} \epsilon^k n^k < 10^{-2k^2} \epsilon n^k$ k -tuples inside $V_{\mathbb{S}}$ have the same colour t .

For $0 \leq i \leq k$ let \mathcal{D}_i be all k -tuples with i points in $V_{\mathbb{S}}$ and $k-i$ points in $V_{\mathbb{S}}$ and let $\theta_i = 10^{-2k^2} (2k2^{2k})^i \epsilon$. We claim that for each i all but at most $\theta_i n^k$ k -tuples of \mathcal{D}_i have colour $t+i \pmod{2}$. Otherwise, choose the smallest i for which this is not true. By the above discussion $i > 0$, and there are at least $\theta_i n^k$ k -tuples in \mathcal{D}_i with colour $1 - (t+i) = t+i-1 \pmod{2}$. Since i was the smallest such index all but at most $\theta_{i-1} n^k$ k -tuples of \mathcal{D}_{i-1} have colour $t+i-1 \pmod{2}$. Let E_{i-1} be the $(k-1)$ -tuples Y with $i-1$ points in $V_{\mathbb{S}}$ and $k-i$ points in $V_{\mathbb{S}}$ for which there are at least $2^{-2k} n$ points $y \in V_{\mathbb{S}}$ such that $Y \cup y$ does not have colour $t+i-1 \pmod{2}$. Then $|E_{i-1}| \leq k\theta_{i-1} n^k / (2^{-2k} n) = \frac{1}{2} \theta_i n^{k-1}$, so at most $\frac{1}{2} \theta_i n^k$ k -tuples contain an element of E_{i-1} . Recall that there are at most $10^{-4k^2} \epsilon^k n^k$ improper k -tuples and at most $10^{-3k^2} \epsilon^{k-1} n^{k-1} \cdot n = 10^{-3k^2} \epsilon^{k-1} n^k$ k -tuples that contain some abnormal $(k-1)$ -tuple. Since $10^{-4k^2} \epsilon^k + 10^{-3k^2} \epsilon^{k-1} < 10^{-2k^2-1} \epsilon < \theta_i/2$ we can find a proper k -tuple $K \in \mathcal{D}_i$ such that K has colour $t+i-1 \pmod{2}$ and for any $(k-1)$ -tuple $Y \subset K$ we have Y normal and $Y \notin E_{i-1}$.

Since $i > 0$, there is $x \in K \cap V_{\mathbb{S}}$. Let $Y = K - x$. Since Y is normal there are at most $2^{-3k} \epsilon n$ vertices y such that $Y \cup y$ is improper, and by definition of E_{i-1}

there are at most $2^{-2k}n$ points $y \in V_s$ such that $Y \cup y$ does not have colour $t+i-1 \pmod{2}$. Since $2^{-3k}\epsilon n + 2^{-2k}n < 2^{-k}(n-k)$ there is $y \in V_s$ such that $Y \cup y$ is proper and has colour $t+i-1 \pmod{2}$. Then $c_Y(x) = c_Y(y) = t+i-1 \pmod{2}$. But $x \in V_{\bar{s}}$ and $y \in V_s$, so $c_Y(x) + 1 - s_1 = c_Y(p_1)$ and $c_Y(y) + s_1 = c_Y(p_1)$, both mod 2. This is a contradiction, so we conclude that all but at most $\theta_i n^k$ k -tuples of \mathcal{D}_i have colour $t+i$.

Now partition V into 2 classes V_1, V_2 so that $V_s \subset V_1, V_{\bar{s}} \subset V_2$, and the other vertices are distributed arbitrarily. Incorrect $2k$ -tuples with respect to this partition belong to the one of the following three groups.

(i) Improperly coloured $2k$ -tuples. There are at most $10^{-5k^2} \epsilon^k n^{2k}$ of those.

(ii) Properly coloured $2k$ -tuples which use at least one vertex not in $V_s \cup V_{\bar{s}}$. There are at most $2^k 2^{-2k} \epsilon n \binom{n}{2k-1} < 2^{-k} \epsilon n^{2k}$ such $2k$ -tuples.

(iii) Properly coloured $2k$ -tuples which contain a k -tuple of \mathcal{D}_i with colour $t+i-1 \pmod{2}$. There are at most $\sum_{i=0}^k \theta_i n^k \binom{n}{k} < \theta_k n^{2k} = 10^{-2k^2} (2k 2^{2k})^k \epsilon n^{2k} < 10^{-k^2} \epsilon n^{2k}$ such $2k$ -tuples.

Therefore all but at most $(10^{-5k^2} \epsilon^k + 2^{-k} \epsilon + 10^{-k^2} \epsilon) n^{2k} < \epsilon n^{2k}$ $2k$ -tuples are correct with respect to this partition. This completes the proof of the theorem. ■

3.3. The Turán number of $\mathcal{C}_3^{(2k)}$

In this subsection we complete the proof of Frankl’s conjecture.

Proof of Theorem 1.1. Let H be a $2k$ -uniform hypergraph on n vertices, which has $e(H) \geq b_{2k}(n)$ and contains no $\mathcal{C}_3^{(2k)}$. By the same argument given in the proof in the case $k=2$ we can assume that H has minimum degree at least $b_{2k}(n) - b_{2k}(n-1)$. Applying Lemma 3.2 gives

$$(9) \quad \delta(H) \geq \frac{1}{2} \binom{n-1}{2k-1}.$$

For convenience of notation we set $\eta = (100k)^{-10^k}$. By Theorem 3.4 there is a partition with all but at most $(\eta/20k)^{2k} n^{2k}$ edges of H being good, i.e., they have odd intersection with both parts. Let $V(H) = V_1 \cup V_2$ be the partition which minimises the number of bad edges. Then every vertex belongs to at least as many good edges as bad edges, or we can move it to the other class of the partition. Recall that, by definition, the number of good $2k$ -tuples with respect to this partition is at most $b_{2k}(n)$. We must have $||V_1| - n/2| < \frac{1}{10} \eta n$

and $||V_2| - n/2| < \frac{1}{10}\eta n$. Otherwise by Lemma 3.1, part (v)

$$e(H) < \frac{1}{2} \binom{n}{2k} - \frac{1}{2} \left(2 \cdot \frac{1}{10}\eta n \right) + \left(10 \cdot \frac{1}{10}\eta \right)^k n^{2k-1} + (\eta/20k)^{2k} n^{2k} < b_{2k}(n),$$

which is a contradiction.

Note that there is no k -tuple of vertices P for which there are both $(10k)^{-k}\eta n^k$ k -tuples Q such that $P \cup Q$ is a good edge and $(10k)^{-k}\eta n^k$ k -tuples R such that $P \cup R$ is a bad edge. Indeed, each such Q and R which are disjoint give a $2k$ -tuple $Q \cup R$ which is good, but cannot be an edge as it would create a $\mathcal{C}_3^{(2k)}$. Moreover, every $2k$ -tuple can be obtained at most $\frac{1}{2} \binom{2k}{k}$ times in this way, and every Q is disjoint from all but at most $k \binom{n}{k-1}$ k -tuples R . Thus at least $(10k)^{-k}\eta n^k \left((10k)^{-k}\eta n^k - k \binom{n}{k-1} \right) / \left(\frac{1}{2} \binom{2k}{k} \right) > 2(\eta/20k)^{2k} n^{2k}$ good $2k$ -tuples are not edges of H , and therefore $e(H) < b_{2k}(n) - 2(\eta/20k)^{2k} n^{2k} + (\eta/20k)^{2k} n^{2k} < b_{2k}(n)$, which is a contradiction.

Claim 3.5. Any vertex of H is contained in at most ηn^{2k-1} bad edges.

Proof. Suppose some vertex a belongs to ηn^{2k-1} bad edges. Call a $(k-1)$ -tuple X good if there are at most $(10k)^{-k}\eta n^k$ k -tuples Q such that $a \cup X \cup Q$ is a bad edge, otherwise call X bad. By the above discussion, for every bad $(k-1)$ -tuple X there are at most $(10k)^{-k}\eta n^k$ k -tuples R such that $a \cup X \cup R$ is a good edge. There are at least ηn^{k-1} bad $(k-1)$ -tuples or we would only have $\eta n^{k-1} \cdot \binom{n}{k} + \left(\binom{n-1}{k-1} - \eta n^{k-1} \right) \cdot (10k)^{-k}\eta n^k < \eta n^{2k-1}$ bad edges through a .

Note that there are at most $\binom{n}{k-1} \cdot (10k)^{-k}\eta n^k$ good edges that contain a and a bad $(k-1)$ -tuple. To see this, we bound the number of such good edges by picking the bad $(k-1)$ -tuple X in at most $\binom{n}{k-1}$ ways and then a k -tuple R such that $a \cup X \cup R$ is a good edge in at most $(10k)^{-k}\eta n^k$ ways (as remarked above). By choice of partition there are at least as many good edges containing a as bad, so by (9) we see that a is in at least $\frac{1}{4} \binom{n-1}{2k-1}$ good edges. It follows that there are at least $\frac{1}{4} \binom{n-1}{2k-1} - (10k)^{-k}\eta n^{2k-1}$ good edges that contain only good $(k-1)$ -tuples. In particular there are at least $\left(\frac{1}{4} \binom{n-1}{2k-1} - (10k)^{-k}\eta n^{2k-1} \right) / \binom{n-1}{k} \geq n^{k-1} / (2k)!$ good $(k-1)$ -tuples.

Suppose there are $\alpha \binom{n}{k-1}$ good $(k-1)$ -tuples, where by the above we see that $(2k)^{-k-1} \leq \alpha \leq 1 - (k-1)!\eta$. We can bound the number of good edges that contain a and do not contain a bad $(k-1)$ -tuple as follows. Given any such edge W containing a we consider ordered triples (X, Y, b) , where X and Y are good $(k-1)$ -tuples, b is a vertex and $X \cup Y \cup b \cup a = W$. Each edge gives rise to $k \binom{2k-1}{k-1}$ such triples. On the other hand we can choose X and Y in at most $\left(\alpha \binom{n}{k-1} \right)^2$ ways; then to make E good b is constrained to lie in some

particular class V_i of the partition, so can be chosen in at most $(\frac{1}{2} + \frac{1}{10}\eta)n$ ways. This shows that the number of such edges W is at most

$$\left(\left(\alpha \binom{n}{k-1} \right)^2 \left(\frac{1}{2} + \frac{1}{10}\eta \right) n \right) / \left(k \binom{2k-1}{k-1} \right) < \left(\alpha^2 + 3 \cdot \frac{1}{10}\eta \right) \frac{1}{2} \binom{n-1}{2k-1}.$$

A similar argument shows that the number of bad edges that contain a and do not contain a good $(k-1)$ -tuple is at most $((1-\alpha)^2 + 3 \cdot \frac{1}{10}\eta) \frac{1}{2} \binom{n-1}{2k-1}$. We showed above that there are at most $\binom{n}{k-1} \cdot (10k)^{-k} \eta n^k$ good edges that contain a and a bad $(k-1)$ -tuple. Also, by definition there are at most $\binom{n}{k-1} \cdot (10k)^{-k} \eta n^k$ bad edges that contain a and a good $(k-1)$ -tuple. Therefore the total number of edges containing a is at most

$$\left(\alpha^2 + (1-\alpha)^2 + 6 \cdot \frac{1}{10}\eta \right) \frac{1}{2} \binom{n-1}{2k-1} + 2 \cdot \binom{n}{k-1} \cdot (10k)^{-k} \eta n^k.$$

From the bounds $(2k)^{-k-1} \leq \alpha \leq 1 - (k-1)!\eta$ we see that this is at most $(\frac{1}{2} - \eta/2) \binom{n-1}{2k-1}$. This contradicts equation (9), so the claim is proved. ■

Now write $|V_1| = n/2 + t$, $|V_2| = n/2 - t$ with $-\frac{1}{10}\eta n < t < \frac{1}{10}\eta n$. By possibly renaming the classes (i.e. replacing t with $-t$) we can assume that $d(n, t) \leq d(n, -t)$. Now any vertex of V_1 belongs to $d(n, t)$ good $2k$ -tuples, and $d(n, t)$ is the minimum degree of $\mathcal{B}^{(2k)}(n, t)$, which is certainly at most the maximum degree of $\mathcal{B}_n^{(2k)}$. From Lemma 3.1, part (iii) we have a bound $d(n, t) < \frac{1}{2} \binom{n-1}{2k-1} + (20k)^k n^{k-1/2}$ but we will only use the weaker bound $d(n, t) < \frac{1}{2} \binom{n-1}{2k-1} + 10^{4k^2} n^{k-1/2}$. Later we will show that this weaker bound also holds for $d(n, -t)$, and then the subsequent argument will apply switching V_1 and V_2 .

Claim 3.6. 1. If a is a vertex of V_1 for which K of the good $2k$ -tuples containing a are not edges then there are at least $K - 10^{4k^2} n^{k-1/2}$ bad edges containing a .

2. If b is a vertex of V_2 for which L of the good $2k$ -tuples containing b are not edges then there are at least $L - (\eta n/5)^{2k-1}$ bad edges containing b .

Proof. 1. By the preceding remarks a belongs to at most $\frac{1}{2} \binom{n-1}{2k-1} + 10^{4k^2} n^{k-1/2}$ good $2k$ -tuples and therefore it belongs to at most $\frac{1}{2} \binom{n-1}{2k-1} + 10^{4k^2} n^{k-1/2} - K$ good edges. Then by equation (9) a belongs to at least $K - 10^{4k^2} n^{k-1/2}$ bad edges.

2. From Lemma 3.1, part (v) b belongs to at most $\frac{1}{2} \binom{n-1}{2k-1} + \left(2 \cdot \frac{1}{10}\eta n - 1 \right)$ good $2k$ -tuples, and the stated bound follows as in (1). ■

Before proving the next claim we make a remark that will be used on several occasions without further comment. Suppose W is a bad edge, so that $|W \cap V_i|$ is even for $i=1,2$. If we partition $W = P \cup Q$ with $|P|=|Q|=k$ then $|P \cap V_i|=|Q \cap V_i| \pmod{2}$ for $i=1,2$. Then for any k -tuple R disjoint from P and Q with $|R \cap V_i|=|P \cap V_i|+1 \pmod{2}$ both $2k$ -tuples $P \cup R$ and $Q \cup R$ are good. We can obtain such a k -tuple $R \subset V - (P \cup Q)$ by picking any $(k-1)$ -tuple, and then another vertex which, because of parity, is constrained to lie in some particular V_i . This counts each k -tuple k times, so the number of choices for R is at least

$$k^{-1} \binom{n-2k}{k-1} \left(\left(\frac{1}{2} - \frac{1}{10} \eta \right) n - 3k \right) > n^k / (3 \cdot k!).$$

Claim 3.7. *Suppose $t \leq k$ and T is a t -tuple of vertices belonging to θn^{2k-t} bad edges, for some $\theta > (20k)^k \eta$. Then any $S \subset T$ with $|S|=t-1$ belongs to at least $(10k)^{-k} \theta n^{2k-t+1}$ good non-edges.*

Proof. Write $T = S \cup v$. Consider a bad edge W containing T and a partition $W = T \cup X \cup Y$, where $|X|=k-1$ and $|Y|=k+1-t$. By the above remark, there are at least $n^k / (3 \cdot k!)$ k -tuples R for which $v \cup X \cup R$ and $S \cup Y \cup R$ are both good $2k$ -tuples. Note that they can't both be edges, or we would have a copy of $C_3^{(2k)}$. Suppose that for at least $\frac{1}{2} \theta n^{2k-t}$ such W there is a partition $W = T \cup X \cup Y$ for which there are at least $n^k / 2(3 \cdot k!)$ k -tuples R for which $v \cup X \cup R$ is a good non-edge. This clearly gives at least $\frac{1}{2} \theta n^{k-1}$ choices for X . Each such non-edge can be partitioned in at most $\binom{2k-1}{k}$ ways in the form $v \cup X \cup R$, so there are at least

$$\binom{2k-1}{k}^{-1} \frac{1}{2} \theta n^{k-1} \frac{n^k}{2(3 \cdot k!)} > (10k)^{-k} \theta n^{2k-1}$$

good non-edges containing v . Now Claim 3.6 shows that there are at least

$$(10k)^{-k} \theta n^{2k-1} - (\eta n / 5)^{2k-1} > (20k)^{-k} \theta n^{2k-1} > \eta n^{2k-1}$$

bad edges containing v , which contradicts Claim 3.5. It follows that for at least $\frac{1}{2} \theta n^{2k-t}$ such W and any partition of $W = T \cup X \cup Y$ we have a good non-edge $S \cup Y \cup R$ for at least $n^k / 2(3 \cdot k!)$ k -tuples R . This gives at least $\frac{1}{2} \theta n^{k+1-t}$ choices for Y . Each such non-edge has at most $\binom{2k-t+1}{k}$ representations as $S \cup Y \cup R$, so there are at least

$$\binom{2k-t+1}{k}^{-1} \frac{1}{2} \theta n^{k+1-t} \frac{n^k}{2(3 \cdot k!)} > (10k)^{-k} \theta n^{2k-t+1}$$

good non-edges containing S . ■

Suppose for the sake of contradiction that there is some bad edge incident with V_1 . Denote the set of bad edges containing some vertex v by $\mathcal{Z}(v)$. Let a be a vertex in V_1 belonging to the maximum number of bad edges and let $Z = |\mathcal{Z}(a)|$. Note that $Z > 0$.

Claim 3.8. *Suppose $t \leq k$, $(20k)^k \eta < \phi < (100k)^{-k}$ and \mathcal{F} is a set of at least $\phi Z n^{-(2k-t)}$ t -tuples containing a such that each $F \in \mathcal{F}$ is contained in at least ϕn^{2k-t} bad edges. Then there are at least $\phi^5 Z n^{-(2k-t+1)}$ $(t-1)$ -tuples containing a each of which is contained in at least $\phi^5 n^{2k-t+1}$ bad edges.*

Proof. Let \mathcal{G} be the set of $(t-1)$ -tuples containing a that are contained in a member of \mathcal{F} . By Claim 3.7 each $G \in \mathcal{G}$ is contained in at least $(10k)^{-k} \phi n^{2k-t+1}$ good non-edges. Each such good non-edge is counted by at most $\binom{2k-1}{t-2}$ different G 's, so there are at least $\binom{2k-1}{t-2}^{-1} |\mathcal{G}| (10k)^{-k} \phi n^{2k-t+1} > (40k)^{-k} |\mathcal{G}| \phi n^{2k-t+1}$ good non-edges containing a . Since $a \in V_1$ Claim 3.6 gives at least

$$(40k)^{-k} |\mathcal{G}| \phi n^{2k-t+1} - 10^{4k^2} n^{k-1/2} > (50k)^{-k} |\mathcal{G}| \phi n^{2k-t+1}$$

bad edges containing a , so by definition of Z we get $|\mathcal{G}| < (50k)^k \phi^{-1} Z \cdot n^{-(2k-t+1)}$. Let $\mathcal{G}_1 \subset \mathcal{G}$ consist of those G that belong to at least $\phi^3 n$ members of \mathcal{F} . Then

$$\phi Z n^{-(2k-t)} \leq |\mathcal{F}| < |\mathcal{G}_1| n + |\mathcal{G}| \phi^3 n < |\mathcal{G}_1| n + (50k)^k \phi^2 Z n^{-(2k-t)}$$

so $|\mathcal{G}_1| > \phi^5 Z n^{-(2k-t-1)}$ with room to spare. For each $G \in \mathcal{G}_1$ there are at least $\phi^3 n$ sets of \mathcal{F} each contributing ϕn^{2k-t} bad edges containing G . Each such bad edge is counted by at most $2k-t+1$ different $F \in \mathcal{F}$, so G belongs to at least $(2k-t+1)^{-1} \phi^3 n \cdot \phi n^{2k-t} > \phi^5 n^{2k-t+1}$ bad edges. ■

Let $\mathcal{Z}_1(a)$ be those bad edges W containing a for which there is some partition into two k -tuples $W = P \cup Q$ with $a \in P$ so that there are at least $n^k/2(3 \cdot k!)$ k -tuples R for which $P \cup R$ is a good non-edge. Let $\mathcal{Z}_2(a) = \mathcal{Z}(a) - \mathcal{Z}_1(a)$, and write $Z_i = |\mathcal{Z}_i(a)|$ for $i = 1, 2$. Then one of Z_1, Z_2 is at least $Z/2$.

Case 1. Suppose $Z_1 \geq Z/2$. Let \mathcal{P} be the (non-empty) set of k -tuples P containing a such that there is some edge $P \cup Q$ in $\mathcal{Z}_1(a)$, and $P \cup R$ is a good non-edge for at least $n^k/2(3k!)$ k -tuples R . Each such good non-edge is counted by at most $\binom{2k-1}{k-1}$ different P 's, so there are at least $\binom{2k-1}{k-1}^{-1} |\mathcal{P}| n^k/2(3 \cdot k!) > (10k)^{-k} |\mathcal{P}| n^k$ good non-edges containing a . Now

Claim 3.6 gives at least $(10k)^{-k}|\mathcal{P}|n^k - 10^{4k^2}n^{k-1/2} > (20k)^{-k}|\mathcal{P}|n^k$ bad edges containing a , so by definition of Z , $|\mathcal{P}| < (20k)^k Zn^{-k}$. On the other hand, let $\mathcal{P}_1 \subset \mathcal{P}$ consist of those P that belong to at least $\frac{1}{10}(20k)^{-k}n^k$ bad edges. Then

$$Z/2 \leq Z_1 < |\mathcal{P}_1|n^k + |\mathcal{P} \setminus \mathcal{P}_1| \frac{1}{10}(20k)^{-k}n^k < |\mathcal{P}_1|n^k + Z/10$$

so $|\mathcal{P}_1| > 0.4 Zn^{-k}$. Now apply **Claim 3.8** $k-1$ times, starting with $t=k$ and $\phi = (100k)^{-k}$. We deduce that a belongs to at least $\phi^{5^{k-1}}n^{2k-1} > \eta n^{2k-1}$ bad edges, which contradicts **Claim 3.5**.

Case 2. Now suppose $Z_2 \geq Z/2$. Note that every bad edge containing a contains at least one other point of V_1 , so there is some $b \in V_1$ belonging to at least Z_2/n edges of $\mathcal{Z}_2(a)$. Fix one such b . Let \mathcal{X} be the set of $(k-1)$ -tuples X for which there exists a $(k-1)$ -tuple Y such that $W = a \cup b \cup X \cup Y$ is an edge of $\mathcal{Z}_2(a)$. By definition of $\mathcal{Z}_2(a)$ for any such partition of W , there are at least $n^k/2(3 \cdot k!)$ k -tuples R such that $b \cup X \cup R$ is a good non-edge. This gives at least $\frac{n^k}{2(3 \cdot k!)}|\mathcal{X}| > (10k)^{-k}|\mathcal{X}|n^k$ good non-edges containing b , and since $b \in V_1$ **Claim 3.6** gives at least $(10k)^{-k}|\mathcal{X}|n^k - 10^{4k^2}n^{k-1/2} > (20k)^{-k}|\mathcal{X}|n^k$ bad edges containing b . Thus, by definition of Z , $|\mathcal{X}| < (20k)^k Zn^{-k}$. Note that each edge in $\mathcal{Z}_2(a)$ that contains b is obtained by picking a pair of $(k-1)$ -tuples in \mathcal{X} , so $Z/(2n) \leq Z_2/n \leq \binom{|\mathcal{X}|}{2} < \frac{1}{2}(20k)^{2k} Z^2 n^{-2k}$. Therefore $Z > (20k)^{-2k}n^{2k-1} > \eta n^{2k-1}$, which contradicts **Claim 3.5**.

We conclude that there are no bad edges incident to the vertices of V_1 , i.e. all bad edges are entirely contained in V_2 . As in the case $k = 2$ this gives a more precise bound on t , defined by $|V_1| = n/2 + t$, $|V_2| = n/2 - t$. If $|t| \geq 20^k \sqrt{n}$ then **Lemma 3.1**, part (iv) gives $d_{2k}(n, t) < \frac{1}{2} \binom{n-1}{2k-1} - 20^k n^{k-1/2}$. This is a contradiction, since the vertices of V_1 only belong to good edges, of which there are at most $d_{2k}(n, t) < \delta(H)$. Therefore $|t| < 20^k \sqrt{n}$. Now **Lemma 3.1**, part (ii) gives

$$d_{2k}(n, -t) < \frac{1}{2} \binom{n-1}{2k-1} + (10(20k)^2)^k n^{k-1/2} < \frac{1}{2} \binom{n-1}{2k-1} + 10^{4k^2} n^{k-1/2}.$$

As we remarked earlier, this bound allows us to repeat the above argument interchanging V_1 and V_2 , so we deduce that there are no bad edges incident with V_2 either, i.e. all edges are good. Then by definition of $b_{2k}(n)$ we have $e(H) \leq b_{2k}(n)$, with equality only when H is a $\mathcal{B}_n^{(2k)}$, so the theorem is proved. ■

4. Hypergraphs without $\mathcal{C}_r^{(4)}$

We recall that $\mathcal{C}_r^{(2k)}$ is the $2k$ -uniform hypergraph obtained by letting P_1, \dots, P_r be pairwise disjoint sets of size k and taking as edges all sets $P_i \cup P_j$ with $i \neq j$. In this section we will be concerned with the case $k=2$ and general r .

Sidorenko [8] showed that the Turán density of $\mathcal{C}_r^{(2k)}$ is at most $\frac{r-2}{r-1}$. This is a consequence of Turán’s theorem applied to an auxiliary graph G constructed from a $2k$ -uniform hypergraph H of order n . The vertices of G are the k -tuples of vertices of H , and two k -tuples P_1, P_2 are adjacent if $P_1 \cup P_2$ is an edge of H . It is easy to see that the graph G has $\binom{n}{k}$ vertices, $\frac{1}{2} \binom{2k}{k} e(H)$ edges and contains no K_r . Thus the upper bound on the number of edges of H follows immediately from Turán’s theorem. The following construction from [8] gives a matching lower bound when r is of the form $2^p + 1$.

Let W be a vector space of dimension p over the field $GF(2)$, i.e. the finite field with 2 elements $\{0, 1\}$. Partition a set of vertices V as $\bigcup_{w \in W} V_w$, $|V_w| = |V|/(r-1)$. Given t and a t -tuple of vertices $X = x_1 \cdots x_t$ with $x_i \in V_{w_i}$ we define $\Sigma X = \sum_1^t w_i$. Define a $2k$ -uniform hypergraph H , where a $2k$ -tuple X is an edge iff $\Sigma X \neq 0$. Observe that this doesn’t contain a copy of $\mathcal{C}_r^{(2k)}$. Indeed, if P_1, \dots, P_r are disjoint k -tuples then there is some $i \neq j$ with $\Sigma P_i = \Sigma P_j$ (by the pigeonhole principle). Then $\Sigma(P_i \cup P_j) = \Sigma P_i + \Sigma P_j = 0$, so $P_i \cup P_j$ is not an edge.

This construction depends essentially on an algebraic structure, which only exists for certain values of r . Perhaps surprisingly, we will show that this is an intrinsic feature of the problem, by proving [Theorem 1.2](#), which gives a stronger upper bound on the Turán density of $\mathcal{C}_r^{(4)}$, when r is not of the form $2^p + 1$. We make no attempt to optimize the constant in this bound.

In addition, our proof of this theorem implies that, for $r = 2^p + 1$, any $\mathcal{C}_r^{(4)}$ -free 4-uniform hypergraph with density $\frac{r-2}{r-1} - o(1)$ looks approximately like Sidorenko’s construction.

Corollary 4.1. *Let $r = 2^p + 1$ be an integer and let W be a p -dimensional vector space over the field $GF(2)$. For every $\epsilon > 0$ there is $\eta > 0$ so that if H is a $\mathcal{C}_r^{(4)}$ -free 4-uniform hypergraph with $e(H) > \frac{r-2}{r-1} \binom{n}{4} - \eta n^4$ then there is a partition of the vertex set as $\bigcup_{w \in W} V_w$ such that all but ϵn^4 edges X of H satisfy $\Sigma X \neq 0$.*

The rest of this section is organized as follows. In the [first subsection](#) we will prove a lemma showing that certain edge-colourings of the complete graph K_s exist only if s is a power of 2. In the [following subsection](#) we will

recall a proof of the Simonovits stability theorem so that we can calculate some explicit constants. The [final subsection](#) contains the proof of [Theorem 1.2](#).

4.1. A lemma on edge-colourings of a complete graph

Lemma 4.2. *Suppose that we have a colouring of the edges of the complete graph K_s in $s-1$ colours, so that every colour is a matching and each subset of 4 vertices spans edges of either 3 or 6 different colours. Then $s = 2^p$ for some integer p .*

Proof. Since the number of colours is $s-1$, every colour is a matching and the total number of edges in K_s is $s(s-1)/2$ it is easy to see that every colour is a perfect matching. Also, if wx and yz are disjoint edges of the same colour, then by hypothesis only 3 different colours appear on $wxyz$, so wy and xz have the same colour, as do xy and wz . Denote the set of colours by $C = \{c_1, \dots, c_{s-1}\}$. We define a binary operation $+$ on C using the following rule. Pick a vertex x . Given c_i and c_j let $e_i = xy_i$ and $e_j = xy_j$ be the edges incident with x with these colours. These edges exist, as each colour is a perfect matching. Define $c_i + c_j$ to be the colour of $y_i y_j$.

To see that this is well-defined, let x' be another vertex and suppose $e'_i = x'y'_i$ has colour c_i and $e'_j = x'y'_j$ has colour c_j . If $y_i = y'_j$ then opposite edges of $xy_j y_i x'$ have the same colours, so $x'y_j$ has colour c_i , i.e. $y_j = y'_i$ and there is nothing to prove. Therefore we can assume that all y_i, y_j, y'_i, y'_j are distinct. Consider the 4-tuple $xx'y_i y'_i$. Since xy_i and xy'_i have the same colour we deduce that xx' and $y_i y'_i$ have the same colour. Similarly xx' and $y_j y'_j$ have the same colour, from which we see that $y_i y'_i$ and $y_j y'_j$ have the same colour. Now looking at $y_i y'_i y_j y'_j$ we see that $y_i y_j$ and $y'_i y'_j$ have the same colour, so $c_i + c_j$ is well-defined.

Let D be a set obtained by adjoining another element called $\mathbf{0}$ to C . Extend $+$ to an operation on D by defining $\mathbf{0} + d = d + \mathbf{0} = d$ and $d + d = \mathbf{0}$ for all $d \in D$. We claim that $(D, +)$ is an abelian group. Note that $+$ is commutative by definition, $\mathbf{0}$ is an identity and inverses exist. It remains to show associativity, i.e. for any d_1, d_2, d_3 we have $(d_1 + d_2) + d_3 = d_1 + (d_2 + d_3)$. This is immediate if any of the d_i are $\mathbf{0}$ or if they are all equal. If $d_1 = d_2 \neq d_3$ then $d_1 + d_2 = \mathbf{0}$ and there is a triangle with colours $d_1, d_3, d_1 + d_3$, so $d_1 + (d_2 + d_3) = d_3$ as required. The same argument applies when $d_2 = d_3 \neq d_1$. If $d_1 = d_3$ then $d_1 + d_2 = d_2 + d_3$ by commutativity, and so $(d_1 + d_2) + d_3 = d_1 + (d_2 + d_3)$ also by commutativity. So we can assume that the d_i are pairwise distinct and non-zero. Pick a vertex x , let xy_1 be the edge of colour d_1 and xy_2 the

edge of colour d_2 . Let y_2z have colour d_3 . We can suppose $z \neq y_1$, otherwise $d_1 + d_2 = d_3$ and $d_2 + d_3 = d_1$ and $(d_1 + d_2) + d_3 = d_1 + (d_2 + d_3) = \mathbf{0}$. Now y_1y_2 has colour $d_1 + d_2$ and xz has colour $d_2 + d_3$. Consider the edge y_1z . From the triangle it forms with x we see that it has colour $d_1 + (d_2 + d_3)$ and from the triangle with y_2 we see that it has colour $(d_1 + d_2) + d_3$. This proves associativity, so D is an abelian group.

Finally, note that every non-zero element has order 2, so D is in fact a vector space over the field with 2 elements. If p is its dimension then $s = |D| = 2^p$. ■

4.2. The Simonovits stability theorem

In this subsection we will recall a proof of the Simonovits stability theorem [10] so that we can calculate some explicit constants. Let $T_s(N)$ be the s -partite Turán graph on N vertices, i.e. a complete s -partite graph with part sizes as equal as possible. Write $t_s(N)$ for the number of edges in $T_s(N)$. Then Turán’s theorem states that any K_{s+1} -free graph on N vertices has at most $t_s(N)$ edges, with equality only for $T_s(N)$. It is easy to show that $\frac{s-1}{s}N^2/2 - s < t_s(N) \leq \frac{s-1}{s}N^2/2$.

Proposition 4.3. *Suppose G is a K_{s+1} -free graph on N vertices with minimum degree $\delta(G) \geq (1 - \frac{1}{s} - \alpha)N$ and $\alpha < 1/s^2$. Then there is a partition of the vertex set of G as $V(G) = U_1 \cup \dots \cup U_s$ with $\sum e(U_i) < \alpha N^2$.*

Proof. By Turán’s theorem G contains a copy of K_s ; let $A = \{a_1, \dots, a_s\}$ be its vertex set. Note that any vertex x not in A has at most $s - 1$ neighbours in A , or we get a K_{s+1} . Let B be those vertices with exactly $s - 1$ neighbours in A , and $C = V(G) - A - B$. Partition $A \cup B$ as $U_1 \cup \dots \cup U_s$ where U_i consists of those vertices adjacent to $A - a_i$. Then there are no edges inside any U_i , as if xy is such an edge then $xy + A - a_i$ forms a K_{s+1} . Distribute the vertices of C arbitrarily among the U_i . Counting edges between A and $V - A$ gives

$$s(\delta(G) - s + 1) \leq e(A, V - A) \leq (s - 1)|B| + (s - 2)|C| = (s - 1)(N - s) - |C|$$

so $|C| \leq (s - 1)N - s\delta(G) \leq \alpha N$. Therefore $\sum e(U_i) < \alpha N^2$. ■

Theorem 4.4. *Suppose G is a K_{s+1} -free graph on N vertices with at least $(\frac{s-1}{2s} - c)N^2$ edges and $c < 1/(4s^4)$. Then there is a partition of the vertex set of G as $V(G) = U_1 \cup \dots \cup U_s$ with $\sum e(U_i) < (2s + 1)\sqrt{c} N^2$.*

Proof. Construct a sequence of graphs $G = G_N, G_{N-1}, \dots$ where if G_m has a vertex of degree at most $(1 - \frac{1}{s} - 2\sqrt{c})m$ then we delete it to get G_{m-1} .

Suppose we can delete $\sqrt{c} N$ vertices by this process and reach a graph $G_{(1-\sqrt{c})N}$. Then $G_{(1-\sqrt{c})N}$ is K_{s+1} -free and has at least

$$\left(\frac{s-1}{2s} - c - \sqrt{c}\left(1 - \frac{1}{s} - 2\sqrt{c}\right)\right) N^2 > \frac{s-1}{2s}(1 - \sqrt{c})^2 N^2$$

edges. This contradicts Turán’s theorem, so the sequence terminates at some G_m with $m \geq (1 - \sqrt{c})N$ and minimum degree at least $(1 - \frac{1}{s} - 2\sqrt{c})m$. By Proposition 4.3 there is a partition $V(G_m) = U_1 \cup \dots \cup U_s$ with $\sum e(U_i) < 2s\sqrt{c} N^2$. Now distribute the $\sqrt{c}N$ deleted vertices arbitrarily among the U_i . Then $\sum e(U_i) < (2s+1)\sqrt{c} N^2$. ■

4.3. Proof of Theorem 1.2

Let V be the vertex set of H . Define a graph G whose vertices are all pairs in V , where the pairs ab and cd are adjacent exactly when $abcd$ is an edge of H . Since H is $C_r^{(4)}$ -free we see that G is K_r -free. Also, each edge of H creates exactly 3 edges in G (corresponding to the 3 ways of breaking a 4-tuple into pairs) so

$$e(G) > 3\left(\frac{r-2}{r-1} - 10^{-33}r^{-70}\right) \binom{n}{4} > \left(\frac{r-2}{2(r-1)} - 10^{-33}r^{-70}\right) N^2,$$

where $N = \binom{n}{2}$.

Applying Theorem 4.4 with $s = r - 1$ gives a partition of the pairs of vertices in V as $\bigcup_1^{r-1} P_i$ with $\sum_1^{r-1} e(P_i) < 10^{-16}r^{-34}N^2$. If there is some P_i with $|P_i| < (\frac{1}{r-1} - 10^{-3}r^{-7})N$ then

$$\begin{aligned} \frac{e(G)}{N^2} &< \frac{\binom{r-2}{2}}{(r-2)^2} \left(\frac{r-2}{r-1} + 10^{-3}r^{-7}\right)^2 \\ &\quad + \left(\frac{1}{r-1} - 10^{-3}r^{-7}\right) \left(\frac{r-2}{r-1} + 10^{-3}r^{-7}\right) + 10^{-16}r^{-34} \\ &< \frac{r-2}{2(r-1)} - 10^{-6}r^{-14}/2 + 10^{-16}r^{-34}. \end{aligned}$$

This is a contradiction so $|P_i| \geq (\frac{1}{r-1} - 10^{-3}r^{-7})N$ for all i . Also if some $|P_i| > (\frac{1}{r-1} + 10^{-3}r^{-6})N$, then there is j such that $|P_j| < (\frac{1}{r-1} - 10^{-3}r^{-7})N$. Therefore for all i

$$(10) \quad \left| |P_i| - \frac{1}{r-1}N \right| \leq 10^{-3}r^{-6}n^2.$$

Note that all but at most $10^{-16}r^{-34}n^4$ edges of H are formed by taking a pair from P_i and a pair from P_j with $i \neq j$. We think of the P_i as a colouring of pairs. A 4-tuple $abcd$ will be called *properly coloured* if either

- (i) $abcd$ is an edge and each of the 3 sets $\{ab, cd\}, \{ac, bd\}, \{ad, bc\}$ contains two pairs with different colours, or
- (ii) $abcd$ is not an edge and each of the 3 sets $\{ab, cd\}, \{ac, bd\}, \{ad, bc\}$ consists of two pairs with the same colour.

An improperly coloured 4-tuple is either an edge that is the union of two pairs of the same colour or a non-edge which is the union of two pairs with different colours. There are at most $10^{-16}r^{-34}N^2$ of the former 4-tuples, and the number of latter is at most

$$\frac{r-2}{2(r-1)}N^2 - (e(G) - 10^{-16}r^{-34}N^2) \leq (10^{-16}r^{-34} + 10^{-33}r^{-70})n^4.$$

Therefore all but $10^{-15}r^{-34}n^4$ 4-tuples are properly coloured. Call a pair ab *bad* if there are at least $10^{-12}r^{-32}n^2$ pairs cd such that $abcd$ is improperly coloured; otherwise call it *good*. Then there are at most $\binom{4}{2}(10^{-15}r^{-34}n^4)/(10^{-12}r^{-32}n^2) < 10^{-2}r^{-2}n^2$ bad pairs.

Consider a graph on V whose edges are the pairs in P_1 . As noted in (10) it has at least $\frac{1}{r-1}N - 10^{-3}r^{-6}n^2$ edges. For vertices a and b in V , let $d(a)$ denote the degree of a and $d(a, b)$ the *codegree* of a and b (i.e. the size of their common neighbourhood). Then

$$\sum_{a, b \in V} d(a, b) = \sum_{c \in V} \binom{d(c)}{2} \geq n \binom{\sum d(c)/n}{2} = n \binom{2|P_1|/n}{2} > \frac{1}{5r^2}nN.$$

Suppose there are at most m pairs (a, b) for which $d(a, b) > \frac{n}{10r^2}$. Then $\frac{1}{5r^2}nN < \sum d(a, b) \leq mn + N\frac{n}{10r^2}$, so $\frac{1}{10r^2}N < m$, i.e. there are at least $\frac{1}{10r^2}\binom{n}{2}$ pairs (a, b) for which $d(a, b) > \frac{n}{10r^2}$. At least one such pair is good, as the number of bad pairs is at most $10^{-2}r^{-2}n^2 < \frac{1}{20r^2}n^2$. Let (a, b) be such a pair and suppose it belongs to P_t .

Let B be the set of pairs cd for which $abcd$ is improperly coloured. Since ab is good we have $|B| \leq 10^{-12}r^{-32}n^2$. Therefore there are at most $|B|\binom{n}{2} < 10^{-12}r^{-32}n^4$ 4-tuples of vertices that contain any pair of B . We will call a 4-tuple *normal* if it is properly coloured and does not contain a pair from B ; otherwise we call it *abnormal*. Then all but at most $10^{-12}r^{-32}n^4 + 10^{-15}r^{-34}n^4 < 10^{-11}r^{-32}n^4$ 4-tuples are normal.

Partition the vertices of $V - ab$ into $(r-1)^2$ sets U_{ij} , where c is in U_{ij} iff $ac \in P_i$ and $bc \in P_j$. Then by the above discussion $|U_{11}| \geq \frac{n}{10r^2}$. Now we claim that for $i \neq j$ we have $|U_{ij}| < 10^{-3}r^{-11}n$. For suppose that $|U_{ij}| \geq$

$10^{-3}r^{-11}n$. Let P_k be the colour that appears most frequently among pairs joining vertices of U_{11} to U_{ij} . Then there are at least $\frac{1}{r-1}|U_{11}||U_{ij}|$ pairs of colour P_k with one endpoint in U_{11} and the other in U_{ij} . Consider the 4-tuples of the form $c_1c_2d_1d_2$, with $c_1, c_2 \in U_{11}$, $d_1, d_2 \in U_{ij}$ and $c_1d_1, c_2d_2 \in P_k$. There are at least

$$\begin{aligned} \left(\frac{1}{r-1}|U_{11}||U_{ij}|\right) \left(\frac{1}{r-1}|U_{11}||U_{ij}| - 2n\right) / 4 &> r^{-2}(10^{-1}r^{-2}n \cdot 10^{-3}r^{-11}n)^2 / 4 \\ &> 10^{-11}r^{-32}n^4 \end{aligned}$$

such 4-tuples, so some $c_1c_2d_1d_2$ is normal. By definition of normality each of its pairs forms a properly coloured 4-tuple with ab . Since ac_1 and bc_2 are in P_1 and ab is in P_t we deduce that c_1c_2 is in P_t as well. Also $ad_1 \in P_i$, $bd_2 \in P_j$ and $i \neq j$, so d_1d_2 cannot be in P_t . But c_1d_1 and c_2d_2 both belong to P_k so $c_1c_2d_1d_2$ is improperly coloured. This contradicts the definition of normality, so we do have $|U_{ij}| < 10^{-3}r^{-11}n$.

For convenience write $U_i = U_{ii}$. Then all but at most $(r-1)^2 10^{-3}r^{-11}n \leq 10^{-3}r^{-9}n$ vertices belong to one of the U_i . Suppose cd is a pair such that $abcd$ is properly coloured. Since ab is a good pair, this is the case for all but at most $10^{-11}r^{-32}n^2$ pairs cd . If c and d both belong to some U_i then ac and bd both have colour i . Since ab has colour t we see that cd has colour t . Similarly, if $c \in U_i$ and $d \in U_j$ with $i \neq j$ we see that cd cannot have colour t .

Let E_i denote the pairs with both endpoints in U_i , so that $|E_i| = \binom{|U_i|}{2}$. By the above discussion, all but at most $10^{-12}r^{-32}n^2$ pairs in $\cup_i E_i$ belong to P_t . Suppose $|U_i| < (\frac{1}{r-1} - 10^{-1}r^{-3})n$ for some i , so that

$$\begin{aligned} \sum |E_i| &> \left(\frac{1}{r-1} - 10^{-1}r^{-3}\right)^2 \frac{n^2}{2} \\ &\quad + (r-2) \left(\frac{1 - 1/(r-1) + 10^{-1}r^{-3} - 10^{-3}r^{-9}}{r-2}\right)^2 \frac{n^2}{2} - O(n) \\ &> \left(\frac{1}{r-1} + 10^{-2}r^{-6}\right) \frac{n^2}{2} - O(n). \end{aligned}$$

By (10), this gives the following contradiction.

$$\begin{aligned} \frac{1}{r-1} \binom{n}{2} + 10^{-3}r^{-6}n^2 &\geq |P_t| \geq \sum_i |E_i| - 10^{-12}r^{-32}n^2 \\ &> \frac{1}{r-1} \frac{n^2}{2} + \frac{10^{-2}r^{-6}}{3} n^2. \end{aligned}$$

Therefore $|U_i| \geq (\frac{1}{r-1} - 10^{-1}r^{-3})n$ for each i .

Let E_{ij} denote the edges with one endpoint in U_i and the other in U_j . We claim that one colour is dominant among these edges, i.e. there is some q such that all but $10^{-2}r^{-4}n^2$ edges of E_{ij} belong to P_q . Indeed, suppose that there are colours q_1 and q_2 for which E_{ij} contains at least $10^{-2}r^{-4}n^2$ edges of colour q_1 and at least $10^{-2}r^{-4}n^2$ edges of colour q_2 . Then there are at least $(10^{-2}r^{-4}n^2)(10^{-2}r^{-4}n^2 - 2n) > 10^{-11}r^{-32}n^4$ 4-tuples $c_1c_2d_1d_2$ with c_1, c_2 in U_i , d_1, d_2 in U_j , c_1d_1 of colour q_1 and c_2d_2 of colour q_2 . At least one such 4-tuple $c_1c_2d_1d_2$ is normal, since there at most $10^{-11}r^{-32}n^4$ abnormal 4-tuples. But then c_1c_2 and d_1d_2 both have colour t , so $c_1c_2d_1d_2$ is improperly coloured, which is a contradiction.

Consider the complete graph K_{r-1} on the vertex set $\{1, \dots, r-1\}$ and colour edge ij with the dominant colour of E_{ij} . We show that this colouring satisfies the hypotheses of Lemma 4.2. First of all we show that colour t doesn't occur in this edge-colouring of K_{r-1} , i.e. there are only $r-2$ colours. Suppose ij has colour t . Then

$$\begin{aligned} & \frac{1}{r-1} \binom{n}{2} + 10^{-3}r^{-6}n^2 \\ & \geq |P_t| \geq \sum_i |E_i| - 10^{-12}r^{-32}n^2 + |U_i||U_j| - 10^{-2}r^{-4}n^2 \\ & \geq (r-1) \binom{(\frac{1}{r-1} - 10^{-1}r^{-3})n}{2} + \left(\left(\frac{1}{r-1} - 10^{-1}r^{-3} \right) n \right)^2 - 10^{-1}r^{-4}n^2 \\ & \geq \frac{1}{r-1} \frac{n^2}{2} + \frac{n^2}{(r-1)^2} - \frac{r+2}{r-1} 10^{-1}r^{-3}n^2 > \frac{1}{r-1} \frac{n^2}{2} + \frac{n^2}{r^2}, \end{aligned}$$

is a contradiction.

Now suppose that some colour ℓ is not a matching, i.e. there are edges ij and ik in K_{r-1} both of colour ℓ . Then all but at most $2 \cdot 10^{-2}r^{-4}n^2$ pairs of $E_{ij} \cup E_{ik}$ have colour ℓ . Consider the 4-tuples of the form c_1c_2de , with $c_1, c_2 \in U_i$, $d \in U_j$ and $e \in U_k$, such that c_1d, c_2d, c_1e, c_2e all have colour ℓ . There are at least

$$\begin{aligned} & \binom{|U_i|}{2} |U_j||U_k| - 2 \cdot 10^{-2}r^{-4}n^2 \binom{n}{2} \\ & \geq \frac{1}{2} \left(\frac{1}{r-1} - 10^{-1}r^{-3} \right)^4 n^4 - O(n^3) - 10^{-2}r^{-4}n^4 > 10^{-11}r^{-32}n^4 \end{aligned}$$

such 4-tuples, so there is one such c_1c_2de which is normal. But then c_1c_2 has colour t and de cannot have colour t , since by normality $abde$ is properly

coloured. Therefore c_1c_2de is improperly coloured. This is a contradiction, so each colour forms a matching.

It remains to show that if some 4 vertices $x_1x_2x_3x_4$ in K_{r-1} do not span 6 different colours then they span only 3 colours. Suppose that x_1x_2 and x_3x_4 have colour α , x_1x_3 has colour β and x_2x_4 has colour γ . Recall that all but at most $10^{-2}r^{-4}n^2$ pairs in $E_{x_ix_j}$ have the corresponding colour of x_ix_j . Consider the 4-tuples in H of the form $c_1c_2c_3c_4$ with $c_i \in U_{x_i}$ such that c_ic_j has the same colour as x_ix_j . There are at least

$$\prod_1^4 |U_{x_i}| - 4 \cdot 10^{-2}r^{-4}n^2 \binom{n}{2} > \left(\frac{1}{r-1} - 10^{-1}r^{-3} \right)^4 n^4 - 2 \cdot 10^{-2}r^{-4}n^4 \\ > 10^{-11}r^{-32}n^4$$

such 4-tuples. Since the the number of abnormal 4-tuples is at most $10^{-11}r^{-32}n^4$, some such $c_1c_2c_3c_4$ should be normal. Then $\beta = \gamma$, or $c_1c_2c_3c_4$ would be improperly coloured. We see that opposite edges of $x_1x_2x_3x_4$ have the same colour. Therefore we can apply [Lemma 4.2](#) with $s = r - 1$ to deduce that $r - 1$ is of the form 2^p . ■

Finally it is not difficult to check that, when $r = 2^p + 1$, the above arguments together with the proof of [Lemma 4.2](#) imply [Corollary 4.1](#).

5. Concluding remarks

Among the various techniques that we used in this paper, the stability approach stands out as one that should be widely applicable in extremal combinatorics. The process of separating the argument into a stability stage and a refinement stage focuses attention on the particular difficulties of each, and often leads to progress where the raw problem has appeared intractable. For recent examples we refer to our proofs of the conjecture of Sós on the Turán number of the Fano plane [4], and a conjecture of Erdős and Rothschild on edge colourings with no monochromatic cliques [1].

Our methods probably apply to $\mathcal{C}_r^{(2k)}$ for general k when r is of the form $2^p + 1$, although the reader who has grappled with the thornier aspects of this paper will note the formidable technical difficulties that would arise. It would be far more interesting to say more about the behaviour of the Turán density of $\mathcal{C}_r^{(2k)}$ for general r . Even $\mathcal{C}_4^{(4)}$ presents an enigma for which there is no obvious plausible conjecture. We find it remarkable that the seemingly similar hypergraphs $\mathcal{C}_3^{(4)}$ and $\mathcal{C}_5^{(4)}$ are actually distinguished from $\mathcal{C}_4^{(4)}$ by a hidden algebraic feature, so are loathe even to speculate on the nature of the best construction for this case.

References

- [1] N. ALON, J. BALOGH, P. KEEVASH and B. SUDAKOV: The number of edge colourings with no monochromatic cliques, *J. London Math. Soc.*, to appear.
- [2] P. FRANKL: Asymptotic solution of a Turán-type problem, *Graphs and Combinatorics* **6** (1990), 223–227.
- [3] Z. FÜREDI: Turán type problems, in: *Surveys in combinatorics*, London Math. Soc. Lecture Note Ser. **166**, Cambridge Univ. Press, Cambridge, 1991, 253–300.
- [4] P. KEEVASH and B. SUDAKOV: The Turán number of the Fano plane, *Combinatorica* **25(5)** (2005), 561–574.
- [5] I. KRASIKOV and S. LITSYN: Survey of binary Krawtchouk polynomials, in: *Codes and association schemes*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci. **56**, Amer. Math. Soc., Providence, RI, 2001, 199–211.
- [6] V. LEVENSHTEN: Krawtchouk polynomials and universal bounds for codes and designs in Hamming spaces, *IEEE Trans. Inform. Theory* **41** (1995), 1303–1321.
- [7] L. LOVÁSZ: *Combinatorial Problems and Exercises*, North-Holland, Amsterdam, 1993.
- [8] A. SIDORENKO: An analytic approach to extremal problems for graphs and hypergraphs, in: *Extremal problems for finite sets*, Bolyai Soc. Math. Stud. **3**, Budapest, 1994, 423–455.
- [9] A. SIDORENKO: What we know and what we do not know about Turán numbers, *Graphs and Combinatorics* **11** (1995), 179–199.
- [10] M. SIMONOVITS: A method for solving extremal problems in graph theory, stability problems; in: *Theory of Graphs (Proc. Colloq. Tihany, 1966)*, Academic Press, New York, and Akad. Kiadó, Budapest, 1968, 279–319.

Peter Keevash

Department of Mathematics
Princeton University
Princeton, NJ 08540
USA

keevash@math.princeton.edu

Benny Sudakov

Department of Mathematics
Princeton University
Princeton, NJ 08540
USA

bsudakov@math.princeton.edu