

Pacific Journal of Mathematics

ON A LEMMA OF BISHOP AND PHELPS

ARNE BRØNSTED

ON A LEMMA OF BISHOP AND PHELPS

ARNE BRØNDSTED

The main purpose of the present note is to establish a theorem on the existence of maximal elements in certain partially ordered uniform spaces. The theorem unifies a lemma of E. Bishop and R. R. Phelps and a number of known extensions of this lemma.

1. In the proof of the fundamental theorem of E. Bishop and R. R. Phelps [1, Theorem 1] on the density of the set of support points of a closed convex subset of a Banach space, a lemma [1, Lemma 1] on the existence of maximal elements in certain partially ordered complete subsets of a normed linear space played a central role. Subsequently, this lemma was extended for various other purposes. Since none of the extensions is sufficiently general to cover all of them, it seems natural to look for a common generalization. In §2 we shall present such a general theorem. Actually, we first prove a theorem (Theorem 1) which is too general to be directly applicable, at least in the present context, and next we apply it to obtain the desired theorem (Theorem 2). The theorems are formulated in terms of uniform structures, due to the facts that completeness is the crucial assumption and that both non-metric topological linear spaces and metric nonlinear spaces are to be covered. The proofs are heavily influenced by known arguments. In §3 we shall discuss the relations of Theorem 2 to the Bishop-Phelps lemma and its extensions. Finally, in §4 we shall give a simplified proof of a recent result of J. Daneš by applying the Bishop-Phelps procedure.

2. Everywhere in the following E is assumed to be a non-empty set. By an *extended real valued function* on E we shall mean a mapping $\varphi: E \rightarrow]-\infty, +\infty]$, not identically $+\infty$. The set of points $x \in E$ such that $\varphi(x) < +\infty$ is denoted $\text{dom } \varphi$. When E is equipped with a uniformity \mathcal{U} , and φ is an extended real valued function on E , we shall say that φ is *inf-complete* when the set of points $x \in E$ such that $\varphi(x) \leq r$ is complete for each real r . Note that this implies lower semi-continuity of φ , and that the converse holds if \mathcal{U} is a complete uniformity. By an *ordering* on E we shall mean a reflexive, asymmetric and transitive relation \leq ; the corresponding irreflexive relation is denoted $<$. For $x \in E$ we shall denote by $S(x, \leq)$ the set of points $y \in E$ such that $x \leq y$. We shall say that an extended real valued function φ on E is *decreasing* resp.

strictly decreasing (with respect to \leq), if $x_1 < x_2$ implies $\varphi(x_1) \geq \varphi(x_2)$ resp. $\varphi(x_1) > \varphi(x_2)$.

All uniformities and topologies considered are assumed to be Hausdorff. All linear spaces considered are assumed to be real.

THEOREM 1. *Let \mathcal{U} be a uniformity on E , let \leq be an ordering on E , and let φ be an extended real valued function on E which is bounded below. Assume that the following conditions are fulfilled:*

- (a) *The set $S(x, \leq)$ is complete for each $x \in E$.*
- (b) *The function φ is decreasing.*
- (c) *For each $U \in \mathcal{U}$ there exists $\delta > 0$ such that $x_1 \leq x_2$ and $\varphi(x_1) - \varphi(x_2) < \delta$ implies $(x_1, x_2) \in U$.*

Then for each $x \in \text{dom } \varphi$ there exists $x_0 \in \text{dom } \varphi$ such that $x \leq x_0$ and x_0 is maximal in (E, \leq) .

Proof. For any given point $x \in \text{dom } \varphi$ there exists by Zorn's lemma a maximal totally ordered subset M of E containing x . We shall index the points of M by the elements of a totally ordered set (I, \leq) so that for $\alpha, \beta \in I$ we have $x_\alpha \leq x_\beta$ if and only if $\alpha \leq \beta$. Since φ is bounded below and decreasing, the net $(\varphi(x_\alpha))_{\alpha \in I}$ converges to some $a \in]-\infty, +\infty[$. Let $U \in \mathcal{U}$ be given, and let $\delta > 0$ be such that (c) is fulfilled. Choose $\alpha \in I$ such that $\varphi(x_\alpha) < a + \delta$. Then for $\alpha < \beta < \gamma$ we have $(x_\beta, x_\gamma) \in U$ by (c). This shows that the net $(x_\alpha)_{\alpha \in I}$ is a Cauchy net, and hence, by (a), it converges to a point x_0 with $x_\alpha \leq x_0$ for all $\alpha \in I$. In particular, $x \leq x_0$. By the nature of M it next follows that x_0 is maximal in (E, \leq) . Finally, (b) implies $x_0 \in \text{dom } \varphi$.

REMARK 1. As is well known, the use of Zorn's lemma may be replaced by an induction argument (involving the axiom of choice) along the following lines. Let $x_1 = x$, let $E_1 = S(x_1, \leq)$ and let $a_1 = \inf \varphi(E_1)$. When x_1, x_2, \dots, x_n have been chosen, let $E_n = S(x_n, \leq)$, let $a_n = \inf \varphi(E_n)$, and choose $x_{n+1} \in E_n$ such that $\varphi(x_{n+1}) \leq a_n + n^{-1}$. In doing so, one obtains sequences $(x_n)_{n \in \mathbb{N}}$ and $(a_n)_{n \in \mathbb{N}}$ such that $x_n \leq x_{n+1}$, $a_n \leq a_{n+1} \leq \varphi(x)$, $a_n \leq \varphi(x_{n+1}) \leq a_n + n^{-1}$. Using these relations and the idea of the proof above, the conclusion follows.

REMARK 2. We add an obvious but useful remark. Let E, \mathcal{U}, φ , and \leq be as assumed in Theorem 1, and let \leq' be an ordering on E which is finer than \leq , i.e., $x_1 \leq' x_2$ implies $x_1 \leq x_2$. Then (b) and (c) are also fulfilled for \leq' . Therefore, if (a) is fulfilled for

\cong' , then the theorem applies to \cong' . Even if (a) does not hold for \cong' , we may still conclude, however, that there exists $x_0 \in \text{dom } \varphi$ which is maximal in (E, \cong') , due to the fact that if x_0 is maximal with respect to \cong , then it is also maximal with respect to \cong' .

In order to obtain the desired theorem, we shall apply Theorem 1 to certain orderings which we shall denote $\cong_{d,\varphi}$. Let E be a set, and let d be a nonnegative extended real valued function on $E \times E$ satisfying the following conditions:

(d) $d(x, y) = 0$ if and only if $x = y$.

(e) $d(x, z) \leq d(x, y) + d(y, z)$.

Furthermore, let φ be an extended real valued function on E . We then define $\cong_{d,\varphi}$ as follows: $x_1 \cong_{d,\varphi} x_2$ if and only if either $x_1 = x_2$, or $x_1, x_2 \in \text{dom } \varphi$ and $d(x_1, x_2) \leq \varphi(x_1) - \varphi(x_2)$. Then it is easily checked that $\cong_{d,\varphi}$ is in fact an ordering and that φ is strictly decreasing with respect to $\cong_{d,\varphi}$, cf. (b). Also, if E is equipped with a uniformity such that φ is inf-complete and the functions $y \rightarrow d(x, y)$, $x \in E$, are lower semi-continuous, then for each $x \in E$ the set $S(x, \cong_{d,\varphi})$ is a closed subset of a complete set, and hence complete, cf. (a). Finally, note that if for each $U \in \mathcal{U}$ there exists $\delta > 0$ such that $d(x, y) < \delta$ implies $(x, y) \in U$, then (c) holds for $\cong_{d,\varphi}$. Therefore, by Theorem 1 we have:

THEOREM 2. *Let \mathcal{U} be a uniformity on E , let d be a non-negative extended real valued function on $E \times E$ satisfying (d) and (e), and let φ be an extended real valued function on E which is bounded below. Assume that the following conditions are fulfilled:*

(f) *The function φ is inf-complete.*

(g) *The functions $y \rightarrow d(x, y)$, $x \in E$, are lower semi-continuous.*

(h) *For each $U \in \mathcal{U}$ there exists $\delta > 0$ such that $d(x, y) < \delta$ implies $(x, y) \in U$.*

Then for each $x \in \text{dom } \varphi$ there exists $x_0 \in \text{dom } \varphi$ such that $x \cong_{d,\varphi} x_0$ and x_0 is maximal in $(E, \cong_{d,\varphi})$, i.e.,

$$\varphi(x_0) \leq \varphi(x) - d(x, x_0),$$

and

$$\varphi(y) > \varphi(x_0) - d(x_0, y)$$

for all $y \in E \setminus \{x_0\}$.

REMARK 3. Note that if \cong' is finer than $\cong_{d,\varphi}$, then under the conditions of Theorem 2 there exists x_0 which is maximal in (E, \cong') , and if in addition the sets $S(x, \cong')$ are closed, then we may obtain $x \cong' x_0$ for any given x ; cf. Remark 2.

3. In this section we shall clarify the relations of Theorem 2 to the Bishop-Phelps lemma and the most general extensions known to the author, namely that of I. Ekeland [5, Théorème 1] for complete metric spaces, and that of R. R. Phelps [6, Lemma 1] for non-metric topological linear spaces.

Note first that if d is a metric on E , then the conditions (d) and (e) are fulfilled. Furthermore, if one takes \mathcal{U} to be the uniformity on E generated by d , then (g) and (h) hold. Hence, if E is a metric space with metric d , and φ is bounded below and inf-complete on E (in particular, if d is a complete metric, and φ is bounded below and lower semi-continuous), then Theorem 2 applies. Adding the obvious remark that φ may be replaced by $k^{-1}\varphi$ for any $k > 0$, one obtains (under the condition stated) for each $x \in \text{dom } \varphi$ an $x_0 \in \text{dom } \varphi$ such that

$$\varphi(x_0) \leq \varphi(x) - kd(x, x_0)$$

and

$$\varphi(y) > \varphi(x_0) - kd(x_0, y)$$

for all $y \in E \setminus \{x_0\}$. This statement is essentially the theorem of I. Ekeland.

To obtain the result of R. R. Phelps, let X be a topological linear space, and let C be a closed, bounded, convex subset of X containing o . Let μ be the gauge function of C , i.e.,

$$\mu(x) = \inf \{ \lambda > 0 \mid x \in \lambda C \}$$

for $x \in X$, and let $d(x, y) = \mu(x - y)$ for $x, y \in X$. Then, as is well known, d maps $X \times X$ into $[0, +\infty]$ (with $d(x, y) = +\infty$ if and only if C does not absorb $x - y$), and the conditions (d) and (e) are fulfilled. Also, if \mathcal{U} is taken to be the uniformity given on X (X being a topological linear space), then (g) and (h) hold; for (g), the closedness of C is essential, for (h), the boundedness is essential. Therefore, if E is a subset of X , and φ is bounded below and inf-complete on E (in particular, if E is complete and φ is bounded below and lower semi-continuous) then Theorem 2 applies. Noting, as before, that φ may be replaced by $k^{-1}\varphi$ for any $k > 0$, one obtains a slight reformulation of the result of R. R. Phelps, stating that (under the conditions above) for each $x \in \text{dom } \varphi$ there exists $x_0 \in \text{dom } \varphi$ such that

$$\varphi(x) \leq \varphi(x_0) - k\mu(x - x_0),$$

and

$$\varphi(y) > \varphi(x_0) - k\mu(x_0 - y)$$

for all $y \in E \setminus \{x_0\}$. (In [6], the set corresponding to the set C above is not assumed to be convex. This condition seems necessary, however, to ensure that the relation considered be an ordering. For all the applications in [6], the appropriate set is in fact convex.)

Note that if one takes $d(x, y) = \|x - y\| = \mu(x - y)$, both of the two cases above covers the case of E being a subset of a normed linear space. In particular, both cases covers the original lemma of Bishop and Phelps which we shall briefly review. Let E be a complete subset of a normed linear space X , let ξ be a nonzero continuous linear functional on X which is bounded above on E , and let $k > 0$. Then the set

$$K(\xi, k) = \{x \in X \mid \|x\| \leq k\langle x, \xi \rangle\}$$

is a closed convex cone in X . (Note that $K(\xi, k) = \{0\}$ when $k\|\xi\| < 1$, and that $\text{int} K(\xi, k) \neq \emptyset$ when $k\|\xi\| > 1$.) Being also proper, the cone determines an ordering \leq on X in the usual manner, and the ordering on E induced by \leq is simply $\leq_{d, \varphi}$ with $d(x, y) = \|x - y\|$ and $\varphi(x) = -k\langle x, \xi \rangle$. Therefore, for each $x \in E$ there exists $x_0 \in E$ such that $x \leq x_0$ and x_0 is maximal in (E, \leq) . This is the lemma of Bishop and Phelps, slightly reformulated. In this connection, note that if \leq' is an ordering on X determined by a convex subcone C of $K(\xi, k)$, then by Remark 3 we may still conclude the existence of a point $x_0 \in E$ such that x_0 is maximal in (E, \leq') ; and if in addition C is closed, we may obtain $x \leq' x_0$ for a given $x \in E$. This covers a result of F. E. Browder [2, Lemma 2].

For a review of applications of the Bishop-Phelps procedure in the case of orderings on Banach spaces generated by convex cones, see R. R. Phelps [8]. In particular, this paper contains a comprehensive bibliography. For applications in more general cases, see e.g. A. Brøndsted and R. T. Rockafellar [3], R. R. Phelps [7] and the papers of Ekeland and Phelps quoted above. See also § 4.

4. By a drop $D(z, r, y)$ in a normed linear space X we shall mean the convex hull of a closed ball $B(z, r)$ and a point y not belonging to the ball. This notion was introduced by J. Daneš who proved the following [4, Drop Theorem]:

THEOREM 3 (J. Daneš). *Let F be a closed subset of a Banach space X , and let z be a point in $X \setminus F$. Let $0 < r < R < \rho$, where $R = \text{dist}(z, F)$. Then there exists a point $x_0 \in \text{bd}F$ such that $\|x_0 - z\| \leq \rho$ and $D(z, r, x_0) \cap F = \{x_0\}$.*

This theorem is related to a theorem of F. E. Browder [2, Theorem 4]. Roughly speaking, Browder proves that a certain drop

may be translated to “support” the appropriate set, i.e., the “peak” is the only point of the drop that belongs to the set, whereas Daneš proves that for certain given z and r there exists a “supporting” drop of the particular form $D(z, r, y)$ for a suitable y . The proof of Browder’s theorem depends on the lemma of Browder referred to in § 3; hence the Bishop-Phelps procedure applies with an ordering defined by a convex cone. Although not explicitly stated, the proof of J. Daneš also contains a sort of a Bishop-Phelps argument. We shall give a considerably clarified version of the proof by applying directly the Bishop-Phelps procedure (with an ordering not generated by a convex cone).

Proof (Theorem 3). Without loss of generality we may assume that $z = o$. We define a relation \leq on the set $E = F \cap B(o, \rho)$ by putting $x_1 \leq x_2$ if and only if $x_2 \in D(o, r, x_1)$. It is easily checked that \leq is in fact an ordering. Let

$$\varphi(x) = \frac{\rho + r}{R - r} \|x\|$$

for $x \in E$. It is claimed that \leq is finer than the ordering $\leq_{d, \varphi}$, where $d(x, y) = \|x - y\|$. Let x_1 and x_2 be points in E such that $x_1 < x_2$. From the definition of a drop it then follows that

$$(1) \quad x_2 = (1 - t)x_1 + tv$$

for appropriate $t \in]0, 1[$ and $v \in B(o, r)$. By (1), we have $\|x_2\| \leq (1 - t)\|x_1\| + t\|v\|$, whence

$$t(\|x_1\| - \|v\|) \leq \|x_1\| - \|x_2\|.$$

On the other hand, we clearly have $R - r \leq \|x_1\| - \|v\|$, and we therefore obtain

$$(2) \quad t \leq (\|x_1\| - \|x_2\|)(R - r)^{-1}.$$

From (1) it also follows that $x_2 - x_1 = t(v - x_1)$. Using this and (2) we get

$$\begin{aligned} \|x_2 - x_1\| &= t\|v - x_1\| \leq t(\|v\| + \|x_1\|) \\ &\leq t(r + \rho) \leq (\rho + r)(R - r)^{-1}(\|x_1\| - \|x_2\|), \end{aligned}$$

as desired. Now, by Theorem 2 (see also § 3), there exists a maximal element x_0 in $(E, \leq_{d, \varphi})$. But then x_0 is also maximal in (E, \leq) , cf. Remark 3. This proves that $\|x_0\| \leq \rho$ and that x_0 is the only point of $D(o, r, x_0)$ belonging to E . But then x_0 is the only point of $D(o, r, x_0)$ belonging to the given set F , since the points in $D(o, r, x_0)$

have norm $\leq \rho$, whereas the points in $F \setminus E$ have norm $> \rho$. Finally, it is obvious that x_0 is a boundary point of F .

Note that since a drop is closed, we may obtain $x_0 \in D(z, r, x)$ for any given $x \in F$ with $\|x - z\| \leq \rho$, cf. Remark 3.

REFERENCES

1. E. Bishop and R. R. Phelps, *The support functionals of a convex set*, Proc. Symp. Pure Math. VII, Convexity, Amer. Math. Soc., (1963), 27-36.
2. F. E. Browder, *Normal solvability and the Fredholm alternative for mappings into infinite dimensional manifolds*, J. Funct. Anal., **8** (1971), 250-274.
3. A. Brøndsted and R. T. Rockafellar, *On the subdifferentiability of convex functions*, Proc. Amer. Math. Soc., **16** (1965), 605-611.
4. J. Daneš, *A geometric theorem useful in nonlinear functional analysis*, Bollettino U.M.I. (4), **6** (1972), 369-375.
5. I. Ekeland, *Sur les problèmes variationnelles*, C. R. Acad. Sci. Paris Sér. A, **275** (1972), 1057-1059.
6. R. R. Phelps, *Support cones and their generalizations*, Proc. Symp. Pure Math. VII, Convexity, Amer. Math. Soc., (1963), 393-401.
7. ———, *Weak* support points of convex sets in E^** , Israel J. Math., **2** (1964), 177-182.
8. ———, *Support cones in Banach spaces and their applications*, Advances in Math., **13** (1974), 1-19.

Received March 23, 1973.

UNIVERSITY OF COPENHAGEN
DENMARK

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, (not dittoed), double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. Items of the bibliography should not be cited there unless absolutely necessary, in which case they must be identified by author and Journal, rather than by item number. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. **39**. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California, 90024.

The Pacific Journal of Mathematics expects the author's institution to pay page charges, and reserves the right to delay publication for nonpayment of charges in case of financial emergency.

100 reprints are provided free for each article, only if page charges have been substantially paid. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$72.00 a year (6 Vols., 12 issues). Special rate: \$36.00 a year to individual members of supporting institutions.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 270, 3-chome Totsuka-cho, Shinjuku-ku, Tokyo 160, Japan.

Copyright © 1973 by Pacific Journal of Mathematics
Manufactured and first issued in Japan

Pacific Journal of Mathematics

Vol. 55, No. 2

October, 1974

Walter Allegretto, <i>On the equivalence of two types of oscillation for elliptic operators</i>	319
Edward Arthur Bertram, <i>A density theorem on the number of conjugacy classes in finite groups</i>	329
Arne Brøndsted, <i>On a lemma of Bishop and Phelps</i>	335
Jacob Burbea, <i>Total positivity and reproducing kernels</i>	343
Ed Dubinsky, <i>Linear Pincherle sequences</i>	361
Benny Dan Evans, <i>Cyclic amalgamations of residually finite groups</i>	371
Barry J. Gardner and Patrick Noble Stewart, <i>A “going down” theorem for certain reflected radicals</i>	381
Jonathan Light Gross and Thomas William Tucker, <i>Quotients of complete graphs: revisiting the Heawood map-coloring problem</i>	391
Sav Roman Harasymiv, <i>Groups of matrices acting on distribution spaces</i>	403
Robert Winship Heath and David John Lutzer, <i>Dugundji extension theorems for linearly ordered spaces</i>	419
Chung-Wu Ho, <i>Deforming p. l. homeomorphisms on a convex polygonal 2-disk</i>	427
Richard Earl Hodel, <i>Metrizability of topological spaces</i>	441
Wilfried Imrich and Mark E. Watkins, <i>On graphical regular representations of cyclic extensions of groups</i>	461
Jozef Krasinkiewicz, <i>Remark on mappings not raising dimension of curves</i>	479
Melven Robert Krom, <i>Infinite games and special Baire space extensions</i>	483
S. Leela, <i>Stability of measure differential equations</i>	489
M. H. Lim, <i>Linear transformations on symmetric spaces</i>	499
Teng-Sun Liu, Arnoud C. M. van Rooij and Ju-Kwei Wang, <i>On some group algebra modules related to Wiener’s algebra M_1</i>	507
Dale Wayne Myers, <i>The back-and-forth isomorphism construction</i>	521
Donovan Harold Van Osdol, <i>Extensions of sheaves of commutative algebras by nontrivial kernels</i>	531
Alan Rahilly, <i>Generalized Hall planes of even order</i>	543
Joylyn Newberry Reed, <i>On completeness and semicompleteness of first countable spaces</i>	553
Alan Schwartz, <i>Generalized convolutions and positive definite functions associated with general orthogonal series</i>	565
Thomas Jerome Scott, <i>Monotonic permutations of chains</i>	583
Eivind Stensholt, <i>An application of Steinberg’s construction of twisted groups</i>	595
Yasuji Takeuchi, <i>On strongly radical extensions</i>	619
William P. Ziemer, <i>Some remarks on harmonic measure in space</i>	629
John Grant, <i>Corrections to: “Automorphisms definable by formulas”</i>	639
Peter Michael Rosenthal, <i>Corrections to: “On an inversion for the general Mehler-Fock transform pair”</i>	640
Carl Clifton Faith, <i>Corrections to: “When are proper cyclics injective”</i>	640