

## ON A POSITIVE TRIGONOMETRIC SUM

RICHARD ASKEY,<sup>1</sup> JAMES FITCH AND GEORGE GASPER<sup>2</sup>

We give a new proof of the following theorem of P. Turán [2]. See [1] for another proof, shorter than Turán's but longer than this proof.

THEOREM. Let  $\sum_{n=1}^N a_n \sin(2n-1)\theta \geq 0$ ,  $0 \leq \theta \leq \pi$ . Then

$$\sum_{n=1}^N \frac{a_n \sin n\phi}{n} > 0, \quad 0 < \phi < \pi,$$

unless all  $a_n = 0$ .

A simple computation shows that

$$\frac{d}{dy} \frac{\sin \alpha y}{\alpha (\sin y)^\alpha} = - \frac{\sin(\alpha - 1)y}{(\sin y)^{\alpha+1}}.$$

Letting  $\alpha = 2n$  and  $y = \phi/2$  we see that

$$\frac{\sin n\phi}{n} = 2 \int_{\phi/2}^{\pi/2} \left( \frac{\sin \phi/2}{\sin \theta} \right)^{2n} \frac{\sin(2n-1)\theta}{\sin \theta} d\theta.$$

Thus

$$\sum_{n=1}^N \frac{a_n \sin n\phi}{n} = 2 \int_{\phi/2}^{\pi/2} \sum_{n=1}^N a_n \left( \frac{\sin \phi/2}{\sin \theta} \right)^{2n} \sin(2n-1)\theta \frac{d\theta}{\sin \theta}.$$

But  $\sum_{n=1}^N a_n r^{2n-1} \sin(2n-1)\theta > 0$ ,  $0 < r < 1$ , if  $\sum_{n=1}^N a_n \sin(2n-1)\theta \geq 0$  and not all  $a_n$  are zero and this completes the proof.

### REFERENCES

1. C. Hyltén-Cavallius, *A positive trigonometrical kernel*, Tolfte Skand. Matematiker kongressen 1953, Lund (1954).
2. P. Turán, *On a trigonometric sum*, Ann. Polon. Math. 25 (1953), 155-161.

UNIVERSITY OF WISCONSIN

Received by the editors February 26, 1968.

<sup>1</sup> Sponsored by the Mathematics Research Center, United States Army, Madison, Wisconsin, under Contract No. DA-31-124-ARO-D-462.

<sup>2</sup> Supported in part by National Science Foundation grant GP-6764.