

D. M. Shaydayeva

On basic concepts of non-commutative topology

Czechoslovak Mathematical Journal, Vol. 34 (1984), No. 3, 378–389

Persistent URL: <http://dml.cz/dmlcz/101963>

Terms of use:

© Institute of Mathematics AS CR, 1984

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON BASIC CONCEPTS OF NON-COMMUTATIVE TOPOLOGY

D. M. SHAYDAYEVA, MOSCOW

(Received May 10, 1982)

A non-commutative generalization of locally compact Hausdorff spaces was independently offered by Ch. A. Akemann [1–3], R. Giles and H. Kummer [8; 9]. They used the description of such topological spaces in terms of the bounded continuous functions algebra as the matter to be extended to the non-abelian situation. In that way a non-commutative or “quantum” topology associated with a C^* -algebra was defined as a certain family of projections in its atomic W^* -envelope, and the C^* -algebra was interpreted as an algebra of “continuous” elements in accordance with this topology.

In the present paper we give an intrinsic axiomatic definition of a general non-commutative topology in terms of the lattice of all projections in an arbitrary atomic W^* -algebra B . The system of axioms connects the properties of non-commutative topology with order, Jordan and C^* -structures on B . For all that two key ideas are pursued: firstly, to generalize to non-abelian case the description of any topology by means of the bounded lower semicontinuous functions cone and, secondly, to provide that the set of “continuous” elements in B be a C^* -algebra. Moreover, we give an effective characterization of compactness and show that a non-commutative topology is locally compact iff it is the Akemann-Giles topology associated with a certain C^* -algebra.

1. Preliminaries. Let us consider, together with any set X , the commutative W^* -algebra $B(X)$ of all complex-valued bounded functions on X . The field of all subsets of X may be naturally identified with the atomic boolean algebra $\text{Pr}B(X)$ of all projections in $B(X)$ (each subset of X is assigned its characteristic function); the points of X are in one to one correspondence with the atoms in the lattice $\text{Pr}B(X)$. Now, if there is a topology τ on X , it may be regarded as some family of projections in $B(X)$. Namely, $\tau = \text{Pr}B(X) \cap L(X)$, where $L(X)$ is the convex cone of all lower semi-continuous bounded functions on X . In particular, if τ is completely regular, let $C(\tau)$ denote the C^* -algebra of all continuous bounded functions on X and let $C(\tau)^m$ denote the set of suprema in $B(X)$ (i.e. pointwise suprema) of all bounded increasing nets of real-valued elements in $C(\tau)$, then $\tau = \text{Pr}B \cap C(\tau)^m$.

It easily follows from the spectral theory that an arbitrary commutative atomic

W^* -algebra M is isomorphic to the W^* -algebra $B(X)$, where X is the set of all minimal projections in M (i.e. atoms in the atomic boolean algebra $\text{Pr}M$). In the light of these facts the lattice $\text{Pr}B$ of all projections in an arbitrary non-commutative atomic W^* -algebra B can be regarded as a non-commutative analogue of the concept of set; minimal projections in B play the role of points, any projection in B , being the supremum of minimal ones, plays the role of a subset. Finally, a “non-commutative topology” arises as the appropriate family of projections in a non-commutative atomic W^* -algebra.

This outlook at a non-commutative generalization of topology was the basis for the Akemann-Giles analogue of a locally compact Hausdorff space. In what follows we construct a non-commutative analogue of a general topological space and show that the Akemann-Giles construction coincides with ours in the locally compact case.

For the general theory of C^* - and W^* -algebras we shall make systematic use of books [6] and [12].

1.1. q -sets. An atomic W^* -algebra B together with the set of all minimal projections therein will be called a q -set, the elements of $\text{Pr}B$ q -subsets and the atoms in $\text{Pr}B$ q -points. Union and intersection of q -sets are to be taken in the lattice $\text{Pr}B$. Two q -sets e and f will be called *disjoint* iff $ef = 0$. (The terminology is lifted from [8]).

Any C^* -algebra A is associated with a q -set as follows. The second conjugate space A^{**} is a W^* -algebra; let Z_A be the supremum of all minimal projections in A^{**} , then Z_A belongs to the center of the algebra A^{**} [1; p. 278]. Set $B_A = Z_A A^{**}$, then B_A is the atomic W^* -algebra and we can consider A as the weakly dense sub-algebra of B_A , since $A \subset A^{**}$ and $A \rightarrow Z_A A$ is an isomorphism (see [3], p. I). If A is abelian with the spectrum X , then the points of X are in one to one correspondence with the minimal non-zero projections of A^{**} so that $B_A \approx B(X)$ and the Gelfand isomorphism give $A \approx C_0(X) \subset B(X)$.

1.2. Notation. For any subset $E \subset B$ put $E^+ = \{a \in E \mid a \geq 0\}$, $E^s = \{a \in E \mid a = a^*\}$, $E_1 = \{a \in E \mid \|a\| \leq 1\}$. Recall that the self-adjoint part B^s of a W^* -algebra B is an ordered space, in which every norm bounded increasing net has a supremum. Let E^m denote the set of suprema in B^s of all norm-bounded increasing nets of elements of E^s ; put $E_m = -(-E)^m$.

1.3. The Akemann-Giles Q -topology. For any C^* -algebra A consider $A \subset B_A$ as above. The family of q -sets $\tau_A = \text{Pr}B_A \cap (A^+)^m$ will be called the *Akemann-Giles q -topology* on B_A . The elements of τ_A are called q -open q -sets, the elements of $1 - \tau_A = \{1 - e \mid e \in \tau_A\}$ are called q -closed. The pair (B_A, τ_A) will be called the q -spectrum of the C^* -algebra A and denote as $q \text{ spec } A$. If A is commutative, then τ_A is the usual topology on the spectrum of A and therefore is Hausdorff and locally compact. In the non-abelian case τ_A has similar properties. Namely, a q -topology τ_A

is Hausdorff: i.e., given disjoint q -points x and y there exist disjoint open q -sets e and f with $x \leq e$ and $y \leq f$ [8; III.6]. After Akemann a q -set $p \in \text{Pr}B_A$ is called q -compact if p is closed and there exists $a \in A_1^+$ with $p \leq a$ [3; II.I and II.5]. The q -topology τ_A is locally compact: i.e., for any q -point x there exist an open q -set e and a compact q -set p with $x \leq e \leq p$ (this follows from [3; III.I]). In the abelian case, when $A \approx C_0(X) \subset B(X)$, these concepts of q -compactness are equivalent to the usual definitions.

1.4. Gelfand-Akemann-Giles theorem. An element $a \in B_A^s$ is called τ_A -continuous if each spectral projection of a which corresponds to an open subset of the real numbers is also an open q -set. A τ_A -continuous element a is called vanishing at ∞ if each spectral projection of a corresponding to a closed subset of the real numbers which do not contain 0 is q -compact [3; I.I and III.3].

Denote by $C(\tau_A)^s$ the set of all τ_A -continuous elements of B_A^s and by $C_0(\tau_A)^s$ the set of all elements vanishing at ∞ . Set $C(\tau_A) = C(\tau_A)^s + iC(\tau_A)^s$ and $C_0(\tau)_A = C_0(\tau_A)^s + iC_0(\tau_A)^s$. The elements of $C(\tau_A)$ and $C_0(\tau_A)$ will also be called τ_A -continuous and vanishing at ∞ , respectively.

Theorem [3; 5; 8]. *A C^* -algebra A is exactly the algebra of all τ_A -continuous elements of B_A vanishing at ∞ , i.e. $A = C_0(\tau_A)$; the set of all continuous elements of B_A coincides with the C^* -algebra of all multipliers of A , i.e. $C(\tau_A) = M(A) \equiv \{b \in B_A \mid Ab + bA \subset A\}$.*

Throughout the whole paper B will always be used to denote an arbitrary atomic W^* -algebra, and $\tau \subset \text{Pr}B$ a family of projections (i.e. q -sets) in B . We start to discuss the individual axioms which will connect the properties of τ with various structures on B .

2. Order axioms. It is easy to see that any Akemann-Giles q -topology contains 0 and 1 and that a union of open q -sets is also open. Akemann has shown that in contradistinction to the usual topology, the intersection of two open q -sets is not necessarily open (see a counterexample in [1]). The first three axioms describe the corresponding properties of the general q -topology.

AXIOM A1. $0, 1 \in \tau$;

AXIOM A2. $(e_\alpha) \subset \tau \Rightarrow \bigvee_\alpha e_\alpha \in \tau$, i.e. "union of open q -sets is open".

AXIOM A3. $e, f \in \tau, [e, f] = 0 \Rightarrow e \wedge f \in \tau$, i.e. "intersection of two commuting open q -sets is open".

Remark that actually axiom A3 is an order condition since $[e, f] = 0 \Leftrightarrow e = (e \wedge f) \vee (e \wedge (1 - f))$. Any Akemann-Giles q -topology satisfies all these axioms [1].

We have noticed in § 1 that in the commutative case a topology of a topological space X may be described algebraically by the equality $\tau = \text{Pr}B(X) \cap L(X)$. Our fourth axiom will be a non-commutative version of this description, therefore we need an appropriate definition of the class of lower semicontinuous (LSC) elements in B .

Definition 2.1. The set $E \subset B^s$ is *lower monotone closed* (LMC) if $E = E^m$; E is *upper monotone closed* (UMC) if $E = E_m$. The minimal LMC set in B^s containing E is called the *lower monotone closure* of E and is denoted by $L(E)$. The *upper monotone closure* of E is similarly defined and denoted by $U(E)$.

Lemma 2.2. *If E is a convex cone in B^s , then $L(E)$ is a convex cone, too. If, besides, $E \supset \mathbb{R} \cdot 1$, then the convex cone $L(E)$ is norm-closed in B^s .*

Proof. Take $\lambda, \mu \in \mathbb{R}^+$ and $a \in E$, then

$$(*) \quad E \subset M(a) \equiv \{b \in L(E) \mid \lambda a + \mu b \in L(E)\} \subset L(E)$$

and the set $M(a)$ is LMC. So $M(a) = L(E)$ and for each $a \in E$ and $b \in L(E)$ we have $\lambda a + \mu b \in L(E)$. This implies that for each $a \in L(E)$ (*) is correct and we get similarly $M(a) = L(E)$. The last equality shows that $L(E)$ is a convex cone. Let $E \supset \mathbb{R} \cdot 1$. If a sequence $(a_n) \subset L(E)$ converges to an element $b \in B^s$ we may suppose that $\|a_n - b\| \leq 2^{-n}$. Then the increasing sequence $\tilde{a}_n = a_n - 2^{-n+1} \cdot 1$ is contained in $L(E)$ and also converges to b . Hence $b = \bigvee_n \tilde{a}_n \in L(E)$ and $L(E)$ is norm-closed.

Put $A^+(\tau) = \{\sum_{i=1}^n \lambda_i e_i \mid \lambda_i \geq 0, e_i \in \tau, n \in \mathbb{N}\}$ and consider $A(\tau) = A^+(\tau) + \mathbb{R} \cdot 1$, the minimal convex cone containing τ and "constants". We now define the class of *lower semicontinuous* (LSC) *elements* in B^s as $L(\tau) = L(A(\tau))$. By Lemma 2.2, $L(\tau)$ is a norm-closed convex cone in B^s . If τ were the usual topology on a set X , $L(\tau)$ would be the class of all lower semicontinuous real-valued functions on X .

Similarly we may define the class $U(\tau)$ of *upper semicontinuous* (USC) *elements* in B^s by setting

$$A^+(1 - \tau) = \left\{ \sum_{i=1}^n \lambda_i f_i \mid \lambda_i \geq 0, f_i \in 1 - \tau \right\},$$

$$A(1 - \tau) = A^+(1 - \tau) + \mathbb{R} \cdot 1 \quad \text{and} \quad U(\tau) = U(A(1 - \tau)).$$

For any subset $E \subset B^s$ we shall denote by \bar{E} the norm-closure of E in B^s .

Lemma 2.3. *If A is a C^* -algebra and τ_A is the Akemann-Giles topology associated with A , then*

$$L(\tau_A) = \bar{A}^m,$$

where $\tilde{A} = A + \mathbb{C} \cdot 1 \subset B_A$ is the C^* -algebra obtained by adjoining the unit 1 of B_A to A .

Proof. We have $A(\tau_A) \subset \tilde{A}^m \subset \bar{A}^m$ since $\tau_A \subset (A^+)^m$. By [4; 3.3] the convex cone \bar{A}^m is LMC, therefore we get $L(\tau_A) \subset \bar{A}^m$. To show the converse inclusion notice that by virtue of the Gelfand-Akemann-Giles theorem (1.4) we have $\tilde{A} \subset C(\tau_A)$ and so by the spectral theory $\tilde{A}^s \subset A(\tau_A)^m$. This implies $\tilde{A}^m \subset L(\tau_A)$, which means $\bar{A}^m \subset L(\tau_A)$ as desired.

The proof of the last lemma was based on the assertion in [4], which was stated

there in terms of A^{**} , but applying theorems [9; 3.8 and 4; 2.6] we get the result in terms of B_A .

From the papers [9; 4] by Pedersen and Akemann we know that there is an isometric map of $\overline{A^m}$ onto the set $L^{\infty}(S(A))$ of all bounded LSC affine functions on the state space $S_A = \{\varphi \in A^{*+} \mid \|\varphi\| = 1\}$ provided with the weak* topology of A^* . Through Lemma 2.3 we may therefore identify the convex cone $L(\tau_A)$ with the cone $L^{\infty}(S(A))$ (see also § 5 below).

The next definition coincides in essence with that in 1.1, and in the commutative case it is the usual definition of continuous functions.

Definition 2.4. An element $a \in B^s$ is called q -continuous, if for each open set $I \subset \mathbb{R}$ the spectral projection $E_a(I)$ belongs to τ , $a = \int \lambda dE_a$ being the spectral representation of a . Let $C(\tau)^s$ denote the set of all q -continuous elements of B^s and $C(\tau) = C(\tau)^s + iC(\tau)^s$. The elements of $C(\tau)$ will also be called q -continuous.

Proposition 2.5. If a family $\tau \subset \text{Pr}B$ satisfies axioms A1–A3, then

$$C(\tau)^s = \bigcup \{C^s \mid C \subset C(\tau), C \text{ is commutative } C^* \text{-subalgebra}\}.$$

Proof. Let $a \in C(\tau)^s$ and let $C^*(1, a)$ be the commutative C^* -subalgebra of B , generated by a and the unit 1. By the Gelfand representation theorem $C^*(1, a)$ is isomorphic to the C^* -algebra $C(\text{Spa})$ of all continuous functions on the spectrum of a , and any $b \in C^*(1, a)^s$ is associated with the continuous real-valued function f_b on Spa with $E_b(I) = E_a(f_b^{-1}(I))$ for each $I \subset \mathbb{R}$. This shows that for any $b \in C^*(1, a)^s$ and any open set $I \subset \mathbb{R}$ the q -set $E_b(I) \in \tau$, which means that $C^*(1, a)^s \subset C(\tau)^s$ and completes the proof.

It follows from the spectral theory that $C(\tau)^s \subset L(\tau) \cap U(\tau)$, i.e. that any continuous element in B^s is both lower and upper semicontinuous. Set

$$Q(\tau)^s = L(\tau) \cap U(\tau) \quad \text{and} \quad Q(\tau) = Q(\tau)^s + iQ(\tau)^s.$$

The elements of $Q(\tau)$ will be called q -quasicontinuous. By Lemma 2.2, $Q(\tau)$ is a norm-closed linear subspace of B . In the commutative case, when τ is the usual topology, $Q(\tau)$ certainly coincides with $C(\tau)$. But it is not so for Akemann-Giles topologies.

Proposition 2.6. Let A be a C^* -algebra and $(B_A, \tau_A) = q \text{ spec } A$. Then

$$Q(\tau_A) = \{x \in B_A \mid a \times b \in A \ \forall a, b \in A\} \equiv Q(A),$$

i.e. $Q(\tau_A)$ is the space of all quasimultiples of A .

Proof. It follows from Lemma 2.3 that $Q(\tau_A) = \overline{A^m} \cap \overline{A_m}$; by [4; 4.1] this intersection is exactly the space $Q(A)$ of all quasimultiples of A in B_A .

By the Gelfand-Akemann-Giles theorem, $C(\tau_A) = M(A)$ where $M(A)$ is the C^* -algebra of all multiples of A in B_A ; in the paper [4] an example is given of a C^* -algebra A for which $Q(A) \neq M(A)$ and $Q(A)$ is not a Jordan algebra.

AXIOM A4. $\tau = \text{Pr}B \cap L(\tau)$.

Definition 2.7. A family $\tau \subset \text{Pr}B$, which obeys the axioms A1–A4, is called a *q-topology*. The elements of τ are *q-open q-sets* the elements of $1 - \tau$ are *q-closed q-sets*. Given any *q-set* $e \in \text{Pr}B$, its *q-closure* is $\bar{e} = \bigwedge \{f \mid f \text{ is } q\text{-closed and } e \leq f\}$; similarly its *q-interior* is ${}^{\circ}e = \bigvee \{g \mid g \text{ is open and } e \geq g\}$. The pair (B, τ) is called the *q-topological space*.

It follows from Lemma 2.3 and [4; 3,6] that any Akemann-Giles *q-topology* satisfies axiom A4. In the commutative case that axiom follows from A1–A3, but if B is non-abelian, A4 is independent of them. Axioms A1–A3 imply neither that $C(\tau)$ is a linear subspace of B nor that it is norm-closed.

Theorem 2.8. *If τ is a q-topology, then*

$$C(\tau)^s = \{a \in B^s \mid a^n \in Q(\tau)^s \text{ for all } n \in \mathbb{N}\}.$$

Proof. If $a \in C(\tau)^s$ then by Proposition 2.5 $a^n \in C(\tau)^s$ for all $n \in \mathbb{N}$ and so $C(\tau)^s \subset \{a \in B^s \mid a^n \in Q(\tau)^s \text{ for all } n \in \mathbb{N}\}$. Conversely, let $b \in B^s$, $b^n \in Q(\tau)^s$ for all $n \in \mathbb{N}$ and denote by $C^*(1, b)$ the C^* -subalgebra of B generated by b and the unit 1. then through the Stone-Weierstrass theorem we get $C^*(1, b)^s \subset Q(\tau)^s$. Since for any open $I \subset \mathbb{R}$ such that $E_b(I) \neq 0$ there exists an increasing sequence $(b_n) \subset C^*(1, b)^+$ with $E_b(I) = \bigvee_n b_n$, it implies that $E_b(I) \in (Q(\tau))^m \subset L(\tau)$. Thus in virtue of axiom A4 $E_b(I)$ is open, whence $b \in C(\tau)^s$.

Corollary 2.9. *If τ is a q-topology then $C(\tau)^s$ is a norm-closed subset of B^s .*

Recall that B^s is a real Jordan algebra and B is a Jordan C^* -algebra (JC^* -algebra) with multiplication $a \circ b = \frac{1}{2}(ab + ba)$.

Proposition 2.10. *Let τ be a q-topology. If $C(\tau)^+$ is a convex cone in B^s then $C(\tau)^s$ is a norm-closed real Jordan subalgebra of B^s and $C(\tau)$ is a JC^* -subalgebra of B .*

Proof. By Proposition 2.5, given $a, b \in C(\tau)^s$ we have $a + \|a\| \cdot 1, b + \|b\| \cdot 1 \in C(\tau)^+$ so $(a + b) + (\|a\| + \|b\|) \cdot 1 \in C(\tau)^+$ whence $a + b \in C(\tau)^s$. Moreover we get again by Proposition 2.5 $a^2 \in C(\tau)^s$ whenever $a \in C(\tau)^s$ and thus $a \circ b = \frac{1}{4}((a + b)^2 - (a - b)^2) \in C(\tau)^s$ whenever $a, b \in C(\tau)^s$. Together with Corollary 2.9 this proves the first assertion. Take a sequence $(a_n) \subset C(\tau)$ which uniformly converges to $b \in B$. Setting $\text{Re } x = \frac{1}{2}(x + x^*)$, $\text{Im } x = \frac{1}{2}(x - x^*) \in B^s$ for any $x \in B$ we see that $\|\text{Re } a_n - \text{Re } b\| \rightarrow 0$ and $\|\text{Im } a_n - \text{Im } b\| \rightarrow 0$. Since $C(\tau)^s$ is closed, we obtain that $\text{Re } b, \text{Im } b \in C(\tau)^s$ which implies $b \in C(\tau)$.

Lemma 2.11. *Let τ be a q-topology. Suppose $X \subset Q(\tau)^+$ is a convex cone satisfying*

- (i) $X \supset \mathbb{R}^+ \cdot 1$; (ii) $a^{1/2} \in Q(\tau)^s$ for any $a \in X$.

Then $X \subset C(\tau)^+$.

Proof. Take $a \in X$. By Theorem 2.8 it suffices to prove that $a^n \in Q(\tau)^+$ whenever $n \in \mathbb{N}$. This assertion is trivial for $n = 1$ and we assume that it has been proved for all values of $n < m$. Consider $\alpha > \|a\|$; since by (i) $1 + \alpha^{-1}a \in X$ we see from (ii)

that the element

$$(1 + \alpha^{-1}a)^{1/2} = 1 + \frac{1}{2}\alpha^{-1}a + \frac{\frac{1}{2}(\frac{1}{2} - 1)}{2!} \alpha^{-2}a^2 + \frac{\frac{1}{2}(\frac{1}{2} - 1)(\frac{1}{2} - 2)}{3!} \alpha^{-3}a^3 + \dots$$

belongs to $Q(\tau)^+$. So a^m is the uniform limit

$$a^m = \lim_{\alpha \rightarrow +\infty} \alpha^m \binom{\frac{1}{2}}{m}^{-1} \left[(1 + \alpha^{-1}a)^{1/2} - \sum_{k=0}^{m-1} \binom{\frac{1}{2}}{k} \alpha^{-k} a^k \right]$$

of elements in $Q(\tau)^s$. Because $Q(\tau)^s$ is closed, we get the lemma.

3. Algebraic regularity of q -topology. Next we introduce a condition connecting τ with the Jordan algebra structure on B .

AXIOM A5. If $a \in A^+(\tau)$, then $a^{1/2} \in L(\tau)$; if $a \in A^+(1 - \tau)$, then $a^{1/2} \in U(\tau)$.

An arbitrary Akemann-Giles q -topology satisfies this axiom. Indeed, if $\tau = \tau_A$ for a C^* -algebra A , then $\tau \in (A^+)^m$ whence $A^+(\tau) \subset (A^+)^m$ and $A^+(\tau)^{1/2} \subset ((A^+)^m)^{1/2}$. Since $(A^+)^{1/2} \subset A^+$ and $t^{1/2}$ is an operator monotone function on B^+ [10] we have $((A^+)^m)^{1/2} \subset (A^+)^m$; applying Lemma 2.3 we see that $(A^+)^m \subset L(\tau)$. From all that we obtain $(A^+(\tau))^{1/2} \subset L(\tau)$. Similarly we can conclude that $(A^+(1 - \tau))^{1/2} \subset U(\tau)$.

Theorem 3.1. *If a q -topology τ satisfies axiom A5, then $C(\tau)^s$ is a norm-closed real Jordan subalgebra of B^s and $C(\tau)$ is a JC^* -subalgebra of B .*

Proof. Let X denote the convex cone $\overline{A^+(\tau) \cap A^+(1 - \tau)}$. In view of Proposition 2.10 it suffices to show that $C(\tau)^+ = X$. The inclusion $C(\tau)^+ \subset X$ follows from the spectral theory. The inverse inclusion is valid since the cone X satisfies all the conditions of Lemma 2.11 (we have $a^{1/2} \in Q(\tau)^s$ whenever $a \in X$ in virtue of axiom A5 and norm-continuity of the operator function $t^{1/2}$).

Axiom A5 may be weakened to get a necessary and sufficient condition for the set $C(\tau)$ to be a JC^* -subalgebra of B . Such a weak variant, being equivalent to the original axiom A5 for completely regular q -topologies (see Definition 3.4 below), will concern only the part of q -topology τ , which can be reproduced by means of $C(\tau)$.

Let us define the *regularization* τ^{reg} of a q -topology τ by setting $\tau^{\text{reg}} = \text{Pr}B \cap \bigcap (C(\tau)^+)^m$ (in general the projections family τ^{reg} need not be a q -topology).

Lemma 3.2. *If $\tau \in \text{Pr}B$ satisfies axioms A1–A3, then $C(\tau) = C(\tau^{\text{reg}})$.*

Proof. Clearly, $C(\tau^{\text{reg}}) \subset C(\tau)$. Conversely, by Proposition 2.5, given $a \in C(\tau)^s$ the C^* -subalgebra $C^*(1, a) \subset C(\tau)$, so for any open $I \subset \mathbb{R}$ the spectral projection $E_a(I)$, being a supremum of an increasing sequence of elements in $C^*(1, a)$, belongs to τ^{reg} . Thus we get $C(\tau)^s \subset C(\tau^{\text{reg}})^s$ and, consequently, $C(\tau) \subset C(\tau^{\text{reg}})$.

AXIOM A5°. If $a \in A^+(\tau^{\text{reg}})$, then $a^{1/2} \in L(\tau)$, if $a \in A^+(1 - \tau^{\text{reg}})$ then $a^{1/2} \in U(\tau)$.

Theorem 3.3. *A q -topology τ satisfies axiom A5° iff $C(\tau)$ is a JC^* -subalgebra of B .*

Proof. Necessity follows from Proposition 2.9 and the equality

$$C(\tau)^+ = \overline{A^+(\tau^{\text{reg}})} \cap \overline{A^+(1 - \tau^{\text{reg}})},$$

which can be easily deduced from Lemma 3.2 and axiom A5° in a manner similar to the proof of Theorem 3.1.

Sufficiency. If $C(\tau)$ is a Jordan algebra, then $C(\tau)^+$ is a convex cone and $A^+(\tau^{\text{reg}}) \subset (C(\tau)^+)^m$. Indeed, whenever $a = \sum_{i=1}^n \lambda_i e_i$, $\lambda_i \geq 0$, $e_i \in \tau^{\text{reg}}$, there exist increasing sequences $(b_{ik})_{k=1}^\infty \subset C(\tau)^+$, $i = 1, 2, \dots, n$, with $e_i = \bigvee_k b_{ik}$ and $a = \bigvee_k c_k$, where $c_k = \sum_{i=1}^n \lambda_i b_{ik}$ is an increasing sequence in $C(\tau)^+$. Now by Proposition 2.5 we have $(C(\tau)^+)^{1/2} \subset C(\tau)^+$ and the operator monotonicity of the function $t^{1/2}$ gives $((C(\tau)^+)^m)^{1/2} \subset (C(\tau)^+)^m \subset L(\tau)$. Similarly we can conclude that $A^+(1 - \tau^{\text{reg}})^{1/2} \subset U(\tau)$. Hence axiom A5° holds.

Definition 3.4. A q -topology τ is called *completely regular*, if for any q -point x disjoint from a closed q -set f there exists an element $a \in C(\tau)_1^+$ with $ax = x$ and $af = 0$ (this means that a takes the value 1 at x and the value 0 on f).

Any Akemann-Giles q -topology is completely regular [8; 4.7].

Theorem 3.5. *If a completely regular q -topology τ satisfies axiom A5°, then $\tau = \tau^{\text{reg}}$.*

Proof. Given any open q -set $e \in \tau$ put

$$I(e) = \{(1 + a)^{-1} \cdot a \mid a \in C(\tau)^+, a \leq \lambda e \text{ for some } \lambda > 0\}.$$

Whenever $(1 + a_i)^{-1} \cdot a_i \in I(e)$, $i = 1, 2$, we have $\tilde{a} = (1 + (a_1 + a_2))^{-1} \cdot (a_1 + a_2) \in I(e)$ and $(1 + a_i)^{-1} \cdot a_i \leq \tilde{a}$ for $i = 1, 2$ (since by [6; 16.8] the function $(1 + t)^{-1}$ is antimonotone and so the function $(1 + t)^{-1} \cdot t = 1 - (1 + t)^{-1}$ is operator monotone). This means that $I(e)$ is a directed set and there is a supremum e_1 of $I(e)$ in B^s . To complete the proof we shall show that $e = e_1$. The implication $a \leq \lambda e \Rightarrow (1 + a)^{-1} \cdot a \leq e$ gives $e_1 \leq e$. Inasmuch τ is completely regular, for any q -point $x \leq e$ there exists $a_x \in C(\tau)^+$ with $a_x x = x$ and $a_x \leq e$. So for each natural n we have $(1 + na_x)^{-1} na_x \in I(e)$ whence $e_1 \geq (1 + na_x)^{-1} na_x \geq (1 + nx)^{-1} \cdot nx$ and $e_1 \geq x$. Finally, we have $e_1 \geq \bigvee_{x \leq e} x = e$ and $e_1 = e$.

Corollary 3.6. *A complete regular q -topology τ satisfies axiom A5 iff the set $C(\tau)$ is a JC^* -subalgebra of B .*

4. Symmetry: the sixth axiom. Every unitary $u \in B$ (i.e. such that $uu^* = 1 = u^*u$) yields an $*$ automorphism $\phi_u : a \rightarrow u^*au$ of the W^* -algebra B , which induces an automorphism of the lattice $\text{Pr}B$ onto itself. If τ is a q -topology in B and a unitary element u is τ -continuous, it is very natural to require ϕ_u to be a “homeomorphism” of τ . Such requirement seems to be independent of axioms A1 – A5 and so it becomes our last axiom.

AXIOM A6. $u^* \tau u \subset \tau$ for any unitary $u \in C(\tau)$.

As a matter of fact we need only the weakened variant of A6 like that in § 3.

AXIOM A6°. $u^* \tau^{reg} u \subset \tau$ for any unitary $u \in C(\tau)$.

Theorem 4.1. *Let τ be a q -topology. Then $C(\tau)$ is a C^* -subalgebra of B iff τ satisfies axioms A5° and A6°.*

Proof. Necessity. In view of Theorem 3.3 we need to check axiom A6° only. Take a unitary $u \in C(\tau)$, then $u^* C(\tau)^+ u = C(\tau)^+$ since φ_u is an $*$ automorphism and $C(\tau)$ is a C^* -algebra. Axiom A6° holds because

$$\begin{aligned} u^* \tau^{reg} u &= u^* ((C(\tau)^+)^n \cap \text{Pr} B) u = \\ &= u^* (C(\tau)^+)^n u \cap \text{Pr} B = (C(\tau)^+)^n \cap \text{Pr} B = \tau^{reg}. \end{aligned}$$

Sufficiency. By Theorem 3.3, $C(\tau)$ is a JC^* -subalgebra of B and we have to prove that $i[a, b] = i(ab - ba) \in C(\tau)^s$ whenever $a, b \in C(\tau)^s$ for this implies that $ab = a \circ b + \frac{1}{2}i[a, b] \in C(\tau)$. For any $t > 0$ consider the unitary element

$$u_t = \exp(itb) \equiv \sum_{n=0}^{\infty} \frac{(it)^n}{n!} b^n \in C(\tau)^s.$$

Then $(1/t)(u_t^* a u_t - a) = a_t$ uniformly converges to $i[a, b]$ as t tends to 0. Since for any unitary $u \in C(\tau)$ we have by the spectral theory, Lemma 3.2 and by virtue of axiom A6°

$$u^* C(\tau)^s u = u^* C(\tau^{reg})^s u \subset C(u^* \tau^{reg} u)^s \subset C(\tau)^s,$$

the elements a_t belong to $C(\tau)^s$ and thus $i[a, b] \in C(\tau)^s$.

Definition 4.2. A q -topology τ is a T_1 q -topology, if for any two disjoint q -points x and y there exists an open q -set $e \in \tau$ with $e \supseteq x$ and $ey = 0$. It also means that any q -point is τ -closed.

Proposition 4.3. *Let a T_1 q -topology τ satisfy axioms A5° and A6°, then τ is completely regular iff $\tau = \tau^{reg}$.*

Proof. By virtue of Theorem 3.5 it is enough to prove sufficiency. Since τ is T_1 , elements of $C(\tau)$ distinguish normal pure states of B and so the C^* -algebra $C(\tau)$ is weakly dense in B . For any $e \in \tau$ consider the C^* -subalgebra $A(e) = \{a \in C(\tau) \mid eae = a\}$. By [8; 4.2 and 4.5] $A(e)$ is weakly dense in the W^* -algebra eBe , so by the transitivity theorem [8; 2.7] for any q -point $x \leq e$ there exists $a \in A(e)_1^+$ (i.e. $a \leq e$) with $ax = x$.

We now define a C^* -topology as a q -topology which obeys axioms A5 and A6. Any Akemann-Giles q -topology is a C^* -topology.

5. Compactness. Ch. Akemann introduced the notion of q -compactness in terms of the C^* -algebra A (see 1.3), but an intrinsic q -topological description of q -compact q -sets has not been given. Nevertheless, Akemann showed in [3] that the intersection

condition for a q -set p (for any decreasing net (q_α) of τ_A -closed q -sets $\forall p \wedge (\bigwedge_{i=1}^n q_{\alpha_i}) \neq 0$ implies $p \wedge (\bigwedge_\alpha q_\alpha) \neq 0$) and the regularity after Effros [7] follow from the q -compactness of p , it being unknown whether these conditions are sufficient. We make use of a multiplicative version of the intersection condition.

Let us define a q -topological space to be a pair (B, τ) where τ is a q -topology. A q -set $p \in \text{Pr}B$ is called *regular* if for any open q -set $e \in \tau$ $\|pe\| = \|\bar{p}e\|$ (\bar{p} is τ -closure of p).

Definition 5.1. Let (B, τ) be a q -topological space. A q -set $p \in \text{Pr}B$ is called *quasi-compact* if for any decreasing net $(b_\alpha) \subset U(\tau)^+$ with $b = \bigwedge_\alpha b_\alpha \in U(\tau)^+$, $\inf \|pb_\alpha p\| = \|pbp\|$. If (B, τ) is a completely regular T_1 q -topological space, then a q -set $p \in \text{Pr}B$ is called *compact* if p is quasicompact and regular. (B, τ) is called a *compact q -space* if the unit 1 is a compact q -set.

Proposition 5.2. *If a completely regular T_1 q -topology satisfies axiom A5, then any compact q -set p is closed.*

Proof. Suppose, on the contrary, that $p \neq \bar{p}$ and consider a q -point x with $x \leq \bar{p} - p$. By Theorem 3.5 there exists an increasing net $(a_\alpha) \subset C(\tau)_1^+$ with $1 - x = \bigvee_\alpha a_\alpha$. Put $b = 1 - a_\alpha$, then $x = \bigwedge_\alpha b_\alpha$, $(b_\alpha) \subset C(\tau)^+ \subset U(\tau)^+$. For all natural n and all α we have $x \leq E_{b_\alpha}((1 - 1/n, \infty)) \equiv e_{\alpha n} \in \tau$ and $b_\alpha \geq ((n - 1)/n) \cdot e_{\alpha n}$. Since p is regular we see that

$$\begin{aligned} 1 &\geq \|pb_\alpha p\| \geq \frac{n-1}{n} \|pe_{\alpha n} p\| = \frac{n-1}{n} \|\bar{p}e_{\alpha n} \bar{p}\| \geq \frac{n-1}{n} \|\bar{p}x \bar{p}\| = \\ &= \frac{n-1}{n} \|x\| = \frac{n-1}{n} \end{aligned}$$

and finally $\|pb_\alpha p\| = 1$. Since p is compact, this implies that $\|\bar{p}x \bar{p}\| = 1$ which contradicts $xp = 0$.

Any Akemann-Giles q -topology is T_1 complete regular [3; III.1 and 8; 3.9] so Definition 5.1 of compact q -sets is applicable.

Theorem 5.3. *Let (B_A, τ_A) be the q -spectrum of a C^* -algebra A . A q -set $p \in \text{Pr}B_A$ is q -compact after Akemann (see 1.3) iff p is compact in the sense of Definition 5.1.*

Proof. In §2 we have mentioned an isomorphism of \bar{A}_m on the cone $U^q(S_A)$. With any q -closed q -set $p \in 1 - \tau_A$ this isomorphism correlates the USC affine function \hat{p} on S_A and the closed face $F(p) = \{\varphi \in S_A \mid \hat{p}(\varphi) = 1\}$. The map $p \mapsto F(p)$ is induced by the Effros-Akemann correspondence between the q -closed q -sets and the $\sigma(A^*, A)$ -closed order ideals of A^* ([7], [2]). So that map is a bijection of $1 - \tau_A$ on the set of all closed faces of S_A and for each $p \in 1 - \tau_A$ and $b \in \bar{A}_m$ we have $\|pbp\| = \max_{\varphi \in F(p)} |\hat{b}(\varphi)|$.

Notice that a q -closed p is q -compact after Akemann iff $F(p)$ is a compact subset of S_A . It may be proved in the same way as Akemann-Urysohn's lemma [3; III.1].

Necessity. It is enough to prove that p is quasicompact. Consider a decreasing net $(b_\alpha) \subset U(\tau)^+$ with $b = \bigwedge_\alpha b_\alpha$ and put $\lambda = \inf_\alpha \|pb_\alpha p\| \geq \|pbp\|$. For each α , $F_\alpha = \{\varphi \in F(p) \mid \hat{b}_\alpha(\varphi) \geq \lambda\} \neq \emptyset$ is a closed subset of the compact set $F(p)$ so $F_0 = \bigcap_x F_\alpha \neq \emptyset$. Take $\tilde{\varphi} \in F_0$, then $\hat{b}(\tilde{\varphi}) = \inf_\alpha \hat{b}_\alpha(\tilde{\varphi}) \geq \lambda$ whence $\|pbp\| = \max_{\varphi \in F(p)} \hat{b}(\varphi) \geq \lambda$.

Sufficiency. If $A \ni 1$, it follows from Proposition 5.2. Let $A \bar{\ni} 1$ and $\tilde{A} = A + \mathcal{C}$. 1 as in 2.3. Denote as $\overline{F(p)} \subset S_{\tilde{A}}$ the $\sigma(\tilde{A}^*, \tilde{A})$ -closure of $F(p)$. Since $S_{\tilde{A}}$ is compact, it is enough to show $F(p) = \overline{F(p)}$. For any $\varphi \in \overline{F(p)}$ we have $\varphi = \varphi_0 + \lambda\varphi_\infty$, where $\varphi_0 \in A^{*+}$, $\lambda \geq 0$, φ_∞ is the unique pure state of \tilde{A} which vanishes on A and $\|\varphi\| = \|\varphi_0\| + |\lambda|$. We shall show $\|\varphi\| = \|\varphi_0\|$, whence $\lambda = 0$ and $\varphi \in F(p)$. Let $(u_\alpha) \subset A^+$ be an increasing approximate unit, then $\|pu_\alpha p - p\| \rightarrow 0$ since $(1 - u_\alpha) \subset C(\tau_A)^+$, $0 = \bigwedge_\alpha (1 - u_\alpha)$ in B^+ and p is compact. Let us choose $u_n = u_{\alpha_n}$ with $u_{n+1} \geq u_n$ and $pu_n p \geq p - 2^{-n}p$, n being natural. Consider the closed subsets $F_n = \{\theta \in \tilde{A}^{*+} \mid \hat{u}_n(\theta) \geq 1 - 2^{-n}\} \subset \tilde{A}^+$; then $F(p) \subset F_n$, hence $\overline{F(p)} \subset F_n$. This means that $\varphi(u_n) = \varphi_0(u_n) \geq 1 - 2^{-n}$. Since n was arbitrary, $\|\varphi_0\| = 1 = \|\varphi\|$ as desired.

Definition 5.4. A completely regular T_1 q -topological space (B, τ) is called *locally compact* if for any q -point x there exists an open q -set $e \geq x$ with the compact q -closure \bar{e} .

Theorem 5.5. A q -topological space (B, τ) is the q -spectrum of a certain C^* -algebra iff τ is a locally compact C^* -topology.

Proof. Necessity. Let A be a C^* -algebra, then τ_A is the complete regular T_2 C^* -topology by the above. For any q -point $x \in B_A$ take $a \in A_1^+$ with $ax = x$, then $e = E_a(\frac{1}{2}, \infty) = E_a([\frac{1}{2}, 1]) \geq x$, $e \in \tau$ and $\bar{e} \leq E_a([\frac{1}{2}, 1]) \leq 2a$. So e is q -compact after Akemann, hence e is compact by Theorem 5.3.

Sufficiency. Let (B, τ) be a locally compact C^* -topological space. Then $C(\tau)$ is the weakly dense C^* -subalgebra of B and $\tau = \text{Pr}B \cap (C^+(\tau))^m$ (see 4.3). Put $(\hat{B}, \hat{\tau}) = q \text{ spec } C(\tau)$. By [8; 3.4] there exists a central projection $z \in \text{Pr}\hat{B}$ with $B \approx z\hat{B}$. Since the identification $B = z\hat{B}$ agrees with the inclusions $C(\tau) \subset B$ and $C(\tau) \subset \hat{B}$, we have

$$\tau = \text{Pr}B \cap (C(\tau)^+)^m = \text{Pr}(z\hat{B}) \cap z(C(\tau)^+)^m = z(\text{Pr}\hat{B} \cap (C(\tau)^+)^m) = z\hat{\tau}.$$

Let us show $z \in \hat{\tau}$. By hypothesis, if $x \leq z$ is a q -point in \hat{B} (hence in B) there exists $e \in \tau$ with $e \geq x$ and \bar{e} q -compact. Besides, there exists $a \in C(\tau)_1^+$ with $ax = x$ and $a \leq e$. Let p be the support of a in \hat{B} . Then $p \in \hat{\tau}$ and $x \leq p$. If we show $p \leq z$, the assertion will follow for then $z = \bigvee \{p \in \hat{\tau} \mid x \leq p \leq z, x \text{ is a } q\text{-point}\}$ which is $\hat{\tau}$ -open. To prove $p \leq z$ we consider any q -point $y \leq 1 - z$ and show $ay = 0$. Indeed, by [8; 3.9 and 4.2] there exists a decreasing net $(b_\alpha) \subset C(\tau)_1^+$, with $y = \bigwedge_\alpha b_\alpha$

in \hat{B} . Then $\|aya\| \leq \inf_{\alpha} \|ab_{\alpha}a\|$ and

$$\|ab_{\alpha}a\| = \|(b_{\alpha}^{1/2}a) * (b_{\alpha}^{1/2}a)\| = \|b^{1/2}a^2b^{1/2}\| \leq \|b^{1/2}\bar{e}b^{1/2}\| = \|\bar{e}b_{\alpha}\bar{e}\|.$$

Since $0 = \bigwedge_{\alpha}(zb_{\alpha})$ we have $\|\bar{e}b_{\alpha}\bar{e}\| \rightarrow 0$, for \bar{e} is a compact q -set. Thus $\|aya\| = 0$, i.e. $ay = 0$. It implies that $py = 0$, hence $p \leq z$. So we have that z is $\hat{\tau}$ -open. Now consider $A = \{a \in C(\tau) \mid az = a\}$. Then by [8; 5.9] $q \text{ spec } A = (z\hat{B}, z\hat{\tau}) = (B, \tau)$, which completes the proof.

Corollary 5.6. *A C^* -topological space (B, τ) is compact iff $(B, \tau) = q \text{ spec } C(\tau)$.*

This last theorem shows that for an arbitrary locally compact C^* -topological space (B, τ) the q -space $(\hat{B}, \hat{\tau}) = q \text{ spec } C(\tau)$ may be described as "the Stone-Čech compactification of (B, τ) ".

References

- [1] C. A. Akemann: The general Stone-Weierstrass problem for C^* -algebras, J. Functional Analysis, 4 (2) (1969), 277—294.
- [2] C. A. Akemann: Left ideal structure of C^* -algebras, J. Functional Analysis, 6 (2) (1970), 306—317.
- [3] C. A. Akemann: A Gelfand representation theory for C^* -algebras, Pacific J. Math., 39 (1) (1971), 1—11.
- [4] C. A. Akemann and G. K. Pedersen: Complications of semicontinuity in C^* -algebra theory, Duke Math. J., 40 (4) (1973), 785—796.
- [5] C. A. Akemann, G. K. Pedersen and J. Tomiyama: Multipliers of C^* -algebras, J. Functional Analysis, 13 (3) (1973), 277—301.
- [6] J. Dixmier: Les C^* -algèbres et leurs représentations, Gauthier-Villars, Paris, 1964.
- [7] E. G. Effros: Order ideals in a C^* -algebra and its dual, Duke Math. J., 30 (1963), 391—412.
- [8] R. Giles and H. Kummer: A non-commutative generalization of topology, Indiana Univ. Math. J. 21 (1) (1971), 91—102.
- [9] R. Giles: Foundations for quantum mechanics, J. Mathematical Phys., 11 (1970), 2139 to 2160.
- [10] G. K. Pedersen: Applications of weak* semicontinuity in C^* -algebra theory, Duke Math. J., 39 (3) (1972), 413—430.
- [11] G. K. Pedersen: Some operator monotone functions, Proc. Amer. Math. Soc., 36 (1972), 309—310.
- [12] S. Sakai: C^* -algebras and W^* -algebras, Springer-Verlag, Berlin—Heidelberg—New York, 1971.

Author's address: University of People's Friendship, Dept. of Math. Analysis, Ordzhonikidze 3, 117923 Moscow, USSR.