

Nelson Merentes; Sergio Rivas

On characterization of the Lipschitzian composition operator between spaces of functions of bounded  $p$ -variation

*Czechoslovak Mathematical Journal*, Vol. 45 (1995), No. 4, 627–637

Persistent URL: <http://dml.cz/dmlcz/128558>

## Terms of use:

© Institute of Mathematics AS CR, 1995

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON CHARACTERIZATION OF THE LIPSCHITZIAN COMPOSITION  
OPERATOR BETWEEN SPACES OF FUNCTIONS  
OF BOUNDED  $p$ -VARIATION

N. MERENTES, S. RIVAS, Caracas

(Received November 18, 1993)

INTRODUCTION

Let  $I = [a, b]$  be an interval,  $f: I \times \mathbb{R} \rightarrow \mathbb{R}$  a fixed two-place function, and  $\mathcal{F}(I)$  the linear space of all functions  $u: I \rightarrow \mathbb{R}$ . The function  $F: \mathcal{F}(I) \rightarrow \mathcal{F}(I)$  given by the formula

$$(Fu)(t) := f(t, u(t)) \quad t \in I, u \in \mathcal{F}(I),$$

is called a composition operator. In [4] it is proved that a composition operator  $F$  maps the space  $\text{Lip}(I)$  of all Lipschitzian function into itself and is globally Lipschitzian if and only if  $f(t, x) = g(t)x + h(t)$ , where  $g, h \in \text{Lip}(I)$ .

This result has been further extended to some other function Banach spaces (see [1–7]). Recently N. Merentes (see [7]) proved an analogous theorem in the space  $RV_p[a, b]$  of functions of bounded  $p$ -variation in the sense of Riesz ( $1 < p < \infty$ ). In the present paper we generalize these results in the case that the composition operator  $F$  is globally Lipschitzian between spaces  $RV_p[a, b]$  and  $RV_q[a, b]$  where  $1 \leq q \leq p$ . On the other hand, if  $1 \leq p < q$ , the composition operator  $F$  is constant.

1. PRELIMINARY RESULTS

Given  $1 \leq p < \infty$  and  $u: [a, b] \rightarrow \mathbb{R}$ , we write

$$V_p(u; \pi) := \sup_{\pi} \sum_{i=1}^n \frac{|u(t_i) - u(t_{i-1})|^p}{|t_i - t_{i-1}|^{p-1}}$$

for the  $p$ -variation of the function  $u$  in the sense of Riesz, where the supremum is taken over all partitions  $\pi: a = t_0 < \dots < t_n = b$  of the interval  $[a, b]$ . By

$RV_p = RV_p[a, b]$  we denote the Banach space of all functions  $u$  on  $[a, b]$  for which the norm

$$\|u\|_p := |u(a)| + (V_p(u; [a, b]))^{\frac{1}{p}}$$

is finite. Usually, one takes  $BV_\infty[a, b]$  as the space  $\text{Lip}[a, b]$  of all Lipschitzian functions on  $[a, b]$  with the norm

$$\|u\|_{\text{Lip}[a, b]} := |u(a)| + \sup_{x \neq y} \frac{|u(x) - u(y)|}{|x - y|}.$$

Moreover, the space  $RV_1[a, b]$  is simply denoted by  $BV[a, b]$  and it is the classical space of functions of bounded variation on  $[a, b]$ .

It is easy to see that if  $p > 1$ , then every function  $u \in RV_p[a, b]$  is continuous. More precisely, the inclusions

$$\text{Lip}[a, b] \subset RV_p[a, b] \subset AC[a, b] \subset BV[a, b] \quad (p > 1)$$

hold, where  $AC[a, b]$  is the space of all absolutely continuous functions.

**Lemma 1** ([8], Riesz). *Let  $1 < p < \infty$  be a fixed number. A function  $u$  fulfills  $u \in RV_p[a, b]$  if and only if  $u \in AC[a, b]$  and  $u' \in L_p[a, b]$ . In that case we also have the equality*

$$V_p(u; [a, b]) = \int_a^b |u'(t)|^p dt.$$

F. Szigeti (see [9], p. 13) proved that the space  $RV_p[a, b]$  ( $1 < p < \infty$ ) is also a Banach algebra.

In [7] it is proved that the composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  maps the space  $RV_p[a, b]$  ( $1 < p < \infty$ ) into itself and is globally Lipschitzian if and only if  $f(t, x) = g(t)x + h(t)$  ( $t \in [a, b]; x \in \mathbb{R}$ ) for some  $g, h \in RV_p[a, b]$ . In the case  $p = 1$ , J. Matkowski and J. Miś (see [6]) proved that the composition operator  $F$ , generated by  $f$ , maps the space  $BV[a, b]$  into itself and satisfies the global Lipschitzian condition if and only if

$$\bar{f}(x, y) = g(x)y + h(x)$$

for two functions  $g, h \in NBV[a, b]$ , where

$$\bar{f}(x, y) = \lim_{\delta \rightarrow 0} f(x - \delta, y) \quad (y \in \mathbb{R})$$

is the left-continuous regularization of  $f$  and  $NBV[a, b]$  is the subspace of all functions  $u \in BV[a, b]$  such that  $u$  is continuous on  $[a, b]$  from the left.

## MAIN RESULTS

In this section we will present a characterization of functions  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  for which the composition operator  $F$  generated by  $f$  maps the space  $RV_p[a, b]$  into the space  $RV_q[a, b]$  ( $1 \leq q \leq p$ ) and is globally Lipschitzian. In the case  $1 \leq p < q$ , the composition operator is constant.

**Theorem 1.** *Let  $p, q$  be real numbers such that  $1 < q \leq p$ . The composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  maps the space  $RV_q[a, b]$  into the space  $RV_p[a, b]$  and is globally Lipschitzian if and only if the function  $f$  satisfies the following conditions:*

a) For all  $t \in [a, b]$  there exists  $M(t) > 0$  such that

$$(1) \quad |f(t, x) - f(t, y)| \leq M(t)|x - y| \quad (x, y \in \mathbb{R}),$$

b)

$$(2) \quad f(t, x) = g(t)x + h(t) \quad (t \in [a, b], x \in \mathbb{R}),$$

where  $g, h \in RV_q[a, b]$ .

*Proof.* Suppose that there exist  $g, h \in RV_q[a, b]$  such that  $f(t, x) = g(t)x + h(t)$  ( $t \in [a, b], x \in \mathbb{R}$ ). Then the composition  $F$  generated by  $f$  is given by

$$(Fu)(t) = g(t)u(t) + h(t) \quad (t \in [a, b], u \in RV_q[a, b]).$$

Since  $F(RV_p[a, b]) \subset RV_q[a, b]$  ( $1 < q \leq p$ ) and  $RV_q[a, b]$  is a Banach algebra, then  $Fu \in RV_q[a, b]$  for all  $u \in RV_p[a, b]$ .

Moreover,

$$\|Fu_1 - Fu_2\|_q \leq \|g\|_q \|u_1 - u_2\|_p \quad (u_1, u_2 \in RV_p[a, b]).$$

Thus, the composition operator  $F$  maps the space  $RV_p[a, b]$  into the space  $RV_q[a, b]$  and is globally Lipschitzian.

Suppose now that  $F: RV_p[a, b] \rightarrow RV_q[a, b]$  ( $1 < q \leq p$ ) is globally Lipschitzian, then there exists a constant  $M > 0$  such that

$$\|Fu_1 - Fu_2\|_q \leq M \|u_1 - u_2\|_p \quad (u_1, u_2 \in RV_p[a, b]).$$

Let  $t \in (a, b]$ . Using the definition of the operator  $F$  and of the norm  $\|\cdot\|_q$  we have

$$(3) \quad |f(t, u_1(t)) - f(t, u_2(t)) - f(a, u_1(a)) + f(a, u_2(a))| \leq M|t - a|^{1-\frac{1}{q}} \|u_1 - u_2\|_p$$

for all  $u_1, u_2 \in RV_p[a, b]$ .

Define a function  $\alpha: [a, b] \rightarrow \mathbb{R}$  by

$$\alpha(\tau) := \begin{cases} \frac{\tau - a}{t - a}, & a \leq \tau \leq t, \\ 1, & t \leq \tau \leq b. \end{cases}$$

We have  $\alpha \in RV_p[a, b]$  and

$$V_p(\alpha; [a, b]) = \frac{1}{|t - a|^{p-1}}.$$

Let us fix  $x, y \in \mathbb{R}$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  ( $i = 1, 2$ ) by

$$(4) \quad u_1(\tau) := x, \quad \tau \in [a, b], \quad u_2(\tau) := \alpha(\tau)(y - x) + x, \quad \tau \in [a, b].$$

The functions  $u_i$  fulfill  $u_i \in RV_p([a, b])$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = (V_p(\alpha; [a, b]))^{\frac{1}{p}} |x - y| = \frac{|x - y|}{|t - a|^{1 - \frac{1}{p}}}.$$

Hence, substituting into the inequality (3) the particular functions  $u_i$  ( $i = 1, 2$ ) defined by (4), we obtain

$$(5) \quad |f(t, x) - f(t, y)| \leq M \frac{|t - a|^{1 - \frac{1}{q}}}{|t - a|^{1 - \frac{1}{p}}} |x - y|$$

for all  $t \in (a, b]$ ,  $x, y \in \mathbb{R}$ .

Now, let  $t = a$ . Define a function  $\beta: [a, b] \rightarrow \mathbb{R}$  by

$$\beta(\tau) := \frac{\tau - a}{b - a} \quad (\tau \in [a, b]).$$

The function  $\beta$  fulfills  $\beta \in RV_p[a, b]$  and

$$V_p(\beta; [a, b]) = \frac{1}{|b - a|^{p-1}}.$$

Let us fix  $x, y \in \mathbb{R}$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  ( $i = 1, 2$ ) by

$$(6) \quad u_1(\tau) := x, \quad \tau \in [a, b], \quad u_2(\tau) := \beta(\tau)(x - y) + y, \quad \tau \in [a, b].$$

The functions  $u_i$  fulfill  $u_i \in RV_p[a, b]$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = \left(1 + (V_p(\beta; [a, b]))^{\frac{1}{p}}\right) |x - y| = \left(1 + \frac{1}{|b - a|^{1 - \frac{1}{p}}}\right) |x - y|.$$

Hence, substituting into the inequality (3) the particular functions  $u_i$  ( $i = 1, 2$ ) defined by (6), we obtain

$$|f(a, x) - f(a, y)| \leq M|b - a|^{1-\frac{1}{q}} \left(1 + \frac{1}{|b - a|^{1-\frac{1}{p}}}\right) |x - y|$$

for all  $x, y \in \mathbb{R}$ .

Define a function  $M: [a, b] \rightarrow \mathbb{R}$  by

$$M(t) := \begin{cases} M \frac{|t - a|^{1-\frac{1}{q}}}{|t - a|^{1-\frac{1}{p}}}, & a < t \leq b, \\ M|b - a|^{1-\frac{1}{q}} \left(1 + \frac{1}{|b - a|^{1-\frac{1}{p}}}\right), & t = a. \end{cases}$$

Hence we have for all  $t \in [a, b]$  that there exists  $M(t) > 0$  such that the inequality (1) holds. Thus for all  $t \in [a, b]$  the function  $f(t, \cdot): \mathbb{R} \rightarrow \mathbb{R}$  is continuous.

Next we shall prove that  $f$  satisfies the equality (2).

Let us fix  $t, t_0 \in [a, b]$  such that  $t_0 < t$ . Since the composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  is globally Lipschitzian between  $RV_p[a, b]$  and  $RV_q[a, b]$  ( $1 < q \leq p$ ), there exists a constant  $M > 0$  such that

$$(7) \quad |f(t, u_1(t)) - f(t, u_2(t)) - f(t_0, u_1(t_0)) + f(t_0, u_2(t_0))| \leq M \|u_1 - u_2\|_p |t - t_0|^{1-\frac{1}{q}}$$

for all  $u_1, u_2 \in RV_p[a, b]$ .

Define a function  $\gamma: [a, b] \rightarrow \mathbb{R}$  by

$$\gamma(\tau) := \begin{cases} \frac{\tau - a}{t_0 - a}, & a \leq \tau \leq t_0, \\ -\frac{\tau - t}{t - t_0}, & t_0 \leq \tau \leq t, \\ 0, & t \leq \tau \leq b. \end{cases}$$

The function  $\gamma$  fulfills  $\gamma \in RV_p[a, b]$ . Let us fix  $x, y \in \mathbb{R}$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  by

$$(8) \quad \begin{aligned} u_1(\tau) &:= \frac{\gamma(t)}{2}x + \left(1 + \frac{\gamma(\tau)}{2}\right)y \quad (\tau \in [a, b]), \\ u_2(\tau) &:= \frac{1 + \gamma(\tau)}{2}x + \frac{1 - \gamma(\tau)}{2}y \quad (\tau \in [a, b]). \end{aligned}$$

The functions  $u_i$  fulfill  $u_i \in RV_p[a, b]$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = \frac{|x - y|}{2}.$$

Hence, substituting into the inequality (7) the particular functions  $u_i$  ( $i = 1, 2$ ) defined by (8), we obtain

$$(9) \quad \left| f(t, y) - f\left(t, \frac{x+y}{2}\right) - f\left(t_0, \frac{x+y}{2}\right) + f(t_0, x) \right| \leq \frac{M}{2} |t - t_0|^{1-\frac{1}{q}} |x - y|.$$

Since  $F$  maps  $RV_p[a, b]$  into  $RV_q[a, b]$  ( $1 < q \leq p$ ), then for all  $x \in \mathbb{R}$  the function  $f(\cdot, x)$  is continuous on  $[a, b]$ . Consequently, letting  $t_0 \uparrow t$  in the inequality (9), we get

$$\left| f(t, y) - f\left(t, \frac{x+y}{2}\right) - f\left(t, \frac{x+y}{2}\right) + f(t, x) \right| = 0$$

for all  $t \in [a, b]$  and  $x, y \in \mathbb{R}$ .

Thus for all  $t \in [a, b]$ ,  $x, y \in \mathbb{R}$ , we have

$$\frac{f(t, x) + f(t, y)}{2} = f\left(t, \frac{x+y}{2}\right).$$

Consequently, for all  $t \in [a, b]$  the function  $f(t, \cdot): \mathbb{R} \rightarrow \mathbb{R}$  satisfies the Jensen equation and since the function  $f(t, \cdot)$  is continuous on  $\mathbb{R}$ , we have that there exist two functions  $g, h: [a, b] \rightarrow \mathbb{R}$  such that

$$f(t, x) = g(t)x + h(t), \quad (t \in [a, b], x \in \mathbb{R}).$$

Since  $h(t) = f(t, 0) = F(0)$ ,  $g(t) = f(t, 1) - f(1, 0) = F(1) - F(0)$  and  $F$  maps  $RV_p[a, b]$  into  $RV_q[a, b]$ , we conclude  $g, h \in RV_q[a, b]$ .  $\square$

**Remark 1.** It is easy to observe that the above theorem remains true if there exist Banach spaces  $(X, \|\cdot\|_X)$  and  $(Y, \|\cdot\|_Y)$  such that  $RV_p[a, b] \hookrightarrow X \hookrightarrow Y \hookrightarrow RV_q[a, b]$  ( $1 < q \leq p$ ) and the composition operator  $F$  maps the space  $X$  into the space  $Y$  and is globally Lipschitzian.

**Theorem 2.** *Let  $p, q$  be real numbers such that  $1 < p < q$ . If the composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  maps the space  $RV_p[a, b]$  into the space  $RV_q[a, b]$  and is globally Lipschitzian, then the function  $f$  satisfies the condition*

$$f(t, x) = f(t, 0) \quad (t \in [a, b], x \in \mathbb{R}).$$

As an immediate consequence of Theorem 2 we obtain that the composition operator  $F$  is constant.

PROOF. Since the composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ , maps the space  $RV_p[a, b]$  into the space  $RV_q[a, b]$  ( $1 < p < q$ ) and is globally Lipschitzian, there exists a constant  $M > 0$  such that

$$\|Fu_1 - Fu_2\|_q \leq M\|u_1 - u_2\|_p \quad (u_1, u_2 \in RV_p[a, b]).$$

Let us fix  $t, t_0 \in [a, b]$  such that  $t_0 < t$ . Using the definitions of the operator  $F$  and of the norm  $\|\cdot\|_q$ , we have

$$(10) \quad \begin{aligned} & |f(t, u_1(t)) - f(t, u_2(t)) - f(t_0, u_1(t_0)) + f(t_0, u_2(t_0))| \\ & \leq M|t - t_0|^{1-\frac{1}{q}}\|u_1 - u_2\|_p \quad (u_1, u_2 \in RV[a, b]). \end{aligned}$$

Define a function  $\alpha: [a, b] \rightarrow \mathbb{R}$  by

$$\alpha(\tau) := \begin{cases} 1, & a \leq \tau \leq t_0, \\ -\frac{\tau - t}{t - t_0}, & t_0 \leq \tau \leq t, \\ 0, & t \leq \tau \leq b. \end{cases}$$

The function  $\alpha$  fulfills  $\alpha \in RV_p[a, b]$  and

$$V_p(\alpha; [a, b]) = \frac{1}{|t - t_0|^{p-1}}.$$

Let us fix  $x \in \mathbb{R}$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  ( $i = 1, 2$ ) by

$$(11) \quad u_1(\tau) := x \quad \tau \in [a, b], \quad u_2(\tau) := \alpha(\tau)x \quad \tau \in [a, b].$$

The functions  $u_i$  fulfill  $u_i \in RV_p[a, b]$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = \frac{|x|}{|t - t_0|^{1-\frac{1}{p}}}.$$

Hence, substituting into the inequality (10) the particular functions  $u_i$  ( $i = 1, 2$ ) defined by (11), we obtain

$$(12) \quad |f(t, x) - f(t, 0)| \leq M \frac{|t - t_0|^{1-\frac{1}{q}}}{|t - t_0|^{1-\frac{1}{p}}}|x|.$$

Since  $q > p$ , letting  $t_0 \uparrow t$  in the inequality (12) we obtain

$$f(t, x) = f(t, 0) \quad (t \in [a, b], x \in \mathbb{R}).$$



Next we shall consider the case when the composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  maps the space  $RV_p[a, b]$  into the space  $BV[a, b]$ . In this case a similar result holds for the left regularization  $f^*: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  of the function  $f$  defined by

$$f^*(t, x) := \begin{cases} \lim_{s \uparrow t} f(s, x), & t \in (a, b], x \in \mathbb{R}, \\ \lim_{s \downarrow a} \lim_{v \uparrow s} f(v, x), & t = a, x \in \mathbb{R}. \end{cases}$$

□

**Theorem 3.** *Let  $p$  be a real number such that  $1 < p < \infty$ . The composition operator  $F$  generated by  $f: [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$  maps the space  $RV_p[a, b]$  into the space  $BV[a, b]$  and if it is globally Lipschitzian, then the function  $f$  satisfies the following conditions:*

a) For each  $t \in [a, b]$  there exists  $M(t) > 0$  such that

$$(13) \quad |f^*(t, x) - f^*(t, y)| \leq M(t)|x - y| \quad (x, y \in \mathbb{R}),$$

b)

$$(14) \quad f^*(t, x) = g(t)x + h(t) \quad (t \in [a, b], x \in \mathbb{R}),$$

where  $g, h \in NBV[a, b]$ .

**Proof.** Let  $t \in [a, b)$  and define a function  $\alpha: [a, b] \rightarrow \mathbb{R}$  by

$$\alpha(t) := \begin{cases} 1, & a \leq \tau \leq t, \\ \frac{\tau - b}{t - b}, & t \leq \tau \leq b. \end{cases}$$

The function  $\alpha$  fulfills  $\alpha \in RV_p[a, b]$  and

$$V_p(\alpha, [a, b]) = \frac{1}{|b - t|^{p-1}}.$$

Let us fix  $x, y \in K$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  ( $i = 1, 2$ ) by

$$(15) \quad u_1(\tau) := x \quad \tau \in [a, b], \quad u_2(\tau) := \alpha(\tau)(y - x) + x, \quad \tau \in [a, b].$$

The functions  $u_i$  fulfil  $u_i \in RV_p[a, b]$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = (V_p(\alpha; [a, b]))^{\frac{1}{p}} |x - y| = \left(1 \frac{1}{|b - t|^{1 - \frac{1}{p}}}\right) |x - y|.$$

Since the composition operator  $F$  is globally Lipschitzian between  $RV_p[a, b]$  and  $BV[a, b]$ , there exists a constant  $M > 0$  such that

$$(16) \quad |f(b, u_1(b)) - f(b, u_2(b)) - f(t, u_1(t)) + f(t, u_2(t))| \leq M \|u_1 - u_2\|_p$$

for all  $u_1, u_2 \in RV_p[a, b]$ .

Hence, substituting into the inequality (16) the particular functions  $u_i$  ( $i = 1, 2$ ) defined by (15) we obtain

$$|f(t, y) - f(t, x)| \leq M \left[ 1 + \frac{1}{|b - t|^{1 - \frac{1}{p}}} \right]$$

for all  $t \in [a, b]$ .

In the case  $t = b$ , by a similar argument as above, we obtain that there exists a constant  $M(b) > 0$  such that

$$|f(b, x) - f(b, y)| \leq M(b)|x - y| \quad (x, y \in \mathbb{R}).$$

Thus, defining a function  $M: [a, b] \rightarrow \mathbb{R}$  by

$$M(t) := \begin{cases} 1 + \frac{1}{|b - t|^{1 - \frac{1}{p}}}, & t \in [a, b), \\ M(b), & t = b, \end{cases}$$

we obtain that for each  $t \in [a, b)$  there exists  $M(t) > 0$  such that

$$(17) \quad |f(t, x) - f(t, y)| \leq M(t)|x - y| \quad (t \in [a, b), x, y \in \mathbb{R}).$$

Hence, passing to the limit in the inequality (17), by the definition of  $f^*$  we have for all  $t \in [a, b]$  that there exists  $M(t) > 0$  such that

$$|f^*(t, x) - f^*(t, y)| \leq M(t)|x - y| \quad (x, y \in \mathbb{R}).$$

Next we shall prove that  $f^*$  satisfies the equality (14).

Let us fix  $t, t_0 \in [a, b]$ ,  $n \in \mathbb{N}$  such that  $t_0 < t$ . Define a partition  $\pi_n$  of the interval  $[t_0, t]$  by  $\pi_n: a < t_0 < t_1 < \dots < t_{2n-1} < t_{2n} = t$ , where

$$t_1 - t_{i-1} = \frac{t - t_0}{2n}, \quad i = 1, 2, \dots, 2n.$$

Since the composition operator  $F$  is globally Lipschitzian between  $RV_p[a, b]$  and  $BV[a, b]$ , there exists a constant  $M > 0$  such that

$$(18) \quad \sum_{i=1}^n |f(t_{2i}, u_1(t_{2i})) - f(t_{2i}, u_2(t_{2i})) - f(t_{2i-1}, u_1(t_{2i-1})) + f(t_{2i-1}, u_2(t_{2i-1}))| \leq M \|u_1 - u_2\|_p \quad (u_1, u_2 \in RV_p[a, b]).$$

Define a function  $\alpha: [a, b] \rightarrow \mathbb{R}$  in the following way:

$$\alpha(\tau) := \begin{cases} 0, & a \leq \tau \leq t_0, \\ \frac{\tau - t_{i-1}}{t_i - t_{i-1}}, & t_{i-1} \leq \tau \leq t_i, \quad i = 1, 3, \dots, 2n-1, \\ -\frac{\tau - t_i}{t_i - t_{i-1}}, & t_{i-1} \leq \tau \leq t_i, \quad i = 2, 4, \dots, 2n, \\ 0, & t \leq \tau \leq b. \end{cases}$$

The function  $\alpha$  fulfils  $\alpha \in RV_p[a, b]$  and

$$V_p(\alpha; [a, b]) = \frac{2^p n^p}{|t - t|^{p-1}}.$$

Let us fix  $x, y \in \mathbb{R}$  and define functions  $u_i: [a, b] \rightarrow \mathbb{R}$  by

$$(19) \quad \begin{aligned} u_1(\tau) &:= \frac{\alpha(\tau)}{2}x + \left(1 - \frac{\alpha(\tau)}{2}\right)y \quad (\tau \in [a, b]), \\ u_2(\tau) &:= \frac{1 + \alpha(\tau)}{2}x + \frac{1 - \alpha(\tau)}{2}y \quad (\tau \in [a, b]). \end{aligned}$$

The functions  $u_i$  fulfil  $u_i \in RV_p[a, b]$  ( $i = 1, 2$ ) and

$$\|u_1 - u_2\|_p = \frac{|x - y|}{2}.$$

Hence, substituting into the inequality (18) the particular functions  $u_i$  ( $i = 1, 2$ ) defined in (19), we obtain

$$(20) \quad \sum_{i=1}^n \left| f(t_{2i}, y) - f\left(t_{2i}, \frac{x+y}{2}\right) - f\left(t_{2i-1}, \frac{x+y}{2}\right) + f(t_{2i-1}, x) \right| \leq M \frac{|x-y|}{2}$$

for all  $x, y \in \mathbb{R}$ .

Since the composition operator  $F$  maps the space  $RV_p[a, b]$  into the space  $BV[a, b]$ , then  $f(\cdot, x) \in BV[a, b]$  for all  $x \in \mathbb{R}$ , thus letting  $t_0 \uparrow t$  in the inequality (20) we get

$$(21) \quad \left| f^*(t, y) - f^*\left(t, \frac{x+y}{2}\right) - f^*\left(t, \frac{x+y}{2}\right) + f^*(t, x) \right| \leq M \frac{|x-y|}{2n}$$

for all  $x, y \in \mathbb{R}$ ,  $n \in \mathbb{N}$ .

Passing to the limit for  $n \rightarrow \infty$  in the inequality (21), we get

$$\frac{f^*(t, y) + f^*(t, x)}{2} = f^*\left(t, \frac{x+y}{2}\right)$$

for all  $t \in [a, b]$ ,  $x, y \in \mathbb{R}$ .

Thus for all  $t \in [a, b]$ , the function  $f^*(t, \cdot): \mathbb{R} \rightarrow \mathbb{R}$  satisfies the Jensen equation and by property (a) of this theorem we get that there exist two functions  $g, h \in NBV[a, b]$  such that

$$f^*(t, x) = g(t)x + h(t) \quad (t \in [a, b], x \in \mathbb{R}).$$

□

**Remark 2.** It is easy to observe that the above theorem remains true if there exists a Banach space  $(X, \|\cdot\|_X)$  such that  $RV_p[a, b] \subset X \subset BV[a, b]$  ( $1 < p < \infty$ ) and the composition operator  $F$  maps the space  $X$  into the space  $BV[a, b]$  and is globally Lipschitzian.

#### References

- [1] *Matkowska A.*: On characterization of Lipschitzian operators of substitution in the class of Hölder functions. *Sci. Bul. Łódz Technical Univ.* 17 (1984), 81–85.
- [2] *Matkowska A., Matkowski J and Merentes N.*: Remark on globally Lipschitzian composition operators. submitted.
- [3] *Matkowski J.*: Form of Lipschitz operators of substitution in Banach spaces of differentiable functions. *Sci. Bull. Łódz Technical Univ.* 17 (1984), 5–10.
- [4] *Matkowski J.*: Functional equation and Nemytskii operator. *Funkcialaj Ekvacioj* 25 (1982), 127–132.
- [5] *Matkowski J. and Merentes N.*: Characterization of globally Lipschitzian composition operators in the Banach space  $BV_p^2[a, b]$ . *Archivum Math.* 28 (1992), 181–186.
- [6] *Matkowski J. and Miś J.*: On a characterization of Lipschitzian operators of substitution in the space  $BV[a, b]$ . *Math. Nachr.* 117 (1984), 155–159.
- [7] *Merentes N.*: On a characterization of Lipschitzian operators of substitution in the space of bounded Reisz  $\varphi$ -variation. *Annales Univ. Sci. Budapest* 34 (1991), 139–144.
- [8] *Riesz F.*: Untersuchungen über Systeme integrierbarer Funktionen. *Math. Annalen* 69 (1910), 449–497.
- [9] *Szigeti F.*: Composition of Sobolev functions and application. *Notas de Matemáticas, Univ. Los Andes, Venezuela* 86 (1987), 1–25.

*Authors' addresses:* N. Merentes, Central University of Venezuela, Caracas, Venezuela;  
S. Rivas, Open National University, Caracas, Venezuela.