

Ján Jakubík

On complete MV -algebras

Czechoslovak Mathematical Journal, Vol. 45 (1995), No. 3, 473–480

Persistent URL: <http://dml.cz/dmlcz/128535>

Terms of use:

© Institute of Mathematics AS CR, 1995

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON COMPLETE MV -ALGEBRAS

JÁN JAKUBÍK, Košice

(Received August 8, 1993)

Though the number of published papers on MV -algebras is rather large (the fundamental source are Chang's articles [1] and [2]), the terminology and notation in this field seem to be far from being unified. We will apply the terminology from [5], [6].

It is well-known that MV -algebras are term equivalent to Wajsberg algebras (called also W -algebras); cf., e.g., Cignoli [3]. Further, MV -algebras are categorically equivalent to bounded commutative BCK -algebras (cf. Mundici [8]); such BCK -algebras were studied by Traczyk [10].

Cignoli [3] studied the structure of MV -algebras which are complete and atomic. His main result is the following theorem:

- (*) ([3], Theorem 2.6.) An MV -algebra is complete and atomic if and only if it is a direct product of finite linearly ordered MV -algebras.

An MV -algebra \mathcal{A} which is a direct product of MV -algebras \mathcal{A}_i ($i \in I$) is complete if and only if all \mathcal{A}_i are complete. Further, a complete linearly ordered MV -algebra is atomic if and only if it is finite (cf. 1.3 below). Thus (*) can be expressed as follows:

- (**) An MV -algebra is complete and atomic if and only if it is a direct product of complete atomic linearly ordered algebras.

Let $\mathcal{A} = (A; \oplus, *, \neg, 0, 1)$ be an MV -algebra. We can introduce lattice operations \vee, \wedge , and hence also the corresponding partial order \leq on A (cf. Section 1 below). Let $0 < x \in A$ and let $\alpha > 1$ be a cardinal. The element x will be called an α -atom of \mathcal{A} if the interval $[0, x]$ is a chain having cardinality α . Hence the notion of the 2-atom coincides with the usual notion of the atom. The MV -algebra \mathcal{A} is said to be α -atomic if for each $0 < y \in A$ there exists an α -atom x of \mathcal{A} with $x \leq y$.

Let R be the additive group of all reals with the natural linear order. For each MV -algebra \mathcal{A} there exists a lattice ordered group G with a strong unit such that \mathcal{A} can be constructed by means of G (cf. $(*_2)$ and $(*_3)$ in Section 1 below). If G is isomorphic to R , then \mathcal{A} will be said to be of type R .

By applying the results of [6] the following will be proved in the present paper:

- (A) Let \mathcal{A} be an MV -algebra and let α be a cardinal.
- (i) \mathcal{A} is complete and α -atomic if and only if it is isomorphic to a direct product of complete α -atomic linearly ordered MV -algebras.
 - (ii) Let $\alpha > 2$. An MV -algebra is complete, α -atomic and linearly ordered if and only if it is of type R .
 - (iii) If \mathcal{A} is a complete α -atomic MV -algebra with $A \neq \{0\}$, then either $\alpha = 2$ or $\alpha = c$ (the cardinality of the continuum).
- (B) Let \mathcal{A} be a complete MV -algebra. Then \mathcal{A} is isomorphic to a direct product $\mathcal{A}_1 \times \mathcal{A}_2 \times \mathcal{A}_3$ such that
- (i) \mathcal{A}_1 is atomic;
 - (ii) \mathcal{A}_2 is c -atomic;
 - (iii) for each cardinal α , there are no α -atoms in \mathcal{A}_3 .

Let us remark that for each infinite cardinal α there exists a non-complete MV -algebra \mathcal{A} such that, whenever x is a nonzero element of A , then x is an α -atom of \mathcal{A} .

1. PRELIMINARIES AND AUXILIARY RESULTS

For the notion of the MV -algebra we introduce the following definition (cf. [5] and [6]):

- (**) An MV -algebra is a system $\mathcal{A} = (A; \oplus, *, \neg, 0, 1)$ (where $\oplus, *$ are binary operations, \neg is a unary operation and $0, 1$ are nullary operations) such that the following identities are satisfied:

- (m₁) $x \oplus (y \oplus z) = (x \oplus y) \oplus z$;
- (m₂) $x \oplus 0 = x$;
- (m₃) $x \oplus y = y \oplus x$;
- (m₄) $x \oplus 1 = 1$;
- (m₅) $\neg\neg x = x$;
- (m₆) $\neg 0 = 1$;
- (m₇) $x \oplus \neg x = 1$;
- (m₈) $\neg(\neg x \oplus y) \oplus y = \neg(x \oplus \neg y) \oplus x$;
- (m₉) $x * y = \neg(\neg x \oplus \neg y)$.

We recall the following results $(*_i)$ ($i = 1, 2, 3$) (for $(*_1)$ cf. [5]; for $(*_2)$ and $(*_3)$ cf. [7] 2.5 and 3.8; cf. also [6], 1.2, 1.3 and 1.4).

$(*_1)$ Let \mathcal{A} be an *MV*-algebra. For each $x, y \in A$ put $x \vee y = (x * \neg y) \oplus y$ and $x \wedge y = \neg(\neg x \vee \neg y)$. Then $\mathcal{L}(\mathcal{A}) = (A; \vee, \wedge)$ is a distributive lattice with the least element 0 and the greatest element 1.

$(*_2)$ Let G be an abelian lattice ordered group with a strong unit u . Let A be the interval $[0, u]$ of G . For each a and b in A we put

$$a \oplus b = (a + b) \wedge u, \quad \neg a = u - a, \quad 1 = u, \quad a * b = \neg(\neg a \oplus \neg b).$$

Then $\mathcal{A} = (A; \oplus, *, \neg, 0, 1)$ is an *MV*-algebra.

If G and \mathcal{A} are as in $(*_2)$, then we put $\mathcal{A} = \mathcal{A}_0(G, u)$.

$(*_3)$ Let \mathcal{A} be an *MV*-algebra. Then there exists an abelian lattice ordered group G with a strong unit u such that $\mathcal{A} = \mathcal{A}_0(G, u)$.

In what follows, \mathcal{A} and G are as in $(*_2)$ and $(*_3)$.

1.1. Lemma. *\mathcal{A} is complete if and only if G is complete.*

Proof. Let \mathcal{A} be complete. Hence the interval $[0, u]$ is complete. The fact that u is a strong unit of G implies that for proving the completeness of G it suffices to verify that for each positive integer n the lattice $[0, nu]$ is complete.

We proceed by induction on n . The case $n = 1$ is trivial. Suppose that $n > 1$ and that the interval $[0, (n - 1)u]$ is complete. Since $[(n - 1)u, nu]$ is isomorphic to $[0, u]$, we obtain that $[(n - 1)u, nu]$ is complete as well.

Let $X = \{x_i\}_{i \in I}$ be a nonempty subset of $[0, nu]$. For each $i \in I$ we put

$$x_i^1 = x_i \wedge (n - 1)u, \quad x_i^2 = x_i \vee (n - 1)u.$$

In view of the assumption the elements

$$x^1 = \bigvee_{i \in I} x_i^1, \quad x^2 = \bigvee_{i \in I} x_i^2$$

exist. For each $i \in I$ the relation

$$x_i = x_i^1 + (x_i^2 - a)$$

is valid, where $a = (n - 1)u$. Put

$$x = x^1 + (x^2 - a).$$

Then $x \geq x_i$ for each $i \in I$. Let $y \in [0, nu], y \geq x_i$ for each $i \in I$. Put $y^1 = y \wedge (n - 1)u, y^2 = y \vee (n - 1)u$. Then $y^1 \geq x_i^1$ and $y^2 \geq x_i^2$ for each $i \in I$. At the same time we have

$$y = y^1 + (y^2 - a).$$

Therefore $y \geq x$. Thus $x = \sup X$ is valid in $[0, nu]$. Similarly we can verify that $\inf X$ does exist in $[0, nu]$. Hence $[0, nu]$ is complete.

The converse implication is obvious. □

1.2. Lemma. *\mathcal{A} is linearly ordered if and only if G is linearly ordered.*

Proof. If G is linearly ordered, then clearly \mathcal{A} is linearly ordered as well. Suppose that G fails to be linearly ordered. Then there are $g_i \in G$ with $0 < g_i$ ($i = 1, 2$), $g_1 \wedge g_2 = 0$. Since u is a strong unit in G we infer that $u_i = g_i \wedge u > 0$ ($i = 1, 2$). We have $u_1 \wedge u_2 = 0$ and $u_1, u_2 \in A$. Hence \mathcal{A} is not linearly ordered. □

Let Z be the additive group of all integers with the usual linear order. It is well-known that if $H \neq \{0\}$ is a complete linearly ordered group, then H is isomorphic either to Z or to R ; hence if $0 < h \in H$, then the interval $[0, h]$ is atomic if and only if $[0, h]$ is finite. Hence $(*_1), (*_2), 1.1$ and 1.2 yield

1.3.1. Corollary. *Let \mathcal{A} be an MV-algebra, $A \neq \{0\}$. Suppose that \mathcal{A} is linearly ordered and complete. Then (i) \mathcal{A} is finite if and only if it is atomic, and (ii) \mathcal{A} is infinite if and only if it is c-atomic.*

1.3.2. Corollary. *Let \mathcal{A} be as in 1.3.1. Then (i) \mathcal{A} is atomic if and only if G is isomorphic to Z ; (ii) \mathcal{A} is c-atomic if and only if it is of type R .*

For each nonempty subset X of a lattice ordered group H we denote

$$X^\delta = \{y \in H : |y| \wedge |x| = 0 \text{ for each } x \in X\}.$$

X^δ will be said to be the polar in H generated by the set X . For a thorough theory of polars in lattice ordered groups cf. Šik [9]. Each polar is a convex ℓ -subgroup of H ; if $0 \in X$ and X is a linearly ordered convex subset of H , then $X^{\delta\delta}$ is linearly ordered as well.

1.4. Lemma. *Let \mathcal{A} be a complete MV-algebra. Let $0 < x \in G$ and suppose that the interval $[0, x]$ is linearly ordered. Then either $[0, x]$ is finite or $[0, x]$ has cardinality c .*

Proof. Put $X = [0, x]$. Then X is, at the same time, an interval of G . Thus $X^{\delta\delta}$ is linearly ordered. According to 1.1, G is complete. Hence by the Riesz Theorem (cf., e.g., Fuchs [4], Chap. V), $X^{\delta\delta}$ is a direct factor of G . Therefore in view of [6], 3.2, $X_1 = X^{\delta\delta} \cap [0, u]$ is a direct factor of \mathcal{A} . Moreover, $X^{\delta\delta}$ is linearly ordered and hence X_1 is linearly ordered as well. Each direct factor of a complete MV -algebra must be complete. Now it suffices to apply 1.3.2. \square

1.5. Corollary. *Let α be a cardinal and let \mathcal{A} be a complete MV -algebra, $A \neq \{0\}$. If \mathcal{A} is α -atomic, then either $\alpha = 2$ or $\alpha = c$.*

The notion of an α -atom of a lattice ordered group can be defined in the same way as in the case of MV -algebras. A lattice ordered group G is said to be α -atomic if for each $g \in G$ with $0 < g$ there exists an α -atom g_1 in G such that $g_1 \leq g$.

By a similar argument as above we obtain

1.5'. Lemma. *Let α be a cardinal and let G be a complete nonzero lattice ordered group. If G is α -atomic, then either $\alpha = 2$ or $\alpha = c$.*

1.6. Example. Let α be an infinite cardinal. Next, let I be a linearly ordered set which is isomorphic to the first ordinal having the power α . For each $i \in I$ let G_i be a linearly ordered group isomorphic to Z . Put $G' = \Gamma_{i \in I} G_i$, where Γ denotes the operation of lexicographic product (cf., e.g., [4]). For $g' \in G'$ and $i \in I$ let g'_i be the component of g' in G_i . Denote $I(g') = \{i \in I : g'_i \neq 0\}$. Let G'' be the subgroup of G consisting of all $g' \in G'$ for which the set $I(g')$ is finite; G'' is linearly ordered by the inherited order. G'' is a non-complete linearly ordered group such that whenever $x, y \in G$ and $x < y$, then the power of the interval $[x, y]$ in G is α . Choose $u \in G''$ with $0 < u$ and let G be the convex ℓ -subgroup of G'' generated by the element u . Hence u is a strong unit in G . Let \mathcal{A} be as in $(*_2)$. Thus each strictly positive element of A is an α -atom in \mathcal{A} .

2. PROOFS OF (A) AND (B)

The assertion (ii) and (iii) of (A) were already proved (cf. 1.3.2 and 1.5); the remaining part of (A) will be proved as follows. The case $A = \{0\}$ being trivial we can suppose that $A \neq \{0\}$.

a) Suppose that an MV -algebra \mathcal{A} is a direct product of MV -algebras \mathcal{A}_i ($i \in I$). Without loss of generality we can suppose that the direct decomposition under consideration is internal (in the sense of [5]). Assume that all \mathcal{A}_i are linearly ordered, complete and α -atomic. For each $x \in A$ and $i \in I$ we denote by x_i the component

of x in \mathcal{A}_i . Let $x > 0$. Then there exists $i \in I$ such that $x_i > 0$. There is an α -atom y^i of \mathcal{A}_i with $y^i \leq x_i$. The element y^i is, at the same time, an α -atom in \mathcal{A} and $y^i \leq x$. Thus \mathcal{A} is α -atomic.

b) Suppose that \mathcal{A} is a complete and α -atomic MV-algebra. Since $A \neq \{0\}$, the set of all α -atoms of \mathcal{A} is nonempty. For each α -atom x of \mathcal{A} let X be as in the proof of 1.4. Hence $X^{\delta\delta}$ is a direct factor of G ; let $\{G_i\}_{i \in I}$ be the set of all $X^{\delta\delta}$ which can be constructed in this way. Each G_i is linearly ordered, complete and α -atomic.

For each $y \in G$ and $i \in I$ let y_i be the component of y in G_i . It is well-known that if $y \geq 0$, then y_i is the greatest element of the set $G_i \cap [0, y]$.

Let $y \in A$. We have $y_i \leq y$ for each $i \in I$; since \mathcal{A} is complete, there exists $y' = \bigvee_{i \in I} y_i$ in A . We shall verify that $y' = y$. By way of contradiction, suppose that $y'' = y - y' > 0$. Then $y'' \in A$ and hence there exists an α -atom x in A such that $x \leq y''$. Thus there is $i(1) \in I$ such that $G_{i(1)} = [0, x]^{\delta\delta}$. Clearly $y''_{i(1)} \geq x_{i(1)} = x > 0$. Since $0 \leq y_{i(1)} \in G_{i(1)} \cap [0, y']$ we obtain $y_{i(1)} \leq y'_{i(1)}$. On the other hand, the relation $y' < y$ gives $y'_{i(1)} \leq y_{i(1)}$. Thus $y_{i(1)} = y'_{i(1)}$. Therefore $y_{i(1)} = y'_{i(1)} + y''_{i(1)} = y_{i(1)} + x > y_{i(1)}$, which is a contradiction. Thus

$$(1) \quad y = \bigvee_{i \in I} y_i.$$

If $i(1)$ and $i(2)$ are distinct elements of I , then $G_{i(1)} \cap G_{i(2)} = \{0\}$. This implies that $y_{i(1)} \wedge y_{i(2)} = 0$.

Let φ be a mapping of A into the direct product $\prod_{i \in I} A_i$ (where $A_i = [0, u_i]$ for each $i \in I$) defined by

$$\varphi(y) = (y_i)_{i \in I}.$$

We consider A_i to be partially ordered by the inherited partial order. Let y and z be elements of A . If $y \leq z$, then clearly $y_i \leq z_i$ for each $i \in I$. Conversely, assume that $y_i \leq z_i$ for each $i \in I$; then we infer from (1) that $y \leq z$. Thus if y and z are distinct, then $\varphi(y)$ and $\varphi(z)$ are distinct as well. Further, let $(t^i) \in \prod_{i \in I} A_i$. There exists $t \in A$ with $t = \bigvee_{i \in I} t^i$. For each $i(1) \in I$ we have

$$t_{i(1)} = t_{i(1)} \wedge t = t_{i(1)} \wedge \left(\bigvee_{i \in I} t^i \right) = t_{i(1)} \wedge t^{i(1)}$$

(since $t_{i(1)} \wedge t^i = 0$ whenever $i \neq i(1)$). Thus $t_{i(1)} \leq t^{i(1)}$. On the other hand, $t^{i(1)} \in G_{i(1)} \cap [0, t]$ and hence $t^{i(1)} \leq t_{i(1)}$. We obtain $t^{i(1)} = t_{i(1)}$ and therefore $\varphi(t) = (t^i)_{i \in I}$. We have verified that φ is an isomorphism of the lattice A onto $\prod_{i \in I} A_i$.

For each $i \in I$ the element u_i is a strong unit in G_i , hence the MV -algebra $\mathcal{A}_i = (A_i; \oplus, *, \neg, 0, u_i)$ exists. From the construction of the isomorphism φ and from [6], 3.5 we infer that φ is, at the same time, an isomorphism of \mathcal{A} onto $\prod_{i \in I} \mathcal{A}_i$. Each \mathcal{A}_i is complete, linearly ordered and α -atomic. This completes the proof of (A).

PROOF OF (B). Let \mathcal{A} be a complete MV -algebra and let G be as above. We denote by X_1 and X_2 the system of all atoms of \mathcal{A} or the system of all c -atoms of \mathcal{A} , respectively. Put $G_i = X_i^{\delta}$ ($i = 1, 2$). By the Riesz Theorem, G_1 and G_2 are direct factors of G . For each $x_1 \in X_1$ and $x_2 \in X_2$ we have $x_1 \wedge x_2 = 0$. This yields that $G_1 \cap G_2 = \{0\}$. Therefore

$$(2) \quad G = G_1 \times G_2 \times G_3,$$

where

$$(3) \quad G_3 = (G_1 \cup G_2)^{\delta}.$$

All G_i ($i = 1, 2, 3$) are complete. It follows from the definition of G_1 that it is atomic; analogously, G_2 is c -atomic. The relation (3) yields that for each cardinal α no α -atom exists in G_3 .

For $i \in \{1, 2, 3\}$ let u_i be the component of u in G_i . We can construct the MV -algebras $\mathcal{A}_i = (A_i; \oplus, *, \neg, 0, u_i)$ for $i = 1, 2, 3$, where $A_i = [0, u_i]$. Then all \mathcal{A}_i are complete, \mathcal{A}_1 is atomic, \mathcal{A}_2 is c -atomic, and for each cardinal α , \mathcal{A}_3 has no α -atoms. Now we can apply [6], Lemma 3.2 (this lemma deals with direct decompositions having two factors, but by an obvious induction we can extend the validity of the lemma to direct decompositions having a finite number of direct factors); from (2) we infer that \mathcal{A} is a direct product of \mathcal{A}_1 , \mathcal{A}_2 and \mathcal{A}_3 . \square

References

- [1] C. C. Chang: Algebraic analysis of many-valued logics. Trans. Amer. Math. Soc. 88 (1958), 467–490.
- [2] C. C. Chang: A new proof of the completeness of the Lukasiewicz axioms. Trans. Amer. Math. Soc. 93 (1959), 74–80.
- [3] R. Cignoli: Complete and atomic algebras of the infinite valued Lukasiewicz logic. Studia Logica 50 (1991), 375–384.
- [4] L. Fuchs: Partially ordered algebraic systems. Pergamon Press, Oxford, 1963.
- [5] D. Gluschkof: Cyclic ordered groups and MV -algebras. Czechoslov. Math. J. 43 (1993), 249–263.
- [6] J. Jakubík: Direct product decompositions of MV -algebras. Czechoslov. Math. J. To appear.
- [7] D. Mundici: Interpretation of AFC^* -algebras in Lukasiewicz sentential calculus. Jour. Functional. Anal. 65 (1986), 15–63.

- [8] *D. Mundici*: *MV*-algebras are categorically equivalent to bounded commutative *BCK*-algebras. *Math. Japonica* 31 (1986), 889–894.
- [9] *F. Šik*: To the theory of lattice ordered groups. *Czechoslov. Math. J.* 6 (1956), 1–25. (In Russian.)
- [10] *T. Traczyk*: On the variety of bounded commutative *BCK*-algebras. *Math. Japonica* 24 (1979), 238–282.

Author's address: Matematický ústav SAV, dislokované pracovisko, Grešákova 6, 040 01 Košice, Slovakia.