

# ON COMPLETELY MONOTONE FUNCTIONS ON $C_+(X)$

J. HOFFMANN-JØRGENSEN and P. RESSEL

## 1. Introduction.

Let  $X$  be a completely regular Hausdorff space, let  $C(X)$  denote the vector space of all bounded realvalued continuous functions on  $X$  and  $M(X)$  the vector space of all real Radon measures on  $X$ . The positive cones in  $C(X)$  and  $M(X)$  are denoted by  $C_+(X)$  and  $M_+(X)$ .

Under pointwise addition the cone  $C_+(X)$  becomes a 2-divisible abelian semigroup in the sense of [1]. As in [1] we define the character semigroup  $\hat{S}$  of  $S := C_+(X)$  by  $\varrho \in \hat{S}$  if and only if  $\varrho: S \rightarrow [0, 1]$  and

$$(1.1) \quad \varrho(0) = 1$$

$$(1.2) \quad \varrho(f+g) = \varrho(f)\varrho(g) \quad \forall f, g \in S.$$

In the topology of pointwise convergence and with pointwise multiplication  $\hat{S}$  becomes a compact topological abelian semigroup.

Let  $L$  denote all functionals  $\lambda: C_+(X) \rightarrow [0, \infty]$  satisfying

$$(1.3) \quad \lambda(0) = 0$$

and

$$(1.4) \quad \lambda(f+g) = \lambda(f) + \lambda(g) \quad \forall f, g \in C_+(X).$$

Each  $\lambda$  satisfying (1.3) and (1.4) is increasing and hence positive homogeneous, i.e.

$$(1.5) \quad \lambda(af) = a\lambda(f) \quad \forall a \geq 0, \forall f \in C_+(X),$$

with the usual conventions  $0 \cdot \infty = 0$  and  $a \cdot \infty = \infty, \forall a > 0$ . Equipped with the topology of pointwise convergence  $L$  becomes compact and

$$\lambda(\cdot) \mapsto e^{-\lambda(\cdot)}$$

is a homeomorphism of  $L$  onto  $\hat{S}$ .

We shall consider the following subsets of  $L$ :

$$L_0 := \{ \lambda \in L \mid \lambda(f) < \infty, \forall f \in C_+(X) \}$$

and, if  $Y \subseteq X$

$$L_Y := \left\{ \lambda \in L_0 \mid \exists \mu \in M_+(Y) \text{ such that } \lambda(f) = \int_Y f d\mu \forall f \in C_+(X) \right\}.$$

Let  $w^*$  be the *weak topology* on  $M(X)$ , that is,  $w^* = \sigma(M(X), C(X))$ , then the map

$$(1.6) \quad \mu \mapsto \int_X \cdot d\mu$$

is a homeomorphism of  $M_+(X)$  onto  $L_X$ .

Let  $X$  denote the Stone-Čech compactification of  $X$  and let  $\tilde{f}$  denote the unique continuous extension of  $f$  to  $\beta X$ , for all  $f \in C(X)$ . Then the map

$$(1.7) \quad \tilde{\mu} \mapsto \int_{\beta X} \tilde{\cdot} d\tilde{\mu}$$

is a homeomorphism of  $(M_+(\beta X), w^*)$  onto  $L_0$ .

A function  $\varphi: C_+(X) \rightarrow \mathbb{R}$  is *completely monotone* if and only if  $\varphi$  is bounded and

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j \varphi(f_i + f_j) \geq 0$$

for all  $n \in \mathbb{N}$ ,  $c_1, \dots, c_n \in \mathbb{R}$  and  $f_1, \dots, f_n \in C_+(X)$ , (cf. [2] and Theorem 4.2 in [1]). From [2] we know that every completely monotone function  $\varphi: C_+(X) \rightarrow \mathbb{R}$  has a unique *representing measure*  $\xi \in M_+(L)$  in the sense that

$$(1.8) \quad \varphi(f) = \int_L e^{-\lambda(f)} d\xi(\lambda), \quad \forall f \in C_+(X).$$

Our aim in the following will be to establish a connection between continuity properties of  $\varphi$  and the concentration of the measure  $\xi$  to some “nice” subsets of  $L$ . A very special result of this type has already been proved in Theorem 6.1 of [1]. There  $X$  is the finite set  $\{1, \dots, p\}$  with discrete topology,  $C_+(X)$  can be identified with  $\mathbb{R}_+^p$  and  $L$  with  $[0, \infty]^p$  and it is shown that a completely monotone functions on  $\mathbb{R}_+^p$  is continuous if and only if the representing measure is concentrated on  $\mathbb{R}_+^p$ .

If we consider the dual pair  $(C(X), M(X))$ , two natural topologies on  $C(X)$  arise, the *weak topology*, denoted by  $w$ , and the *Mackey topology*, which we

shall denote by  $m$ . We shall need two further topologies. First we define the  $L_1$ -topology  $\tau$  on  $C(X)$  by the family of seminorms

$$(1.9) \quad r_{K, \sigma}(f) := \int_K \left| \int_X f d\mu \right| d\sigma(\mu)$$

where  $K$  runs through all  $w^*$ -compact, uniformly tight subsets of  $M(X)$  and  $\sigma$  runs through  $M_+(M(X), w^*)$ . There is a simpler description of this topology, but first we need a lemma:

LEMMA 1.1. *Let  $\sigma \in M_+(M(X), w^*)$  and suppose that the function  $\mu \mapsto |\mu|(X)$  is  $\sigma$ -integrable, then*

$$\lambda(A) := \int_{M(X)} |\mu|(A) d\sigma(\mu), \quad A \in \mathcal{B}(X)$$

is a positive finite  $\tau$ -smooth measure on  $(X, \mathcal{B}(X))$ , and for every bounded Borel functions  $g$  on  $X$  we have

$$\int_X g d\lambda = \int_{M(X)} \left( \int_X g d|\mu| \right) d\sigma(\mu).$$

If  $\sigma$  satisfies

$$\sigma(M(X)) = \sup \{ \sigma(K) \mid K \text{ uniformly tight and closed} \}$$

then  $\lambda$  is a Radon measure on  $X$ .

REMARK.  $\mathcal{B}(X)$  of course denotes the Borel  $\sigma$ -algebra of  $X$ . The notion of  $\tau$ -smoothness may be found in [7, p. XII], and from P 16 p. XIII in [7] it follows that if  $X$  can be homeomorphically embedded as a universally measurable subset of a compact space  $Y$ , then every  $\tau$ -smooth finite measure on  $X$  is a Radon measure (e.g. if  $X$  is analytic, or if  $X$  is  $\sigma$ -compact, or if  $X$  is locally compact or if  $X$  is complete in the sense of Čech). From Proposition 1 in [5] we know that the function  $\mu \mapsto |\mu|(A)$  is Borel on  $M(X)$  for every Borel set  $A \subseteq X$ , it is lower semicontinuous if  $A$  is open.

PROOF OF LEMMA 1.1. In the first part we only need to show  $\tau$ -smoothness of  $\lambda$ . Let a collection of open sets  $G_\alpha \subseteq X$  filter up to  $G$ . Then the lower semicontinuous functions  $\mu \mapsto |\mu|(G_\alpha)$  filter up to  $\mu \mapsto |\mu|(G)$  and applying P 15 of [7] we get  $\lambda(G) = \sup \lambda(G_\alpha)$ .

The second part is proved in a straightforward manner, taking into account that  $\lambda$  is inner regular w.r.t. the closed subsets of  $X$ , cf. P 16 in [7].

COROLLARY 1.2. *The  $L_1$ -topology  $\tau$  on  $C(X)$  is generated by the seminorms*

$$q_\mu(f) := \int_X |f| d\mu$$

where  $\mu$  runs through  $M_+(X)$ .

PROOF. Let  $K$  be a  $w^*$ -compact and uniformly tight subset of  $M(X)$  and  $\sigma$  a positive finite Radon measure on  $M(X)$ , then

$$\mu(A) := \int_K |\nu|(A) d\sigma(\nu) \quad A \in \mathcal{B}(X)$$

is a finite positive Radon measure on  $X$  by Lemma 1.1, and

$$r_{K,\sigma}(f) \leq \int_K \left( \int_X |f| d\nu \right) d\sigma(\nu) = \int_X |f| d\mu.$$

If  $\mu \in M_+(X)$ , then there exists a measurable function  $\alpha: X \rightarrow [0, 1]$  such that  $\{\alpha \geq \varepsilon\}$  is compact for all  $\varepsilon > 0$ , and

$$\int_X \frac{1}{\alpha} d\mu < \infty.$$

Let  $\psi(x) := \alpha(x)\delta_x$ , where  $\delta_x$  is the one point measure in  $x$ , then  $\psi$  is a Borel map from  $X$  into  $M_+(X)$ ,  $\psi(X)$  is uniformly tight and  $K := \overline{\psi(X)}$  is therefore  $w^*$ -compact and uniformly tight. Let  $d\lambda := (1/\alpha)d\mu$  and  $\sigma := \lambda \circ \psi^{-1}$ , then  $\sigma$  is a finite positive Radon measure on  $M(X)$ , and

$$\begin{aligned} q_\mu(f) &= \int_X |f| d\mu = \int_X \alpha(x)|f(x)| d\lambda(x) \\ &= \int_X \left| \int_X f(y) d(\psi(x))(y) \right| d\lambda(x) = \int_K \left| \int_X f d\nu \right| d(\lambda \circ \psi^{-1})(\nu) \\ &= \int_K \left| \int f d\nu \right| d\sigma(\nu). \end{aligned}$$

This shows that  $\{q_\mu\}$  and  $\{r_{K,\sigma}\}$  generate the same topology.

We shall need a fourth topology on  $C(X)$ . This is the so-called *strict topology* on  $C(X)$ , which we denote by  $\beta$ . The strict topology is generated by the seminorms

$$p_\alpha(f) := \|\alpha f\|_X = \sup_{x \in X} |\alpha(x)f(x)|$$

where  $\alpha$  runs through all bounded measurable functions on  $X$  which vanish at infinity, i.e.  $\{|\alpha| \geq \varepsilon\}$  is relatively compact for all  $\varepsilon > 0$ . This topology was first introduced by C. R. Buck for locally compact spaces and later generalized by many authors to general completely regular Hausdorff spaces (see e.g. [4]).

From Theorem 2 in [4] we know that  $w \subseteq \beta \subseteq m$ , and from Corollary 1.2 it follows easily that  $w \subseteq \tau \subseteq \beta$ , therefore we have

$$(1.10) \quad w \subseteq \tau \subseteq \beta \subseteq m$$

and we shall leave to the reader to prove that  $\tau \neq w$  and  $\tau \neq \beta$  if  $X$  is infinite.

From Theorem 3 in [6] one easily deduces the following form of Riesz' representation theorem:

**THEOREM 1.3.** (D. Pollard and F. Topsøe [6]). *Let  $\lambda: C_+(X) \rightarrow [0, \infty[$  be additive and suppose that  $\lambda$  satisfies*

$$(1.3.1) \quad \forall \varepsilon > 0 \exists \delta > 0 \exists C \text{ compact } \subseteq X \text{ such that } \lambda(f) \leq \varepsilon \text{ whenever } 0 \leq f \leq 1 \text{ and } f \leq \delta \text{ on } C.$$

*Then there exists a unique measure  $\mu \in M_+(X)$  representing  $\lambda$ , that is,*

$$\lambda(f) = \int_X f d\mu, \quad \forall f \in C_+(X).$$

**REMARK.** Note that (1.3.1) holds if there exists  $\{f_n\} \subseteq C_+(X)$  with the following two properties:

$$(1.3.2) \quad \{f_n \leq 1\} \text{ is compact for all } n \geq 1,$$

$$(1.3.3) \quad \lim_{n \rightarrow \infty} \lambda(f_n) = 0.$$

## 2. Concentration of $L_0$ or $L_X$ .

Let  $X$  be a completely regular Hausdorff space and  $\varphi$  a completely monotone function on  $C_+(X)$ , with representing measure  $\xi$ . We shall give necessary and sufficient conditions for  $\xi$  to be concentrated on  $L_0$  or  $L_X$ . First we need a measurability lemma:

**LEMMA 2.1.** *If  $Y$  is a Borel subset of the Stone-Čech compactification  $\beta X$ , then  $L_Y$  is a Borel subset of  $L$ . The subset  $L_0$  is open in  $L$ .*

**PROOF.** From  $L_0 = \{\lambda \in L \mid \lambda(1) < \infty\}$  follows that  $L_0$  is open. Identifying  $L_0$  with  $M_+(\beta X)$ , see (1.7), we get

$$L_Y = \{\tilde{\mu} \in M_+(\beta X) \mid \tilde{\mu}(\beta X \setminus Y) = 0\}$$

and from the fact that  $\tilde{\mu} \mapsto \tilde{\mu}(\beta X \setminus Y)$  is Borel on  $M_+(\beta X)$  by Proposition 1 in [5] we deduce that  $L_Y$  is Borel in  $L_0$ , hence in  $L$ .

**THEOREM 2.2.** *Let  $\varphi: C_+(X) \rightarrow \mathbb{R}$  be completely monotone with representing measure  $\xi$  (see (1.8.)). Then the following 3 statements are equivalent:*

$$(2.2.1) \quad \xi(L \setminus L_0) = 0,$$

$$(2.2.2) \quad \lim_{t \rightarrow 0} \varphi(tf) = \varphi(0), \quad \forall f \in C_+(X),$$

$$(2.2.3) \quad \lim_{t \rightarrow 0} \varphi(t) = \varphi(0),$$

where  $t$  also denotes the constant function equal to  $t$ .

**PROOF.** (2.2.1)  $\Rightarrow$  (2.2.2): Let  $f \in C_+(X)$  and define  $F_t(\lambda) := e^{-t\lambda f}$ . Then

$$0 \leq F_t(\lambda) \leq 1 \quad \text{and} \quad \lim_{t \rightarrow 0} F_t(\lambda) = 1$$

for all  $\lambda \in L_0$ . Hence the assumption implies that

$$\varphi(tf) = \int_{L_0} F_t(\lambda) d\xi(\lambda) \rightarrow \xi(L_0) = \varphi(0),$$

as  $t$  tends to zero.

(2.2.2)  $\Rightarrow$  (2.2.3): Obvious.

(2.2.3)  $\Rightarrow$  (2.2.1): Let  $F_t(\lambda)$  be defined as above but with  $f$  replaced by the constant 1. If  $\lambda \in L \setminus L_0$  then  $\lambda(1) = \infty$ , therefore we get

$$\lim_{t \rightarrow 0} F_t(\lambda) = 1_{L_0}(\lambda) \quad \text{for all } \lambda \in L.$$

Hence by assumption

$$\xi(L) = \varphi(0) = \lim_{t \rightarrow 0} \int_L F_t(\lambda) d\xi(\lambda) = \xi(L_0)$$

and so  $\xi(L \setminus L_0) = 0$ .

**THEOREM 2.3.** *Let  $\varphi: C_+(X) \rightarrow \mathbb{R}$  be a completely monotone function with representing measure  $\xi$ , and let  $\varrho$  be a topology on  $C(X)$  satisfying  $\tau \subseteq \varrho \subseteq \beta$ . Then the following 6 statements are equivalent:*

$$(2.3.1) \quad \exists Y \text{ } \sigma\text{-compact } \subseteq X \text{ such that } \xi(L \setminus L_Y) = 0,$$

$$(2.3.2) \quad \xi(L) = \sup \{ \xi(K) \mid K \subseteq M_+(X) \text{ compact and uniformly tight} \},$$

$$(2.3.3) \quad \varphi \text{ is uniformly } \varrho\text{-continuous,}$$

$$(2.3.4) \quad \varphi \text{ is } \varrho\text{-continuous at } 0,$$

(2.3.5)  $\varphi|B$  is  $\varrho$ -continuous at 0, where  $B = \{f \in C(X) \mid 0 \leq f \leq 1\}$ ,

(2.3.6)  $\forall \varepsilon > 0 \exists \delta > 0 \exists C$  compact  $\subseteq X$  such that  $\varphi(0) - \varphi(f) \leq \varepsilon$  whenever  $f \in B$  and  $f \leq \delta$  on  $C$ .

Note that we identify  $M_+(X)$  with  $L_X$  in (2.3.2) (see (1.6)).

PROOF. (2.3.1)  $\Rightarrow$  (2.3.2): First we note that  $L_Y \subseteq L_X \subseteq L_0$  and therefore  $\xi(L \setminus L_0) = 0$ . Let  $\{C_n\}$  be compact sets in  $X$  with  $C_1 = \emptyset$  and  $C_n \uparrow Y$ ; then we may define

$$F_n(v) := v(\beta X \setminus C_n) \quad \text{for } v \in L_0 = M_+(\beta X).$$

Then  $F_n: L_0 \rightarrow [0, \infty[$  is Borel and  $\lim_{n \rightarrow \infty} F_n(v) = 0$  for all  $v \in L_Y$ . From Lemma 2.1 we know that  $L_Y \in \mathcal{B}(L)$ , and since  $\xi(L \setminus L_Y) = 0$  by assumption, we have that  $F_n \rightarrow 0$  a.e.  $[\xi]$ . Hence by Egoroff's theorem we can find for any given  $\varepsilon > 0$  a sequence  $a_1 \geq a_2 \geq \dots \geq 0$  of positive numbers such that

$$\lim_{n \rightarrow \infty} a_n = 0,$$

$$\xi(\{v \mid F_n(v) \leq a_n, \forall n \geq 1\}) \geq \xi(L) - \varepsilon.$$

Now since  $F_n$  is lower semi-continuous on  $L_0$  (see Proposition 1 in [5]) and  $F_1(v) = v(\beta X)$ , we have that

$$K := \{v \in L_0 \mid F_n(v) \leq a_n, \forall n \geq 1\}$$

is a compact uniformly tight subset of  $M_+(X)$  with  $\xi(K) \geq \xi(L) - \varepsilon$ . Hence (2.3.2) holds.

(2.3.2)  $\Rightarrow$  (2.3.3): Given  $\varepsilon > 0$  choose  $K \subseteq M_+(X)$  compact and uniformly tight so that  $\xi(K) \geq \xi(L) - \varepsilon/4$ . We claim that  $|\varphi(f) - \varphi(g)| < \varepsilon$  whenever  $r_{K, \xi}(|f - g|) < \varepsilon/2$  (see (1.9) for the definition of  $r_{K, \xi}$ ). In fact, if  $r_{K, \xi}(|f - g|) < \varepsilon/2$  for two functions  $f, g \in C_+(X)$ , then

$$\begin{aligned} |\varphi(f) - \varphi(g)| &\leq \varphi(f) - \varphi(f \vee g) + \varphi(g) - \varphi(f \vee g) \\ &= \int_L (e^{-\lambda f} - e^{-\lambda(f \vee g)}) d\xi(\lambda) + \int_L (e^{-\lambda g} - e^{-\lambda(f \vee g)}) d\xi(\lambda) \\ &\leq \int_L [1 - e^{-\lambda(f \vee g - f)}] d\xi(\lambda) + \int_L [1 - e^{-\lambda(f \vee g - g)}] d\xi(\lambda) \\ &\leq \frac{\varepsilon}{2} + \int_K \lambda(f \vee g - f) d\xi(\lambda) + \int_K \lambda(f \vee g - g) d\xi(\lambda) \\ &= \frac{\varepsilon}{2} + \int_K \lambda(2(f \vee g) - f - g) d\xi(\lambda) \end{aligned}$$

$$\begin{aligned}
&= \frac{\varepsilon}{2} + \int_K \lambda(|f-g|) d\xi(\lambda) \\
&= \frac{\varepsilon}{2} + r_{K,\xi}(|f-g|) < \varepsilon.
\end{aligned}$$

This shows that  $\varphi$  is uniformly  $\tau$ -continuous. (2.3.3) follows because  $\tau$  is weaker than  $\varrho$  by assumption.

(2.3.3)  $\Rightarrow$  (2.3.4)  $\Rightarrow$  (2.3.5): Obvious.

(2.3.5)  $\Rightarrow$  (2.3.6): Since  $\varrho$  is weaker than  $\beta$ , we have that  $\varphi|B$  is  $\beta$ -continuous at 0. Let  $\varkappa$  be the topology on  $C(X)$  of uniform convergence on compact subsets of  $X$ , then by Proposition 1 in [4],  $\varkappa$  coincides with  $\beta$  on  $B$ . Hence  $\varphi|B$  is  $\varkappa$ -continuous at 0 and this evidently implies (2.3.6).

(2.3.6)  $\Rightarrow$  (2.3.1): First we note that (2.3.6) implies that  $\lim_{t \rightarrow 0} \varphi(t) = \varphi(0)$ , therefore  $\xi(L \setminus L_0) = 0$  by Theorem 2.1. Now let

$$M_n := \{v \in L_0 \mid v(1) \leq n\}.$$

Let  $f \in C_+(X)$  and define

$$F_t(\lambda) := \frac{1}{t}(1 - e^{-t\lambda(f)}) \quad \text{for } t > 0, \lambda \in L_0.$$

Then we have

$$\begin{aligned}
\lim_{t \rightarrow 0} F_t(\lambda) &= \lambda(f) \quad \forall \lambda \in L_0 \\
0 &\leq F_t(\lambda) \leq \lambda(f) \leq \|f\|_X \lambda(1)
\end{aligned}$$

and this implies that

$$\mu_n(f) := \int_{M_n} \lambda(f) d\xi(\lambda) = \lim_{t \rightarrow 0} \int_{M_n} F_t(\lambda) d\xi(\lambda)$$

for all  $f \in C_+(X)$ . Let for  $A \in \mathcal{B}(\beta X)$

$$\tilde{\mu}_n(A) := \int_{M_n} \lambda(A) d\xi(\lambda)$$

then by Lemma 1.1  $\tilde{\mu}_n$  is a positive Radon measure on  $\beta X$  with

$$\mu_n(f) = \int_{\beta X} \tilde{f} d\tilde{\mu}_n \quad \forall f \in C(X).$$

Now we use the elementary inequality

$$x \leq (1+a)(1 - e^{-x}) \quad \text{for } 0 \leq x \leq a$$



to conclude that

$$F_t(\lambda) \leq \lambda(f) \leq (n+1)(1 - e^{-\lambda(f)})$$

for  $f \in B$  and  $\lambda \in M_n$ . Hence we get

$$\mu_n(f) \leq (n+1) \int_{M_n} (1 - e^{-\lambda(f)}) d\xi(\lambda) \leq (n+1)(\varphi(0) - \varphi(f))$$

for all  $f \in B$ . The assumption (2.3.6) now implies that  $\mu_n$  satisfies (1.3.1) and by Theorem 1.3 we have that  $X$  is  $\tilde{\mu}_n$ -measurable and  $\tilde{\mu}_n(\beta X \setminus X) = 0$ .

Hence we can find a  $\sigma$ -compact subset  $Y \subseteq X$  such that  $\tilde{\mu}_n(\beta X \setminus Y) = 0$  for all  $n \geq 1$ . But then we have

$$\xi(\{\lambda \in M_n \mid \lambda(\beta X \setminus Y) > 0\}) = 0 \quad \forall n \geq 1$$

and since  $M_n \uparrow L_0$  and  $\xi(L \setminus L_0) = 0$ , we finally get

$$\xi(L \setminus L_Y) = \xi(L_0 \setminus L_Y) = \xi(\{\lambda \in L_0 \mid \lambda(\beta X \setminus Y) > 0\}) = 0$$

which proves our theorem.

### 3. The Lévy continuity theorem on $M_+(X)$ .

Let  $X$  be a completely regular Hausdorff space. Then  $M_+(M_+(X))$  denotes the set of positive finite Radon measures on  $(M_+(X), w^*)$ , and  $M_t(M_+(X))$  denotes the set of all  $\sigma \in M_+(M_+(X))$  satisfying

$$(3.1) \quad \sigma(M_+(X)) = \sup \{ \sigma(K) \mid K \text{ compact and uniformly tight} \}.$$

Note that  $M_t(M_+(X)) = M_+(M_+(X))$  if  $X$  is semi-Radonian (see Theorem 10 in [5]).

If  $\sigma \in M_+(M_+(X))$ , then we define its *Laplace transform*  $\hat{\sigma}$  by

$$\hat{\sigma}(f) := \int_{M_+(X)} \exp\left(-\int_X f \, dv\right) d\sigma(v)$$

for  $f \in C_+(X)$ . Note that  $\hat{\sigma}$  is completely monotone on  $C_+(X)$ . If  $X$  is  $\sigma$ -compact, then the set of all Laplace transforms of measures on  $M_+(X)$  is characterised by those completely monotone functions  $\varphi$  on  $C_+(X)$  which satisfy one of the continuity properties (2.3.3)–(2.3.6) stated in Theorem 2.3.

We shall consider  $M_+(M_+(X))$  and  $M_t(M_+(X))$  equipped with their weak topologies, coming from the space  $C(M_+(X), w^*)$ . Let  $\psi$  denote the map  $M_+(X) \rightarrow L$  given by (1.6), and let

$$\Psi(\sigma) := \sigma \circ \psi^{-1}$$

be the corresponding map from  $M_+(M_+(X))$  to  $M_+(L)$ . It is easily checked that

(3.2)  $\Psi$  is a homeomorphism of  $M_+(M_+(X))$  onto

$$M_X(L) := \{ \xi \in M_+(L) \mid \xi^*(L \setminus L_X) = 0 \}$$

(see e.g. Corollary 9 in [3, p. 244]).

**THEOREM 3.1.** *Let  $\{\sigma_\alpha\}$  be a net in  $M_+(M_+(X))$  satisfying*

(3.1.1)  $\sup_\alpha \sigma_\alpha(M_+(X)) < \infty,$

(3.1.2)  $\hat{\sigma}_\alpha(f) \rightarrow \varphi(f)$  for all  $f \in C_+(X)$ , where  $\varphi|B$  is  $\beta$ -continuous at 0,  
 $B := \{f \in C(X) \mid 0 \leq f \leq 1\}.$

Then there exists a measure  $\sigma \in M_t(M_+(X))$  whose Laplace transform is  $\varphi$  and  $\sigma_\alpha \rightarrow \sigma$  weakly.

**PROOF.** Let  $A := \sup_\alpha \sigma_\alpha(M_+(X))$  and let

$$M_A := \{ \xi \in M_+(L) \mid \xi(L) \leq A \}.$$

Then  $\xi_\alpha := \Psi(\sigma_\alpha) \in M_A$  for all  $\alpha$ , and  $M_A$  is a compact subset of  $M_+(L)$ . If  $\xi$  is a limit point of  $\{\xi_\alpha\}$ , then

$$\begin{aligned} \xi(f) &= \int_L e^{-\lambda(f)} d\xi(\lambda) = \lim_\alpha \int_L e^{-\lambda(f)} d\xi_\alpha(\lambda) \\ &= \lim_\alpha \hat{\sigma}_\alpha(f) = \varphi(f). \end{aligned}$$

Hence  $\xi$  is a completely monotone function on  $C_+(X)$ , with representing measure  $\xi$ . Since a measure on  $L$  is uniquely determined by its Laplace transform (see Corollary 2.5 of [1]), we find that  $\{\xi_\alpha\}$  admits at most one limit point in  $M_+(L)$ . Hence  $\xi = \lim_\alpha \xi_\alpha$  exists and  $\xi = \varphi$ .

Then by (3.1.2) and Theorem 2.3 we conclude that  $\xi = \Psi(\sigma)$  for some  $\sigma \in M_t(M_+(X))$ , and  $\hat{\sigma} = \varphi$ . Therefore by (3.2) we find that  $\sigma_\alpha \rightarrow \sigma$  weakly.

**THEOREM 3.2.** *Let  $\mathcal{X}$  be a subset of  $M_+^1(M_+(X))$ , the probability Radon measures on  $M_+(X)$ . Let again  $B := \{f \in C_+(X) \mid 0 \leq f \leq 1\}$ . Then we have*

- (i) *If  $\{\hat{\sigma}|B \mid \sigma \in \mathcal{X}\}$  is  $\beta$ -equicontinuous at 0, then  $\mathcal{X}$  is a relatively compact subset of  $M_t(M_+(X))$ .*
- (ii) *If  $\mathcal{X}$  is uniformly tight and  $X$  is a Prohorov space (see e.g. [5] for this notion), then  $\{\hat{\sigma}|B \mid \sigma \in \mathcal{X}\}$  is  $\beta$ -equicontinuous at 0.*

**PROOF.** (i). Follows immediately from Theorem 3.1.

(ii). Let  $\varepsilon > 0$  be given. There exists by assumption a compact set  $K \subseteq M_+(X)$  such that

$$\sup \{ \sigma(M_+(X) \setminus K) \mid \sigma \in \mathcal{X} \} < \frac{\varepsilon}{3}.$$

$X$  being a Prohorov space we can find a compact subset  $C$  of  $X$  such that

$$\sup \{ \nu(X \setminus C) \mid \nu \in K \} < \frac{\varepsilon}{3}.$$

From the compactness of  $K$  we deduce that  $A := \sup \{ \nu(X) \mid \nu \in K \}$  is finite. Now suppose that  $f \in B$ ,  $f|_C < \varepsilon/3A$  and  $\sigma \in \mathcal{X}$ . Then we get

$$1 - \hat{\sigma}(f) = \int_{M_+(X)} (1 - e^{-\lambda(f)}) d\sigma(\lambda) \leq \frac{\varepsilon}{3} + \int_K (1 - e^{-\lambda(f)}) d\sigma(\lambda)$$

and for  $\lambda \in K$

$$1 - e^{-\lambda(f)} \leq \lambda(f) \leq \frac{\varepsilon}{3} + \int_C f d\lambda \leq \frac{2\varepsilon}{3}$$

hence  $1 - \hat{\sigma}(f) \leq \varepsilon$ , showing the required  $\beta$ -equicontinuity at 0.

The next theorem might be more useful for applications. Note that if  $\sigma \in M_+(M_+(X))$ , then its Laplace transform is defined in a natural way on all non-negative Borel functions on  $X$ , in particular on Borel subsets of  $X$ .

**THEOREM 3.3.** *Let  $\mathcal{X} \subseteq M_+^1(M_+(X))$  satisfy the following two conditions*

$$(3.3.1) \quad \limsup_{A \rightarrow \infty} \sigma(\{ \nu \in M_+(X) \mid \nu(C) > A \}) = 0, \quad \forall C \subseteq X \text{ compact},$$

$$(3.3.2) \quad \limsup_C \limsup_{\sigma \in \mathcal{X}} (1 - \hat{\sigma}(X \setminus C)) = 0,$$

where the limit is taken along the net of compact subsets of  $X$ . Then  $\mathcal{X}$  is a relatively compact subset of  $M_+(M_+(X))$ .

**PROOF.** Let  $0 < \varepsilon < 1$ ,  $0 < \delta < 1$ ; then  $1 - e^{-\delta} \geq \frac{1}{2}\delta$  and

$$\begin{aligned} 1 - \hat{\sigma}(X \setminus C) &= \int_{M_+(X)} (1 - e^{-\lambda(X \setminus C)}) d\sigma(\lambda) \\ &\geq \int_{\{ \lambda \mid \lambda(X \setminus C) \geq \frac{1}{2}\delta \}} (1 - e^{-\lambda(X \setminus C)}) d\sigma(\lambda) \geq \frac{\delta}{4} \sigma \left( \left\{ \lambda \mid \lambda(X \setminus C) \geq \frac{\delta}{2} \right\} \right). \end{aligned}$$

By (3.3.2) there exists a compact set  $C \subseteq X$  such that

$$\sup_{\sigma \in \mathcal{X}} \frac{\delta}{4} \sigma \left( \left\{ \lambda \mid \lambda(X \setminus C) \geq \frac{\delta}{2} \right\} \right) \leq \frac{\varepsilon \delta}{24}$$

hence

$$\inf_{\sigma \in \mathcal{X}} \sigma \left( \left\{ \lambda \mid \lambda(X \setminus C) < \frac{\delta}{2} \right\} \right) \geq 1 - \frac{\varepsilon}{6}$$

and applying (3.3.1) we find  $A \in \mathbf{R}$  such that

$$\inf_{\sigma \in \mathcal{X}} \sigma(\{\lambda \mid \lambda(C) \leq A\}) \geq 1 - \frac{\varepsilon}{6}.$$

Putting

$$L_1 := \left\{ \lambda \in M_+(X) \mid \lambda(X \setminus C) < \frac{\delta}{2} \text{ and } \lambda(C) \leq A \right\}$$

we have

$$\inf_{\sigma \in \mathcal{X}} \sigma(L_1) \geq 1 - \frac{\varepsilon}{3}.$$

Now let  $f \in C_+(X)$ ,  $0 \leq f \leq 1$  and  $\sup_{x \in C} f(x) < \varepsilon/3A$ . Then for any  $\sigma \in \mathcal{X}$  we get

$$1 - \hat{\sigma}(f) = \int_{M_+(X)} (1 - e^{-\lambda(f)}) d\sigma(\lambda) \leq \frac{\varepsilon}{3} + \int_{L_1} \lambda(f) d\sigma(\lambda)$$

and for  $\lambda \in L_1$

$$\lambda(f) = \int_X f d\lambda \leq \frac{\delta}{2} + \int_C f d\lambda \leq \frac{\delta}{2} + \frac{\varepsilon}{3}.$$

Hence, choosing  $\delta = \frac{2}{3}\varepsilon$ ,  $1 - \hat{\sigma}(f) \leq \varepsilon$ , proving  $\beta$ -equicontinuity of  $\{\hat{\sigma} \mid \sigma \in \mathcal{X}\}$  at 0. From Theorem 3.2 we get the desired conclusion.

#### 4. Completely alternating functions on $C_+(X)$ .

A class of functions on  $C_+(X)$ , closely connected to completely monotone functions, is that of completely alternating (or alternating of infinite order) functions. A function  $\psi: C_+(X) \rightarrow [0, \infty]$  is *completely alternating* if and only if

$$\sum_{i=1}^n \sum_{j=1}^n c_i c_j \psi(f_i + f_j) \leq 0$$

for all  $n \in \mathbf{N}$ ,  $f_1, \dots, f_n \in C_+(X)$  and  $c_1, \dots, c_n \in \mathbf{R}$  such that  $\sum_{i=1}^n c_i = 0$ , (see [1, Proposition 3.2 and Theorem 4.2]). One of the main results in [1] was the ‘‘Lévy–Khinchin’’-representation for completely alternating functions (Theorem 3.7 in [1]). This uniquely determined representation has the form

$$\psi(f) = c + h(f) + \int_{L \setminus \{0\}} (1 - e^{-\lambda(f)}) d\xi_0(\lambda)$$

where  $c \in [0, \infty[$ ,  $h: C_+(X) \rightarrow [0, \infty[$  is additive and  $\xi_0$  is a non-negative Radon-measure on  $L \setminus \{0\}$ . Observing that

$$L \setminus \{0\} = \{\lambda \in L \mid \lambda(1) > 0\}$$

we can write this representation in the following form

$$(4.1) \quad \psi(f) = c + \int_{\beta X} \tilde{f} d\kappa + \int_{L \setminus \{0\}} \frac{1 - e^{-\lambda(f)}}{1 - e^{-\lambda(1)}} d\xi(\lambda)$$

where  $\kappa \in M_+(\beta X)$ ,  $\xi \in M_+(L)$  and

$$\Delta_1 \psi(f) := \psi(f+1) - \psi(f) = \int_L e^{-\lambda(f)} d\xi(\lambda),$$

cf. the proof of Theorem 3.7 in [1].

Note that each completely alternating function  $\psi$  on  $C_+(X)$  satisfies the inequalities

$$(4.2) \quad \alpha \psi(f) \leq \psi(\alpha f) \quad \forall f \in C_+(X), \forall \alpha \in [0, 1]$$

$$(4.3) \quad \psi(\beta f) \leq \beta \psi(f) \quad \forall f \in C_+(X), \forall \beta \in [1, \infty[.$$

This follows from (4.1) and from the fact that

$$1 - e^{-\alpha \lambda} \geq \alpha(1 - e^{-\lambda}) \quad \forall \lambda \geq 0, \forall \alpha \in [0, 1]$$

$$1 - e^{-\beta \lambda} \leq \beta(1 - e^{-\lambda}) \quad \forall \lambda \geq 0, \forall \beta \in [1, \infty[$$

which is easily established using Cauchy’s mean value theorem. Another important property is subadditivity

$$(4.4) \quad \psi(f+g) \leq \psi(f) + \psi(g) \quad \forall f, g \in C_+(X),$$

cf. Proposition 3.5 in [1].

**THEOREM 4.1.** *Let the completely alternating function  $\psi: C_+(X) \rightarrow [0, \infty[$  have the representation (4.1). Then  $\xi(L \setminus L_0) = 0$  if and only if  $\lim_{t \rightarrow 0} \psi(t) = \psi(0)$ .*

PROOF. We may and do assume  $\psi(0)=c=0$ . Suppose that  $\lim_{t \rightarrow 0} \psi(t)=0$ . By (4.4)

$$0 \leq \psi(t+1) - \psi(1) \leq \psi(t) \rightarrow 0 \quad \text{as } t \rightarrow 0,$$

hence

$$\Delta_1 \psi(t) = \psi(t+1) - \psi(t) \rightarrow \psi(1) = \Delta_1 \psi(0)$$

and we get  $\xi(L \setminus L_0)=0$  from Theorem 2.2. The other direction follows immediately from (4.1).

THEOREM 4.2. Let  $\psi: C_+(X) \rightarrow [0, \infty[$  be a completely alternating function with the representation (4.1); let  $\varrho$  be any topology on  $C(X)$  satisfying  $\tau \subseteq \varrho \subseteq \beta$  and put

$$B := \{f \in C(X) \mid 0 \leq f \leq 1\}.$$

Then the following 5 statements are equivalent:

(4.2.1)  $\exists Y$   $\sigma$ -compact  $\subseteq X$  such that  $\kappa(X \setminus Y)=0$  and  $\xi(L \setminus L_Y)=0$

(4.2.2)  $\psi|_B$  is  $\tau$ -continuous at 0

(4.2.3)  $\psi|_B$  is uniformly  $\varrho$ -continuous

(4.2.4)  $\psi|_B$  is  $\beta$ -continuous at 0

(4.2.5)  $\forall \varepsilon > 0 \exists \delta > 0 \exists C$  compact  $\subseteq X$  such that  $\psi(f) - \psi(0) \leq \varepsilon$  whenever  $f \in B$  and  $f \leq \delta$  on  $C$ .

PROOF. Again we assume  $\psi(0)=c=0$ .

(4.2.1)  $\Rightarrow$  (4.2.2): The function  $f \mapsto \int_{\beta X} \tilde{f} d\kappa$  is  $\tau$ -continuous because  $\kappa \in M_+(X)$ , cf. Corollary 1.2. By Theorem 2.2 there exist compact uniformly tight subsets  $K_n \subseteq L_Y \setminus \{0\}$  such that  $\xi(L \setminus K_n) \leq 1/n$ . We define

$$\psi_n(f) := \int_{K_n} \frac{1 - e^{-\lambda(f)}}{1 - e^{-\lambda(1)}} d\xi(\lambda), \quad f \in C_+(X), n \in \mathbb{N}.$$

By Theorem 2.3  $\{\psi_n\}$  is a sequence of  $\tau$ -continuous completely alternating functions. Now

$$\sup_{f \in B} \frac{1 - e^{-\lambda(f)}}{1 - e^{-\lambda(1)}} \leq 1 \quad \text{for all } \lambda \in L$$

implying that  $\psi_n$  converges uniformly to  $\psi(f) - \int f d\kappa$  on  $B$ . Hence  $\psi|_B$  is  $\tau$ -continuous.

(4.2.2)  $\Rightarrow$  (4.2.3): If  $f, g$  belong to  $C_+(X)$  then applying the subadditivity (4.4) we have

$$\begin{aligned} |\psi(f) - \psi(g)| &\leq \psi(f \vee g) - \psi(f) + \psi(f \vee g) - \psi(g) \\ &\leq \psi((f-g)^+) + \psi((g-f)^+). \end{aligned}$$

If now  $\psi|B$  is  $\tau$ -continuous at 0, then  $\psi|B$  is uniformly  $\tau$ -continuous, as one can see immediately from the definition of the  $\tau$ -topology.

(4.2.3)  $\Rightarrow$  (4.2.4): Obvious.

(4.2.4)  $\Rightarrow$  (4.2.5): This can be seen as in Theorem 2.3.

(4.2.5)  $\Rightarrow$  (4.2.1): Let  $\varepsilon > 0$  and choose  $\delta > 0$ ,  $C \subseteq X$  compact such that  $\psi(f) < \varepsilon$  for all  $f \in B$ ,  $f \leq \delta$  on  $C$ . For those  $f$  we get

$$\Delta_1 \psi(0) - \Delta_1 \psi(f) = \psi(1) - \psi(f+1) + \psi(f) \leq \psi(f) < \varepsilon$$

hence there exists by Theorem 2.3 a  $\sigma$ -compact subset  $Y_1 \subseteq X$  such that  $\xi(L \setminus L_{Y_1}) = 0$ . The function  $f \mapsto \int_{\beta X} \tilde{f} d\kappa$  has of course also the continuity property of (4.2.5), hence by Theorem 1.3.,  $\kappa$  belongs to  $M_+(X)$  and therefore is concentrated on a  $\sigma$ -compact subset  $Y_2 \subseteq X$ . The union  $Y := Y_1 \cup Y_2$  fulfills condition (4.2.1).

#### REFERENCES

1. C. Berg, J. P. R. Christensen and P. Ressel, *Positive definite functions on abelian semigroups*, Math. Ann. (to appear).
2. G. Choquet, *Theory of capacities*, Ann. Inst. Fourier (Grenoble), 5 (1954), 131–295.
3. J. Hoffmann-Jørgensen, *The theory of analytic spaces*, Århus University, Various Publications Series No. 10 (1970).
4. J. Hoffmann-Jørgensen, *A generalization of the strict topology*, Math. Scand. 30 (1972), 313–323.
5. J. Hoffmann-Jørgensen, *Weak compactness and tightness of subsets of  $M(X)$* , Math. Scand. 31 (1972), 127–150.
6. D. Pollard and F. Topsøe, *A unified approach to Riesz type representation theorems*, Studia Math. 54 (1975), 173–190.
7. F. Topsøe, *Topology and measure*, Lecture Notes in Mathematics 133, Springer-Verlag, Berlin · Heidelberg · New York, 1970.

UNIVERSITY OF ÅRHUS  
DENMARK

AND

UNIVERSITY OF FREIBURG/BR.  
WEST-GERMANY