

On construction of recursion operators from Lax representation

Metin Gürses, Atalay Karasu, and Vladimir V. Sokolov

Citation: *J. Math. Phys.* **40**, 6473 (1999); doi: 10.1063/1.533102

View online: <http://dx.doi.org/10.1063/1.533102>

View Table of Contents: <http://jmp.aip.org/resource/1/JMAPAQ/v40/i12>

Published by the [American Institute of Physics](#).

Related Articles

Conditional stability theorem for the one dimensional Klein-Gordon equation
J. Math. Phys. **52**, 112703 (2011)

Solitary wave evolution in a magnetized inhomogeneous plasma under the effect of ionization
Phys. Plasmas **18**, 102116 (2011)

Modulational instability of ion acoustic waves in e-p-i plasmas with electrons and positrons following a q-nonextensive distribution
Phys. Plasmas **18**, 102313 (2011)

On polygonal relative equilibria in the N-vortex problem
J. Math. Phys. **52**, 103101 (2011)

Global existence and blow-up phenomena for a weakly dissipative periodic 2-component Camassa-Holm system
J. Math. Phys. **52**, 103701 (2011)

Additional information on *J. Math. Phys.*

Journal Homepage: <http://jmp.aip.org/>

Journal Information: http://jmp.aip.org/about/about_the_journal

Top downloads: http://jmp.aip.org/features/most_downloaded

Information for Authors: <http://jmp.aip.org/authors>

ADVERTISEMENT



AIPAdvances

Submit Now

Explore AIP's new
open-access journal

- Article-level metrics now available
- Join the conversation! Rate & comment on articles

On construction of recursion operators from Lax representation

Metin Gürses^{a)}

*Department of Mathematics, Faculty of Sciences, Bilkent University,
06533 Ankara—Turkey*

Atalay Karasu

*Department of Physics, Faculty of Arts and Sciences, Middle East Technical University,
06531 Ankara—Turkey*

Vladimir V. Sokolov

Landau Institute, Moscow, 117940 Russia

(Received 28 June 1999; accepted for publication 16 August 1999)

In this work we develop a general procedure for constructing the recursion operators for nonlinear integrable equations admitting Lax representation. Several new examples are given. In particular, we find the recursion operators for some KdV-type systems of integrable equations. © 1999 American Institute of Physics. [S0022-2488(99)03212-0]

I. INTRODUCTION

It is well known that most of the integrable nonlinear partial differential equations,

$$u_t = F(t, x, u, u_x, \dots, u_{nx}), \quad (1)$$

admit a Lax representation,

$$L_t = [A, L], \quad (2)$$

so that the inverse scattering method is applicable. The generalized symmetries¹ of (1) have also Lax representations with the same L operator,

$$L_{t_n} = [A_n, L], \quad n \geq 1. \quad (3)$$

The recursion operator \mathcal{R} , satisfying the equation (see Ref. 2)

$$\mathcal{R}_t + [D_F, \mathcal{R}] = 0, \quad (4)$$

where D_F is the Frechét derivative of the function F , generates symmetries of (1) starting from the simplest ones. In general, \mathcal{R} is a nonlocal operator (a pseudodifferential operator).

The construction of the recursion operator of a given integrable system (1) is not an easy task. Several works are devoted to this subject. Among these works, most of the authors use (4) for the construction of the recursion operator.^{3–8} There are several difficulties in this direct approach. The main problems are the choices of the order of \mathcal{R} and the structure of the nonlocal terms. This is an approach having no relation with the Lax representation (2).

On the other hand, some of the authors used Lax representation for this purpose. Most of these works are related to the squared eigenfunctions of the Lax operator^{9–13} and are based on finding an eigenvalue equation for the squared eigenfunctions of the Lax operator. The operator corresponding to this eigenvalue equation turns out to be the adjoint of the recursion operator.

^{a)}Electronic mail: gurses@fen.bilkent.edu.tr

There is an alternative use of the Lax representation to construct recursion operators. This approach is based on the explicit construction of the A_n operators (3). It was first used by Symes,¹⁴ Adler¹⁵ (see also Dorfman–Fokas,¹⁶ Fokas–Gel’fand¹⁷) and Antonowicz–Fordy.^{18,19} Although these authors use the Lax representation in different ways, their approach is basically the same. Symes and Adler use the Gel’fand–Dikii²⁰ construction of the A_n operators. On the other hand, Antonowicz–Fordy determines these operators from integrability condition (3) and by using an ansatz for A_n . Their basic aim is to determine the Hamiltonian operators θ_1 and θ_2 ²¹ of the equations under consideration. The recursion operator is simply given by $\mathcal{R} = \theta_2 \theta_1^{-1}$. Their approach is based on some explicit formulas for coefficients of the A_n operator. This is necessary to find the Hamiltonian operators θ_1 and θ_2 , and it seems that this approach is quite effective to determine the bi-Hamiltonian structure for the simple cases but it becomes more complicated when the L -operator has a sophisticated structure.

If one is interested only in the determination of the recursion operator \mathcal{R} , we shall show in this work that it is possible to succeed this without any concrete information of the coefficients of A_n operators. We use only an ansatz $\tilde{A} = \mathcal{P}A + R$ that relates A_n operators for different n . Here \mathcal{P} is some operator that commutes with the L operator and R is the remainder.

We follow this basic idea, partially used by Symes,¹⁴ Adler,¹⁵ Shabat and Sokolov,²² and establish an extremely simple, effective, and algorithmic method for the construction of recursion operators when the Lax representation (2) is given.²³

In the next section we consider the case where L is a scalar operator. We first consider the case where L is a differential operator and then the case where it is a pseudodifferential operator. In each case we present our method, discuss the reductions, and give examples for illustrations. In Sec. III we consider Lax operator taking values in a Lie algebra. We give our method both for the general case and also for the reductions. We give one example for each case in the text. Several additional examples are given in the Appendices A, B, and C corresponding to all different cases.

II. SCALAR LAX REPRESENTATIONS

First we consider equations with the scalar Lax representations of the form

$$L_t = [A, L], \quad (5)$$

where L is, in general, a pseudodifferential operator of order m and A is a differential operator whose coefficients are functions of x and t .

The different choice of operators A for a given L leads to a hierarchy of nonlinear systems (3). It is well known that one can define operators A_n by the following formula:²⁰

$$A_n = (L^{n/m})_+, \quad (6)$$

where $L^{n/m}$ is a pseudodifferential series of the form $L^{n/m} = \sum_{-\infty}^n v_i D^i$ and $(L^{n/m})_+ = \sum_{i=0}^n v_i D^i$. Here v_i are some concrete functions depending on the coefficients of L and D is the total derivative with respect to x .

In Refs. 25 and 26 the relationships between the Kac–Moody algebras and special types of scalar differential and pseudodifferential operators L were established. All corresponding integrable systems are Hamiltonian ones. For most of them a second Hamiltonian structure is not known up to now.

In this section and Appendices A, B, and C we consider the simplest systems from Refs. 25 and 26 as examples and find their recursion operators. In the sequel these examples will be referred to as Drinfeld–Sokolov (DS) systems. It is interesting to note that in all these examples the order of the recursion operator is equal to the Coxeter number of the corresponding Kac–Moody algebra.

A. Gel'fand–Dikii systems

In this section we shall consider the case where L is a differential operator,

$$L = D^m + u_{m-2}D^{m-2} + \dots + u_0, \tag{7}$$

where $u_i, i=0,1,\dots,m-2$ are functions of x, t . In the framework of Ref. 25, this corresponds to the Kac–Moody algebras of the type $A_{m-1}^{(1)}$.

To show that (3) is equivalent to a system of $(m-1)$ evolution equations with respect to u_i one can use the following standard reasoning. Set

$$L^{n/m} = (L^{n/m})_+ + (L^{n/m})_-, \tag{8}$$

where $(L^{n/m})_+$ is the differential part of the series $L^{n/m}$ and $(L^{n/m})_-$ is a series of order ≤ -1 . Since $[L, L^{n/m}] = 0$ we have

$$[(L^{n/m})_+, L] = [L, (L^{n/m})_-]. \tag{9}$$

The left-hand side of (9) is a differential operator, but the right side is a series of order $\leq n-2$. Thus, both sides of (3) are differential operators of order $\leq n-2$ and it is equivalent to a system of evolution equations for the dependent variables $u_i, i=0,1,\dots,m-2$. This system can be obtained by comparing the coefficients of D^i , where $0,\dots,m-2$ in (3).

Since $L^{(n+m)/m} = LL^{n/m}$, then we have

$$A_{m+n} = (LL^{n/m})_+ = L(L^{n/m})_+ + (L(L^{n/m})_-)_+, \tag{10}$$

which leads directly to

$$L_{t_{n+m}} = [A_{n+m}, L] = LL_{t_n} + [(L(L^{n/m})_-)_+, L]. \tag{11}$$

The above equation (11) has been given also by Symes¹⁴ (see also Adler's paper¹⁵). In his work Symes expressed the coefficients of the both parts of (11), in a rather complicated way, in terms of some finite set of coefficients of the resolvent for an L operator. That allows him to express $L_{t_{n+m}}$ in terms of L_{t_n} . This relation gives directly the recursion operator. He gave explicit formulas for the cases $m=2$ and $m=3$.

In this section we shall show that in order to construct the recursion operator it suffices to know only that

$$L_{t_{n+m}} = LL_{t_n} + [R_n, L]. \tag{12}$$

Obviously, it follows from the following.

Proposition 1: For any n ,

$$A_{n+m} = LA_n + R_n, \tag{13}$$

where R_n is a differential operator of order $\leq m-1$.

Proof: The relation (13) coincides with (10) if we put

$$R_n = (L(L^{n/m})_-)_+. \tag{14}$$

Since $(L^{n/m})_-$ is a series of order ≤ -1 , then $\text{ord}(R_n) \leq m-1$.

Remark 1: It follows from the formula

$$A_{n+m} = (L^{n/m}L)_+ = (L^{n/m})_+L + ((L^{n/m})_-L)_+, \tag{15}$$

that

$$A_{n+m} = A_n L + \bar{R}_n, \tag{16}$$

and

$$L_{t_{n+m}} = L_{t_n} L + [L, \bar{R}_n], \tag{17}$$

where \bar{R}_n is a differential operator of order $\leq m - 1$.

To find the recursion operator we can simply equate the coefficients of different powers of D in (12). It is easy to see that in this comparison of the coefficients of D^i , $i = 2m - 2, \dots, m - 1$ we determine R_n in terms of the coefficients of operators L and L_{t_n} . It is important that the resulting formulas turn out to be linear in the coefficients of L_{t_n} . The remaining coefficients of D^i , $i = m - 2, \dots, 0$ in (12) give us the relation

$$\begin{pmatrix} u_0 \\ \cdot \\ \cdot \\ \cdot \\ u_{m-2} \end{pmatrix}_{t_{n+m}} = \mathcal{R} \begin{pmatrix} u_0 \\ \cdot \\ \cdot \\ \cdot \\ u_{m-2} \end{pmatrix}_{t_n}, \tag{18}$$

where \mathcal{R} is a recursion operator. Instead of (12) one can use (17). The corresponding recursion operators coincide.

Example 1. KdV equation: The KdV equation,

$$u_t = \frac{1}{4}(u_{3x} + 6uu_x), \tag{19}$$

has a Lax representation with

$$L = D^2 + u, \quad A = (L^{3/2})_+. \tag{20}$$

Since in this case $L_{t_{n+2}} = u_{t_{n+2}} \equiv u_{n+2}$ and $L_{t_n} = u_{t_n} \equiv u_n$, the main relation (12) takes the form

$$u_{n+2} = (D^2 + u) \cdot u_n + [R_n, L], \tag{21}$$

with $R_n = a_n D + b_n$.

Now if we equate successively to zero the coefficients of D^2 , D , and D^0 in the above equation, we obtain

$$a_n = \frac{1}{2} D^{-1}(u_n), \quad b_n = \frac{3}{4} u_n,$$

and

$$u_{n+2} = (\frac{1}{4} D^2 + u + \frac{1}{2} u_x D^{-1}) u_n,$$

that gives the standard recursion operator for the KdV equation,

$$\mathcal{R} = \frac{1}{4} D^2 + u + \frac{1}{2} u_x D^{-1}. \tag{22}$$

In the same way one can find a recursion operator for the Boussinesq equation (see Appendix A).

B. Symmetric and skew-symmetric reductions of a differential Lax operator

The standard reductions of the Gel'fand-Dikii systems are given by the conditions $L^* = L$ or $L^* = -L$. Here $*$ denotes the adjoint operation defined as follows. Let L be a differential operator,

$L = \sum a_i D^i$. Its adjoint L^* is given by $L^* = \sum (-D)^i \cdot a_i$. It is easy to see that if $L^* = L$ then $m = \text{ord}(L)$ must be an even integer. For the case $L^* = -L$, it must be an odd integer.

It is well known that for both reductions all possible A_n are defined by (6), where n takes odd integer values. This condition provides that $(A_n)^* = -A_n$ that is necessary for (3) to be compatible.

If $L^* = L$, the formula $A_{n+m} = (LL^{n/m})_+ = (L^{(n+m)/m})_+$ gives a correct A_n operator since $n + m$ is an odd integer. Thus, in this case Proposition 1 remains valid and the recursion operator can be found from (12) or (17).

On the other hand, if $L^* = -L$ then both integers m and n are odd and hence their sum $m + n$ is an even integer. This means that $(L^{(n+m)/m})_+$ cannot be taken as an A_n operator. In this (skew adjoint) case we must take

$$A_{n+2m} = (L^{(n+2m)/m})_+ = (L^2 L^{n/m})_+,$$

to find the recursion operator. Following the proof of Proposition 1 we obtain Proposition 2.

Proposition 2: If $L^* = -L$ then

$$A_{n+2m} = L^2 A_n + R_n, \tag{23}$$

where $\text{ord}(R_n) < 2 \text{ord}(L)$. It follows from (23) that

$$L_{t_{n+2m}} = L^2 L_{t_n} + [R_n, L]. \tag{24}$$

Remark 2: Instead of (23) we can use the ansatz

$$A_{n+2m} = L A_n L + \tilde{R}_n, \tag{25}$$

or

$$A_{n+2m} = A_n L^2 + \tilde{\tilde{R}}_n. \tag{26}$$

The recursion operators obtained by the utility of (23), (25), and (26) all coincide.

In the works,^{25,26} more general reductions $L^\dagger = \pm L$ were also considered. Here $L^\dagger = K L^* K^{-1}$, where K is a given differential operator, such that $L K^{-1}$ is a differential operator. In this general reductions, as well, possible A_n operators are given by (6), with n being an odd integer. Propositions 1 and 2 are valid for this general symmetric and skew-symmetric cases and hence one can use Eqs. (12), (24) accordingly to obtain the recursion operators.

Example 2. Kupershmidt equation: This equation,

$$u_t = u_{5x} + 10uu_{3x} + 25u_x u_{2x} + 20u^2 u_x, \tag{27}$$

has the Lax pair

$$L = D^3 + 2uD + u_x, \quad A = (L^{5/3})_+. \tag{28}$$

In this case $L^* = -L$; therefore we use Eq. (24) with

$$\tilde{R}_n = a_n D^5 + b_n D^4 + c_n D^3 + d_n D^2 + e_n D + f_n. \tag{29}$$

By equating the coefficients of powers of D in (24), we obtain

$$\begin{aligned} a_n &= \frac{2}{3} D^{-1}(u_n), & b_n &= \frac{11}{3} u_n, & c_n &= \frac{1}{9} (20u D^{-1}(u_n) + 73u_{n,x}), \\ d_n &= \frac{1}{3} (10u_x D^{-1}(u_n) + 22uu_n + 27u_{n,2x}), \end{aligned}$$

$$\begin{aligned}
 e_n &= \frac{1}{27}(70u_{2x}D^{-1}(u_n) - 2D^{-1}(u_{2x}u_n) + 40u^2D^{-1}(u_n) - 8D^{-1}(u^2u_n) \\
 &\quad + 134u_{n,3x} + 212uu_{n,x} + 184u_xu_n), \\
 f_{n,x} &= \frac{1}{27}(20u_{4x}D^{-1}(u_n) + 74u_{3x}u_n + 126u_{2x}u_{n,x} + 40uu_{2x}D^{-1}(u_n) + 40u_x^2D^{-1}(u_n) \\
 &\quad + 136u_xu_{n,2x} + 27uu_xu_n + 28u_{n,5x} + 64uu_{n,3x} + 16u^2u_{n,x}),
 \end{aligned}$$

and the recursion operator for the Kupershmidt equation:

$$\begin{aligned}
 \mathcal{R} &= D^6 + 12uD^4 + 36u_xD^3 + (49u_{2x} + 36u^2)D^2 + 5(7u_{3x} + 24uu_x)D + 13u_{4x} + 82uu_{2x} + 69u_x^2 \\
 &\quad + 32u^3 + 2u_xD^{-1}(u_{2x} + 4u^2) + 2(u_{5x} + 10uu_{3x} + 25u_xu_{2x} + 20u^2u_x)D^{-1}. \tag{30}
 \end{aligned}$$

C. Pseudodifferential Lax operator

In this section we generalize our scheme to the case of pseudodifferential Lax operators. The only difference is that in formulas like (13) and (23) the R_n operator also becomes a pseudodifferential operator.

It follows from these formulas that the structure of the nonlocal terms in R_n is, in general, similar to the nonlocal terms in L since A_{n+m} and A_n are differential operators.

For skew-symmetric case, A_n may be defined by either (23) or (25), or (26). In the pseudodifferential case they are not equivalent, in the sense that the nonlocal part of R_n depends on which ansatz we choose. For illustration, let us consider the case $L = MD^{-1}$, where M is a differential operator. The following lemma shows that if $L^\dagger = L$ or $L^\dagger = -L$, where

$$L^\dagger = DL^*D^{-1}, \tag{31}$$

then the formulas (13) and (25) are much suitable then (16), (23), and (26).

Lemma: Let $L^\dagger = \epsilon L$, where $\epsilon = \pm 1$. Then

$$R_n = D^{m-1} + \dots + a_0, \quad \text{for } \epsilon = 1, \tag{32}$$

where R_n is defined by (13), and

$$\tilde{R}_n = D^{2m-1} + \dots + a_{-1}D^{-1}, \quad \text{for } \epsilon = -1, \tag{33}$$

where \tilde{R}_n is defined by (25).

Proof: If $L = MD^{-1}$ then $L^\dagger = \epsilon L$ implies $M^* = -\epsilon M$. It is easy to show that $(L^{1/m})^\dagger = -L^{1/m}$. Hence $(L^{n/m})^\dagger = -L^{n/m}$ for an odd integer n . Define now a series K_n by

$$L^{n/m} = DK_n.$$

It is easy to prove that $K_n^* = K_n$. Since $K_n = (K_n)_+ + (K_n)_-$ and $(K_n)^* = K_n$, we have

$$(K_n)_+^* = (K_n)_+, \quad (K_n)_-^* = (K_n)_-.$$

From the last formula it follows that $\text{ord}(K_n)_- \leq -2$, which leads to an important result,

$$A_n = (L^{n/m})_+ = D(K_n)_+.$$

This implies that

$$LA_n = M(K_n)_+ \tag{34}$$

is a differential operator. Now using (34) in (13) and (25) for the cases $\epsilon = 1$ and $\epsilon = -1$, respectively, we find the ansatz for A_n given by (32) and (33).

Example 3 ($\epsilon = -1$): It is known that the KdV equation has, besides the standard Lax representation, the following Lax pair:

$$L = (D^2 + u)D^{-1}, \quad A = (L^3)_+ \tag{35}$$

The L operator satisfies the reduction $L^\dagger = -L$. According to the formula (33) we have

$$\bar{R}_n = a_n D + b_n + c_n D^{-1}.$$

It follows from (25) that

$$a_n = D^{-1}(u_n), \quad b_n = u_n, \quad c_n = -u_{n,x} - uD^{-1}(u_n).$$

The remaining equation in (25) gives the recursion operator

$$\mathcal{R} = D^2 + 4u + 2u_x D^{-1} \tag{36}$$

Example 4 ($\epsilon = 1$). *DSIII system*: The DSIII system^{25,26} is given by

$$\begin{aligned} u_t &= -u_{3x} + 6uu_x + 6v_x, \\ v_t &= 2v_{3x} - 6uv_x. \end{aligned} \tag{37}$$

The nonlocal Lax representation for this system is

$$\begin{aligned} L &= (D^5 - 2uD^3 - 2D^3u - 2Dw - 2wD)D^{-1}, \\ A &= (L^{3/4})_+, \end{aligned} \tag{38}$$

where $w = v - u_{2x}$. Since $L^\dagger = L$ we can use (32), which gives us

$$R_n = a_n D^3 + b_n D^2 + c_n D + d_n \tag{39}$$

By equating the coefficients of the powers of D in (25), we obtain

$$\begin{aligned} a_n &= D^{-1}(u_n), \quad b_n = 4u_n, \\ c_n &= \frac{1}{2}(-6uD^{-1}(u_n) + 11u_{n,x} + 2D^{-1}(uu_n) + 2D^{-1}(v_n)), \\ d_{n,x} &= -\frac{1}{2}(6u_{2x}D^{-1}(u_n) + 10u_xu_n - 5u_{n,3x} + 4uu_{n,x} - 6v_{n,x}). \end{aligned}$$

The recursion operator of the DSIII is found as

$$\mathcal{R} = \begin{pmatrix} \mathcal{R}_0^0 & \mathcal{R}_1^0 \\ \mathcal{R}_0^1 & \mathcal{R}_1^1 \end{pmatrix}, \tag{40}$$

with

$$\begin{aligned} \mathcal{R}_0^0 &= D^4 - 8uD^2 - 12u_xD - 8u_{2x} + 16u^2 + 16v + (-2u_{3x} + 12uu_x + 12v_x)D^{-1} + 4u_xD^{-1}u, \\ \mathcal{R}_1^0 &= -10D^2 + 8u + 4u_xD^{-1}, \\ \mathcal{R}_0^1 &= 10v_xD + 12v_{2x} + (4v_{3x} - 12uv_x)D^{-1} + 4v_xD^{-1}u, \\ \mathcal{R}_1^1 &= -4D^4 + 16uD^2 + 8u_xD + 16v + 4v_xD^{-1}. \end{aligned} \tag{41}$$

This recursion operator has recently been given in Ref. 6.

III. MATRIX L OPERATOR OF THE FIRST ORDER

In this section we demonstrate how our approach, given in the previous sections, can be generalized to the case where L is a matrix operator of the form

$$L = D_x + \lambda a + q(x, t). \tag{42}$$

A. General case

Let us consider the Lax operator (42), where q and a belong to a Lie algebra g and λ is the spectral parameter. The constant element a is supposed to be such that

$$g = \text{Ker}(\text{ad}_a) \oplus \text{Im}(\text{ad}_a). \tag{43}$$

First, let us recall the procedure²⁵ of constructing the A operators for the Lax operator (42).

Proposition 3: There exist unique series,

$$u = u_{-1}\lambda^{-1} + u_{-2}\lambda^{-2} + \dots, \quad u_i \in \text{Im}(\text{ad}_a), \tag{44}$$

$$h = h_0 + h_{-1}\lambda^{-1} + h_{-2}\lambda^{-2} + \dots, \quad h_i \in \text{Ker}(\text{ad}_a), \tag{45}$$

such that

$$e^{\text{ad}_u}(L) = L + [u, L] + \frac{1}{2}[u, [u, L]] + \dots = D_x + a\lambda + h. \tag{46}$$

Let b be a constant element of g such that $[b, \text{Ker}(\text{ad}_a)] = \{0\}$. It follows from (45) that $[b\lambda^n, D_x + a\lambda + h] = 0$. Hence $[\Phi_{b,n}, L] = 0$, where

$$\Phi_{b,n} = e^{-\text{ad}_u}(b\lambda^n). \tag{47}$$

Then the corresponding A operator of the form

$$A_{b,n} = b\lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_0, \tag{48}$$

is defined by the formula

$$A_{b,n} = (\Phi_{b,n})_+, \tag{49}$$

where

$$(\sum_{-\infty}^n \alpha_i \lambda^i)_+ = \sum_0^n \alpha_i \lambda^i. \tag{50}$$

According to (47),

$$\Phi_{b,n+1} = \lambda \Phi_{b,n}. \tag{51}$$

Hence

$$A_{b,n+1} = (\lambda \Phi_{b,n})_+ = \lambda (\Phi_{b,n})_+ + (\lambda (\Phi_{b,n})_-)_+. \tag{52}$$

The last formula shows that

$$A_{b,n+1} = \lambda A_{b,n} + R_n, \quad R_n \in g, \tag{53}$$

where R_n does not depend on λ . Substituting (53) into the Lax equation $L_{t_{n+1}} = [A_{b,n+1}, L]$, we get

$$L_{t_{n+1}} = \lambda L_{t_n} + [R_n, L]. \tag{54}$$

Using the ansatz (54), one can easily find the corresponding recursion operator.

Example 5: The system

$$\begin{aligned} u_t &= -\frac{1}{2}u_{xx} + u^2v, \\ v_t &= \frac{1}{2}v_{xx} - v^2u, \end{aligned} \tag{55}$$

is equivalent to the nonlinear Schrödinger equation, has a Lax operator

$$L = D + \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \lambda + \begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}. \tag{56}$$

The Lie algebra g in this example coincides with $sl(2)$.

Using (54) with

$$R_n = \begin{pmatrix} a_n & b_n \\ c_n & -a_n \end{pmatrix},$$

we find that

$$\begin{aligned} a_n &= \frac{1}{2}D^{-1}(vu_n + uv_n), \\ b_n &= \frac{1}{2}u_n, \quad c_n = -\frac{1}{2}v_n, \end{aligned}$$

and the recursion operator of the system (55) is given by

$$\mathcal{R} = \begin{pmatrix} -\frac{1}{2}D + uD^{-1}v & uD^{-1}u \\ -vD^{-1}v & \frac{1}{2}D - vD^{-1}u \end{pmatrix}. \tag{57}$$

B. Reductions in matrix case

In the general case considered in the previous section the A_n operators belong to the Lie algebra,

$$\mathfrak{a}_+ = \{ \sum_{i=0}^{\kappa} a_i \lambda^i, \quad a_i \in g, \quad \kappa \in Z_+ \}, \tag{58}$$

that is a subalgebra of the Lie algebra,

$$\mathfrak{a} = \{ \sum_{i=-\infty}^{\kappa} a_i \lambda^i, \quad a_i \in g, \quad \kappa \in Z \}. \tag{59}$$

A standard σ reduction is defined by any automorphism σ of the Lie algebra g of finite order κ . Because $\sigma^\kappa = \text{Id}$, the eigenvalues of σ are $\epsilon^i, i=0, \dots, \kappa-1$, where ϵ is a primitive κ root of unity.

Let g_i be an eigenspace corresponding to eigenvalue ϵ^i . Then the following reduction $a_j \in g_i$, where $i = j \pmod{\kappa}$ in (58) and (59) is compatible with Eqs. (3). Note that according to this definition $a \in g_1$, and the potential $q(x, t)$ in (42) belongs to g_0 or, the same, satisfies $\sigma(q) = q$.

It is easy to see that, to satisfy such a reduction, we must use the ansatz

$$A_{b,n+\kappa} = \lambda^\kappa A_{b,n} + R_n, \tag{60}$$

where

$$R_n = r_{\kappa-1} \lambda^{\kappa-1} + \dots + r_0, \quad r_i \in g_i. \tag{61}$$

Further generalizations are associated with modifications of sign “+” in (50), which corresponds to the simplest decomposition of algebra \mathfrak{a} into the direct sum of two subalgebras,

$$\mathfrak{a} = \mathfrak{a}_+ \oplus \mathfrak{a}_-, \tag{62}$$

where \mathfrak{a}_+ is given by (58) and

$$\mathfrak{a}_- = \{ \sum_{-\infty}^{-1} a_i \lambda^i, a_i \in g \}. \tag{63}$$

The sign “+” in (50) is the projection of onto \mathfrak{a}_+ parallel to \mathfrak{a}_- . If we have a different decomposition (62), then the construction from Proposition 3 is also valid, but we have the following condition:

$$R_n \in \mathfrak{a}_+ \cap \lambda \mathfrak{a}_-, \tag{64}$$

instead of $R_n \in g$. If we also have the σ reduction, we must use the most general ansatz (60), where

$$R_n \in \mathfrak{a}_+ \cap \lambda^k \mathfrak{a}_-. \tag{65}$$

Example 6: Let us consider the following equation:

$$u_t = \frac{1}{4} u_{xxx} - \frac{3}{8} u_{xx} u + \frac{3}{8} u u_{xx} - \frac{3}{8} u u_x u, \tag{66}$$

where u is a square matrix of arbitrary size, or more generally, u belongs to an arbitrary associative algebra \mathcal{K} . This equation has a Lax representation with

$$L = D + \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \lambda + \begin{pmatrix} u & 0 \\ 0 & 0 \end{pmatrix}. \tag{67}$$

Here $\mathbf{1}$ is the unity of \mathcal{K} . The reduction (67) can be described as follows (see Ref. 27). The Lie algebra g is the algebra of all 2×2 matrices with entries belonging to \mathcal{K} . The automorphism σ is defined by

$$\sigma(X) = TXT^{-1}, \tag{68}$$

where

$$T = \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix}.$$

Obviously $\sigma^2 = \text{Id}$ and eigenvalues of σ are 1 and -1 . The corresponding eigenspaces are

$$g_0 = \left\{ \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix} \right\}, \quad g_1 = \left\{ \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix} \right\}, \tag{69}$$

and therefore the coefficients a_i in (59) have the following structure:

$$a_{2j} = \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix}, \quad a_{2j+1} = \begin{pmatrix} 0 & * \\ * & 0 \end{pmatrix}. \tag{70}$$

The subalgebra \mathfrak{a}_+ is given by (58), where the coefficients have the structure (70) and, additionally,

$$a_0 = \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}.$$

The subalgebra \mathfrak{a}_- has the following form:

$$a_- = \sum_{-\infty}^0 a_i \lambda^i, \tag{71}$$

where a_0 is of the form

$$a_0 = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}, \quad \alpha \in \mathcal{K}.$$

The A operator for (66) is given by formula $A = (\Phi_{a,3})_+$ [see (49)], where

$$a = \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix},$$

and “+” means the projection onto a_+ parallel to a_- .

According to (65), R_n is of the form

$$R_n = \begin{pmatrix} a_n & 0 \\ 0 & a_n \end{pmatrix} \lambda^2 + \begin{pmatrix} 0 & b_n \\ c_n & 0 \end{pmatrix} \lambda + \begin{pmatrix} d_n & 0 \\ 0 & 0 \end{pmatrix}. \tag{72}$$

It follows from

$$L_{t_{n+2}} = \lambda^2 L_{t_n} + [R_n, L], \tag{73}$$

that

$$\begin{aligned} u_n - a_{n,x} + [a_n, u] + b_n - c_n &= 0 & c_n - b_n - a_{n,x} &= 0, \\ d_n - b_{n,x} - u b_n &= 0, & d_n + c_{n,x} - c_n u &= 0, \\ u_{n+2} &= -d_{n,x} + [d_x, u]. \end{aligned}$$

Finding a_n , b_n , c_n , and d_n from this system, we obtain the following recursion operator:

$$\mathcal{R} = -(D + \text{ad}_u)(-D + R_u)(2D + \text{ad}_u)^{-1}(D + L_u)D(2D + \text{ad}_u)^{-1}, \tag{74}$$

where R_u and L_u are the operators of right and left multiplications by u , respectively.

Note that in the commutative case (66) coincides with the modified KdV equation. It is easy to verify that (74) becomes the standard recursion operator of a modified KdV equation. All factors in (74) have to be regarded as operators acting on a (noncommutative) polynomial depending on u, u_x, u_{xx}, \dots .

IV. CONCLUSION

In this work we devoted ourselves in the construction of recursion operators when the Lax representation is given. We have shown that our approach can be easily generalized to all cases where the L operator is a polynomial of λ . It would be interesting to generalize it for the cases of more complicated λ dependence of L as well as for the cases of 2 + 1-dimensional equations, Toda-type lattices, and ordinary differential equations.

ACKNOWLEDGMENTS

We would like to thank Dr. Jing Ping Wang for reading the manuscript and pointing out some misprints. This work is partially supported by the Scientific and Technical Research Council of Turkey (TUBITAK). M. G. is a member of the Turkish Academy of Sciences (TUBA). V. S. is supported by Russian Foundation of Basic Researches (RFBR) Grant No. 99-01-00294 and INTAS.

APPENDIX A: EXAMPLE TO SEC. II A

The Boussinesq equation,

$$u_{tt} = -\frac{1}{3}(u_{4x} + 2(u^2)_{2x}), \quad (\text{A1})$$

can be expressed in the form of a pair of first-order evolution equations,

$$\begin{aligned} u_t &= v_x, \\ v_t &= -\frac{1}{3}(u_{3x} + 8uu_x). \end{aligned} \quad (\text{A2})$$

This system has a Lax pair,

$$L = D^3 + 2uD + u_x + v, \quad A = (L^{2/3})_+. \quad (\text{A3})$$

To construct the recursion operator for this system, we use Eq. (12) with the differential operator,

$$R_n = a_n D^2 + b_n D + c_n.$$

By equating the coefficients of the powers of D in (12), we find

$$\begin{aligned} a_n &= \frac{2}{3}D^{-1}(u_n), \quad b_n = \frac{1}{3}(5u_n + D^{-1}(v_n)), \\ c_n &= \frac{1}{9}(6v_n + 8uD^{-1}(u_n) + 10u_{n,x}), \end{aligned}$$

and after that we obtain the recursion operator of the form (40) for (A2) with

$$\begin{aligned} \mathcal{R}_0^0 &= 3v + 2v_x D^{-1}, \\ \mathcal{R}_1^0 &= D^2 + 2u + u_x D^{-1}, \\ \mathcal{R}_0^1 &= -\left(\frac{1}{3}D^4 + \frac{10}{3}uD^2 + 5u_x D + 3u_{2x} + \frac{16}{3}u^2 + \left(\frac{2}{3}u_{3x} + \frac{16}{3}uu_x\right)D^{-1}\right), \\ \mathcal{R}_1^1 &= 3v + v_x D^{-1}. \end{aligned} \quad (\text{A4})$$

APPENDIX B: EXAMPLES TO SEC. II B**1. Sawada–Kotera equation**

The Lax pair for the Sawada–Kotera equation,²⁸

$$u_t = u_{5x} + 5uu_{3x} + 5u_x u_{2x} + 5u^2 u_x, \quad (\text{B1})$$

is given by

$$L = D^3 + uD, \quad A = (L^{5/3})_+. \quad (\text{B2})$$

In this example, $L^\dagger = -L$, where $L^\dagger = D^{-1}L^*D$ and L is skew-symmetric, then we use (24). The operator \tilde{R}_n has the same form as (29), with the coefficients given by

$$\begin{aligned} a_n &= \frac{1}{3}D^{-1}(u_n), \quad b_n = \frac{5}{3}u_n, \quad c_n = \frac{1}{9}(5uD^{-1}(u_n) + 29u_{n,x}), \\ d_n &= \frac{1}{9}(5u_x D^{-1}(u_n) + 14uu_n + 26u_{n,2x}), \end{aligned}$$

$$e_n = \frac{1}{27}(10u_{2x}D^{-1}(u_n) - 2D^{-1}(u_{2x}u_n) - D^{-1}(u^2u_n) + 5u^2D^{-1}(u_n) + 28u_{n,3x} + 32uu_{n,x} + 32u_xu_n),$$

$$f_n = 0.$$

The recursion operator is given as

$$\mathcal{R} = D^6 + 6uD^4 + 9u_xD^3 + (9u^2 + 11u_{2x})D^2 + (10u_{3x} + 21uu_x)D + 5u_{4x} + 16uu_{2x} + 6u_x^2 + 4u^3 + (u_{5x} + 5uu_{3x} + 5u_xu_{2x} + 5u^2u_x)D^{-1} + u_xD^{-1}(u^2 + 2u_{2x}). \tag{B3}$$

2. DSI system

The DSI system,^{25,26}

$$u_t = 3vv_x,$$

$$v_t = 2v_{3x} + 2uv_x + vu_x, \tag{B4}$$

has a Lax representation with

$$L = [D^3 + (u + v)D + \frac{1}{2}(u + v)_x][D^3 + (u - v)D + \frac{1}{2}(u - v)_x],$$

$$A = (L^{1/2})_+. \tag{B5}$$

Here R_n is a differential operator of order 5, and since L is symmetric we again use Eq. (12). The expressions for the coefficients of the operator R_n are very long and complicated. Hence we do not display them here. We find that the recursion operator \mathcal{R} of this system is of the form (40), where

$$\begin{aligned} \mathcal{R}_0^0 &= -4D^6 - 24uD^4 - 27u_xD^3 + 2(-49u_{2x} - 18u^2 + 42v^2)D^2 + 10(-7u_{3x} - 12uu_x + 30vv_x)D \\ &\quad - 26u_{4x} - 82uu_{2x} - 69u_x^2 + 222vv_x + 141v_x^2 - 16u^3 + 48v^2u \\ &\quad + 2(-2u_{5x} - 10uu_{3x} - 25u_xu_{2x} - 10u^2u_x + 15v^2u_x + 30vv_{3x} + 45v_xv_{2x} + 30uv_{v_x})D^{-1} \\ &\quad + 2u_xD^{-1}(3v^2 - 2u^2 - u_{2x}), \\ \mathcal{R}_1^0 &= 168vD^4 + 204vD^3 + 6(21v_{2x} + 32uv)D^2 + 6(40vu_x + 7v_{3x} + 22uv_x)D \\ &\quad + 6(13vu_{2x} + 10u_xv_x + v_{4x} + 5uv_{2x} + 4vu^2 + 12v^3) + 108vv_xD^{-1}v + 2u_xD^{-1}(6uv + 9v_{2x}), \\ \mathcal{R}_0^1 &= 56vD^4 + 268v_xD^3 + 2(243v_{2x} + 32uv)D^2 + 2(36vu_x + 219v_{3x} + 106uv_x)D \\ &\quad + 2(27vu_{2x} + 92u_xv_x + 99v_{ax} + 99uv_{2x} + 4vu^2 + 12v^3) + 2(10vu_{3x} + 35u_{2x}v_x + 45u_xv_{2x} \\ &\quad + 10uvu_x + 18v_{5x} + 30uv_{3x} + 10u^2v_x + 15v^2v_x)D^{-1} + 2v_xD^{-1}(3v^2 - 2u^2 - u_{2x}), \\ \mathcal{R}_1^1 &= 108D^6 + 216uD^4 + 432u_xD^3 + 6(81u_{2x} + 18u^2 + 22v^2)D^2 + 6(45u_{3x} + 36uu_x + 70vv_x)D \\ &\quad + 3(18u_{4x} + 18uu_{2x} + 9u_x^2 + 98vv_{2x} + 67v_x^2 + 32uv^2) + 36(2v_{3x} + 2v_xu + vu_x)D^{-1}v \\ &\quad + 2v_xD^{-1}(6uv + 9v_{2x}). \end{aligned} \tag{B6}$$

3. DSII system

The DSII system,^{25,26}

$$\begin{aligned} u_t &= 3v_x, \\ v_t &= -2(v_{3x} + uv_x + vu_x), \end{aligned} \tag{B7}$$

has a Lax representation with

$$\begin{aligned} L &= (D^5 + uD^3 + D^3u + (v + \frac{1}{2}u^2)D + D(v + \frac{1}{2}u^2))D, \\ A &= (L^{1/2})_+. \end{aligned} \tag{B8}$$

Since L is symmetric we again use Eq. (12). In this case the operator R_n is given as follows:

$$R_n = a_n D^5 + b_n D^4 + c_n D^3 + d_n D^2 + e_n D, \tag{B9}$$

where

$$\begin{aligned} a_n &= \frac{1}{3}D^{-1}(u_n), \quad b_n = \frac{5}{3}u_n, \\ c_n &= \frac{1}{9}[5uD^{-1}(u_n) + 3D^{-1}(v_n) + 29u_{n,x}], \\ d_n &= \frac{1}{9}[5u_x D^{-1}(u_n) + 26u_{n,2x} + 14uu_n + 12v_n], \\ e_n &= \frac{1}{27}[5(2u_{2x} + u^2 + 3v)D^{-1}(u_n) - 3D^{-1}(vu_n + uv_n) + 9uD^{-1}(v_n) \\ &\quad - 2D^{-1}(u_{2x}u_n + \frac{1}{2}u^2u_n) + 54u_xu_n + 28u_{n,3x} + 32(uu_{n,x} - u_nu_x) + 42v_{n,x}]. \end{aligned}$$

The recursion operator (40) for the system can be found as²⁹

$$\begin{aligned} \mathcal{R}_0^0 &= -D^6 - 6uD^4 - 9u_xD^3 - (11u_{2x} + 9u^2 + 42v)D^2 + (-10u_{3x} - 21uu_x - 30v_x)D \\ &\quad - 5u_{4x} - 16uu_{2x} - 6u_x^2 - 60v_{2x} - 4u^3 - 24vu + (-u_{5x} - 5uu_{3x} - 5u_xu_{2x} \\ &\quad - 5u^2u_x - 15vu_x - 15v_{3x} - 15uv_x)D^{-1} - u_xD^{-1}(2u_{2x} + u^2 + 3v), \\ \mathcal{R}_1^0 &= -42D^4 - 48uD^2 - 87u_xD - 6(7u_{2x} + u^2 - 6v) + 27v_xD^{-1} - 3u_xD^{-1}u, \\ \mathcal{R}_0^1 &= 28vD^4 + 106v_xD^3 + (165v_{2x} + 32uv)D^2 + (54vu_x + 132v_{3x} + 74v_xu)D + 30vu_{2x} + 79u_xv_x \\ &\quad + 54v_{4x} + 57uv_{2x} + 4u^2v - 24v^2 + (10vu_{3x} + 25v_xu_{2x} + 30u_xv_{2x} + 10uvu_x + 9v_{5x} + 15uv_{3x} \\ &\quad + 5u^2v_x - 15vv_x)D^{-1} - v_xD^{-1}(3v + u^2 + 2u_{2x}), \\ \mathcal{R}_1^1 &= 27D^6 + 54uD^4 + 135u_xD^3 + 3(54u_{2x} + 9u^2 - 22v)D^2 + 3(36u_{3x} + 27uu_x - 28v_x)D \\ &\quad + 3(9u_{4x} + 9uu_{2x} + 9u_x^2 - 21v_{2x} - 16vu) - 18(v_{3x} + u_xv + v_xu)D^{-1} - 3v_xD^{-1}u. \end{aligned} \tag{B10}$$

4. DSIV system

The DSIV system,^{25,26} which is also known as the Hirota–Satsuma system,^{30,31}

$$\begin{aligned} u_t &= \frac{1}{2}u_{3x} + 3uu_x - 6vv_x, \\ v_t &= -v_{3x} - 3uv_x, \end{aligned} \tag{B11}$$

has Lax representation with

$$L = (D^2 + u + v)(D^2 + u - v), \quad A = (L^{3/4})_+. \tag{B12}$$

Since the operator L is symmetric we use Eq. (12). In this case the operator R_n has the same form as (39), with coefficients given by

$$\begin{aligned}
 a_n &= \frac{1}{2}D^{-1}(u_n), \quad b_n = \frac{7}{4}u_n - \frac{1}{2}v_n, \\
 c_n &= \frac{1}{8}[6uD^{-1}(u_n) + 2D^{-1}(uu_n) - 4D^{-1}(vv_n) + 17u_{n,x} - 12v_{n,x}], \\
 d_{n,x} &= \frac{1}{16}[6u_{2x}D^{-1}(u_n) - 12v_{2x}D^{-1}(u_n) + 30u_xu_n - 8u_xv_n + 24uu_{n,x} \\
 &\quad + 15u_{n,3x} - 12v_xv_n - 8uv_{n,x} - 20vv_{n,x} - 28v_{n,3x}].
 \end{aligned}$$

The recursion operator (40) for the given system is

$$\begin{aligned}
 \mathcal{R}_0^0 &= \frac{1}{4}D^4 + 2uD^2 + 3u_xD + 2u_{2x} + 4(u^2 - v^2) + (3uu_x - 6vv_x + \frac{1}{2}u_{3x})D^{-1} + u_xD^{-1}u, \\
 \mathcal{R}_1^0 &= -5vD^2 - 4v_xD - v_{2x} - 4uv - 2u_xD^{-1}v, \\
 \mathcal{R}_0^1 &= -\frac{5}{2}v_xD - 3v_{2x} - (v_{3x} + 3uv_x)D^{-1} + v_xD^{-1}u, \\
 \mathcal{R}_1^1 &= -D^4 - 4uD^2 - 2u_xD - 4v^2 - 2v_xD^{-1}v.
 \end{aligned} \tag{B13}$$

5. N=3 Hirota–Satsuma system

This system is given by²⁹

$$\begin{aligned}
 u_t &= \frac{1}{4}u_{3x} + 3uu_x + 3(-v^2 + w)_x, \\
 v_t &= -\frac{1}{2}v_{3x} - 3uv_x, \\
 w_t &= -\frac{1}{2}w_{3x} - 3uw_x.
 \end{aligned} \tag{B14}$$

This is an example for the $N=3$ system that covers some other $N=2$ systems as special cases. For instance, letting $w=0$, we get DSIV and letting $v=0$ we get DSIII systems.

The corresponding Lax pair is

$$L = (D^2 + 2u - 2v)(D^2 + 2u + 2v) + 4w, \quad A = (L^{3/4})_+. \tag{B15}$$

In this case the operator L is symmetric and hence R_n has the same form as (39), with the coefficients

$$\begin{aligned}
 a_n &= D^{-1}(u_n), \quad b_n = \frac{7}{2}u_n + v_n, \\
 c_n &= \frac{1}{4}[12uD^{-1}(u_n) + 4D^{-1}(uu_n + w_n - 2vv_n) + 17u_{n,x} + 12v_{n,x}], \\
 d_{n,x} &= \frac{1}{8}[12u_{2x}D^{-1}(u_n) + 24v_{2x}D^{-1}(u_n) + 60u_xu_n + 16u_xv_n + 15u_{n,3x} + 48uu_{n,x} + 24v_xu_n \\
 &\quad - 40v_xv_n + 20v_{n,3x} + 16vv_{n,x} + 20w_{n,x}].
 \end{aligned}$$

The recursion operator is given by

$$\mathcal{R} = \begin{pmatrix} \mathcal{R}_0^0 & \mathcal{R}_1^0 & \mathcal{R}_2^0 \\ \mathcal{R}_0^1 & \mathcal{R}_1^1 & \mathcal{R}_2^1 \\ \mathcal{R}_0^2 & \mathcal{R}_1^2 & \mathcal{R}_2^2 \end{pmatrix}, \tag{B16}$$

where

$$\begin{aligned}
 \mathcal{R}_0^0 &= \frac{1}{4}D^4 + 4uD^2 + 6u_xD + 4(u_{2x} + 4u^2 - 4v^2 + 4w) \\
 &\quad + 4(\frac{1}{4}u_{3x} + 3uu_x - 6v_{v_x} + 3w_x)D^{-1} + 4u_xD^{-1}u, \\
 \mathcal{R}_1^0 &= -2(5vD^2 + 4v_xD + v_{2x} + 8uv + 4u_xD^{-1}v), \\
 \mathcal{R}_2^0 &= 5D^2 + 8u + 4u_xD^{-1}, \\
 \mathcal{R}_0^1 &= -5u_xD - 6v_{2x} - 2(v_{3x} + 6v_xu)D^{-1} + 4v_xD^{-1}u, \\
 \mathcal{R}_1^1 &= -D^4 - 8uD^2 - 4u_xD + 8(8w - 2v^2) - 8v_xD^{-1}v - 8D^{-1}w_x, \\
 \mathcal{R}_2^1 &= 4(v_xD^{-1} + 2D^{-1}v_x), \\
 \mathcal{R}_0^2 &= -5w_xD - 6w_{2x} - 2(v_{3x} + 6w_xu)D^{-1} + 4w_xD^{-1}u. \\
 \mathcal{R}_1^2 &= -16vD^{-1}w_x - 8w_xD^{-1}v, \\
 \mathcal{R}_2^2 &= -D^4 - 8uD^2 - 4u_xD + 16(w - v^2) + 4w_xD^{-1} + 16vD^{-1}v_x. \tag{B17}
 \end{aligned}$$

APPENDIX C: EXAMPLES TO SEC. III

1. Non-Abelian Schrödinger equation

This is the system given by

$$\begin{aligned}
 u_t &= -\frac{1}{2}u_{xx} + uvu, \\
 v_t &= \frac{1}{2}v_{xx} + vu v,
 \end{aligned} \tag{C1}$$

where u and v belong to \mathcal{K} (see Example 6 for the notations). The Lax operator of (C1) is given by

$$L = D + \begin{pmatrix} \mathbf{1} & 0 \\ 0 & -\mathbf{1} \end{pmatrix} \lambda + \begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix}. \tag{C2}$$

The corresponding formula (54) reduces to

$$\begin{pmatrix} 0 & u_{n+1} \\ v_{n+1} & 0 \end{pmatrix} = \lambda \begin{pmatrix} 0 & u_n \\ v_n & 0 \end{pmatrix} + [R_n, L], \tag{C3}$$

where

$$R_n = \begin{pmatrix} a_n & b_n \\ c_n & -a_n \end{pmatrix}. \tag{C4}$$

The formula (C3) gives us both a_n, b_n, c_n and the recursion operator \mathcal{R} . They are given by

$$a_n = \frac{1}{2}D^{-1}(u_n v + u v_n), \quad b_n = \frac{1}{2}u_n, \quad c_n = -\frac{1}{2}u_n, \tag{C5}$$

$$\mathcal{R} = \frac{1}{2} \begin{pmatrix} -D + R_u D^{-1} R_v + L_u D^{-1} L_v & R_u D^{-1} L_u + L_u D^{-1} R_u \\ -L_v D^{-1} R_v - R_v D^{-1} L_v & D - R_v D^{-1} R_u - L_v D^{-1} L_u \end{pmatrix}. \tag{C6}$$

2. Non-Abelian modified KdV equation

The standard non-Abelian modified KdV equation is given by

$$u_t = \frac{1}{4}u_{xxx} - \frac{3}{4}u_x u^2 - \frac{3}{4}u^2 u_x. \tag{C7}$$

The Lax representation of this equation is given

$$L = D + \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \lambda + \begin{pmatrix} u & 0 \\ 0 & -u \end{pmatrix}. \tag{C8}$$

The recursion operator \mathcal{R} can be found from (60) and (61). In our case the automorphism σ is the same as in Example 6, and formulas (60) and (61) give us

$$\begin{pmatrix} 0 & u_{n+1} \\ v_{n+1} & 0 \end{pmatrix} = \lambda^2 \begin{pmatrix} 0 & u_n \\ v_n & 0 \end{pmatrix} + [R_n, L], \tag{C9}$$

where

$$R_n = \begin{pmatrix} 0 & a_n \\ b_n & 0 \end{pmatrix} \lambda + \begin{pmatrix} c_n & 0 \\ 0 & d_n \end{pmatrix}. \tag{C10}$$

Using (C9) we find a_n, b_n, c_n, d_n from the following:

$$\begin{aligned} b_n - a_n &= u_n, & -a_{n,x} - a_n u - u a_n + c_n - d_n &= 0, \\ -b_{n,x} + b_n u + u b_n + d_n - c_n &= 0, & d_{n,x} + c_{n,x} &= [c_n - d_n, u], \\ u_{n+1} &= d_{n,x} + [d_n, u]. \end{aligned}$$

The resulting recursion operator is given by

$$\mathcal{R} = \frac{1}{4}(D - \text{ad}_u \cdot D^{-1} \cdot \text{ad}_u)(D - (L_u + R_u)D^{-1}(L_u + R_u)). \tag{C11}$$

¹P. J. Olver, *Applications of Lie Groups to Differential Equations*, Graduate Texts in Mathematics, 2nd ed. (Springer-Verlag, New York, 1993), Vol. 107.
²P. J. Olver, J. Math. Phys. **18**, 1212 (1977).
³M. Gürses and A. Karasu, J. Math. Phys. **36**, 3485 (1995).
⁴M. Gürses and A. Karasu, Phys. Lett. A **214**, 21 (1996).
⁵M. Gürses and A. Karasu, J. Math. Phys. **39**, 2103 (1998).
⁶M. Gürses and A. Karasu, Phys. Lett. A **251**, 247 (1999).
⁷S. I. Svinolupov, Theor. Math. Phys. **87**, 391 (1991).
⁸J. Krasil'shchik, Contemp. Math. **219**, 121 (1998).
⁹A. S. Fokas and R. L. Anderson, J. Math. Phys. **23**, 1066 (1982).
¹⁰A. S. Fokas, Stud. Appl. Math. **77**, 253 (1987).
¹¹A. P. Fordy and J. Gibbons, J. Math. Phys. **22**, 1170 (1980).
¹²P. M. Santini and A. S. Fokas, Commun. Math. Phys. **115**, 375 (1988).
¹³A. S. Fokas and P. M. Santini, Commun. Math. Phys. **116**, 449 (1988).
¹⁴W. Symes, J. Math. Phys. **20**, 721 (1979).
¹⁵M. Adler, Invent. Math. **50**, 219 (1979).
¹⁶I. Ya. Dorfman and A. S. Fokas, J. Math. Phys. **33**, 2504 (1992).
¹⁷A. S. Fokas and I. M. Gel'fand, in *Important Developments in Soliton Theory*, Springer Series in Nonlinear Dynamics, edited by A. S. Fokas and V. E. Zakharov (Springer-Verlag, Berlin, 1993), pp. 259–282.
¹⁸M. Antonowicz and A. P. Fordy, in *Nonlinear Evolution Equations and Dynamical Systems (NEEDS'87)*, edited by J. Leon (World Scientific, Singapore, 1988), pp. 145–160.
¹⁹M. Antonowicz and A. P. Fordy, in *Soliton Theory: A Survey of Results*, edited by A. P. Fordy (Manchester University Press, Manchester, England, 1990). See also the related references therein.
²⁰I. M. Gel'fand and L. A. Dikii, Funct. Anal. Appl. **10**, 13 (1976).
²¹F. Magri, J. Math. Phys. **19**, 1156 (1978).
²²V. V. Sokolov, Sov. Math. Dokl. **30**, 44 (1984).

- ²³Here we note that in 1980 that Shabat and Sokolov independently found the recursion operator for the Sawada–Kotera equation. This result was published in Ref. 24. In Ref. 22, Sokolov found the recursion operator for the Krichever–Novikov equation.
- ²⁴N. H. Ibragimov, *Transformation Groups Applied to Mathematical Physics* (Reidel, Boston, 1985).
- ²⁵V. G. Drinfeld and V. V. Sokolov, *J. Sov. Math.* **30**, 1975 (1985).
- ²⁶V. G. Drinfeld and V. V. Sokolov, *Proc. Sobolev Sem. Novosibirsk* **2**, 5 (1981) (in Russian).
- ²⁷I. Z. Golubchik and V. V. Sokolov, *Theor. Math. Phys.* **112**, 1097 (1997).
- ²⁸K. Sawada and T. Kotera, *Prog. Theor. Phys.* **51**, 1355 (1974).
- ²⁹J. Springael, X. B. Hu, and I. Loris, *J. Phys. Soc. Jpn.* **65**, 1222 (1996).
- ³⁰J. Satsuma and R. Hirota, *J. Phys. Soc. Jpn.* **51**, 3390 (1982).
- ³¹G. Wilson, *Phys. Lett. A* **89**, 332 (1982).