

RESEARCH

Open Access



# On continuity of the fractional derivative of the time-fractional semilinear pseudo-parabolic systems

Erdal Karapinar<sup>1,2,3</sup>, Ho Duy Binh<sup>4,5,1</sup>, Nguyen Hoang Luc<sup>1</sup> and Nguyen Huu Can<sup>6\*</sup> 

\*Correspondence:  
[nguyenhuucan@tdtu.edu.vn](mailto:nguyenhuucan@tdtu.edu.vn)  
<sup>1</sup>Applied Analysis Research Group,  
Faculty of Mathematics and  
Statistics, Ton Duc Thang University,  
Ho Chi Minh City, Vietnam  
Full list of author information is  
available at the end of the article

## Abstract

In this work, we study an initial value problem for a system of nonlinear parabolic pseudo equations with Caputo fractional derivative. Here, we discuss the continuity which is related to a fractional order derivative. To overcome some of the difficulties of this problem, we need to evaluate the relevant quantities of the Mittag-Leffler function by constants independent of the derivative order. Moreover, we present an example to illustrate the theory.

**MSC:** 35K55; 35K70; 35K92; 47A52; 47J06

**Keywords:** Initial value problem; Caputo derivative; Nonlinear fractional pseudo-parabolic equation systems; Regularity

## 1 Introduction

In recent decades, fractional calculus has had many applications in various fields such as mechanic, biological, physical science, and applied science. This topic has increasingly asserted its role in the field of applied mathematics, especially the subjects are investigating the properties of the concept of derivative. Specifically, the situations with integer order in PDEs are not working well. Therefore, PDEs with fractional derivatives are a generalization equation with integer-order partial derivatives and a strong theoretical and practical interest. There have been many authors researching this field, for example, [1–15]. According to our search results, the extended results for a coupled nonlinear fractional pseudo-parabolic equation are still limited. This is the great impetus that motivated us to study the following model. In this paper, we extend the coupled system and consider the

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

following coupled nonlinear fractional pseudo-parabolic equation:

$$\begin{cases} \partial_t^\alpha(u(x,t) - a\Delta u(x,t)) - \Delta u(x,t) = \mathcal{F}(u,v), & (x,t) \in \Omega \times (0,T), \\ \partial_t^\alpha(v(x,t) - a\Delta v(x,t)) - \Delta v(x,t) = \mathcal{G}(u,v), & (x,t) \in \Omega \times (0,T), \\ u(x,t) = v(x,t) = 0, & x \in \partial\Omega, t \in (0,T], \\ u_t(x,0) = v_t(x,0) = 0, & x \in \partial\Omega, \\ u(x,0) = f(x), & x \in \Omega, \\ v(x,0) = g(x), & x \in \Omega, \end{cases} \quad (1.1)$$

where  $T, a$  are positive numbers and  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 1$ , is an open bounded domain with a smooth boundary  $\partial\Omega$ . Note that  $\mathbb{L}^2(\Omega)$ ,  $\mathbb{H}_0^1(\Omega)$ ,  $\mathbb{H}^2(\Omega)$  denote the usual Sobolev spaces. The symmetric uniform elliptic operator  $\Delta : \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)$  is defined by

$$\Delta u(x) = - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left( \Delta_{ij}(x) \frac{\partial}{\partial x_j} u(x) \right) + l(x)u(x,t), \quad x \in \overline{\Omega}.$$

With assumption  $l(x) \in C(\overline{\Omega}, [0, \infty))$ ,  $\Delta_{ij} \in C^1(\overline{\Omega})$ ,  $\Delta_{ij} = \Delta_{ji}$ ,  $1 \leq i, j \leq n$ , and there exists a positive constant  $\tilde{\Delta} > 0$  for  $x \in \overline{\Omega}$ ,  $z = (z_1, z_2, \dots, z_n) \subset \mathbb{R}^n$ , such that

$$\tilde{\Delta} \sum_{i=1}^n z_i^2 \leq \sum_{1 \leq i, j \leq n} z_i \Delta_{ij}(x) z_j,$$

see e.g. [16]. The constant  $\alpha \in (1, 2)$  is the fractional order and  $\partial_t^\alpha$  denotes the left-sided Caputo fractional derivative involving  $t$ , which is defined by

$$\partial_t^\alpha u(x,t) := \frac{1}{\Gamma(2-\alpha)} \int_0^t (t-\eta)^{1-\alpha} \frac{\partial u^2}{\partial \eta}(x, \eta) d\eta,$$

where  $\Gamma$  is the gamma function.

If  $\alpha = 2$ , then  $\partial_t^2$  is interpreted as a derivative of normal time. The second equation in the (1.1) called the fractional pseudo-parabolic equation has many practical applications, for example, the permeability of a homogeneous liquid through a cracked rock [17], one-way propagation of nonlinear dispersion long waves [18–20], and populations of [21] populations. Let  $\Omega$  be an open and bounded domain  $\mathbb{R}^n$  with the boundary  $\partial\Omega$ . The functions  $\mathcal{F}, \mathcal{G}, f, g$  satisfy some assumptions to be specified later.

In practice, many problems with space-time fraction equations are based on fractional parameters, that is, the order of fractions. However, these fractional parameters were not known during the modeling process. Therefore, the continuity of the solution on these parameters is very important for modeling purposes. To the best of our knowledge, there have been no results investigating continuity related to the fractional order of a system of pseudo-nonlinear parabolic equations.

This article is organized as follows. Part 2 provides some basic and preliminary definitions. In Part 3, we give the formula of a mild solution and some lemmas that may be related to the next section. In Part 4, we apply the results of Part 3 to establish the existence, uniqueness, and continuity of the solution to problem (1.1) in fractional order. Finally, we give an example to test the theory.

## 2 Preliminaries

### 2.1 Stability on the parameters of the Mittag-Leffler function

Consider the Mittag-Leffler function, which is defined by

$$E_{\alpha,\beta}(\xi) = \sum_{n=1}^{\infty} \frac{\xi^n}{\Gamma(n\alpha + \beta)} \quad \text{for } \xi \in \mathbb{C}, \alpha > 0, \text{ and } \beta \in \mathbb{R}.$$

We call to mind the following lemmas (see for example [3, 22, 23]), which will be useful for the main analysis of Sect. 3 and Sect. 4.

**Lemma 2.1** *If  $1 < \alpha < 2$ , then, for all  $\xi > 0$ , where  $M$  is a positive constant depending only on  $\alpha$ ,*

$$|E_{\alpha,1}(-\xi)| \leq M, \quad |E_{\alpha,\alpha}(-\xi)| \leq M.$$

Now, we have the following lemmas.

**Lemma 2.2** *Let  $\lambda > 0$  and  $1 < \alpha < 2$ . Then, for all  $\xi > 0$ , the following identities hold:*

$$\begin{aligned} \partial_{\xi} E_{\alpha,1}(-\lambda \xi^{\alpha}) &= -\lambda \xi^{\alpha-1} E_{\alpha,\alpha}(-\lambda \xi^{\alpha}), \\ \partial_{\xi} (\xi^{\alpha-1} E_{\alpha,\alpha}(-\lambda \xi^{\alpha})) &= -\xi^{\alpha-2} E_{\alpha,\alpha-1}(-\lambda \xi^{\alpha}). \end{aligned} \quad (2.1)$$

*Proof* Apply Lemma 2.2 in [24]. □

**Lemma 2.3** (see [25]) *Let  $1 < \alpha < 2$ . If  $T$  is large enough, then*

$$E_{\alpha,1}(-\lambda_j T^{\alpha}) \neq 0 \quad (2.2)$$

for all  $j \in \mathbb{N}$ , then there exist two constants  $m_{\alpha}$  and  $M_{\alpha}$  such that

$$\frac{m_{\alpha}}{1 + \lambda_j T^{\alpha}} \leq |E_{\alpha,1}(-\lambda_j T^{\alpha})| \leq \frac{M_{\alpha}}{1 + \lambda_j T^{\alpha}}. \quad (2.3)$$

From Lemma 2.3 in [26], we have the following lemmas.

**Lemma 2.4** *Let  $1 < \alpha_1 < \alpha_2 < 2$  and  $\alpha \in (\alpha_1, \alpha_2)$ . There exist two positive constants  $M_1, M_2$ , and  $M_3$  which just rely upon  $\alpha_1, \alpha_2$  such that, for any  $\xi \geq 0$ , we get*

$$\begin{aligned} \frac{M_1(\alpha_1, \alpha_2)}{1 + \xi} &\leq |E_{\alpha,1}(-\xi)| \leq \frac{M_2(\alpha_1, \alpha_2)}{1 + \xi} \\ \text{if } \xi \text{ is large enough and } |E_{\alpha,\alpha}(-\xi)| &\leq \frac{M_3(\alpha_1, \alpha_2)}{1 + \xi}. \end{aligned} \quad (2.4)$$

**Lemma 2.5** *Let  $0 < \alpha_1 < \alpha < \alpha' < \alpha_2$  and  $0 < \xi \leq T$ . For any  $\epsilon > 0$  independent of  $\alpha$ , there always exists  $M_{\epsilon}$  such that*

$$|\xi^{\alpha} - \xi^{\alpha'}| \leq \max(T^{\alpha_2+2\epsilon}, 1) M_{\epsilon} (\alpha' - \alpha)^{\epsilon} \xi^{\alpha-\epsilon}. \quad (2.5)$$

*Proof* See Lemma 3.2 in [24].  $\square$

Using Lemmas 3.3 and 3.4, Section 3 in [24], we have the following lemmas.

**Lemma 2.6** Assume that  $1 < \alpha_1 < \alpha < \alpha' < \alpha_2 < 2$  and  $\epsilon > 0$ . Then there exists a positive constant  $A(\alpha_1, \alpha_2, \epsilon, v_0, T)$

$$\begin{aligned} & |E_{\alpha,1}(-\lambda_j \xi^\alpha) - E_{\alpha',1}(-\lambda_j \xi^{\alpha'})| \\ & \leq A(\alpha_1, \alpha_2, \epsilon, v_0, T) \lambda_j^{v_0-1} \xi^{-\alpha_2(1-v_0)-\epsilon} [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \end{aligned} \quad (2.6)$$

for any  $0 \leq v_0 \leq 1$  and  $0 < \xi \leq T$ .

*Proof* See Lemma 3.3 for stability on parameters of the Mittag-Leffler function in [24].  $\square$

**Lemma 2.7** Assume that  $1 < \alpha_1 < \alpha < \alpha' < \alpha_2 < 2$ . For any  $0 \leq v_0 \leq 1$  and  $\epsilon > 0$ , there exists a positive constant  $B(\alpha_1, \alpha_2, \epsilon, v_0, T)$

$$\begin{aligned} & |\xi^{\alpha-1} E_{\alpha,\alpha}(-\lambda_j t^\alpha) - \xi^{\alpha'-1} E_{\alpha',\alpha'}(-\lambda_j \xi^{\alpha'})| \\ & \leq B(\alpha_1, \alpha_2, \epsilon, v_0, T) \lambda_j^{v_0-1} \xi^{\alpha_1 v_0 - \epsilon - 1} [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]. \end{aligned} \quad (2.7)$$

*Proof* Use Lemma 3.4 from Section 3 in [24].  $\square$

## 2.2 Some Sobolev spaces

In this section, we present some appropriate Sobolev space. Let the operator  $\Delta$  be considered on  $\mathbb{L}^2(\Omega)$  with domain  $\mathbb{H}^2(\Omega) \cap \mathbb{H}_0^1(\Omega)$ . Then the spectrum of  $\Delta$  is a non-diminishing arrangement of positive real numbers  $\{\lambda_j\}_{j \geq 1}$  which satisfy that  $\lim_{j \rightarrow \infty} \lambda_j = \infty$ . Let us denote by  $\{\varphi_j\}_{j \geq 1}$  in  $\mathbb{H}^2(\Omega) \cap \mathbb{H}_0^1(\Omega)$  the set of orthonormal eigenfunctions of  $\Delta$ , which means that  $\Delta \varphi_j = \lambda_j \varphi_j$ . The sequence forms an orthonormal basis of  $\mathbb{L}^2(\Omega)$ , see e.g. [27]. For all  $\gamma \geq 0$ , the operator  $\Delta^\gamma$  has the following representation:

$$\Delta^\gamma v := \sum_{j=1}^{\infty} \langle v, \varphi_j \rangle \lambda_j^\gamma \varphi_j, \quad v \in \mathbb{D}(\Delta^\gamma) = \left\{ v \in \mathbb{L}^2(\Omega) : \sum_{j=1}^{\infty} |\langle v, \varphi_j \rangle|^2 \lambda_j^{2\gamma} < \infty \right\}.$$

The domain  $\mathbb{D}(\Delta^\gamma)$  is Banach spaces equipped with the norm

$$\|v\|_{\mathbb{D}(\Delta^\gamma)}^2 := \sum_{j=1}^{\infty} \lambda_j^{2\gamma} |\langle v, \varphi_j \rangle|^2. \quad (2.8)$$

If  $\gamma = 1$ , we have  $\mathbb{D}(\Delta^1) = \mathbb{H}^2(\Omega)$ .

For a given number  $\gamma \geq 0$ , the Hilbert space

$$\mathcal{H}^\gamma(\Omega) = \left\{ v \in \mathbb{L}^2(\Omega) : \sum_{j=1}^{\infty} |\langle v, \varphi_j \rangle|^2 \lambda_j^{2\gamma} < \infty \right\} \quad (2.9)$$

is endowed with the norm as follows:

$$\|v\|_{\mathcal{H}^\gamma}^2 := \sum_{j=1}^{\infty} \lambda_j^{2\gamma} |\langle v, \varphi_j \rangle|^2. \quad (2.10)$$

If  $\gamma = 0$ , then  $\mathcal{H}^0(\Omega) = \mathbb{L}^2(\Omega)$ . We identified a norm for  $w(u, v) \in \mathcal{H}^\gamma(\Omega) = \mathcal{H}^\gamma(\Omega) \times \mathcal{H}^\gamma(\Omega)$  as follows:

$$\|w\|_{\mathcal{H}^\gamma(\Omega)} = \sqrt{\|u\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v\|_{\mathcal{H}^\gamma(\Omega)}^2}. \quad (2.11)$$

Let us denote by  $\mathbb{C}((0, T]; \mathcal{H}^\gamma(\Omega))$  a space of all continuous functions with the map  $(0, T] \rightarrow \mathcal{H}^\gamma(\Omega)$ . For a given number  $0 < \beta < 1$ , we define by  $\mathbb{C}^\beta((0, T]; \mathcal{H}^\gamma(\Omega))$  such that

$$\sup_{0 < t \leq T} t^\beta \|f(t)\|_{\mathcal{H}^\gamma(\Omega)} < \infty; \quad f \in \mathbb{C}((0, T]; \mathcal{H}^\gamma(\Omega)),$$

in which (see [26])

$$\|f\|_{\mathbb{C}^\beta((0, T]; \mathcal{H}^\gamma(\Omega))} := \sup_{0 < t \leq T} t^\beta \|f\|_{\mathcal{H}^\gamma(\Omega)}.$$

The product space  $\mathcal{C}^\beta(0, T, \mathcal{H}^\gamma(\Omega)) = \mathbb{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega)) \times \mathbb{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega))$  is also a Banach space endowed with the norm

$$\|w\|_{\mathcal{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega))} = \sqrt{\|u\|_{\mathbb{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega))}^2 + \|v\|_{\mathbb{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega))}^2}$$

for  $w = (u, v) \in \mathcal{C}^\beta((0, T], \mathcal{H}^\gamma(\Omega))$ . For a given positive real number  $p$ ,  $\mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$  is a Banach space with the norm

$$\|f\|_{\mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))} := \text{ess sup } e^{-pt} \|f\|_{\mathcal{H}^\gamma(\Omega)}. \quad (2.12)$$

The product space  $\mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega)) = \mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega)) \times \mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$  is a Banach space, and we also identified a norm for  $w = (u, v) \in \mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$  as follows:

$$\|w\|_{\mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))} = \sqrt{\|u\|_{\mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))}^2 + \|v\|_{\mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))}^2}.$$

### 3 Relevant notations and a representation of solution

In perception of spectral decomposition

$$u(x, t) = \sum_{j=1}^{\infty} \langle u(\cdot, t), \varphi_j(\cdot) \rangle \varphi_j(x) \quad \text{and} \quad v(x, t) = \sum_{j=1}^{\infty} \langle v(\cdot, t), \varphi_j(\cdot) \rangle \varphi_j(x). \quad (3.1)$$

We can transform the first two equations of (1.1) into

$$\begin{cases} \langle \partial_t^\alpha(u(\cdot, t), \varphi_j) + a \langle \partial_t^\alpha \Delta u(\cdot, t), \varphi_j \rangle + \langle \Delta u(\cdot, t), \varphi_j \rangle = \langle \mathcal{F}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle, \\ \langle \partial_t^\alpha(v(\cdot, t), \varphi_j) + a \langle \partial_t^\alpha \Delta v(\cdot, t), \varphi_j \rangle + \langle \Delta v(\cdot, t), \varphi_j \rangle = \langle \mathcal{G}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle. \end{cases} \quad (3.2)$$

Using the formula  $\Delta \varphi_j = \lambda_j \varphi_j$ , we obtain

$$\begin{cases} (1 + a\lambda_j) \langle \partial_t^\alpha(u(\cdot, t), \varphi_j) + \lambda_j \langle u(\cdot, t), \varphi_j \rangle = \langle \mathcal{F}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle, \\ (1 + a\lambda_j) \langle \partial_t^\alpha(v(\cdot, t), \varphi_j) + \lambda_j \langle v(\cdot, t), \varphi_j \rangle = \langle \mathcal{G}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle. \end{cases} \quad (3.3)$$

The theory of fractional ordinary differential equations (see [3, 22, 23]) gives a unique function  $u_j, v_j$  as follows:

$$\begin{cases} u_j(t) = E_{\alpha,1}\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right)f_j \\ \quad + \frac{1}{1+a\lambda_j} \int_0^t [(t-s)^{\alpha-1} E_{\alpha,\alpha}\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right)\mathcal{F}_j(s)] ds, \\ v_j(t) = E_{\alpha,1}\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right)g_j \\ \quad + \frac{1}{1+a\lambda_j} \int_0^t [(t-s)^{\alpha-1} E_{\alpha,\alpha}\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right)\mathcal{G}_j(s)] ds. \end{cases} \quad (3.4)$$

Here, we denote  $f_j := \langle f, \varphi_j \rangle$ ,  $g_j := \langle g, \varphi_j \rangle$ ,  $\mathcal{F}_j := \langle \mathcal{F}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle$  and  $\mathcal{G}_j := \langle \mathcal{G}(u(\cdot, t), v(\cdot, t)), \varphi_j \rangle$ . Hence solution (1.1) can be described as by Fourier series (3.1) and then given by

$$\begin{cases} u(x, t) = \sum_{j=1}^{\infty} E_{\alpha,1}\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right)f_j \varphi_j \\ \quad + \sum_{j=1}^{\infty} \frac{1}{1+a\lambda_j} [\int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right)\mathcal{F}_j(s) ds] \varphi_j, \\ v(x, t) = \sum_{j=1}^{\infty} E_{\alpha,1}\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right)g_j \varphi_j \\ \quad + \sum_{j=1}^{\infty} \frac{1}{1+a\lambda_j} [\int_0^t (t-s)^{\alpha-1} E_{\alpha,\alpha}\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right)\mathcal{G}_j(s) ds] \varphi_j. \end{cases} \quad (3.5)$$

It is obvious to see that the mild solution (1.1) is given by

$$\begin{cases} u_\alpha(\cdot, t) = \mathcal{P}_{\Delta,\alpha}(t)f + \int_0^t \mathcal{Q}_{\Delta,\alpha}(t-s)\mathcal{F}(u, v)(\cdot, s) ds, \\ v_\alpha(\cdot, t) = \mathcal{P}_{\Delta,\alpha}(t)g + \int_0^t \mathcal{Q}_{\Delta,\alpha}(t-s)\mathcal{G}(u, v)(\cdot, s) ds, \end{cases} \quad (3.6)$$

where

$$\begin{aligned} \mathcal{P}_{\Delta,\alpha}(t)h &:= \sum_{j=1}^{\infty} E_{\alpha,1}\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right) \langle h, \varphi_j \rangle \varphi_j, \\ \mathcal{Q}_{\Delta,\alpha}(t-s)h &:= \sum_{j=1}^{\infty} \frac{1}{1+a\lambda_j} (t-s)^{\alpha-1} E_{\alpha,\alpha}\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right) \langle h, \varphi_j \rangle \varphi_j. \end{aligned}$$

Therefore, with  $1 < \alpha < \alpha' < 2$ , we also get

$$\begin{cases} u_{\alpha'}(\cdot, t) = \mathcal{P}_{\Delta,\alpha'}(t)f + \int_0^t \mathcal{Q}_{\Delta,\alpha'}(t-s)\mathcal{F}(u, v)(\cdot, s) ds, \\ v_{\alpha'}(\cdot, t) = \mathcal{P}_{\Delta,\alpha'}(t)g + \int_0^t \mathcal{Q}_{\Delta,\alpha'}(t-s)\mathcal{G}(u, v)(\cdot, s) ds. \end{cases} \quad (3.7)$$

Next, we give several lemmas related to Sect. 4 as follows.

**Lemma 3.1** *Let  $1 < \alpha_1 < \alpha < \alpha_2 < 2$ ,  $\gamma \geq 0$ , and  $w \in \mathcal{H}^\gamma(\Omega)$ . The following inequalities hold:*

$$\|\mathcal{P}_{\Delta,\alpha}(t)w\|_{\mathcal{H}^\gamma(\Omega)} \leq \overline{M}_2(\alpha_1, \alpha_2, a, \mu_0) t^{-\alpha_1 \mu_0} \|w\|_{\mathcal{H}^\gamma(\Omega)}, \quad (3.8)$$

$$\|\mathcal{Q}_\alpha(t-s)w\|_{\mathcal{H}^\gamma(\Omega)} \leq \overline{M}_3(\alpha_1, \alpha_2, a, \mu_0) (t-s)^{\alpha_1 - 1 - \alpha_1 \mu_0} \|w\|_{\mathcal{H}^\gamma(\Omega)}, \quad (3.9)$$

where  $\mu_0$  is a positive number satisfying  $0 < \mu_0 < 1$ .

*Proof* Using Lemma 2.4, we get

$$\begin{aligned}
\|\mathcal{P}_{\Delta,\alpha}(t)\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}^2 &= \sum_{j=1}^{\infty} \lambda_j^{2\gamma} E_{\alpha,1}^2\left(\frac{-\lambda_j t^\alpha}{1+a\lambda_j}\right) |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq \sum_{j=1}^{\infty} \lambda_j^{2\gamma} \left(\frac{M_2(\alpha_1, \alpha_2)}{1+\frac{\lambda_j t^\alpha}{1+a\lambda_j}}\right)^2 |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq \sum_{j=1}^{\infty} \lambda_j^{2\gamma} \frac{M_2^2(\alpha_1, \alpha_2)}{(1+\frac{\lambda_j t^\alpha}{1+a\lambda_j})^{\mu_0}} |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq \sum_{j=1}^{\infty} \lambda_j^{2\gamma} M_2^2(\alpha_1, \alpha_2) \left(\frac{\lambda_j t^\alpha}{1+a\lambda_j}\right)^{-2\mu_0} |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq M_2^2(\alpha_1, \alpha_2) t^{-2\alpha\mu_0} (\alpha + \lambda_1^{-1})^{2\mu_0} \sum_{j=1}^{\infty} \lambda_j^{2\gamma} |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq M_2^2(\alpha_1, \alpha_2) t^{-2\alpha_1\mu_0} (\lambda_1^{-1} + a)^{2\mu_0} \|\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}^2.
\end{aligned}$$

Therefore, with  $\overline{M}_2(\alpha_1, \alpha_2, a, \mu_0) := M_2(\alpha_1, \alpha_2)(\lambda_1^{-1} + a)^{\mu_0}$ , we have the following estimate:

$$\|\mathcal{P}_{\Delta,\alpha}(t)\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)} \leq \overline{M}_2(\alpha_1, \alpha_2, a, \mu_0) t^{-\alpha_1\mu_0} \|\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}. \quad (3.10)$$

Likewise, utilizing Lemma 2.4, we can get the following estimation:

$$\begin{aligned}
\|\mathcal{Q}_{\Delta,\alpha}(t-s)\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}^2 &= \sum_{j=1}^{\infty} \frac{\lambda_j^{2\gamma}}{(1+a\lambda_j)^2} (t-s)^{2(\alpha-1)} E_{\alpha,\alpha}^2\left(\frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j}\right) |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq \sum_{j=1}^{\infty} \frac{\lambda_j^{2\gamma}}{(1+a\lambda_j)^2} (t-s)^{2(\alpha-1)} \left(\frac{M_3(\alpha_1, \alpha_2)}{1+\frac{\lambda_j(t-s)^\alpha}{1+a\lambda_j}}\right)^2 |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq (t-s)^{2(\alpha-1-\alpha\mu_0)} M_3^2(\alpha_1, \alpha_2) \sum_{j=1}^{\infty} \lambda_j^{2\gamma} \left(\frac{1}{\lambda_j} + a\right)^{2(\mu_0-1)} \lambda_j^{-2} |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq (t-s)^{2(\alpha-1-\alpha\mu_0)} M_3^2(\alpha_1, \alpha_2) (\lambda_1^{-1} + a)^{2(\mu_0-1)} \lambda_1^{-2} \sum_{j=1}^{\infty} \lambda_j^{2\gamma} |\langle \mathbf{w}, \varphi_j \rangle|^2 \\
&\leq (t-s)^{2(\alpha_1-1-\alpha_1\mu_0)} M_3^2(\alpha_1, \alpha_2) (\lambda_1^{-1} + a)^{2(\mu_0-1)} \lambda_1^{-2} \|\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}^2.
\end{aligned} \quad (3.11)$$

Therefore, we deduce

$$\|\mathcal{Q}_\alpha(t-s)\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)} \leq \overline{M}_3(\alpha_1, \alpha_2, a, \mu_0) (t-s)^{\alpha_1-1-\alpha_1\mu_0} \|\mathbf{w}\|_{\mathcal{H}^\gamma(\Omega)}, \quad (3.12)$$

where  $\overline{M}_3(\alpha_1, \alpha_2, a, \mu_0) := M_3(\alpha_1, \alpha_2)(\lambda_1^{-1} + a)^{\mu_0-1} \lambda_1^{-1}$ .

Thus, we complete the proof of Lemma 3.1.  $\square$

**Lemma 3.2** Let  $1 < \alpha_1 < \alpha < \alpha' < \alpha_2 < 2$ ,  $\gamma \geq 0$  with  $0 \leq v_0 \leq 1$  and  $w \in \mathcal{H}^\gamma(\Omega)$ . The following inequalities hold:

$$\begin{aligned} & \|[\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]w\|_{\mathcal{H}^\gamma(\Omega)} \\ & \leq \bar{A}(\alpha_1, \alpha_2, \epsilon, v_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]t^{-\alpha_2(1-v_0)-\epsilon}\|w\|_{\mathcal{H}^\gamma(\Omega)}. \end{aligned} \quad (3.13)$$

Besides, we have

$$\begin{aligned} & \|[\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \alpha}(t-s)]w\|_{\mathcal{H}^\gamma(\Omega)} \\ & \leq \bar{B}(\alpha_1, \alpha_2, \epsilon, v_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)](t-s)^{\alpha_1 v_0 - \epsilon - 1}\|w\|_{\mathcal{H}^\gamma(\Omega)}. \end{aligned}$$

*Proof* We get

$$\begin{aligned} & \|[\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]w\|_{\mathbb{H}^\gamma(\Omega)}^2 \\ & = \sum_{j=1}^{\infty} \lambda_j^{2\gamma} \left[ E_{\alpha', 1} \left( \frac{-\lambda_j t^{\alpha'}}{1+a\lambda_j} \right) - E_{\alpha, 1} \left( \frac{-\lambda_j t^\alpha}{1+a\lambda_j} \right) \right]^2 |\langle w, \varphi_j \rangle|^2. \end{aligned}$$

By using Lemma 2.6, we obtain the following estimates:

$$\begin{aligned} & \|[\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]w\|_{\mathcal{H}^\alpha(\Omega)}^2 \\ & \leq \sum_{j=1}^{\infty} \lambda_j^{2\gamma} A^2(\alpha_1, \alpha_2, \epsilon, v_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\ & \quad \times \left( \frac{\lambda_j}{1+a\lambda_j} \right)^{2(v_0-1)} t^{-2\alpha_2(1-v_0)-2\epsilon} |\langle w, \varphi_j \rangle|^2 \\ & \leq A^2(\alpha_1, \alpha_2, \epsilon, v_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\ & \quad \times t^{-2\alpha_2(1-v_0)-2\epsilon} (\lambda_1^{-1} + a)^{2(1-v_0)} \sum_{j=1}^{\infty} \lambda_j^{2\gamma} |\langle w, \varphi_j \rangle|^2 \\ & \leq A^2(\alpha_1, \alpha_2, \epsilon, v_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\ & \quad \times t^{-2\alpha_2(1-v_0)-2\epsilon} (\lambda_1^{-1} + a)^{2(1-v_0)} \|w\|_{\mathcal{H}^\gamma(\Omega)}^2. \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} & \|[\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]w\|_{\mathcal{H}^\gamma(\Omega)} \\ & \leq \bar{A}(\alpha_1, \alpha_2, \epsilon, v_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]t^{-\alpha_2(1-v_0)-\epsilon}\|w\|_{\mathcal{H}^\gamma(\Omega)}, \end{aligned}$$

where  $\bar{A}(\alpha_1, \alpha_2, \epsilon, v_0, a, T) := A(\alpha_1, \alpha_2, \epsilon, v_0, T)(\lambda_1^{-1} + a)^{1-v_0}$ .

Similarly, by applying Lemma 2.7, we also get

$$\begin{aligned} & \|[\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \alpha}(t-s)]w\|_{\mathcal{H}^\gamma(\Omega)}^2 \\ & = \sum_{j=1}^{\infty} \frac{\lambda_j^{2\gamma}}{(1+a\lambda_j)^2} \left[ (t-s)^{\alpha'-1} E_{\alpha', \alpha'} \left( \frac{-\lambda_j(t-s)^{\alpha'}}{1+a\lambda_j} \right) \right] \end{aligned}$$

$$\begin{aligned}
& - (t-s)^{\alpha-1} E_{\alpha,\alpha} \left( \frac{-\lambda_j(t-s)^\alpha}{1+a\lambda_j} \right)^2 |\langle w, \varphi_j \rangle|^2 \\
& \leq \sum_{j=1}^{\infty} B^2(\alpha_1, \alpha_2, \epsilon, v_0, T) \left( \frac{\lambda_j}{1+a\lambda_j} \right)^{2(v_0-1)} (t-s)^{2\alpha_1 v_0 - 2\epsilon - 2} \\
& \quad \times [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \frac{\lambda_j^{2\gamma}}{(1+a\lambda_j)^2} |\langle w, \varphi_j \rangle|^2 \\
& \leq B^2(\alpha_1, \alpha_2, \epsilon, v_0, T) [(\alpha' - \alpha_2)^\epsilon + (\alpha' - \alpha)]^2 \lambda_1^{-2} (t-s)^{2\alpha_1 v_0 - 2\epsilon - 2} \\
& \quad \times \sum_{j=1}^{\infty} \left( \frac{1}{\lambda_j} + a \right)^{-2v_0} \lambda_j^{2\gamma} |\langle w, \varphi_j \rangle|^2 \\
& \leq B^2(\alpha_1, \alpha_2, \epsilon, v_0, T) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \lambda_1^{-2} a^{-2v_0} (t-s)^{2\alpha_1 v_0 - 2\epsilon - 2} \|w\|_{\mathcal{H}^\gamma(\Omega)}^2.
\end{aligned}$$

We denote  $\bar{B}(\alpha_1, \alpha_2, \epsilon, v_0, a, T) := B(\alpha_1, \alpha_2, \epsilon, v_0, T) \lambda_1^{-1} a^{-v_0}$ . Therefore, we obtain

$$\begin{aligned}
& \|[\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \omega}(t-s)]w\|_{\mathcal{H}^\gamma(\Omega)} \\
& \leq \bar{B}(\alpha_1, \alpha_2, \epsilon, v_0, a, T) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] (t-s)^{\alpha_1 v_0 - \epsilon - 1} \|w\|_{\mathcal{H}^\gamma(\Omega)}. \tag{3.14}
\end{aligned}$$

We get all estimates of Lemma 3.2. This completes the proof.  $\square$

#### 4 Stability of the fractional order of problem (1.1)

In this section, we are interested in studying the existence of a mild solution and the continuous dependence of the solution of problem (1.1) with input (the fractional-order  $\alpha, \alpha'$  and the initial condition  $f, g$ ). We assume that  $\mathcal{F}, \mathcal{G}$  satisfy the following assumptions:

(S.1)

$$\|\mathcal{F}(u, v)(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \leq C_1 (1 + \|u(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} + \|v(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}), \tag{4.1}$$

$$\|\mathcal{G}(u, v)(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \leq C_2 (1 + \|u(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} + \|v(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}), \tag{4.2}$$

where  $(u, v) \in \mathcal{H}^\gamma(\Omega) = \mathcal{H}^\gamma(\Omega) \times \mathcal{H}^\gamma(\Omega)$ .

(S.2)

$$\|\mathcal{F}(u_1, v_1)(\cdot, t) - \mathcal{F}(u_2, v_2)(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \leq K_1 (\|u_1 - u_2\|_{\mathcal{H}^\gamma(\Omega)} + \|v_1 - v_2\|_{\mathcal{H}^\gamma(\Omega)}), \tag{4.3}$$

$$\|\mathcal{G}(u_1, v_1)(\cdot, t) - \mathcal{G}(u_2, v_2)(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \leq K_2 (\|u_1 - u_2\|_{\mathcal{H}^\gamma(\Omega)} + \|v_1 - v_2\|_{\mathcal{H}^\gamma(\Omega)}), \tag{4.4}$$

where  $(u_1, v_1) \in \mathcal{H}^\gamma(\Omega), (u_2, v_2) \in \mathcal{H}^\gamma(\Omega)$ .

**Definition 4.1**  $w = (u(\cdot, t), v(\cdot, t)) \in \mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega)) = \mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega)) \times \mathbb{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$  is called a mild solution of problem (1.1) if it satisfies system (3.6).

**Theorem 4.1** Let  $w_0(f, g) \in \mathcal{H}^\gamma(\Omega)$ . Assume that  $1 < \alpha_1 < \alpha < \alpha' < \alpha_2 < 2$  and  $0 < v_0 < 1$ . The nonlinear integral equation (1.1) has a unique solution  $w(u, v) \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$ . Let  $w_\alpha \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$  and  $w_{\alpha'} \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$  be two solutions of (1.1) with

*fractional order  $\alpha$  and  $\alpha'$ , respectively. If there exist numbers  $\mu_0, \epsilon$  satisfying  $0 < \epsilon < \min(\frac{1}{2} - \alpha_2 + \alpha_2 v_0, \alpha_1 v_0 - \frac{1}{2})$  and  $0 < \mu_0 < 1 - \frac{1}{2\alpha_1}$ , then*

$$\begin{aligned} & \| \mathbf{w} \|_{\mathcal{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^\gamma(\Omega))} \\ & \leq \sqrt{\mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \| w_0 \|_{\mathcal{H}^\gamma(\Omega)})} \exp\left(\frac{\mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T)T}{2}\right), \end{aligned} \quad (4.5)$$

and

$$\begin{aligned} & \| \mathbf{w}_{\alpha'} - \mathbf{w}_\alpha \|_{\mathcal{C}^{\alpha_2(1-v_0)+\epsilon}((0, T], \mathcal{H}^\gamma(\Omega))} \\ & \leq \sqrt{\mathcal{M}_{\mu_0, v_0}^{\mathcal{C}_1, \mathcal{C}_2}(\alpha_1, \alpha_2, \epsilon, a, T, \| w_0 \|_{\mathcal{H}^\gamma(\Omega)})} [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \\ & \times \exp\left(\frac{\mathcal{M}_{\mu_0, v_0}^{\mathcal{K}_1, \mathcal{K}_2}(\alpha_1, \alpha_2, \epsilon, a, T)T}{2}\right). \end{aligned} \quad (4.6)$$

*Proof of Theorem 4.1* We divide the proof into three parts.

*Part 1.* The existence and uniqueness of the solution of the nonlinear fractional pseudo-parabolic equation systems (1.1). For  $\mathbf{w} \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$ , we consider the following function  $\mathcal{H}\mathbf{w} := (\mathcal{H}_{\Delta, \alpha} u(\cdot, t), \mathcal{H}_{\Delta, \alpha} v(\cdot, t))$ , where

$$\begin{cases} \mathcal{H}_{\Delta, \alpha} u(\cdot, t) = \mathcal{P}_{\Delta, \alpha}(t)f + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)\mathcal{F}(u, v)(\cdot, s)ds, \\ \mathcal{H}_{\Delta, \alpha} v(\cdot, t) = \mathcal{P}_{\Delta, \alpha}(t)g + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)\mathcal{G}(u, v)(\cdot, s)ds. \end{cases} \quad (4.7)$$

Let  $\mathbf{w}_1(u_1(\cdot, t), v_1(\cdot, t)) \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$ ,  $\mathbf{w}_2(u_2(\cdot, t), v_2(\cdot, t)) \in \mathcal{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))$ , we have

$$\begin{aligned} \mathcal{H}_{\Delta, \alpha} u_1(\cdot, t) - \mathcal{H}_{\Delta, \alpha} u_2(\cdot, t) &= \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)[\mathcal{F}(u_1, v_1)(\cdot, s) - \mathcal{F}(u_2, v_2)(\cdot, s)]ds, \\ \mathcal{H}_{\Delta, \alpha} v_1(\cdot, t) - \mathcal{H}_{\Delta, \alpha} v_2(\cdot, t) &= \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)[\mathcal{G}(u_1, v_1)(\cdot, s) - \mathcal{G}(u_2, v_2)(\cdot, s)]ds. \end{aligned}$$

With  $p > 0$ , we get the following estimate:

$$\begin{aligned} & \| e^{-pt}(\mathcal{H}_{\Delta, \alpha} u_1(\cdot, t) - \mathcal{H}_{\Delta, \alpha} u_2(\cdot, t)) \|_{\mathcal{H}^\gamma(\Omega)} \\ & = \left\| \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)e^{-pt}[\mathcal{F}(u_1, v_1)(\cdot, s) - \mathcal{F}(u_2, v_2)(\cdot, s)]ds \right\|_{\mathcal{H}^\gamma(\Omega)}. \end{aligned}$$

Applying Lemma 3.1, we get the following estimate:

$$\begin{aligned} & \| e^{-pt}(\mathcal{H}_{\Delta, \alpha} u_1(\cdot, t) - \mathcal{H}_{\Delta, \alpha} u_2(\cdot, t)) \|_{\mathcal{H}^\gamma(\Omega)}^2 \\ & \leq \left[ \int_0^t \overline{M}_3(\alpha_1, \alpha_2, a, \mu_0)(t-s)^{\alpha_1-1-\alpha_1\mu_0}e^{-p(t-s)} \right. \\ & \quad \times \left. [e^{-ps}\|\mathcal{F}(u_1, v_1)(\cdot, s) - \mathcal{F}(u_2, v_2)(\cdot, s)\|_{\mathcal{H}^\alpha(\Omega)}]ds \right]^2 \end{aligned}$$

$$\begin{aligned} &\leq \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \left[ \int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0} e^{-p(t-s)} \right. \\ &\quad \times \left. \left[ e^{-ps} \|\mathcal{F}(u_1, v_1)(\cdot, s) - \mathcal{F}(u_2, v_2)(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)} \right] ds \right]^2. \end{aligned}$$

Using (4.3) and the inequality  $(m+n)^2 \leq 2(m^2 + n^2)$ , we obtain

$$\begin{aligned} &\|e^{-pt} (\mathcal{H}_{\Delta, \alpha} u_1(\cdot, t) - \mathcal{H}_{\Delta, \alpha} u_2(\cdot, t))\|_{\mathcal{H}^\gamma(\Omega)}^2 \\ &\leq \mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \left[ \int_0^t e^{-ps} (\|u_1 - u_2\|_{\mathcal{H}^\gamma(\Omega)} + \|v_1 - v_2\|_{\mathcal{H}^\gamma(\Omega)}) \right. \\ &\quad \times \left. (t-s)^{\alpha_1-1-\alpha_1\mu_0} e^{-p(t-s)} ds \right]^2. \end{aligned} \quad (4.8)$$

Using the inequality  $(m+n)^2 \leq 2(m^2 + n^2)$ , we deduce that

$$\begin{aligned} &\|\mathcal{H}_{\Delta, \alpha} u_1 - \mathcal{H}_{\Delta, \alpha} u_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \\ &\leq \mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) (\|u_1 - u_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))} + \|v_1 - v_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))})^2 \\ &\quad \times \left[ \int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0} e^{-p(t-s)} ds \right]^2 \\ &\leq 2\mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) (\|u_1 - u_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 + \|v_1 - v_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2) \\ &\quad \times \left[ \int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0} e^{-p(t-s)} ds \right]^2 \\ &\leq 2\mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathscr{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \\ &\quad \times \left[ \int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0} e^{-p(t-s)} ds \right]^2. \end{aligned}$$

Applying Hölder's inequality with assumption  $\mu_0 < 1 - \frac{1}{2\alpha_1}$ , we obtain

$$\begin{aligned} &\|\mathcal{H}_{\Delta, \alpha} u_1 - \mathcal{H}_{\Delta, \alpha} u_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \\ &\leq 2\mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathscr{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \\ &\quad \times \int_0^t (t-s)^{2\alpha_1-2-2\alpha_1\mu_0} ds \int_0^t e^{-2p(t-s)} ds \\ &\leq 2\mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathscr{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \frac{t^{2\alpha_1-1-2\alpha_1\mu_0}}{2\alpha_1-1-2\alpha_1\mu_0} \frac{1-e^{-2pt}}{2p}. \end{aligned} \quad (4.9)$$

Hence

$$\begin{aligned} &\|\mathcal{H}_{\Delta, \alpha} u_1 - \mathcal{H}_{\Delta, \alpha} u_2\|_{\mathbb{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2 \\ &\leq \frac{2\mathcal{K}_1^2 \overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0)}{2p(2\alpha_1-2\alpha_1\mu_0-1)} T^{2\alpha_1-2\alpha_1\mu_0-1} (1-e^{-2pT}) \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathscr{L}_p^\infty(0, T; \mathcal{H}^\gamma(\Omega))}^2. \end{aligned} \quad (4.10)$$

We can obtain a similar estimate

$$\begin{aligned} & \|\mathcal{H}_{\Delta,\alpha}v_1 - \mathcal{H}_{\Delta,\alpha}v_2\|_{\mathbb{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}^2 \\ & \leq \frac{2\mathcal{K}_2^2\overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0)}{2p(2\alpha_1 - 2\alpha_1\mu_0 - 1)} T^{2\alpha_1 - 2\alpha_1\mu_0 - 1} (1 - e^{-2pT}) \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}^2. \end{aligned} \quad (4.11)$$

From (4.10)–(4.11), we find that

$$\begin{aligned} & \|\mathcal{H}\mathbf{w}_1 - \mathcal{H}\mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))} \\ & = \sqrt{\|\mathcal{H}u_1 - \mathcal{H}u_2\|_{\mathbb{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}^2 + \|\mathcal{H}v_1 - \mathcal{H}v_2\|_{\mathbb{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}^2} \\ & \leq \sqrt{\frac{2(\mathcal{K}_1^2 + \mathcal{K}_2^2)\overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0)}{2p(2\alpha_1 - 2\alpha_1\mu_0 - 1)} T^{2\alpha_1 - \alpha_1\mu_0 - 1} (1 - e^{-2pT})} \\ & \quad \times \|\mathbf{w}_1 - \mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}. \end{aligned} \quad (4.12)$$

If  $\mathbf{w}_2 = 0$ , then for any  $\mathbf{w} \in \mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$

$$\begin{aligned} & \|\mathcal{H}\mathbf{w}\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))} \\ & \leq \|\mathcal{H}\mathbf{w} - \mathcal{H}\mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))} + \|\mathcal{H}\mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))} \\ & \leq \sqrt{\frac{2(\mathcal{K}_1^2 + \mathcal{K}_2^2)\overline{M}_3^2(\alpha_1, \alpha_2, a, \mu_0)}{2p(2\alpha_1 - 2\alpha_1\mu_0 - 1)} T^{2\alpha_1 - \alpha_1\mu_0 - 1} (1 - e^{-2pT})} \|\mathbf{w}\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))} \\ & \quad + \|\mathcal{H}\mathbf{w}_2\|_{\mathcal{L}_p^\infty(0,T,\mathcal{H}^\gamma(\Omega))}. \end{aligned} \quad (4.13)$$

Note that  $\mathcal{H}\mathbf{w}_2 := (\mathcal{H}_{\Delta,\alpha}u_2(\cdot, t), \mathcal{H}_{\Delta,\alpha}v_2(\cdot, t))$ , where  $u_2 = v_2 = 0$  and

$$\begin{cases} \mathcal{H}_{\Delta,\alpha}u_2(\cdot, t) = \mathcal{P}_{\Delta,\alpha}(t)f, \\ \mathcal{H}_{\Delta,\alpha}v_2(\cdot, t) = \mathcal{P}_{\Delta,\alpha}(t)g. \end{cases} \quad (4.14)$$

By applying Lemma 3.1, we can deduce that  $\mathcal{H}\mathbf{w}_2 \in \mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$ . Therefore, we conclude that if any  $\mathbf{w} \in \mathcal{L}_p^\infty(0, T, \mathcal{H}^\gamma(\Omega))$ , then  $\mathcal{H}\mathbf{w}$  is bounded.

*Part 2.* From (3.6) and applying the inequality  $(a+b)^2 \leq 2(a^2 + b^2)$ , we have the following estimate:

$$\|u(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \leq 2\|\mathcal{P}_{\Delta,\alpha}(t)f\|_{\mathcal{H}^\gamma(\Omega)}^2 + 2\left\|\int_0^t \mathcal{Q}_{\Delta,\alpha}(t-s)\mathcal{F}(u, v)(\cdot, s) ds\right\|_{\mathcal{H}^\gamma(\Omega)}^2.$$

Using Lemma 3.1, we get

$$\begin{aligned} \|u(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 & \leq 2\overline{M}_2(\alpha_1, \alpha_2, a, \mu_0)t^{-2\alpha_1\mu_0}\|f\|_{\mathcal{H}^\gamma(\Omega)}^2 \\ & \quad + 2\left(\int_0^t \overline{M}_3(\alpha_1, \alpha_2, a, \mu_0)(t-s)^{\alpha_1-1-\alpha_1\mu_0}\|\mathcal{F}(u, v)(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)} ds\right)^2. \end{aligned}$$

Multiplying both sides by  $t^{2\alpha_1\mu_0}$  and using Hölder's inequality with assumption (4.1), we can find that

$$\begin{aligned} & \left( t^{2\alpha_1\mu_0} \|u(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \right)^2 \\ & \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 + 2\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1\mu_0} \\ & \quad \times \left( \int_0^t (1 + \|u(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)} + \|v(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}) s^{\alpha_1\mu_0} s^{-\alpha_1\mu_0} (t-s)^{\alpha_1-1-\alpha_1\mu_0} ds \right)^2 \\ & \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 + 2\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1\mu_0} \\ & \quad \times \int_0^t (1 + \|u(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)} + \|v(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)})^2 s^{2\alpha_1\mu_0} ds \int_0^t s^{-2\alpha_1\mu_0} (t-s)^{2\alpha_1-2-2\alpha_1\mu_0} ds. \end{aligned}$$

Using the beta function property  $\int_0^t s^{\theta_1-1} (t-s)^{\vartheta_1-1} ds = t^{\theta_1+\vartheta_1-1} \mathbf{B}(\theta_1, \vartheta_1)$ ,  $\theta_1 > 0$ ,  $\vartheta_1 > 0$  with assumption  $0 < \mu_0 < 1 - \frac{1}{2\alpha_1}$ , then  $2\alpha_1 - 1 - 2\alpha_1\mu_0 > 0$  and  $-2\alpha_1\mu_0 + 1 > 0$ , we obtain

$$\begin{aligned} & \left( t^{2\alpha_1\mu_0} \|\mathbf{u}(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \right)^2 \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 \\ & \quad + 2\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1(1-\mu_0)} \mathbf{B}(\theta_1, \vartheta_1) \\ & \quad \times \int_0^t (1 + \|u(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)} + \|v(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)})^2 s^{2\alpha_1\mu_0} ds, \end{aligned}$$

where  $\theta_1 := -2\alpha_1\mu_0 + 1$ ,  $\vartheta_1 := 2\alpha_1 - 1 - 2\alpha_1\mu_0$ . Using the inequality  $(m+n+p)^2 \leq 3(m^2 + n^2 + p^2)$ , we get

$$\begin{aligned} & \left( t^{2\alpha_1\mu_0} \|u(\cdot, t)\|_{\mathbb{H}^\eta(\Omega)} \right)^2 \\ & \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 \\ & \quad + 6\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1(1-\mu_0)} \mathbf{B}(\theta_1, \vartheta_1) \\ & \quad \times \int_0^t (1 + \|u(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2) s^{2\alpha_1\mu_0} ds \\ & \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 + \frac{6\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1\mu_0+1} \mathbf{B}(\theta_1, \vartheta_1)}{2\alpha_1\mu_0 + 1} \\ & \quad + 6\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1(1-\mu_0)} \mathbf{B}(\theta_1, \vartheta_1) \\ & \quad \times \int_0^t (\|u(\cdot, s)\|_{\mathbb{H}^\eta(\Omega)}^2 + \|v(\cdot, s)\|_{\mathbb{H}^\eta(\Omega)}^2) s^{2\alpha_1\mu_0} ds. \tag{4.15} \end{aligned}$$

Similarly, we can also obtain

$$\begin{aligned} & \left( t^{2\alpha_1\mu_0} \|v(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)} \right)^2 \\ & \leq 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|g\|_{\mathcal{H}^\gamma(\Omega)}^2 + \frac{6\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1\mu_0+1} \mathbf{B}(\theta_1, \vartheta_1)}{2\alpha_1\mu_0 + 1} \\ & \quad + 6\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_1(1-\mu_0)} \mathbf{B}(\theta_1, \vartheta_1) \\ & \quad \times \int_0^t (\|u(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2) s^{2\alpha_1\mu_0} ds. \tag{4.16} \end{aligned}$$

From (4.15) and (4.16), we arrive at

$$\begin{aligned} & (\|u(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}^2) t^{2\alpha_1 \mu_0} \\ & \leq \mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) \\ & \quad + \mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T) \int_0^t (\|u(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2) s^{2\alpha_1 \mu_0} ds, \end{aligned} \quad (4.17)$$

where

$$\begin{aligned} \mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) & := 2\bar{M}_2(\alpha_1, \alpha_2, a, \mu_0) \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}^2 \\ & \quad + \frac{12\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1 \mu_0 + 1} \mathbf{B}(\theta_1, \vartheta_1)}{2\alpha_1 \mu_0 + 1}, \\ \mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T) & := 12\bar{M}_3(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1(1-\mu_0)} \mathbf{B}(\theta_1, \vartheta_1). \end{aligned}$$

Applying  $\|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}^2 = \|f\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|g\|_{\mathcal{H}^{\gamma}(\Omega)}^2$  and Gronwall's inequality, we get

$$\begin{aligned} & \|u\|_{\mathbb{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^{\gamma}(\Omega))}^2 + \|v\|_{\mathbb{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^{\gamma}(\Omega))}^2 \\ & \leq \mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) \exp(\mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T)T). \end{aligned}$$

Therefore, we get

$$\begin{aligned} & \|\mathbf{w}\|_{\mathcal{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^{\gamma}(\Omega))} \\ & = \sqrt{\|u\|_{\mathbb{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^{\gamma}(\Omega))}^2 + \|v\|_{\mathbb{C}^{\alpha_1 \mu_0}((0, T], \mathcal{H}^{\gamma}(\Omega))}^2} \\ & \leq \sqrt{\mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)})} \exp\left(\frac{\mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T)T}{2}\right). \end{aligned} \quad (4.18)$$

*Part 3.* From equation (3.6), we then obtain

$$\begin{cases} u_{\alpha}(\cdot, t) = \mathcal{P}_{\Delta, \alpha}(t)f + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)\mathcal{F}(u_{\alpha}, v_{\alpha})(\cdot, s) ds, \\ v_{\alpha}(\cdot, t) = \mathcal{P}_{\Delta, \alpha}(t)g + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)\mathcal{G}(u_{\alpha}, v_{\alpha})(\cdot, s) ds, \end{cases} \quad (4.19)$$

and

$$\begin{cases} u_{\alpha'}(\cdot, t) = \mathcal{P}_{\Delta, \alpha'}(t)f + \int_0^t \mathcal{Q}_{\Delta, \alpha'}(t-s)\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) ds, \\ v_{\alpha'}(\cdot, t) = \mathcal{P}_{\Delta, \alpha'}(t)g + \int_0^t \mathcal{Q}_{\Delta, \alpha'}(t-s)\mathcal{G}(u_{\alpha'}, v_{\alpha'})(\cdot, s) ds. \end{cases} \quad (4.20)$$

Using (4.19) and (4.20), we get

$$\begin{aligned} & u_{\alpha'}(\cdot, t) - u_{\alpha}(\cdot, t) \\ & = [\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]f + \int_0^t [\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \alpha}(t-s)]\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) ds \\ & \quad + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)[\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) - \mathcal{F}(u_{\alpha}, v_{\alpha})(\cdot, s)] ds, \end{aligned} \quad (4.21)$$

$$\begin{aligned}
& v_{\alpha'}(\cdot, t) - v_{\alpha}(\cdot, t) \\
&= [\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]g + \int_0^t [\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \alpha}(t-s)]\mathcal{G}(u_{\alpha'}, v_{\alpha'})(\cdot, s) ds \\
&\quad + \int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)[\mathcal{G}(u_{\alpha'}, v_{\alpha'})(\cdot, s) - \mathcal{G}(u_{\alpha}, v_{\alpha})(\cdot, s)] ds. \tag{4.22}
\end{aligned}$$

Applying the inequality  $(m+n+p)^2 \leq 3(m^2 + n^2 + p^2)$ , we get the following estimate:

$$\begin{aligned}
& \|u_{\alpha'}(\cdot, t) - u_{\alpha}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 \\
&\leq 3\|[\mathcal{P}_{\Delta, \alpha'}(t) - \mathcal{P}_{\Delta, \alpha}(t)]f\|_{\mathcal{H}^{\gamma}(\Omega)}^2 \\
&\quad + 3\left\|\int_0^t [\mathcal{Q}_{\Delta, \alpha'}(t-s) - \mathcal{Q}_{\Delta, \alpha}(t-s)]\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) ds\right\|_{\mathcal{H}^{\gamma}(\Omega)}^2 \\
&\quad + 3\left\|\int_0^t \mathcal{Q}_{\Delta, \alpha}(t-s)[\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) - \mathcal{F}(u_{\alpha}, v_{\alpha})(\cdot, s)] ds\right\|_{\mathcal{H}^{\gamma}(\Omega)}^2. \tag{4.23}
\end{aligned}$$

Using Lemmas 3.1 and 3.2, we obtain

$$\begin{aligned}
& \|u_{\alpha'}(\cdot, t) - u_{\alpha}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 \\
&\leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)]t^{-\alpha_2(1-\nu_0)-\epsilon}\|f\|_{\mathcal{H}^{\gamma}(\Omega)})^2 \\
&\quad + 3\bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)] \\
&\quad \times \left(\int_0^t (t-s)^{\alpha_1\nu_0-\epsilon-1}\|\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)} ds\right)^2 \\
&\quad + 3\bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) \\
&\quad \times \left(\int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0}\|\mathcal{F}(u_{\alpha'}, v_{\alpha'})(\cdot, s) - \mathcal{F}(u_{\alpha}, v_{\alpha})(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)} ds\right)^2. \tag{4.24}
\end{aligned}$$

Using assumptions (4.1) and (4.3), we obtain

$$\begin{aligned}
& \|u_{\alpha'}(\cdot, t) - u_{\alpha}(\cdot, t)\|_{\mathbb{H}^{\eta}(\Omega)}^2 \\
&\leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)]t^{-\alpha_2(1-\nu_0)-\epsilon}\|f\|_{\mathcal{H}^{\gamma}(\Omega)})^2 \\
&\quad + 3C_1^2\bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)]^2I_1 \\
&\quad + 3K_1^2\bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0)I_2, \tag{4.25}
\end{aligned}$$

where

$$I_1 := \left(\int_0^t (t-s)^{\alpha_1\nu_0-\epsilon-1}(1 + \|u_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}) ds\right)^2, \tag{4.26}$$

$$I_2 := \left(\int_0^t (t-s)^{\alpha_1-1-\alpha_1\mu_0}(\|u_{\alpha'} - u_{\alpha}\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'} - v_{\alpha}\|_{\mathcal{H}^{\gamma}(\Omega)}) ds\right)^2. \tag{4.27}$$

Multiplying both sides by  $t^{2\alpha_2(1-\nu_0)+2\epsilon}$ , we get

$$\begin{aligned} & t^{2\alpha_2(1-\nu_0)+2\epsilon} \|u_{\alpha'}(\cdot, t) - u_{\alpha}(\cdot, t)\|_{\mathbb{H}^{\eta}(\Omega)}^2 \\ & \leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)] \|f\|_{\mathcal{H}^{\gamma}(\Omega)})^2 \\ & \quad + 3\mathcal{K}_1^2 \bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_2(1-\mu_0)+2\epsilon} I_2 \\ & \quad + 3\mathcal{C}_1^2 \bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^{\epsilon} + (\alpha' - \alpha)]^2 t^{2\alpha_2(1-\nu_0)+2\epsilon} I_1. \end{aligned} \quad (4.28)$$

Now we estimate  $I_1$ , from (4.26) we get

$$\begin{aligned} I_1 &= \left( \int_0^t (t-s)^{\alpha_1\nu_0-\epsilon-1} s^{-\alpha_1\mu_0} \right. \\ &\quad \times \left. s^{\alpha_1\mu_0} (1 + \|u_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)}) ds \right)^2. \end{aligned}$$

We assume that  $0 < \epsilon < \alpha_1\nu_0 - \frac{1}{2}$  and  $\mu_0 < \frac{1}{2\alpha_1}$ , then  $2\alpha_1\nu_0 - 2\epsilon - 1 > 0$ ,  $1 - 2\alpha_1\mu_0 > 0$ . Using Hölder's inequality and the properties of beta function, we obtain that

$$\begin{aligned} I_1 &\leq \int_0^t (t-s)^{2\alpha_1\nu_0-2\epsilon-2} s^{-2\alpha_1\mu_0} ds \\ &\quad \times \int_0^t s^{2\alpha_1\mu_0} (1 + \|u_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)})^2 ds \\ &\leq t^{2\alpha_1\nu_0-2\alpha_1\mu_0-2\epsilon} \mathbf{B}(\theta_2, \vartheta_2) \\ &\quad \times \int_0^t s^{2\alpha_1\mu_0} (1 + \|u_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'}(\cdot, t)\|_{\mathcal{H}^{\gamma}(\Omega)})^2 ds, \end{aligned} \quad (4.29)$$

where  $\theta_2 := 2\alpha_1\nu_0 - 2\epsilon - 1$ ,  $\vartheta_2 := 1 - 2\alpha_1\mu_0$ .

Applying the inequality  $(m+n+p)^2 \leq 3(m^2 + n^2 + p^2)$ , we can deduce

$$\begin{aligned} I_1 &\leq 3t^{2\alpha_1\nu_0-2\alpha_1\mu_0-2\epsilon} \mathbf{B}(\theta_2, \vartheta_2) \int_0^t s^{2\alpha_1\mu_0} (1 + \|u_{\alpha'}\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v_{\alpha'}\|_{\mathcal{H}^{\gamma}(\Omega)}^2) ds \\ &\leq 3t^{2\alpha_1\nu_0-2\alpha_1\mu_0-2\epsilon} \mathbf{B}(\theta_2, \vartheta_2) \\ &\quad \times \int_0^t (s^{2\alpha_1\mu_0} + \|u_{\alpha'}\|_{\mathcal{C}^{\alpha_1\mu_0}((0,T], \mathcal{H}^{\gamma}(\Omega))}^2 + \|v_{\alpha'}\|_{\mathcal{C}^{\alpha_1\mu_0}((0,T], \mathcal{H}^{\gamma}(\Omega))}^2) ds. \end{aligned} \quad (4.30)$$

Therefore, from (4.18), we have the following estimate:

$$\begin{aligned} I_1 &\leq \frac{3t^{2\alpha_1\nu_0+1-2\epsilon} \mathbf{B}(\theta_2, \vartheta_2)}{2\alpha_1\mu_0 + 1} \\ &\quad + 3t^{2\alpha_1\nu_0-2\alpha_1\mu_0-2\epsilon+1} \mathbf{B}(\theta_2, \vartheta_2) \|w_{\alpha'}\|_{\mathcal{C}^{\alpha_1\mu_0}((0,T], \mathcal{H}^{\gamma}(\Omega))}^2 \\ &\leq \frac{3T^{2\alpha_1\nu_0+1-2\epsilon} \mathbf{B}(\theta_2, \vartheta_2)}{2\alpha_1\mu_0 + 1} + 3T^{2\alpha_1\nu_0-2\alpha_1\mu_0-2\epsilon+1} \mathbf{B}(\theta_2, \vartheta_2) \\ &\quad \times \mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) \exp(\mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T)T). \end{aligned}$$

To facilitate the calculation, we set

$$\begin{aligned} \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, v_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) \\ := \frac{3T^{2\alpha_1 v_0 + 1 - 2\epsilon} \mathbf{B}(\theta_2, \vartheta_2)}{2\alpha_1 \mu_0 + 1} \\ + 3T^{2\alpha_1 v_0 - 2\alpha_1 \mu_0 - 2\epsilon + 1} \mathbf{B}(\theta_2, \vartheta_2) \\ \times \mathcal{M}_1(\alpha_1, \alpha_2, a, \mu_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}) \exp(\mathcal{M}_2(\alpha_1, \alpha_2, a, \mu_0, T)T). \end{aligned}$$

Hence

$$I_1 \leq \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, v_0, T, \|w_0\|_{\mathcal{H}^{\gamma}(\Omega)}). \quad (4.31)$$

Next, estimate  $I_2$ . From (4.27), applying Hölder's inequality, we deduce

$$\begin{aligned} I_2 &= \left( \int_0^t (t-s)^{\alpha_1 - 1 - \alpha_1 \mu_0} s^{-\alpha_2(1-v_0)-\epsilon} \right. \\ &\quad \times s^{\alpha_2(1-v_0)+\epsilon} (\|u_{\alpha'} - u_\alpha\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'} - v_\alpha\|_{\mathcal{H}^{\gamma}(\Omega)}) ds \Big)^2 \\ &\leq \int_0^t (t-s)^{2\alpha_1 - 2 - 2\alpha_1 \mu_0} s^{-2\alpha_2(1-v_0)-2\epsilon} ds \\ &\quad \times \int_0^t s^{2\alpha_2(1-v_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)} + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)})^2 ds \\ &\leq \int_0^t (t-s)^{2\alpha_1 - 2 - 2\alpha_1 \mu_0} s^{-2\alpha_2(1-v_0)-2\epsilon} ds \\ &\quad \times 2 \int_0^t s^{2\alpha_2(1-v_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2) ds. \end{aligned}$$

Using the beta function property with assumption  $\mu_0 < 1 - \frac{1}{2\alpha_1}$  and  $0 < \epsilon < \min(\frac{1}{2} - \alpha_2 + \alpha_2 v_0, \alpha_1 v_0 - \frac{1}{2})$ , we obtain

$$\begin{aligned} I_2 &\leq 2t^{2\alpha_1 - 2\alpha_1 \mu_0 - 2\alpha_2 + 2\alpha_2 v_0 - 2\epsilon} \mathbf{B}(\theta_3, \vartheta_3) \\ &\quad \times \int_0^t s^{2\alpha_2(1-v_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2) ds, \end{aligned}$$

where

$$\theta_3 := -2\alpha_1 - 1 - 2\alpha_1 \mu_0, \quad \vartheta_3 := 1 - 2\alpha_1(1 - v_0) - 2\epsilon.$$

And so, we get

$$\begin{aligned} I_2 &\leq 2t^{2\alpha_1 - 2\alpha_1 \mu_0 - 2\alpha_2 + 2\alpha_2 v_0 - 2\epsilon} \mathbf{B}(\theta_3, \vartheta_3) \\ &\quad \times \int_0^t s^{2\alpha_2(1-v_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^{\gamma}(\Omega)}^2) ds. \quad (4.32) \end{aligned}$$

From (4.28)–(4.32), we obtain

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \|u_{\alpha'}(\cdot, t) - u_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \\
& \leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \|f\|_{\mathcal{H}^\gamma(\Omega)})^2 \\
& \quad + 3C_1^2 \bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 t^{2\alpha_2(1-\nu_0)+2\epsilon} \\
& \quad + 6K_1^2 \bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) t^{2\alpha_2(1-\mu_0)+2\epsilon} 2t^{2\alpha_1-2\alpha_1\mu_0-2\alpha_2+2\alpha_2\nu_0-2\epsilon} \mathbf{B}(\theta_3, \vartheta_3) \\
& \quad \times \int_0^t s^{2\alpha_2(1-\nu_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2) ds. \quad (4.33)
\end{aligned}$$

Hence, we get the following estimate:

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \|u_{\alpha'}(\cdot, t) - u_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \\
& \leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \|f\|_{\mathcal{H}^\gamma(\Omega)})^2 \\
& \quad + 3C_1^2 \bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 T^{2\alpha_2(1-\nu_0)+2\epsilon} \\
& \quad + 6K_1^2 \bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1(1-\mu_0)+2\alpha_2(\nu_0-\mu_0)} \mathbf{B}(\theta_3, \vartheta_3) \\
& \quad \times \int_0^t s^{2\alpha_2(1-\nu_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2) ds. \quad (4.34)
\end{aligned}$$

In the same way as above, we obtain

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \|v_{\alpha'}(\cdot, t) - v_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \\
& \leq 3(\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \|g\|_{\mathcal{H}^\gamma(\Omega)})^2 \\
& \quad + 3C_2^2 \bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, T)[(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 T^{2\alpha_2(1-\nu_0)+2\epsilon} \\
& \quad + 6K_2^2 \bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1(1-\mu_0)+2\alpha_2(\nu_0-\mu_0)} \mathbf{B}(\theta_3, \vartheta_3) \\
& \quad \times \int_0^t s^{2\alpha_2(1-\nu_0)+2\epsilon} (\|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2) ds. \quad (4.35)
\end{aligned}$$

For simplicity to some math formulas, one should put

$$\begin{aligned}
& \mathcal{M}_{\mu_0, \nu_0}^{\mathcal{C}_1, \mathcal{C}_2}(\alpha_1, \alpha_2, \epsilon, a, T, \|w_0\|_{\mathcal{H}^\gamma(\Omega)}) \\
& := 3\bar{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \|w_0\|_{\mathcal{H}^\gamma(\Omega)}^2 \\
& \quad + 3(C_1^2 + C_2^2) \bar{B}^2(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, T) T^{2\alpha_2(1-\nu_0)+2\epsilon}
\end{aligned}$$

and

$$\begin{aligned}
& \mathcal{M}_{\mu_0, \nu_0}^{\mathcal{K}_1, \mathcal{K}_2}(\alpha_1, \alpha_2, \epsilon, a, T) \\
& := 3(K_1^2 + K_2^2) \bar{M}_3^2(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1(1-\mu_0)+2\alpha_2(\nu_0-\mu_0)} \mathbf{B}(\theta_3, \vartheta_3).
\end{aligned}$$

Combining estimates (4.33) and (4.35), we have

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \left( \|u_{\alpha'}(\cdot, t) - u_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, t) - v_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) \\
& \leq 3 \left( \overline{A}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \right)^2 \left( \|f\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|g\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) \\
& \quad + 3(C_1^2 + C_2^2) \overline{B}(\alpha_1, \alpha_2, \epsilon, \nu_0, a, T) \\
& \quad \times \mathcal{M}(\alpha_1, \alpha_2, a, \mu_0, T) T^{2\alpha_2(1-\nu_0)+2\epsilon} [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\
& \quad + 3(K_1^2 + K_2^2) \overline{M}_3(\alpha_1, \alpha_2, a, \mu_0) T^{2\alpha_1(1-\mu_0)+2\alpha_2(\nu_0-\mu_0)} \mathbf{B}(\theta_3, \vartheta_3) \\
& \quad \times \int_0^t s^{2\alpha_2(1-\nu_0)+2\epsilon} \left( \|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) ds. \quad (4.36)
\end{aligned}$$

Hence, we get the following estimate:

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \left( \|u_{\alpha'}(\cdot, t) - u_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, t) - v_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) \\
& \leq \mathcal{M}_{\mu_0, \nu_0}^{C_1, C_2}(\alpha_1, \alpha_2, \epsilon, a, T, \|w_0\|_{\mathcal{H}^\gamma(\Omega)}) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\
& \quad + \mathcal{M}_{\mu_0, \nu_0}^{K_1, K_2}(\alpha_1, \alpha_2, \epsilon, a, T) \\
& \quad \times \int_0^t s^{2\alpha_2(1-\nu_0)+2\epsilon} \left( \|u_{\alpha'}(\cdot, s) - u_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, s) - v_\alpha(\cdot, s)\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) ds. \quad (4.37)
\end{aligned}$$

Applying Gronwall's inequality, we have the following estimate:

$$\begin{aligned}
& t^{2\alpha_2(1-\nu_0)+2\epsilon} \left( \|u_{\alpha'}(\cdot, t) - u_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 + \|v_{\alpha'}(\cdot, t) - v_\alpha(\cdot, t)\|_{\mathcal{H}^\gamma(\Omega)}^2 \right) \\
& \leq \mathcal{M}_{\mu_0, \nu_0}^{C_1, C_2}(\alpha_1, \alpha_2, \epsilon, a, T, \|w_0\|_{\mathcal{H}^\gamma(\Omega)}) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\
& \quad \times \exp(\mathcal{M}_{\mu_0, \nu_0}^{K_1, K_2}(\alpha_1, \alpha_2, \epsilon, a, T)t). \quad (4.38)
\end{aligned}$$

Therefore, we obtain that

$$\begin{aligned}
& \|u_{\alpha'} - u_\alpha\|_{\mathcal{C}^{\alpha_2(1-\nu_0)+\epsilon}((0, T], \mathcal{H}^\gamma(\Omega))}^2 + \|v_{\alpha'} - v_\alpha\|_{\mathcal{C}^{\alpha_2(1-\nu_0)+\epsilon}((0, T], \mathcal{H}^\gamma(\Omega))}^2 \\
& \leq \mathcal{M}_{\mu_0, \nu_0}^{C_1, C_2}(\alpha_1, \alpha_2, \epsilon, a, T, \|w_0\|_{\mathcal{H}^\gamma(\Omega)}) [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)]^2 \\
& \quad \times \exp(\mathcal{M}_{\mu_0, \nu_0}^{K_1, K_2}(\alpha_1, \alpha_2, \epsilon, a, T)T). \quad (4.39)
\end{aligned}$$

Finally, we obtain

$$\begin{aligned}
& \|\mathbf{w}_{\alpha'} - \mathbf{w}_\alpha\|_{\mathcal{C}^{\alpha_2(1-\nu_0)+\epsilon}((0, T], \mathcal{H}^\gamma(\Omega))} \\
& \leq \sqrt{\mathcal{M}_{\mu_0, \nu_0}^{C_1, C_2}(\alpha_1, \alpha_2, \epsilon, a, T, \|w_0\|_{\mathcal{H}^\gamma(\Omega)})} [(\alpha' - \alpha)^\epsilon + (\alpha' - \alpha)] \\
& \quad \times \exp\left(\frac{\mathcal{M}_{\mu_0, \nu_0}^{K_1, K_2}(\alpha_1, \alpha_2, \epsilon, a, T)T}{2}\right). \quad (4.40)
\end{aligned}$$

This completes the proof.  $\square$

## 5 Numerical results

In this section, we show an example which shows the effectiveness of our method. Let the operator  $-\Delta$  on the domain  $\Omega = (0, \pi)$  with the Dirichlet boundary condition and  $t \in [0, 1]$ ,  $\alpha = 1$ , we have the eigenvalues of  $-\Delta$  given by  $\lambda_j = j^2$  ( $j \in \mathbb{Z}^+$ ) and  $\varphi_j(x) = \sqrt{\frac{2}{\pi}} \sin(jx)$ , respectively. We consider the problem to find  $(u(x, t), v(x, t))$  as follows:

$$\begin{cases} \partial_t^\alpha(u(x, t) + \partial_{xx}^2 u(x, t)) + \partial_{xx}^2 u(x, t) = \mathcal{F}(u, v, x, t), & (x, t) \in (0, \pi) \times (0, 1), \\ \partial_t^\alpha(v(x, t) + \partial_{xx}^2 v(x, t)) + \partial_{xx}^2 v(x, t) = \mathcal{G}(u, v, x, t), & (x, t) \in (0, \pi) \times (0, 1), \end{cases} \quad (5.1)$$

where  $(x, t) \in (0, \pi) \times (0, 1)$ , the source functions are given by

$$\begin{cases} \mathcal{F}(u, v, x, t) = v^2 - 4u - \frac{6t^{2-\alpha} \sin(2x)}{\Gamma(3-\alpha)} + (\frac{t^6}{2} - t^3 + \frac{1}{2})(\cos(8x) - 1), \\ \mathcal{G}(u, v, x, t) = u^2 - 16v - \frac{90t^{3-\alpha} \sin(4x)}{\Gamma(4-\alpha)} - (\frac{t^4}{2} - t^2 + \frac{1}{2})(1 - \cos(4x)), \end{cases} \quad (5.2)$$

and the Dirichlet boundary condition as follows:

$$\begin{cases} u(0, t) = u(\pi, t) = 0, & t \in (0, 1), \\ v(0, t) = v(\pi, t) = 0, & t \in (0, 1). \end{cases} \quad (5.3)$$

Assume that the values of  $u, v$  at the initial time  $t = 0$  are given by

$$\begin{cases} u(x, 0) = f(x) := -\sin(2x), & u_t(x, 0) = 0, \quad x \in (0, \pi), \\ v(x, 0) = g(x) := \sin(4x), & v_t(x, 0) = 0, \quad x \in (0, \pi). \end{cases} \quad (5.4)$$

Then the exact solution of problem (5.1)–(5.4) is given by

$$\begin{cases} u(x, t) = (t^2 - 1) \sin(2x), \\ v(x, t) = (1 - t^3) \sin(3x). \end{cases}$$

At the discretization level, a uniform grid of mesh-points  $(x_m, t_n)$  is used to discretize the space and time intervals

$$x_m = \frac{(m-1)\pi}{h_x}, \quad m = \overline{1, h_x + 1}, \quad \text{and} \quad t_n = \frac{n-1}{h_t}, \quad n = \overline{1, h_t + 1}.$$

Next, by using Simpson's rule of numerical integration, we have the following approximate integration of  $z \in L^2(0, \pi)$ :

$$\int_0^\pi z(s) ds \approx \frac{\Delta s}{3} z(s_1) + \frac{2\Delta s}{3} \sum_{l=1}^{(N+1)/2-1} z(s_{2l}) + \frac{4\Delta s}{3} \sum_{l=1}^{(N+1)/2} z(s_{2l-1}) + \frac{\Delta s}{3} z(s_{N+1}).$$

In code Matlab, we have the solution of problem (5.1) which can be written in a matrix form as follows:

$$U = \begin{bmatrix} u(x_1, t_1) & u(x_2, t_1) & u(x_3, t_1) & \cdots & u(x_{m+1}, t_1) \\ u(x_1, t_2) & u(x_2, t_2) & u(x_3, t_2) & \cdots & u(x_{m+1}, t_2) \\ u(x_1, t_3) & u(x_2, t_3) & u(x_3, t_3) & \cdots & u(x_{m+1}, t_3) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ u(x_1, t_{n+1}) & u(x_2, t_{n+1}) & u(x_3, t_{n+1}) & \cdots & u(x_{m+1}, t_{n+1}) \end{bmatrix}_{(n+1) \times (m+1)},$$

and

$$V = \begin{bmatrix} v(x_1, t_1) & v(x_2, t_1) & v(x_3, t_1) & \cdots & v(x_{m+1}, t_1) \\ v(x_1, t_2) & v(x_2, t_2) & v(x_3, t_2) & \cdots & v(x_{m+1}, t_2) \\ v(x_1, t_3) & v(x_2, t_3) & v(x_3, t_3) & \cdots & v(x_{m+1}, t_3) \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v(x_1, t_{n+1}) & v(x_2, t_{n+1}) & v(x_3, t_{n+1}) & \cdots & v(x_{m+1}, t_{n+1}) \end{bmatrix}_{(n+1) \times (m+1)},$$

where the numerical solutions are presented in the following ( $J$  is a truncation parameter of series):

$$\begin{aligned} & u(x_m, t_n) \\ &= \frac{2}{\pi} \sum_{j=1}^J E_{\alpha,1} \left( \frac{-j^2 t_n^\alpha}{1+j^2} \right) \int_0^\pi f(\xi) \sin(j\xi) d\xi \sin(jx_m) \\ &+ \frac{2}{\pi} \sum_{j=1}^J \frac{1}{1+j^2} \int_0^{t_n} (t_n - s)^{\alpha-1} E_{\alpha,\alpha} \left( \frac{-j^2(t_n-s)^\alpha}{1+j^2} \right) \int_0^\pi \mathcal{F}(\xi) \sin(j\xi) d\xi ds \sin(jx_m) \end{aligned}$$

and

$$\begin{aligned} & v(x_m, t_n) \\ &= \frac{2}{\pi} \sum_{j=1}^J E_{\alpha,1} \left( \frac{-j^2 t_n^\alpha}{1+j^2} \right) \int_0^\pi g(\xi) \sin(j\xi) d\xi \sin(jx_m) \\ &+ \frac{2}{\pi} \sum_{j=1}^J \frac{1}{1+j^2} \int_0^{t_n} (t_n - s)^{\alpha-1} E_{\alpha,\alpha} \left( \frac{-j^2(t_n-s)^\alpha}{1+j^2} \right) \int_0^\pi \mathcal{G}(\xi) \sin(j\xi) d\xi ds \sin(jx_m). \end{aligned}$$

By fixing  $t$ , we consider the following estimations with fractional derivative orders  $\alpha$  and  $\alpha^*$  to compare the regularity of the solution, which are given by

$$\text{ESTU}_\alpha^{\alpha^*}(t) = \left( \sum_{m=1}^{h_x+1} |u_\alpha(x_m, t) - u_{\alpha^*}(x_m, t)|^2 / (h_x + 1) \right)^{1/2}, \quad (5.5)$$

$$\text{ESTV}_\alpha^{\alpha^*}(t) = \left( \sum_{m=1}^{h_x+1} |v_\alpha(x_m, t) - v_{\alpha^*}(x_m, t)|^2 / (h_x + 1) \right)^{1/2}, \quad (5.6)$$

where  $(u_{\alpha^*}, v_{\alpha^*})$  and  $(u_\alpha, v_\alpha)$  are the solutions in the case  $\alpha^*$  and  $\alpha$ , respectively.

**Table 1** The error estimates at  $t = 0.2$  for  $\alpha = 1.1, \alpha^* \in \{1.11, 1.12, 1.13, 1.14, 1.15\}$ 



$t = 0.2$	$\alpha^* = 1.11$	$\alpha^* = 1.12$	$\alpha^* = 1.13$	$\alpha^* = 1.14$	$\alpha^* = 1.15$
ESTU $_{\alpha}^{\alpha^*}(0.2)$	0.05274	0.00283	0.00027	2.67385e-05	3.23890e-06
ESTV $_{\alpha}^{\alpha^*}(0.2)$	0.05903	0.00178	0.00019	2.25538e-05	1.91779e-06

**Table 2** The error estimates at  $t = 0.4$  for  $\alpha = 1.5, \alpha^* \in \{1.51, 1.52, 1.53, 1.54, 1.55\}$ 


$t = 0.4$	$\alpha^* = 1.51$	$\alpha^* = 1.52$	$\alpha^* = 1.53$	$\alpha^* = 1.54$	$\alpha^* = 1.55$
ESTU $_{\alpha}^{\alpha^*}(0.4)$	0.09677	0.00563	0.00073	7.57786e-05	6.87516e-06
ESTV $_{\alpha}^{\alpha^*}(0.4)$	0.16414	0.06821	0.06635	0.066176	0.0642495

**Table 3** The error estimates at  $t = 0.8$  for  $\alpha = 1.8, \alpha^* \in \{1.81, 1.82, 1.83, 1.84, 1.85\}$ 

$t = 0.8$	$\alpha^* = 1.81$	$\alpha^* = 1.82$	$\alpha^* = 1.83$	$\alpha^* = 1.84$	$\alpha^* = 1.85$
ESTU $_{\alpha}^{\alpha^*}(0.8)$	0.01754	0.00266	0.00020	2.05406e-05	1.75455e-06
ESTV $_{\alpha}^{\alpha^*}(0.8)$	0.01971	0.00251	0.00029	2.41501e-05	3.50684e-06

**Figure 1** The solutions at  $t = 0.2$  for  $\alpha = 1.1, \alpha^* \in \{1.11, 1.12, 1.13, 1.14, 1.15\}$ **Figure 2** The solutions at  $t = 0.4$  for  $\alpha = 1.5, \alpha^* \in \{1.51, 1.52, 1.53, 1.54, 1.55\}$ 

Tables 1–3 present the error estimates between the solutions for  $\alpha$  and  $\alpha^*$ . It clearly shows that the solution for  $\alpha^*$  converges to the solution for  $\alpha$  as the deflection of the fractional order tends to zero. For a more intuitive look, we can see the graphs of the solutions in Figs. 1, 2, and 3.



**Figure 3** The solutions at  $t = 0.8$  for  $\alpha = 1.8$ ,  $\alpha^* \in \{1.81, 1.82, 1.83, 1.84, 1.85\}$

## 6 Conclusion

In this paper, we considered the initial value problem for a system of nonlinear parabolic pseudo equations with Caputo fractional derivative. Here, we discuss the continuity which is related to a fractional order derivative. To illustrate the theoretical results, we gave an example in some cases. In the future work, we will expand the problem to the case of other derivative definitions such as Riemann–Liouville and conformable. Especially, the operators affecting the spatial variable of two equations are different, this is an open and difficult problem.

### Acknowledgements

This research is funded by Thu Dau Mot University, Binh Duong, Vietnam. The authors would like to thank the reviewers and editor for their constructive comments and valuable suggestions that improve the quality of this paper.

### Funding

Not applicable.

### Availability of data and materials

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

The authors declare that the study was realized in collaboration with the same responsibility. All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

### Author details

<sup>1</sup>Division of Applied Mathematics, Thu Dau Mot University, Binh Duong Province, Vietnam. <sup>2</sup>Department of Mathematics, Cankaya University, Mimar Sinan Caddesi, 06790, Ankara, Turkey. <sup>3</sup>Department of Medical Research, China Medical University, Hsueh-Shih Road, Taichung, 40402, Taiwan. <sup>4</sup>Department of Mathematics and Computer Science, University of Science, Ho Chi Minh City, Vietnam. <sup>5</sup>Vietnam National University, Ho Chi Minh City, Vietnam. <sup>6</sup>Applied Analysis Research Group, Faculty of Mathematics and Statistics, Ton Duc Thang University, Ho Chi Minh City, Vietnam.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 13 August 2020 Accepted: 12 January 2021 Published online: 23 January 2021

### References

- Atangana, A., Baleanu, D.: New fractional derivatives with nonlocal and non-singular kernel: theory and application to heat transfer model. *Therm. Sci.* **20**(2), 763–769 (2016). <https://doi.org/10.2298/TSCI160111018A>
- Sun, H.G., Zhang, Y., Baleanu, D., Chen, W., Chen, Y.Q.: A new collection of real world applications of fractional calculus in science and engineering. *Commun. Nonlinear Sci. Numer. Simul.* **64**, 213–231 (2018)
- Podlubny, I.: Fractional Differential Equations. Academic Press, London (1999)
- Tuan, N.H., Baleanu, D., Thach, T.N., O'Regan, D., Can, N.H.: Final value problem for nonlinear time fractional reaction-diffusion equation with discrete data. *J. Comput. Appl. Math.* **376**, 112883 (2020) 25 pp

5. Luc, N.H., Huynh, L.N., Baleanu, D., Can, N.H.: Identifying the space source term problem for a generalization of the fractional diffusion equation with hyper-Bessel operator. *Adv. Differ. Equ.* **2020**, Paper No. 261, 23 pp. (2020)
6. Ngoc, T.B., Baleanu, D., Duc, L.M., Tuan, N.H.: Regularity results for fractional diffusion equations involving fractional derivative with Mittag-Leffler kernel. *Math. Methods Appl. Sci.* **43**(12) 7208–7226 (2020)
7. Kiryakova, V.: Generalized Fractional Calculus and Applications. Pitman Research Notes in Mathematics, vol. 301. Longman, Harlow (1994)
8. Chen, W., Holm, S.: Physical interpretation of fractional diffusion-wave equation via lossy media obeying frequency power law (2003). <https://arxiv.org/abs/math-ph/0303040>. preprint
9. Yang, X.J., Baleanu, D., Srivastava, H.M.: Local Fractional Integral Transforms and Their Applications. Academic Press, New York (2015)
10. Zhou, Y.: Basic Theory of Fractional Differential Equations. World Scientific, Singapore (2014)
11. Agarwal, P., Nieto, J.J., Luo, M.J.: Extended Riemann-Liouville type fractional derivative operator with applications. *Open Math.* **15**(1), 1667–1681 (2017)
12. Fan, W., Liu, F., Jiang, X., Turner, I.: A novel unstructured mesh finite element method for solving the time-space fractional wave equation on a two-dimensional irregular convex domain. *Fract. Calc. Appl. Anal.* **20**(2), 352–383 (2017)
13. Wang, R., Li, Y., Wang, B.: Random dynamics of fractional nonclassical diffusion equations driven by colored noise. *Discrete Contin. Dyn. Syst.* **39**(7), 4091–4126 (2019)
14. Wang, R., Shi, L., Wang, B., Wang, B.: Asymptotic behavior of fractional nonclassical diffusion equations driven by nonlinear colored noise on  $\mathbb{R}^N$ . *Nonlinearity* **32**, 4524–4556 (2019)
15. Au, V.V., Jafari, H., Hammouch, Z., Tuan, N.H.: On a final value problem for a nonlinear fractional pseudo-parabolic equation. *Electron. Res. Arch.* **29**(1), 1709–1734 (2021)
16. Sakamoto, K., Yamamoto, M.: Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems. *J. Math. Anal. Appl.* **382**, 426–447 (2011)
17. Barenblat, G.I., Kochina, I.: Basic concepts in the theory of seepage of homogeneous liquids in fissured rock. *J. Appl. Math. Mech.* **24**(5), 1286–1303 (1960)
18. Benjamin, T.B., Bona, J.L., Mahony, J.J.: Model equations for long waves in nonlinear dispersive systems. *Philos. Trans. R. Soc. Lond. Ser. A* **272**, 47–78 (1972)
19. Ting, W.T.: Certain non-steady flows of second-order fluids. *Arch. Ration. Mech. Anal.* **14**, 1–26 (1963)
20. Ngoc, T.B., Tuan, N.H., O'Regan, D.: Existence and uniqueness of mild solutions for a final value problem for nonlinear fractional diffusion systems. *Commun. Nonlinear Sci. Numer. Simul.* **78**, 104882 (2019)
21. Padron, V.: Effect of aggregation on population recovery modeled by a forward-backward pseudoparabolic equation. *Trans. Am. Math. Soc.* **356**, 2739–2756 (2004)
22. Samko, S.G., Kilbas, A.A., Marichev, O.I.: Fractional Integrals and Derivatives, Theory and Applications. Gordon and Breach Science, Naukai Tekhnika, Minsk (1987)
23. Diethelm, K.: The Analysis of Fractional Differential Equations. Springer, Berlin (2010)
24. Tuan, N.H., O'Regan, D., Ngoc, T.B.: Continuity with respect to fractional order of the time fractional diffusion-wave equation
25. Wei, T., Zhang, Y.: The backward problem for a time-fractional diffusion-wave equation in a bounded domain. *Comput. Math. Appl.* **75**(10), 3632–3648 (2018)
26. Dang, D.T., Nane, E., Nguyen, D.M., Tuan, N.H.: Continuity of solutions of a class of fractional equations. *Potential Anal.* **49**, 423–478 (2018)
27. Kato, T.: Perturbation Theory for Linear Operators. Springer, Berlin (1995)

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)