

## ON COVERINGS AND HYPERALGEBRAS OF AFFINE ALGEBRAIC GROUPS

BY

MITSUHIRO TAKEUCHI

**ABSTRACT.** Over an algebraically closed field of characteristic zero, the universal group covering of a connected affine algebraic group, if such exists, can be constructed canonically from its Lie algebra only. In particular the isomorphism classes of simply connected affine algebraic groups are in 1-1 correspondence with the isomorphism classes of finite dimensional Lie algebras of some sort. We shall consider the counterpart of these results (due to Hochschild) in case of a positive characteristic, replacing the Lie algebra by the "hyperalgebra". We show that the universal group covering of a connected affine algebraic group scheme can be constructed canonically from its hyperalgebra only and hence, in particular, that the category of simply connected affine algebraic group schemes is equivalent to a subcategory of the category of hyperalgebras of finite type which contains all the semisimple hyperalgebras.

**Introduction.** Let  $k$  be an arbitrary field of arbitrary characteristic. Let  $\mathcal{G}$  and  $\mathcal{H}$  be connected affine algebraic  $k$ -group schemes. If  $\eta: \mathcal{H} \rightarrow \mathcal{G}$  is an epimorphism of  $k$ -group schemes whose kernel  $\text{Ker}(\eta)$  (in the category of  $k$ -group schemes) is a *finite etale*  $k$ -group scheme, then the pair  $(\mathcal{H}, \eta)$  is called an *etale group covering* of  $\mathcal{G}$ . The  $k$ -group scheme  $\mathcal{G}$  is *simply connected* (or (SC)), if it has no nontrivial etale group covering. An etale group covering  $\gamma: \mathcal{G}^* \rightarrow \mathcal{G}$  is called a *universal group covering* of  $\mathcal{G}$ , if the  $k$ -group scheme  $\mathcal{G}^*$  is (SC). Such a universal group covering  $(\mathcal{G}^*, \gamma)$ , if it exists, should satisfy the following universal mapping property and hence will be determined uniquely up to a unique isomorphism.

For each group covering  $(\mathcal{H}, \eta)$  of  $\mathcal{G}$ , there exists a unique homomorphism of  $k$ -group schemes  $\eta^*: \mathcal{G}^* \rightarrow \mathcal{H}$  with  $\eta \circ \eta^* = \gamma$ .

The purpose of this article is to generalize the following result of Hochschild to the case of arbitrary perfect ground field of arbitrary characteristic:

---

Received by the editors March 22, 1974.

AMS (MOS) subject classifications (1970). Primary 14L15.

Key words and phrases. Group scheme, Hopf algebra, hyperalgebra, group covering, simply connected.

Copyright © 1975, American Mathematical Society

**Theorem (Hochschild [2]).** *If  $k$  is an algebraically closed field of characteristic zero, the following statements hold.*

(a) *A connected affine algebraic  $k$ -group scheme  $\mathfrak{G}$  has a universal group covering if and only if the radical of  $\mathfrak{G}$  is unipotent.*

(b) *Then the universal group covering  $\mathfrak{G}^*$  of  $\mathfrak{G}$  can be constructed canonically from its Lie algebra  $L = \text{Lie}(\mathfrak{G})$  only.*

(c) *In particular, the isomorphism classes of (SC)  $k$ -group schemes are in 1-1 correspondence with the isomorphism classes of finite dimensional Lie algebras  $L$  over  $k$  whose radical  $A$  is nilpotent.*

More precisely, let  $L$  be a finite dimensional Lie algebra over  $k$ . The universal enveloping algebra  $U(L)$  of  $L$  has a unique Hopf algebra structure having  $L$  as primitive elements. Its dual Hopf algebra  $H(L) = U(L)^0$  (see §0.1 for definition) is a commutative (but not always finitely generated) domain as an algebra. Let  $A$  be the radical of  $L$  and denote by  $\langle A \rangle$  the ideal of  $U(L)$  generated by  $A$ . Those elements of  $H(L)$  which annihilate some power of  $\langle A \rangle$  form a finitely generated sub-Hopf algebra of  $H(L)$  denoted by  $B(L)$ . Let  $\mathfrak{G}(L) = \text{Spec}(B(L))$  denote the corresponding affine algebraic  $k$ -group scheme. Suppose that the radical  $A$  is nilpotent. Then the  $k$ -group scheme  $\mathfrak{G}(L)$  is (SC) and its Lie algebra  $\text{Lie}(\mathfrak{G}(L))$  is canonically isomorphic with  $L$ . If in particular  $L = [L, L]$ , then the radical  $A$  is automatically nilpotent and we have  $B(L) = H(L)$ . Now let  $\mathfrak{G}$  be a connected affine algebraic  $k$ -group scheme whose radical is unipotent. Let  $\mathcal{O}(\mathfrak{G})$  denote its affine Hopf algebra. The radical  $A$  of the Lie algebra  $L = \text{Lie}(\mathfrak{G})$  is then nilpotent and the image of the canonical injective homomorphism  $\mathcal{O}(\mathfrak{G}) \hookrightarrow U(L)^0 = H(L)$  is contained in  $B(L)$ . Therefore there results a canonical epimorphism of  $k$ -group schemes  $\gamma: \mathfrak{G}(L) \rightarrow \mathfrak{G}$  which proves to be a universal group covering of  $\mathfrak{G}$ . In particular every (SC)  $k$ -group scheme is of the form  $\mathfrak{G}(L)$  with a uniquely determined finite dimensional Lie algebra  $L$  whose radical  $A$  is nilpotent.

When the characteristic of the field is positive, the hyperalgebra plays the same role as the Lie algebra does in case of characteristic zero. The hyperalgebra  $\text{hy}(\mathfrak{G})$  of an affine algebraic  $k$ -group scheme  $\mathfrak{G}$  is by definition the irreducible component containing 1 of the dual Hopf algebra  $\mathcal{O}(\mathfrak{G})^0$  of the affine Hopf algebra  $\mathcal{O}(\mathfrak{G})$ . Takeuchi [T<sub>I</sub>], [T<sub>II</sub>] develops the theory of hyperalgebras of algebraic groups which is completely analogous to the classical theory of Lie algebras of algebraic groups over a field of characteristic zero. The theory of hyperalgebras is briefly summarized in §0.3 for convenience of the reader. We have been able to characterize the (SC)  $k$ -group schemes by their hyperalgebra as follows:

**Theorem.** *Suppose the base field  $k$  is perfect with a positive characteristic  $p$ . For each connected affine algebraic (not necessarily smooth)  $k$ -group scheme  $\mathcal{G}$ , the following conditions are equivalent to each other:*

(i)  $\mathcal{G}$  is (SC).

(ii) *The affine Hopf algebra  $\mathcal{O}(\mathcal{G})$  is canonically isomorphic with the dual Hopf algebra  $\text{hy}(\mathcal{G})^0$  of the hyperalgebra  $\text{hy}(\mathcal{G})$ .*

(iii) *For each locally algebraic (not necessarily affine)  $k$ -group scheme  $\mathcal{H}$ , the map*

$$\text{Hom}_{k\text{-gr}}(\mathcal{G}, \mathcal{H}) \rightarrow \text{Hopf}_k(\text{hy}(\mathcal{G}), \text{hy}(\mathcal{H}))$$

*which sends each  $f \in \text{Hom}_{k\text{-gr}}(\mathcal{G}, \mathcal{H})$  to the induced homomorphism of hyperalgebras,  $\text{hy}(f) \in \text{Hopf}_k(\text{hy}(\mathcal{G}), \text{hy}(\mathcal{H}))$ , is bijjective.*

From this theorem, it follows that if a connected affine algebraic  $k$ -group scheme  $\mathcal{G}$  has a universal group covering  $(\mathcal{G}^*, \gamma)$ , then the affine Hopf algebra  $\mathcal{O}(\mathcal{G}^*)$  is canonically isomorphic with the dual Hopf algebra  $\text{hy}(\mathcal{G})^0$  of  $\text{hy}(\mathcal{G})$ , because  $\text{hy}(\gamma): \text{hy}(\mathcal{G}^*) \xrightarrow{\simeq} \text{hy}(\mathcal{G})$  and  $\mathcal{O}(\mathcal{G}^*) \xrightarrow{\simeq} \text{hy}(\mathcal{G}^*)^0$ . This means that the dual Hopf algebra  $\text{hy}(\mathcal{G})^0$  is *finitely generated*, the corresponding  $k$ -group scheme  $\text{Spec}(\text{hy}(\mathcal{G})^0)$  is (SC), the hyperalgebra of  $\text{Spec}(\text{hy}(\mathcal{G})^0)$  is canonically isomorphic with  $\text{hy}(\mathcal{G})$  and that the canonical homomorphism of  $k$ -group schemes  $\text{Spec}(\text{hy}(\mathcal{G})^0) \rightarrow \mathcal{G}$  is a universal group covering of  $\mathcal{G}$ . Thus the universal group covering of  $\mathcal{G}$ , if it exists, can be canonically constructed from its *hyperalgebra*  $\text{hy}(\mathcal{G})$  only. On the other hand, we have

**Theorem.** *Suppose  $k$  is perfect and  $p > 0$ . If  $\mathcal{G}$  is an (SC) affine algebraic  $k$ -group scheme, then  $\mathcal{G}/[\mathcal{G}, \mathcal{G}]$  is a finite  $k$ -group scheme, that is its affine Hopf algebra is finite dimensional. If in particular  $\mathcal{G}$  is smooth and (SC), then  $\mathcal{G} = [\mathcal{G}, \mathcal{G}]$  and the radical of  $\mathcal{G}$  is unipotent.*

Thus our (SC)  $k$ -group schemes in case of a positive characteristic correspond with the  $k$ -group schemes  $\mathcal{G}(L)$  with  $L = [L, L]$  in case of characteristic zero. In particular the 'if' part of the statement (a) of the theorem of Hochschild does *not* hold in case of a positive characteristic as it stands. (The additive  $k$ -group scheme  $\mathcal{G}_a$  is (SC) if  $p = 0$  but not if  $p > 0$ .) But if we replace the condition (SC) by the following concept of simply connectedness relative to  $p$ , then statement (a) does hold in any characteristic (cf. Miyanishi [4] also).

Let  $p^*$  denote the *characteristic exponent* ( $= \text{Max}(1, p)$ ) of  $k$ . Let  $\mathcal{H}$  and  $\mathcal{G}$  be connected affine algebraic  $k$ -group schemes. An epimorphism of  $k$ -group schemes  $\eta: \mathcal{H} \rightarrow \mathcal{G}$  is called a  *$p$ -etale group covering* of  $\mathcal{G}$  if the kernel

$\text{Spec}(\eta)$  of  $\eta$  is a finite étale  $k$ -group scheme whose order (= the dimension of the affine ring over  $k$ ) is relatively prime to  $p^*$ . The  $k$ -group scheme  $\mathcal{G}$  is called  *$p$ -simply connected* (or  $(\text{SC})_p$ ) if it has no nontrivial  $p$ -étale group covering. By a  *$p$ -universal group covering* of  $\mathcal{G}$  we mean a  $p$ -étale group covering  $(\mathcal{G}^*, \gamma)$  of  $\mathcal{G}$  with  $\mathcal{G}^* (\text{SC})_p$ . Such a  $p$ -universal group covering of  $\mathcal{G}$ , if it exists, satisfies the same universal mapping property as the usual universal group covering of  $\mathcal{G}$  (where of course the  $p$ -étale group coverings must take the place of the usual étale group coverings) and hence is uniquely determined up to a unique isomorphism. If  $p = 0$ , being  $(\text{SC})_0$  is equivalent to being  $(\text{SC})$ .

**Theorem.** *If  $k$  is perfect, a connected smooth affine algebraic  $k$ -group scheme  $\mathcal{G}$  has a  $p$ -universal group covering if and only if the radical of  $\mathcal{G}$  is unipotent.*

In order to obtain the  $p$ -universal group covering of  $\mathcal{G}$  whose radical is unipotent, we must first treat the *semisimple*  $k$ -group schemes. Indeed we shall show that every connected semisimple  $k$ -group scheme has a universal group covering which is at the same time a  $p$ -universal group covering. Hence if  $\mathcal{G}_u$  denotes the unipotent radical of  $\mathcal{G}$ , the quotient group  $\mathcal{G}/\mathcal{G}_u$ , which is semisimple by assumption, has a  $p$ -universal group covering  $(\mathcal{G}/\mathcal{G}_u)^*$ . If we pull it back along  $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{G}_u$ , then we obtain a  $p$ -universal group covering  $\mathcal{G}^* = \mathcal{G} \times_{\mathcal{G}/\mathcal{G}_u} (\mathcal{G}/\mathcal{G}_u)^*$  of  $\mathcal{G}$ .

The case of connected semisimple  $k$ -group schemes goes as follows. First we consider the case where  $k$  is *algebraically closed*. It is well known that all connected semisimple  $k$ -group schemes are then described up to isomorphisms by their *root system* (cf. Satake [5], etc.). Let  $\mathcal{G} = \mathcal{G}(X, \nabla)$  denote the connected semisimple  $k$ -group scheme (i.e., the Chevalley  $k$ -group scheme) determined by the root system  $(X, \nabla)$ . Let  $X_0 = \{\nabla\}_{\mathbb{Z}}$  be the  $\mathbb{Z}$ -submodule of  $X$  generated by  $\nabla$  and  $X^0$  be the *weight module* of  $(X, \nabla)$ , that is

$$X^0 = \{x \in X_{\mathbb{Q}} \mid \langle \nabla^*, x \rangle \in \mathbb{Z}\}$$

where  $\nabla^*$  denotes the *coroot system* of  $(X, \nabla)$ . We have canonical inclusions  $X_0 \subset X \subset X^0$ . Traditionally the Chevalley  $k$ -group scheme  $\mathcal{G}$  is called "simply connected" if  $X = X^0$  and adjoint if  $X = X_0$ . But the simply connectedness in this sense is *not* equivalent with our  $(\text{SC})$ -ness. That is:

**Theorem.** *Suppose  $k$  is algebraically closed. Determine the subgroup  $\bar{X}$  of  $X^0$  containing  $X$  by the following condition:*

$$(p^*, [\bar{X}: X]) = 1 \quad \text{and} \quad [X^0: \bar{X}] = \text{a power of } p^*.$$

(If  $p = 0$ , then  $\bar{X} = X^0$ .) We have then:

(a)  $\mathbb{G} = \mathbb{G}(X, \nabla)$  is (SC) if and only if  $X = \bar{X}$ .

(b) The natural inclusion of reduced root systems  $(X, \nabla) \hookrightarrow (\bar{X}, \nabla)$  induces an isogeny of Chevalley  $k$ -group schemes  $\gamma: \mathbb{G}(\bar{X}, \nabla) \rightarrow \mathbb{G}(X, \nabla)$  (uniquely determined up to inner automorphisms by  $k$ -rational points of the maximal torus of  $\mathbb{G}(\bar{X}, \nabla)$ ) which is a universal group covering, as well as a  $p$ -universal group covering, of  $\mathbb{G}$ .

Thus all "simply connected" Chevalley  $k$ -group schemes are (SC) in our sense but the converse does not hold. For instance even the adjoint Chevalley  $k$ -group scheme  $\mathfrak{B}\mathbb{G}_n = \mathbb{G}_n / \mu_n$  is (SC) when  $n$  is a power of  $p^*$ . Miyanishi [4] seems to have missed these circumstances. (Look at the "proof" of [4, Lemma 4].)

Coming back to the case where  $k$  is only perfect, if  $\mathbb{G}$  is a connected semisimple affine algebraic  $k$ -group scheme, then  $\mathbb{G} \otimes \bar{k}$  (where  $\bar{k}$  = the algebraic closure of  $k$ ), which is semisimple, has the universal group covering  $(\mathbb{G} \otimes \bar{k})^*$ . We shall prove that the  $\bar{k}$ -group scheme  $(\mathbb{G} \otimes \bar{k})^*$  has a " $k$ -form"  $\bar{\mathbb{G}}$  such that the epimorphism  $(\mathbb{G} \otimes \bar{k})^* \rightarrow \mathbb{G} \otimes \bar{k}$  is "defined over  $k$ ". Then  $\bar{\mathbb{G}}$  is easily seen to be a universal, as well as a  $p$ -universal, group covering of  $\mathbb{G}$ . In other words the dual Hopf algebra  $\text{hy}(\bar{\mathbb{G}})^0$  is finitely generated and the hyperalgebra of  $\bar{\mathbb{G}} = \text{Spec}(\text{hy}(\bar{\mathbb{G}})^0)$ , which is "the" universal group covering of  $\mathbb{G}$ , is canonically isomorphic with  $\text{hy}(\mathbb{G})$ . It is clear that the hyperalgebra  $\text{hy}(\mathbb{G})$  is smooth and *semisimple*, in the sense of having no "radical". Conversely we can prove

**Theorem.** *Let  $J$  be a smooth semisimple hyperalgebra over  $k$  of finite type where  $k$  is perfect. Then  $J$  is the hyperalgebra of some (SC) semisimple  $k$ -group scheme  $\mathbb{G}$  which is uniquely determined, that is  $\mathbb{G} = \text{Spec}(J^0)$ . Thus the category of (SC) semisimple  $k$ -group schemes is equivalent to the category of smooth semisimple finite type hyperalgebras over  $k$ .*

Finally we shall conclude this paper by giving an example of (SC)  $k$ -group schemes which are *not reductive*.

This article is based on the theories of Hopf algebras, group schemes and hyperalgebras. They are prepared in §0.

To avoid confusion, all  $k$ -group schemes are denoted by German letters, but all Hopf algebras by Latin letters.

§0. Preliminaries. Throughout the paper  $k$  denotes a fixed ground field of characteristic  $p$ . The *characteristic exponent* of  $k$  is denoted by  $p^* = \text{Max}(1, p)$ . If  $V$  is a  $k$ -vector space, the dual space  $\text{Hom}_k(V, k)$  is denoted

by  $V^*$ . A subset  $T$  of  $V^*$  is *dense* if  $T^\perp = \{v \in V^* \mid \langle T, v \rangle = 0\} = 0$ . A subspace  $W$  of  $V$  is *cofinite* if  $V/W$  is finite dimensional.

For each homomorphism of fields  $\phi: k \rightarrow K$  and each  $k$ -vector space  $V$ ,  $V \otimes_\phi K$  denotes the scalar extension  $V \otimes_k K$ . In particular  $V^{(p)}$  means the scalar extension  $V \otimes_f k$ , where  $f: k \rightarrow k, \lambda \mapsto \lambda^{p^*}$ . Inductively  $V^{(p^n)} = (V^{(p^{n-1})})^{(p)}$ .

0.1. Concerning coalgebras and Hopf algebras we freely use the notation and the terminology of Sweedler [6]. The structure maps of a  $k$ -coalgebra  $C$  will generally be denoted by  $\Delta: C \rightarrow C \otimes C$  and  $\epsilon: C \rightarrow k$ .  $C^+$  means  $\text{Ker}(\epsilon)$ . The "sigma" notation  $\Delta(x) = \sum_{(x)} x_{(1)} \otimes x_{(2)}$  for  $x \in C$  will be used. The  $k$ -coalgebra  $C$  is *irreducible* if any two nonzero subcoalgebras of  $C$  have nonzero intersection [6, §8.0]. A maximal irreducible subcoalgebra of  $C$  is called an *irreducible component* of  $C$ . The  $k$ -coalgebra  $C$  is *cocommutative* if  $\sum_{(x)} x_{(1)} \otimes x_{(2)} = \sum_{(x)} x_{(2)} \otimes x_{(1)}$  for all  $x \in C$ .

The dual space  $C^*$  becomes a  $k$ -algebra which is called the *dual algebra* of  $C$ , if the product is defined by  $f * g = (f \otimes g) \circ \Delta$ . It is commutative if  $C$  is cocommutative.

Conversely, for each  $k$ -algebra  $A$ , the *dual coalgebra*  $A^0$  is defined to be  $\varinjlim_I (A/I)^*$ , where  $I$  runs through all the *cofinite twosided ideals* of  $A$  [6, §6.0]. The functors  $C \mapsto C^*$  and  $A \mapsto A^0$  are *adjoint* to each other in the sense:

$$\text{Alg}_k(A, C^*) \simeq \text{Coalg}_k(C, A^0).$$

A subspace  $I$  of a  $k$ -coalgebra  $C$  is a *coideal* if  $\epsilon(I) = 0$  and  $\Delta(I) \subset I \otimes C + C \otimes I$ . The quotient space  $C/I$  then has a natural coalgebra structure. If  $f: C \rightarrow D$  is a homomorphism of  $k$ -coalgebras, then the kernel  $\text{Ker}(f)$  is a coideal of  $C$  and the map  $f$  factors through  $C \rightarrow C/\text{Ker}(f)$ .

The structure maps of a *Hopf algebra*  $H$  over  $k$  will generally be denoted by  $\Delta: H \rightarrow H \otimes H$ ,  $m: H \otimes H \rightarrow H$ ,  $\epsilon: H \rightarrow k$ ,  $u: k \rightarrow H$  and  $S: H \rightarrow H$  (the antipode). The dual coalgebra  $H^0$  of the  $k$ -algebra  $H$  is a subalgebra of the dual algebra  $H^*$  of the  $k$ -coalgebra  $H$  and is stable under the map  ${}^tS: H^* \rightarrow H^*, f \mapsto f \circ S$ . The induced algebra structure makes  $H^0$  a Hopf algebra with the antipode  ${}^tS|_{H^0}$ , called the *dual Hopf algebra* of  $H$  [6, §6.2]. The functor  $H \mapsto H^0$  is *selfadjoint* in the following sense:

$$\text{Hopf}_k(H, K^0) \simeq \text{Hopf}_k(K, H^0)$$

for all  $k$ -Hopf algebras  $H$  and  $K$ .

The *bracket product* in a  $k$ -Hopf algebra  $H$  is defined by

$$[x, y] = \sum_{(x,y)} x_{(1)}y_{(1)}S(x_{(2)})S(y_{(2)})$$

for  $x, y \in H$ . If  $K$  and  $J$  are sub-Hopf algebras of  $H$ ,  $[K, J]$  denotes the subalgebra of  $H$  generated by the elements  $[x, y]$  with  $x \in K$  and  $y \in J$ . If  $H$  is cocommutative, this is a subbialgebra of  $H$ .

A subspace  $I$  of  $H$  is a Hopf ideal if it is a coideal and a twosided ideal of  $H$  and  $S(I) \subset I$ . The quotient space  $H/I$  then has a natural Hopf algebra structure. If  $f: H \rightarrow H'$  is a homomorphism of Hopf algebras, then the kernel  $\text{Ker}(f)$  is a Hopf ideal of  $H$  and the map  $f$  factors through  $H \rightarrow H/I$ .

If  $K$  is a sub-Hopf algebra of  $H$ , then  $HK^+$  is a coideal of  $H$ . The quotient coalgebra  $H/HK^+$  is denoted by  $H//K$ . For each coideal  $I$  of  $H$ , we put  $l(I) = \{x \in H \mid \Delta(x) - 1 \otimes x \in I \otimes H\}$ . A sub-Hopf algebra  $K$  of  $H$  is normal if  $\sum_{(x)} x_{(1)}yS(x_{(2)}) \in K$  for all  $x \in H$  and  $y \in K$  or equivalently if  $[H, K] \subset K$ . A Hopf ideal  $J$  of  $H$  is normal if  $\sum_{(x)} x_{(1)}S(x_{(3)}) \otimes x_{(2)} \in H \otimes J$  for all  $x \in J$ . In some cases we have a "bijective correspondence" between some class of sub-Hopf algebras and some class of coideals of  $H$  as follows:

0.1.1. Proposition ([7] or [DG, III, §3, n°7]). Let  $H$  be a commutative Hopf algebra.

- (a) If  $K$  is a sub-Hopf algebra of  $H$ , then  $H$  is faithfully flat over  $K$  and  $I = HK^+$  is a normal Hopf ideal of  $H$ . If  $H$  is finitely generated, then so is  $K$ .
- (b) If  $I$  is a normal Hopf ideal of  $H$ , then  $l(I)$  is a sub-Hopf algebra of  $H$ .
- (c) The correspondences  $K \mapsto HK^+$  and  $I \mapsto l(I)$  establish a bijection between the sets of sub-Hopf algebras and of normal Hopf ideals of  $H$ .

0.1.2. Proposition [T<sub>II</sub>, 5.4.2.1, 4.2.2.1, 5.4.2.5, 5.4.3.7]. Let  $H$  be a cocommutative Hopf algebra.

- (a) If  $K$  is a sub-Hopf algebra of  $H$ , then  $H$  is a left and a right faithfully flat  $K$ -module [7] and an injective cogenerator in the category of right  $H//K$ -comodules.  $HK^+$  is clearly a coideal and a left ideal of  $H$ .
- (b) If  $I$  is a coideal and a left ideal of  $H$ , then  $l(I)$  is a sub-Hopf algebra of  $H$ .
- (c) The correspondences  $K \mapsto HK^+$  and  $I \mapsto l(I)$  establish a bijection between the sets of sub-Hopf algebras of  $H$  and of coideal-left-ideals of  $H$ .
- (d) The coideal  $HK^+$  is a Hopf ideal of  $H$  if and only if  $K$  is normal in  $H$ . In particular the bijective correspondence of (c) induces a bijection between the sets of normal sub-Hopf algebras of  $H$  and of Hopf ideals of  $H$ .

If  $f: H \rightarrow H'$  is a homomorphism of Hopf algebras, the Hopf kernel  $\text{Hopf-ker}(f)$  is defined to be the largest sub-Hopf algebra  $K$  of  $H$  with  $K^+ \subset \text{Ker}(f)$ . This equals  $l(\text{Ker}(f))$  when  $H$  is cocommutative or when  $H$  is com-

mutative and the Hopf ideal  $\text{Ker}(f)$  is normal in  $H$ .

Suppose  $p > 0$ . The *Verschiebung* map of a cocommutative  $k$ -coalgebra  $C$ ,  $V_C: C \rightarrow C^{(p)}$  is a unique  $k$ -coalgebra map such that the composite

$$C^{*(p)} \xrightarrow{\text{cano}} C^{(p)*} \xrightarrow{V} C^*$$

sends each element  $X \otimes \lambda$  (with  $X \in C^*$  and  $\lambda \in k$ ) to  $X^p \lambda$  [T<sub>I</sub>, L9.1], [T<sub>II</sub>, 5.5.3.1]. If  $H$  is a cocommutative Hopf algebra, the *Verschiebung* map  $V_H: H \rightarrow H^{(p)}$  is a Hopf algebra map.

0.2. Concerning the basic theory of algebraic schemes and algebraic groups we refer the reader to [DG]. Here we briefly recall some of the fundamental relations between *affine algebraic* group schemes and commutative Hopf algebras.

A  $k$ -group scheme  $\mathcal{G}$  is *affine algebraic* if it is represented by some finitely generated commutative Hopf algebra  $A$ , that is  $\mathcal{G} = \text{Spec}(A)$ . The Hopf algebra  $A$ , which is uniquely determined by  $\mathcal{G}$ , is denoted by  $\mathcal{O}(\mathcal{G})$ . The following well-known relations between  $\mathcal{G} \leftrightarrow A$  are of particular importance:

- $\mathcal{G}$  is *smooth*  $\iff A^{(p)}$  is reduced,
- $\mathcal{G}$  is *connected*  $\iff A$  has no idempotents other than 0 and 1,
- $\mathcal{G}$  is *unipotent*  $\iff A$  is irreducible as a coalgebra,
- $\mathcal{G}$  is *finite*  $\iff [A: k] < \infty$ ,
- $\mathcal{G}$  is *etale*  $\iff A$  is a finite product of finite separable extensions of  $k$ .

The *additive* and the *multiplicative*  $k$ -group schemes  $\mathcal{G}_a$  and  $\mathcal{G}_m$  are defined by

$$\begin{aligned} \mathcal{G}_a &= \text{Spec}(k[T]), & \Delta(T) &= T \otimes 1 + 1 \otimes T, \\ \mathcal{G}_m &= \text{Spec}(k[X, X^{-1}]), & \Delta(X) &= X \otimes X. \end{aligned}$$

For each finitely generated abelian group  $\Gamma$ , the *diagonalizable*  $k$ -group scheme  $\mathcal{D}(\Gamma)$  is defined by

$$\mathcal{D}(\Gamma) = \text{Spec}(k[\Gamma]), \quad \Delta(\gamma) = \gamma \otimes \gamma \quad \text{for all } \gamma \in \Gamma.$$

If  $K/k$  is an extension of fields, the scalar extension  $\mathcal{G} \otimes_k K$ , where  $\mathcal{G} = \text{Spec}(A)$ , is  $\text{Spec}_K(A \otimes_k K)$ .

The  $k$ -group schemes we shall treat are not necessarily smooth. In particular "finite connected"  $k$ -group schemes as follow will be taken into our consideration also:

$$\begin{aligned} {}_q\alpha_k &= \text{Spec}(k[T]/T^q), & \Delta(T) &= T \otimes 1 + 1 \otimes T, \\ {}_q\mu_k &= \text{Spec}(k[X]/(X^q - 1)), & \Delta(X) &= X \otimes X, \end{aligned}$$

where  $q =$  some power of  $p^*$ . The terms such as kernel, center, quotient, etc.



should be always interpreted in the sense of affine algebraic  $k$ -group schemes (not in the sense of  $\bar{k}$ -rational points).

Let  $\mathcal{G} = \text{Spec}(A)$  be an affine algebraic  $k$ -group scheme. The  $k$ -group scheme of the form  $\mathcal{G}'' = \text{Spec}(B)$  with  $B$  a sub-Hopf algebra of  $A$ , is called a *quotient  $k$ -group scheme* of  $\mathcal{G}$ . The  $k$ -group scheme of the form  $\mathcal{H} = \text{Spec}(A/I)$  with  $I$  a Hopf ideal of  $A$ , is called a *closed subgroup scheme* of  $\mathcal{G}$ . The closed subgroup scheme  $\mathcal{H}$  is *normal* in  $\mathcal{G}$  if so is the Hopf ideal  $I$  in the sense of §0.1. It follows from 0.1.1 (c) that there is a natural bijective correspondence between the sets of normal closed subgroup schemes and of quotient group schemes of  $\mathcal{G}$ . The quotient group scheme of  $\mathcal{G}$  associated with a normal closed subgroup scheme  $\mathcal{N}$  of  $\mathcal{G}$  is denoted by  $\mathcal{G}/\mathcal{N}$ . If  $\mathfrak{f}: \mathcal{G} \rightarrow \mathcal{G}'$  is a homomorphism of affine algebraic  $k$ -group schemes, then the normal closed subgroup scheme of  $\mathcal{G}$  corresponding to the image of the induced Hopf algebra map  $\mathcal{O}(\mathfrak{f}): \mathcal{O}(\mathcal{G}') \rightarrow \mathcal{O}(\mathcal{G}) = A$ , which is a sub-Hopf algebra of  $A$ , is called the *kernel* of  $\mathfrak{f}$  and denoted by  $\mathcal{Ker}(\mathfrak{f})$ . The map  $\mathfrak{f}$  factors through  $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{Ker}(\mathfrak{f})$ , via which the quotient group scheme  $\mathcal{G}/\mathcal{Ker}(\mathfrak{f})$  can be viewed as a closed subgroup scheme of  $\mathcal{G}'$  naturally. The map  $\mathfrak{f}$  is an *epimorphism* (resp. a *monomorphism*) if  $\mathcal{G}/\mathcal{Ker}(\mathfrak{f}) \simeq \mathcal{G}'$  (resp.  $\mathcal{Ker}(\mathfrak{f}) = (e)$ ) or equivalently if the Hopf algebra map  $\mathcal{O}(\mathfrak{f})$  is injective (resp. surjective). A sequence  $1 \rightarrow \mathcal{N} \xrightarrow{i} \mathcal{G} \xrightarrow{\mathfrak{p}} \mathcal{G}'' \rightarrow 1$  of affine algebraic  $k$ -group schemes is *exact* if  $i: \mathcal{N} \simeq \mathcal{Ker}(\mathfrak{p})$  and  $\mathfrak{p}: \mathcal{G}/\mathcal{N} \simeq \mathcal{G}''$ .

If  $\mathcal{H}$  is a closed subgroup scheme of  $\mathcal{G}$ , we denote by  $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$  and  $\mathcal{C}_{\mathcal{G}}(\mathcal{H})$  the *normalizer* and the *centralizer* of  $\mathcal{H}$  in  $\mathcal{G}$  respectively [DG, II, §1, n°3]. They are closed subgroup schemes of  $\mathcal{G}$  [DG, II, §1, 3.7]. In particular so is the *centre*  $\text{cent}(\mathcal{G})$ .

The *derived group*  $[\mathcal{G}, \mathcal{G}]$  is defined to be the kernel of the projection  $\mathcal{G} \rightarrow \text{Spec}(B)$ , where  $B$  denotes the *largest cocommutative* sub-Hopf algebra of  $\mathcal{O}(\mathcal{G})$ . If  $\mathcal{G}$  is smooth, this definition coincides with [DG, II, §5, 4.8].

Let  $\mathcal{G}$  be a connected smooth affine algebraic  $k$ -group scheme. When  $k$  is *perfect*, we define the *radical*  $\text{rad}(\mathcal{G})$  (resp. the *unipotent radical*  $\mathcal{G}_u$ ) of  $\mathcal{G}$  to be the largest normal connected smooth solvable (resp. unipotent) closed subgroup scheme of  $\mathcal{G}$ . If  $\bar{k}$  denotes the algebraic closure of  $k$ , then  $\text{rad}(\mathcal{G}) \otimes_k \bar{k}$  (resp.  $\mathcal{G}_u \otimes_k \bar{k}$ ) is the radical (resp. the unipotent radical) of  $\mathcal{G} \otimes_k \bar{k}$  in the usual sense. The smooth connected  $k$ -group scheme  $\mathcal{G}$  is *semisimple* (resp. *reductive*) if  $\text{rad}(\mathcal{G}) = (e)$  (resp.  $\mathcal{G}_u = (e)$ ).

Suppose  $p > 0$  in the rest of this §0.2. The *Frobenius map*  $\mathfrak{F}: \mathcal{G} \rightarrow \mathcal{G}^{(p)}$  of an affine algebraic  $k$ -group scheme  $\mathcal{G} = \text{Spec}(A)$  corresponds to the Hopf algebra map  $A^{(p)} \rightarrow A$ ,  $a \otimes \lambda \mapsto a^p \lambda$  with  $a \in A$  and  $\lambda \in k$ . The  $k$ -group

scheme  $\mathfrak{G}$  is smooth (resp. etale) if and only if the Frobenius map  $\mathfrak{F}$  is an epimorphism (resp. a monomorphism).

When  $\mathfrak{G}$  is commutative (or equivalently if  $\mathcal{O}(\mathfrak{G}) = A$  is cocommutative), the Verschiebung map of  $\mathfrak{G}$ ,  $\mathfrak{V}_{\mathfrak{G}}: \mathfrak{G}^{(p)} \rightarrow \mathfrak{G}$  [DG, IV, §3, n°6] corresponds to the Verschiebung map of  $A$ ,  $V_A: A \rightarrow A^{(p)}$  (defined in §0.1).

Let  $k[F]$  denote the noncommutative polynomial ring over  $k$  defined by  $F\lambda = \lambda^p F$  for all  $\lambda \in k$ . The category of commutative affine algebraic  $k$ -group schemes killed by the Verschiebung map is antiequivalent to the category of finitely generated left  $k[F]$ -modules [DG, IV, §3, 6.6]. We denote by  $\mathfrak{U}(M)$  the  $k$ -group scheme determined by a left  $k[F]$ -module  $M$ . If we view  $M$  as a commutative  $p$ -Lie algebra over  $k$ ,  $U^{[p]}(M)$ , the universal enveloping algebra of  $M$ , has a unique Hopf algebra structure having  $M$  as primitive elements (i.e.,  $\Delta(x) = x \otimes 1 + 1 \otimes x$  for all  $x \in M$ ) and we have  $\mathfrak{U}(M) = \text{Spec}(U^{[p]}(M))$  with this Hopf algebra structure. We have the following equivalence relations:

$$\mathfrak{U}(M) \text{ is smooth} \Leftrightarrow 0 \rightarrow M^{(p)} \xrightarrow{F} M,$$

$$\mathfrak{U}(M) \text{ is finite} \Leftrightarrow [M: k] < \infty,$$

$$\mathfrak{U}(M) \text{ is etale} \Leftrightarrow M = kFM,$$

$$\mathfrak{U}(M) \text{ is connected} \Leftrightarrow \text{each torsion element of } M \text{ is killed by some power of } F.$$

If  $k$  is algebraically closed, we have further

$$\mathfrak{U}(M) \text{ is etale} \Leftrightarrow M \simeq (k[F]/F - 1)^s,$$

$$\mathfrak{U}(M) \text{ is connected smooth} \Leftrightarrow M \simeq k[F]^r$$

[DG, IV, §3, 6.1].

0.3. As a final preliminary, we briefly recall the theory of *hyperalgebras* developed in  $[T_1]$ ,  $[T_{11}]$ .

A *hyperalgebra* means an irreducible cocommutative Hopf algebra. The Lie algebra  $\text{Lie}(J)$  of a hyperalgebra  $J$  is by definition the primitive elements of  $J$ ,  $P(J) = \{x \in J \mid \Delta(x) = x \otimes 1 + 1 \otimes x\}$ . When  $\text{Lie}(J)$  is finite dimensional,  $J$  is called of *finite type*. If so is  $J$ , the dual algebra  $J^*$  is noetherian  $[T_1, 1.4.1]$  and has a finite Krull dimension which cannot exceed  $[\text{Lie}(J): k]$ . We define the Krull dimension  $\dim(J)$  to be the Krull dimension of  $J^*$   $[T_1, 1.4.4]$ . When the equality  $\dim J = [\text{Lie}(J): k]$  holds, the finite type hyperalgebra  $J$  is called *smooth* or of *Birkhoff-Witt type*  $[T_1, 1.6.1]$ . This is equivalent to saying  $J \simeq B(U)$  as a coalgebra, for some finite dimensional  $k$ -vector space  $U$  with the notation of [6, §12.2]. If  $p = 0$  all hyperalgebras are smooth and if  $p > 0$ ,  $J$  is smooth if and only if the Verschiebung map  $V_J: J \rightarrow J^{(p)}$  is surjective  $[T_1, 1.9.4]$ . From this it follows easily that an arbitrary finite type hyperalgebra  $J$  has the largest smooth subhyperalgebra  $J_{sm}$  called the *smooth part* of  $J$  if  $k$  is perfect  $[T_1, 1.9.5]$ . Then the quotient coalgebra  $J//J_{sm}$  is

finite dimensional over  $k$  and we have  $\dim(J) = \dim(J_{sm}) = [\text{Lie}(J_{sm}): k]$  [T<sub>II</sub>, 5.5.3.8].

Among other things very important and useful is the fact that if  $p > 0$  any finite type hyperalgebra  $J$  is the union of its finite dimensional normal subhyperalgebras [T<sub>II</sub>, 5.5.3.7, 5.5.3.9].

Let  $\mathcal{G}$  be a locally algebraic (not necessarily affine)  $k$ -group scheme [DG, II, §5]. Let  $\mathcal{O}_e$  denote the stalk over the unit  $e$  of the structure sheaf  $\mathcal{O}_{\mathcal{G}}$ . The multiplication  $\mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}$  defines naturally a multiplication of the dual coalgebra  $(\mathcal{O}_e)^0$  and makes it a finite type hyperalgebra [T<sub>I</sub>, 3.1.4, 3.3.1] which is denoted by  $\text{hy}(\mathcal{G})$  and called the hyperalgebra of  $\mathcal{G}$ .

If  $\mathcal{G}$  is affine, then  $\text{hy}(\mathcal{G})$  coincides with the irreducible component containing 1 of the dual Hopf algebra  $\mathcal{O}(\mathcal{G})^0$ . In particular, since the functor  $H \mapsto H^0$  is selfadjoint (§0.1), the inclusion  $\text{hy}(\mathcal{G}) \hookrightarrow \mathcal{O}(\mathcal{G})^0$  corresponds to a natural Hopf algebra map  $\mathcal{O}(\mathcal{G}) \rightarrow \text{hy}(\mathcal{G})^0$  which is not necessarily injective.

A homomorphism of locally algebraic  $k$ -group schemes  $\mathfrak{f}: \mathcal{G} \rightarrow \mathcal{G}'$  induces clearly a homomorphism of hyperalgebras  $\text{hy}(\mathfrak{f}): \text{hy}(\mathcal{G}) \rightarrow \text{hy}(\mathcal{G}')$ . If  $\mathcal{G}$  is affine, it is easy to see that the natural map  $\mathcal{O}(\mathcal{G}) \rightarrow \text{hy}(\mathcal{G})^0$  is bijective if and only if the natural map  $\text{Hom}_{k\text{-gr}}(\mathcal{G}, \mathcal{G}') \rightarrow \text{Hopf}_k(\text{hy}(\mathcal{G}), \text{hy}(\mathcal{G}'))$  which sends each map  $\mathfrak{f}$  to the induced map  $\text{hy}(\mathfrak{f})$  is bijective for all affine algebraic  $k$ -group schemes  $\mathcal{G}'$ .

The functor  $\mathcal{G} \mapsto \text{hy}(\mathcal{G})$  has many interesting properties which are similar and reduce to the properties of the functor  $\mathcal{G} \mapsto \text{Lie}(\mathcal{G})$  in case  $p = 0$ :

0.3.1. **Proposition.** Let  $\mathcal{G}$  be a locally algebraic  $k$ -group scheme.

(a) [T<sub>I</sub>, 3.1.8, 3.3.1]. The Lie algebra of  $\mathcal{G}$ ,  $\text{Lie}(\mathcal{G})$  [DG, II, §4, n°1], equals  $\text{Lie}(\text{hy}(\mathcal{G}))$ . The dimension of  $\mathcal{G}$ ,  $\dim \mathcal{G}$  [DG, II, §5, 1.3], equals  $\dim(\text{hy}(\mathcal{G}))$ .

(b) [T<sub>I</sub>, 3.1.7]. If  $K/k$  is an arbitrary extension of fields, then  $\text{hy}(\mathcal{G}) \otimes_k K$  equals the  $K$ -hyperalgebra  $\text{hy}_K(\mathcal{G} \otimes_k K)$  of the locally algebraic  $K$ -group scheme  $\mathcal{G} \otimes_k K$ .

(c) [T<sub>I</sub>, 2.2.9]. If  $p > 0$ , the Frobenius map  $\mathfrak{F}_{\mathcal{G}}: \mathcal{G} \rightarrow \mathcal{G}^{(p)}$  induces the Verschiebung map  $V: \text{hy}(\mathcal{G}) \rightarrow \text{hy}(\mathcal{G})^{(p)}$ .

(d) [T<sub>I</sub>, 3.3.5, 3.3.11].  $\mathcal{G}$  is smooth if and only if  $\text{hy}(\mathcal{G})$  is. When  $k$  is perfect, let  $\mathcal{G}_{red}$  denote the reduced part of  $\mathcal{G}$  [DG, II, §5, 2.3]. Then the hyperalgebra  $\text{hy}(\mathcal{G}_{red})$  equals the smooth part  $\text{hy}(\mathcal{G})_{sm}$  of  $\text{hy}(\mathcal{G})$ .

(e) [T<sub>I</sub>, 3.3.3].  $\mathcal{G}$  is étale if and only if  $\text{hy}(\mathcal{G}) = k$ .

(f) [T<sub>I</sub>, 3.3.6]. A subgroup scheme  $\mathcal{H}$  of  $\mathcal{G}$  is open if and only if  $\text{hy}(\mathcal{H}) = \text{hy}(\mathcal{G})$ . In particular if  $\mathcal{G}$  is connected, then  $\mathcal{H} = \mathcal{G}$  if and only if  $\text{hy}(\mathcal{H}) = \text{hy}(\mathcal{G})$ .

(g) If  $\mathcal{G}$  is affine, the canonical map  $\mathcal{O}(\mathcal{G}) \rightarrow \text{hy}(\mathcal{G})^0$  is injective if and only if  $\mathcal{G}$  is connected.

(h) If  $p > 0$  and if  $\mathcal{G}$  is affine connected, then  $\mathcal{G}$  is unipotent if and only if each element of  $\text{hy}(\mathcal{G})^+$  is nilpotent.

**0.3.2. Proposition.** Let  $f: \mathcal{G} \rightarrow \mathcal{G}'$  be a homomorphism of locally algebraic  $k$ -group schemes.

(a) [ $T_1$ , 3.1.5]. The hyperalgebra of the kernel  $\text{Ker}(f)$  equals the Hopf kernel of  $\text{hy}(f)$ . In particular if  $f$  is a monomorphism, then  $\text{hy}(f)$  is injective.

(b) [ $T_1$ , 3.3.2]. If both  $\mathcal{G}$  and  $\mathcal{G}'$  are algebraic, there is a unique closed subgroup scheme  $f(\mathcal{G})^\sim$  of  $\mathcal{G}'$ , called the image-subgroup of  $f$ , such that the induced map  $f: \mathcal{G} \rightarrow f(\mathcal{G})^\sim$  is faithfully flat [DG, III, §3, 5.2, 2.6, II, §5.5.1]. Then we have

$$\text{hy}(f(\mathcal{G})^\sim) = \text{Im}(\text{hy}(f)) = \text{hy}(\mathcal{G}) // \text{Hopf-kern}(\text{hy}(f)).$$

(c) [ $T_1$ , 3.3.3]. The map  $f$  is nonramified [DG, I, §4, 3.2] if and only if  $\text{hy}(f)$  is injective.

(d) [ $T_1$ , 3.3.4]. The map  $f$  is flat if and only if  $\text{hy}(f)$  is surjective.

(e) [ $T_1$ , 3.3.7]. If  $\mathcal{G}'$  is connected and  $\mathcal{G}$  is algebraic, then  $f(\mathcal{G})^\sim = \mathcal{G}'$  if and only if  $\text{hy}(f)$  is surjective.

(f) [ $T_1$ , 3.3.9]. Let  $\mathcal{H}$  and  $\mathcal{R}$  be subgroup schemes of  $\mathcal{G}$ . If  $\mathcal{H}$  is connected, then  $\mathcal{H} \subset \mathcal{R}$  if and only if  $\text{hy}(\mathcal{H}) \subset \text{hy}(\mathcal{R})$ . In particular the correspondence  $\mathcal{H} \mapsto \text{hy}(\mathcal{H})$  from the set of connected (and hence closed) subgroup schemes of  $\mathcal{G}$  into the set of subhyperalgebras of  $\text{hy}(\mathcal{G})$  is injective.

(g) [ $T_1$ , 3.3.10]. Let  $f': \mathcal{G} \rightarrow \mathcal{G}'$  be another homomorphism. If  $\mathcal{G}$  is connected (and hence algebraic), then  $f = f'$  if and only if  $\text{hy}(f) = \text{hy}(f')$ .

Let  $\mathcal{G}$  be a locally algebraic  $k$ -group scheme. The inner automorphism action  $\mathcal{G} \times \mathcal{G} \rightarrow \mathcal{G}, (g, h) \mapsto ghg^{-1}$  induces a linear representation  $\mathcal{U}\mathfrak{b}: \mathcal{G} \rightarrow \mathcal{U}\mathfrak{L}(\text{hy}(\mathcal{G}))$ , called the adjoint representation of  $\mathcal{G}$  [ $T_1$ , 3.1.6, 3.4.13]. We view as usual each  $k$ -group scheme as a  $k$ -group functor, i.e., a functor from the category  $M_k$  of  $k$ -models to the category of groups [DG, I, §1, n°4].

**0.3.3. Proposition.** Let  $\mathcal{H}$  be a closed subgroup scheme of a locally algebraic  $k$ -group scheme  $\mathcal{G}$ .  $\mathcal{N}_{\mathcal{G}}(\mathcal{H})$  and  $\mathcal{C}_{\mathcal{G}}(\mathcal{H})$  denote the normalizer and the centralizer of  $\mathcal{H}$  in  $\mathcal{G}$ .

(a) [ $T_1$ , 3.4.13].  $\text{hy}(\mathcal{N}_{\mathcal{G}}(\mathcal{H}))$  (resp.  $\text{hy}(\mathcal{C}_{\mathcal{G}}(\mathcal{H}))$ ) is the largest subcoalgebra  $C$  (resp.  $D$ ) of  $\text{hy}(\mathcal{G})$  satisfying

$$\sum_{(c)} \mathcal{U}\mathfrak{b}(h)(c_{(1)})\mathcal{S}(c_{(2)}) \in R \otimes \text{hy}(\mathcal{H})$$

$$\left( \text{resp. } \sum_{(d)} \mathcal{U}\mathfrak{b}(h)(d_{(1)})\mathcal{S}(d_{(2)}) \in R \otimes k \right)$$

for all  $R \in \mathbf{M}_k$ ,  $b \in \mathfrak{H}(R)$  and  $c \in C$  (resp.  $d \in D$ ), where  $S$  denotes the antipode of  $\text{hy}(\mathfrak{G})$ .

(b) [ $T_1$ , 3.4.15]. If  $\mathfrak{G}$  is connected (and hence algebraic) then  $\mathfrak{H}$  is normal (resp. central) in  $\mathfrak{G}$  if and only if

$$\sum_{(a)} \mathfrak{U}\mathfrak{b}(h)(a_{(1)})S(a_{(2)}) \in R \otimes \text{hy}(\mathfrak{H})$$

$$\left( \text{resp. } \sum_{(a)} \mathfrak{U}\mathfrak{b}(h)(a_{(1)})S(a_{(2)}) \in R \otimes k \right)$$

for all  $R \in \mathbf{M}_k$ ,  $b \in \mathfrak{H}(R)$  and  $a \in \text{hy}(\mathfrak{G})$ .

(c) [ $T_1$ , 3.4a.5]. If  $\mathfrak{H}$  is connected, then  $\mathfrak{H}$  is normal (resp. central) in  $\mathfrak{G}$  if and only if  $\text{hy}(\mathfrak{H})$  is  $\mathfrak{U}\mathfrak{b}(\mathfrak{G})$ -stable (resp.  $\mathfrak{U}\mathfrak{b}(\mathfrak{G})$  acts trivially on  $\text{hy}(\mathfrak{H})$ ).

(d) [ $T_1$ , 3.4.15]. If  $\mathfrak{H}$  and  $\mathfrak{G}$  are both connected, then  $\mathfrak{H}$  is normal (resp. central) in  $\mathfrak{G}$  if and only if so is the subhyperalgebra  $\text{hy}(\mathfrak{H})$  in  $\text{hy}(\mathfrak{G})$ . (A sub-Hopf algebra  $J$  of a Hopf algebra  $H$  is central if  $[H, J] = k$ .)

Let  $\mathfrak{G}$  be a locally algebraic  $k$ -group scheme. A subhyperalgebra  $J$  of  $\text{hy}(\mathfrak{G})$  is *algebraic* (or *closed*) if there is a subgroup scheme  $\mathfrak{H}$  of  $\mathfrak{G}$  with  $J = \text{hy}(\mathfrak{H})$ . It follows from 0.3.2(f) that we can take then a *unique connected* (and hence closed algebraic) one as  $\mathfrak{H}$ . For each subhyperalgebra  $J$  of  $\text{hy}(\mathfrak{G})$  there is a unique *smallest* algebraic subhyperalgebra  $A(J)$  of  $\text{hy}(\mathfrak{G})$  containing  $J$  [ $T_1$ , §3.6] called the *algebraic hull* of  $J$ .

**0.3.4. Proposition.** Let  $\mathfrak{G}$  be a locally algebraic  $k$ -group scheme and  $J$  a subhyperalgebra of  $\text{hy}(\mathfrak{G})$ .

(a) If  $[J: k] < \infty$ , then  $J$  is closed.

(b) Let  $K$  be a subhyperalgebra of  $J$ . If  $J//K$  is finite dimensional, then  $K$  is closed in  $\text{hy}(\mathfrak{G})$  if and only if so is  $J$ .

(c) Let  $l/k$  be an arbitrary extension of fields. Then  $J$  is closed in  $\text{hy}(\mathfrak{G})$  if and only if so is  $J \otimes l$  in  $\text{hy}(\mathfrak{G}) \otimes l = \text{hy}_l(\mathfrak{G}_l)$ .

(d) [ $T_1$ , 3.6.2]. Let  $A$  be a subalgebra and  $C$  a subcoalgebra of  $\text{hy}(\mathfrak{G})$  such that  $AC \subset C$ . If  $[J, C] \subset A$ , then  $[A(J), C] \subset A$ .

(e) [ $T_1$ , 3.6.2]. Let  $K$  be a subhyperalgebra of  $J$ . If  $K$  is normal in  $J$ , then so is  $K$  in  $A(J)$ .

(f) [ $T_1$ , 3.6.3]. We have  $[J, J] = [A(J), A(J)]$  (which is closed in  $\text{hy}(\mathfrak{G})$  by the following (g)). In particular the quotient hyperalgebra  $A(J)//J$  is abelian.

(g) If  $J_1$  and  $J_2$  are closed subhyperalgebras of  $\text{hy}(\mathfrak{G})$ , then the commutator subhyperalgebra  $[J_1, J_2]$  is also closed in  $\text{hy}(\mathfrak{G})$ .

(h) [T<sub>1</sub>, 3.5.6]. If  $\mathcal{G}$  is connected smooth, then  $\text{hy}([\mathcal{G}, \mathcal{G}]) = [\text{hy}(\mathcal{G}), \text{hy}(\mathcal{G})]$ .

(i) If  $\mathcal{G}$  is connected affine, then  $\text{hy}([\mathcal{G}, \mathcal{G}]) = [\text{hy}(\mathcal{G}), \text{hy}(\mathcal{G})]$ .

The proof of (a), (b), (c), (f), (g) and (i) above will be published elsewhere. In particular if  $\mathcal{H}$  and  $\mathcal{R}$  are *connected* subgroup schemes of a locally algebraic  $k$ -group scheme  $\mathcal{G}$ , we can define the *commutator* subgroup  $[\mathcal{H}, \mathcal{R}]$  to be the *unique connected* subgroup of  $\mathcal{G}$  with  $\text{hy}([\mathcal{H}, \mathcal{R}]) = [\text{hy}(\mathcal{H}), \text{hy}(\mathcal{R})]$ . This definition generalizes [DG, II, §5, 4.9].

1. The (SC) (or  $(\text{SC})_p$ )  $k$ -group schemes. We have defined in the introduction the concepts of etale group covering, universal group covering, (SC) or  $(\text{SC})_p$   $k$ -group scheme, etc.

1.1. **Proposition.** Let  $\mathcal{G}$  and  $\mathcal{H}$  be connected affine algebraic  $k$ -group schemes and  $\eta: \mathcal{H} \rightarrow \mathcal{G}$  a homomorphism of  $k$ -group schemes. Then  $(\mathcal{H}, \eta)$  is an etale group covering of  $\mathcal{G}$  if and only if  $\text{hy}(\eta): \text{hy}(\mathcal{H}) \simeq \text{hy}(\mathcal{G})$ .

This follows directly from 0.3.2(c) and (e).

1.2. **Proposition.** Let  $\mathcal{G}$  be a connected affine algebraic  $k$ -group scheme and  $\mathcal{R}$  a normal connected closed subgroup scheme of  $\mathcal{G}$ . (a) If  $\mathcal{G}$  is (SC) (resp.  $(\text{SC})_p$ ), then so is  $\mathcal{G}/\mathcal{R}$ . (b) If  $\mathcal{G}/\mathcal{R}$  and  $\mathcal{R}$  are both (SC) (resp.  $(\text{SC})_p$ ), then so is  $\mathcal{G}$ .

The proof is easy.

1.3. **Remark.** All finite connected  $k$ -group schemes are clearly (SC).

1.4. **Remark.** The multiplicative group  $\mathcal{G}_m$  is not  $(\text{SC})_p$ , in view of the canonical extension  $1 \rightarrow \mu_n \rightarrow \mathcal{G}_m \rightarrow \mathcal{G}_m \rightarrow 1$  for all  $n$  relatively prime to  $p^*$ . More generally if  $\mathcal{H}$  is a  $k$ -torus, that is  $\mathcal{H} \otimes \bar{k} \simeq (\mathcal{G}_m \otimes \bar{k})^r$  for some  $r \in \mathbb{N}$ , then  $\mathcal{H}$  is not  $(\text{SC})_p$  unless  $r = 0$ . Indeed if  $\mathcal{R}$  denotes the kernel of the morphism  $\mathcal{H} \rightarrow \mathcal{H}, x \mapsto x^n$ , then  $\mathcal{R} \otimes \bar{k} \simeq (\mu_n \otimes \bar{k})^r$  clearly and so  $\mathcal{R}$  is finite etale with the order  $n^r$  which is prime to  $p^*$  if  $n$  is so chosen. The additive group  $\mathcal{G}_a$  is not (SC) when  $p > 0$ , in view of the Artin-Schreier extension

$$1 \rightarrow (\mathbb{Z}/p\mathbb{Z})_k \rightarrow \mathcal{G}_a \xrightarrow{\mathfrak{F}-1} \mathcal{G}_a \rightarrow 1$$

where  $\mathfrak{F}$  denotes the Frobenius map.

In general, each affine algebraic  $k$ -group scheme  $\mathcal{G}$  has a normal closed connected finite subgroup scheme  $\mathcal{N}$  such that  $\mathcal{G}/\mathcal{N}$  is smooth [DG, III, §3, 6.10]. Since  $\mathcal{N}$  is always (SC), it follows that  $\mathcal{G}$  is (SC) (resp.  $(\text{SC})_p$ ) if and

only if  $\mathcal{G}/\mathcal{N}$  is. If  $\mathcal{G}/\mathcal{N}$  has a ( $p$ -) universal group covering, then pulling it back along  $\mathcal{G} \rightarrow \mathcal{G}/\mathcal{N}$ , we obtain a ( $p$ -) universal group covering of  $\mathcal{G}$ . Hence all consideration concerning group coverings reduces to the case of smooth  $k$ -group schemes.

**1.5. Proposition.** *Connected affine algebraic unipotent  $k$ -group schemes are  $(SC)_p$ .*

**Proof.** We shall consider the case  $p > 0$  first. Let  $\mathcal{G}$  be connected unipotent  $\neq (e)$ . Since  $\mathcal{G} \neq [\mathcal{G}, \mathcal{G}]$ , it suffices to prove that  $\mathcal{G}/[\mathcal{G}, \mathcal{G}]$  is  $(SC)_p$ , by the induction hypothesis. Thus we can assume  $\mathcal{G}$  is commutative. Let  $\mathfrak{V}_{\mathcal{G}}: \mathcal{G}^{(p)} \rightarrow \mathcal{G}$  denote the Verschiebung map of  $\mathcal{G}$ . Since  $\mathcal{G} \neq \mathfrak{V}_{\mathcal{G}}(\mathcal{G}^{(p)})$ , we can assume that  $\mathfrak{V}_{\mathcal{G}} = 0$ , by the induction hypothesis again. There is then a finitely generated left  $k[F]$ -module  $M$  with  $\mathcal{G} \simeq \mathcal{U}(M)$  (§0.2). Let  $1 \rightarrow \mathfrak{R} \rightarrow \mathfrak{H} \rightarrow \mathcal{U}(M) \rightarrow 1$  be a  $p$ -etale group covering of  $\mathcal{U}(M)$ . Since  $\text{hy}(\mathfrak{H}) \simeq \text{hy}(\mathcal{U}(M))$ , it follows from 0.3.1(h) that  $\mathfrak{H}$  is unipotent and from 0.3.1(g) that  $\mathfrak{H}$  is commutative. Since  $\text{hy}(\mathfrak{V}_{\mathfrak{H}})$  is trivial, it follows from 0.3.2(g) that  $\mathfrak{V}_{\mathfrak{H}} = 0$ . In particular we have  $\mathfrak{V}_{\mathfrak{R}} = 0$  and so  $\mathfrak{R} \simeq \mathcal{U}(N)$  for some finite dimensional (over  $k$ ) left  $k[F]$ -module  $N$ . The order of  $\mathfrak{R}$  ( $= [U^{[p]}(N): k]$ ) is a power of  $p$ . But since the order of  $\mathfrak{R}$  is relatively prime to  $p$ , it follows that  $\mathfrak{R} = (e)$ . Hence  $\mathcal{U}(M)$  is  $(SC)_p$ .

Next suppose that  $p = 0$ . Since each affine algebraic unipotent  $k$ -group scheme has a central series of closed subgroup schemes each of whose quotients is isomorphic to  $\mathcal{G}_a$  [DG, IV, §2, 3.9, 4.1], it suffices to show that  $\mathcal{G}_a$  is  $(SC)_0$ . But since any etale group covering of  $\mathcal{G}_a$  is clearly 1-dimensional unipotent and hence isomorphic to  $\mathcal{G}_a$  [DG, IV, §2, 2.10], the claim follows from the fact that  $\mathcal{G}_a$  has no nontrivial etale subgroup scheme [DG, IV, §2, 1.1]. Q.E.D.

**1.6. Theorem.** *If  $k$  is perfect, a connected smooth affine algebraic  $k$ -group scheme  $\mathcal{G}$  is  $(SC)_p$  if and only if the radical  $\text{rad}(\mathcal{G})$  is unipotent and that  $\bar{\mathcal{G}} = \mathcal{G}/\text{rad}(\mathcal{G})$  is  $(SC)$ .*

**Proof.** The 'if' part follows from Propositions 1.5 and 1.2. Suppose  $\mathcal{G}$  is  $(SC)_p$ . Let  $\mathcal{G}_u$  denote the unipotent radical of  $\mathcal{G}$ . Since  $\bar{\mathcal{G}} = \mathcal{G}/\mathcal{G}_u$  is  $(SC)_p$  and reductive, it follows that  $\bar{\mathcal{G}}/[\bar{\mathcal{G}}, \bar{\mathcal{G}}]$  is an  $(SC)_p$  torus. But because any nontrivial  $k$ -torus is not  $(SC)_p$ , it follows that  $\bar{\mathcal{G}} = [\bar{\mathcal{G}}, \bar{\mathcal{G}}]$  or equivalently that  $\bar{\mathcal{G}}$  is semisimple. Since we show in §3 that all connected  $(SC)_p$  semisimple  $k$ -group schemes are  $(SC)$ , it follows that  $\bar{\mathcal{G}}$  is  $(SC)$ . Q.E.D.

**1.7. Theorem.** *If  $k$  is perfect, a connected smooth affine algebraic  $k$ -*

group scheme  $\mathbb{G}$  has a  $p$ -universal group covering if and only if the radical  $\text{rab}(\mathbb{G})$  is unipotent.

**Proof.** The 'only if' part follows from Theorem 1.6. Suppose conversely that the radical  $\text{rab}(\mathbb{G})$  is unipotent. The semisimple  $k$ -group scheme  $\overline{\mathbb{G}} = \mathbb{G}/\text{rab}(\mathbb{G})$  has a  $p$ -universal group covering  $\overline{\mathbb{G}}^*$  as we shall see in §3. The pull-back  $\mathbb{G}^* = \mathbb{G} \times_{\overline{\mathbb{G}}} \overline{\mathbb{G}}^*$  then is a  $p$ -universal group covering of  $\mathbb{G}$  by Theorem 1.6. Q.E.D.

1.8. Lemma. Let  $\mathbb{G}$  be a connected affine algebraic  $k$ -group scheme and consider the following three conditions:

(a)  $\mathbb{G}$  is (SC).

(b) For each locally algebraic (not necessarily affine)  $k$ -group scheme  $\mathbb{H}$ , the natural map

$$\text{Hom}_{k\text{-gr}}(\mathbb{G}, \mathbb{H}) \rightarrow \text{Hopf}_k(\text{hy}(\mathbb{G}), \text{hy}(\mathbb{H})), \quad \mathbb{f} \mapsto \text{hy}(\mathbb{f})$$

is bijjective.

(c) The canonical map  $\mathcal{O}(\mathbb{G}) \rightarrow \text{hy}(\mathbb{G})^0$  is bijjective.

We have then an implication (b)  $\Rightarrow$  (c)  $\Rightarrow$  (a). If further  $\mathbb{G}/[\mathbb{G}, \mathbb{G}]$  is finite, then (a)  $\Rightarrow$  (b).

**Proof.** We pointed out in §0.3 that condition (c) is equivalent to the bijectivity of the maps of (b) for all affine algebraic  $k$ -group schemes  $\mathbb{H}$ . In particular we have (b)  $\Rightarrow$  (c). Let  $\gamma: \overline{\mathbb{G}} \rightarrow \mathbb{G}$  be an etale group covering. Since  $\text{hy}(\gamma)$  is isomorphic, if (c) is valid, there is a unique homomorphism  $\sigma: \overline{\mathbb{G}} \rightarrow \mathbb{G}$  with  $\text{hy}(\sigma) = \text{hy}(\gamma)^{-1}$ . Then  $\gamma \circ \sigma = \text{id}$ , because  $\text{hy}(\gamma \circ \sigma) = \text{id}$ . Since  $\sigma$  is an epimorphism of affine algebraic  $k$ -group schemes, it follows that  $\sigma \circ \gamma = \text{id}$ . Hence  $\gamma$  is an isomorphism, so we have (c)  $\Rightarrow$  (a). Finally suppose that  $\mathbb{G}$  is (SC) with  $\mathbb{G}/[\mathbb{G}, \mathbb{G}]$  finite. Let  $\mathbb{H}$  be an arbitrary locally algebraic  $k$ -group scheme. Let  $\omega: \text{hy}(\mathbb{G}) \rightarrow \text{hy}(\mathbb{H})$  be a homomorphism of hyperalgebras. The composite

$$\psi: \text{hy}(\mathbb{G}) \xrightarrow{\Delta} \text{hy}(\mathbb{G}) \otimes \text{hy}(\mathbb{G}) \xrightarrow{1 \otimes \omega} \text{hy}(\mathbb{G}) \otimes \text{hy}(\mathbb{H}) = \text{hy}(\mathbb{G} \times \mathbb{H})$$

is an injective homomorphism of hyperalgebras. Put  $\mathbb{R} = \mathbb{G} \times \mathbb{H}$ ,  $J = \text{Im}(\psi)$  and  $J' = [J, J]$ . Then  $J'$  is closed in  $\text{hy}(\mathbb{R})$  by 0.3.4(f) and  $J/J' \simeq \text{hy}(\mathbb{G}/[\mathbb{G}, \mathbb{G}])$  (0.3.2(b)) is finite dimensional by assumption and hence  $J$  is closed in  $\text{hy}(\mathbb{G} \times \mathbb{H})$  by 0.3.4(b). There exists a unique connected closed subgroup scheme  $\mathbb{G}^*$  of  $\mathbb{G} \times \mathbb{H}$  with  $\text{hy}(\mathbb{G}^*) = J$ . The projection  $\text{pr}_1: \mathbb{G}^* \rightarrow \mathbb{G}$ , which is then an etale group covering of  $\mathbb{G}$ , is an isomorphism because  $\mathbb{G}$  is (SC). The composite

$$\mathbb{f}: \mathbb{G} \xrightarrow{(\text{pr}_1)^{-1}} \mathbb{G}^* \xrightarrow{\text{pr}_2} \mathbb{H}$$



is easily seen to be a unique homomorphism of  $k$ -group schemes with  $\text{hy}(\mathfrak{f}) = \omega$ . Q.E.D.

**1.9. Theorem.** *If  $k$  is perfect and  $p > 0$ , a connected affine algebraic  $k$ -group scheme  $\mathcal{G}$  is (SC) if and only if  $\mathcal{O}(\mathcal{G}) \simeq \text{hy}(\mathcal{G})^0$ . Then  $\mathcal{G}/[\mathcal{G}, \mathcal{G}]$  is finite.*

**Proof.** By virtue of Lemma 1.8, it suffices to show that  $\mathcal{G}/[\mathcal{G}, \mathcal{G}]$  is finite when  $\mathcal{G}$  is (SC). Suppose  $\mathcal{G}$  is (SC) and  $p > 0$ . We can assume that  $\mathcal{G}$  is smooth. Let  $\mathcal{G}_u$  denote the unipotent radical of  $\mathcal{G}$ . Since  $\mathcal{G}/\mathcal{G}_u$  is semisimple (1.6), it follows that  $\mathcal{G} = [\mathcal{G}, \mathcal{G}]\mathcal{G}_u$ . Hence  $\mathcal{G}/[\mathcal{G}, \mathcal{G}]$  is (SC), unipotent and smooth. Since it has a central series of closed connected subgroups each of whose quotients is isomorphic with  $\mathcal{G}_a$  [DG, IV, §2, 3.9], it follows from Remark 1.4 that  $\mathcal{G} = [\mathcal{G}, \mathcal{G}]$ . Q.E.D.

**2. The case of Chevalley group schemes.** In this section we assume  $k$  to be algebraically closed and consider the problem of group coverings for connected semisimple  $k$ -group schemes. A connected smooth affine algebraic  $k$ -group scheme is semisimple if its radical is (e). Let  $\mathcal{G}$  be a connected semisimple  $k$ -group scheme with a maximal torus  $\mathfrak{T}$ . The character group  $X = X(\mathfrak{T}) = \text{Hom}_{k\text{-gr}}(\mathfrak{T}, \mathbb{G}_m)$  is a free  $\mathbb{Z}$ -module of finite rank and has a natural root system  $\nabla$  in it. For each root  $\alpha \in \nabla$ , the associated coroot  $\alpha^*$  is uniquely determined in the space  $\hat{X} = \text{Hom}_{\mathbb{Z}}(X, \mathbb{Z})$ . Let  $\nabla^*$  denote the set of coroots  $\alpha^*$ ,  $\alpha \in \nabla$ . We then have the following subgroups of  $X_{\mathbb{Q}} = X \otimes_{\mathbb{Z}} \mathbb{Q}$ :

$$X_0 = \{\nabla\}_{\mathbb{Z}} \subset X \subset X^0 = \{\nabla^*\}_{\mathbb{Z}} = \{x \in X_{\mathbb{Q}} \mid \langle \nabla^*, x \rangle \subset \mathbb{Z}\}.$$

The group  $X^0$  is called the weight module of  $(X, \nabla)$ . The quotient group  $X^0/X_0$  is finite. See Iwahori [3, Vol. 2, p. 58] for the table of  $X^0/X_0$  for the irreducible root systems. For any subgroup  $Y$  between  $X_0$  and  $X^0$ , the pair  $(Y, \nabla)$  is a reduced root system, say, in the sense of Satake [5, p. 44] and if we identify  $Y_{\mathbb{Q}} = X_{\mathbb{Q}}$ , then  $Y$  has the same coroot system as  $X$  and we have  $Y_0 = X_0$  and  $Y^0 = X^0$ .

Let  $\mathfrak{f}: \mathcal{G} \rightarrow \mathcal{G}'$  be an "isogeny" of connected affine algebraic  $k$ -group schemes, by which we mean an epimorphism of affine algebraic  $k$ -group schemes whose kernel  $\text{Ker}(\mathfrak{f})$  is a finite  $k$ -group scheme. If  $\mathcal{G}'$  is smooth and  $\text{Ker}(\mathfrak{f})$  is etale, then  $\mathcal{G}$  is smooth too. Suppose that  $\mathcal{G}$  and  $\mathcal{G}'$  are both smooth. If one of  $\mathcal{G}$  and  $\mathcal{G}'$  is reductive (resp. semisimple), then so is the other. Hence suppose that both  $\mathcal{G}$  and  $\mathcal{G}'$  are semisimple. If  $\mathfrak{T}$  is a maximal torus of  $\mathcal{G}$ , then  $\mathfrak{T}' = \mathfrak{f}(\mathfrak{T})$  is a maximal torus of  $\mathcal{G}'$  and the restricted isogeny  $\mathfrak{f}: \mathfrak{T} \rightarrow \mathfrak{T}'$  induces an injection of abelian groups  $\mathfrak{f}: X' = X(\mathfrak{T}) \hookrightarrow X = X(\mathfrak{T}')$ . This is a special homomorphism in the following sense:

(i)  $[X: {}^t\mathfrak{f}(X')] < \infty$ .

(ii) If  $\nabla$  and  $\nabla'$  denote the root systems of  $(\mathfrak{G}, \mathfrak{T})$  and  $(\mathfrak{G}', \mathfrak{T}')$  respectively, then there are a bijection  $\beta: \nabla \xrightarrow{\sim} \nabla'$  and a family  $(q_\alpha)_{\alpha \in \nabla}$  of powers of  $p^*$  with  ${}^t\mathfrak{f}(\beta(\alpha)) = q_\alpha \alpha$  for all  $\alpha \in \nabla$ .

The famous uniqueness theorem of Chevalley [5, p. 53] tells us conversely that each special homomorphism  $\psi: X' \hookrightarrow X$  comes from an isogeny  $\mathfrak{f}: \mathfrak{G} \rightarrow \mathfrak{G}'$  such that  $\mathfrak{f}(\mathfrak{T}) = \mathfrak{T}'$ , which is determined uniquely up to inner automorphisms by the  $k$ -rational points of  $\mathfrak{T}$ . On the other hand for each reduced root system  $(X, \nabla)$  there exists a connected semisimple  $k$ -group scheme  $(\mathfrak{G}, \mathfrak{T})$ , determined uniquely up to isomorphisms, having  $(X, \nabla)$  as its root system (the existence theorem of Chevalley). The connected semisimple  $k$ -group scheme determined by the root system  $(X, \nabla)$  is denoted by  $\mathfrak{G}(X, \nabla)$ .

An isogeny of connected semisimple  $k$ -group schemes  $\mathfrak{f}: \mathfrak{G} \rightarrow \mathfrak{G}'$  (or the corresponding special homomorphism  ${}^t\mathfrak{f}: X' \hookrightarrow X$ ) is called *standard* if all the indices  $q_\alpha$  are equal to 1. The following facts concerning standard isogenies are well known. Recall that  $\mathfrak{D}(\Gamma)$  denotes the *diagonalizable*  $k$ -group scheme represented by a finitely generated  $\mathbb{Z}$ -module  $\Gamma$  (§0.2).

**2.1. Lemma.** *Let  $\mathfrak{f}: \mathfrak{G} \rightarrow \mathfrak{G}'$  be a standard isogeny of connected semisimple  $k$ -group schemes,  $\mathfrak{T}$  a maximal torus of  $\mathfrak{G}$ ,  $\mathfrak{T}' = \mathfrak{f}(\mathfrak{T})$ ,  $X = X(\mathfrak{T})$  and  $X' = X(\mathfrak{T}')$ . Via the injection  ${}^t\mathfrak{f}: X' \hookrightarrow X$ , we view  $X'$  as a subgroup of  $X$ .*

(1)  $\mathfrak{Rer}(\mathfrak{f}) = \mathfrak{Rer}(\mathfrak{f}|_{\mathfrak{T}}) = \mathfrak{D}(X/X')$ .

(2)  $\mathfrak{Rer}(\mathfrak{f}) \subset \mathfrak{Cent}(\mathfrak{G})$ .

(3)  $\mathfrak{Cent}(\mathfrak{G}) = \mathfrak{D}(X/X_0)$ .

(4)  $\mathfrak{Rer}(\mathfrak{f})$  is etale if and only if  $(p^*, [X: X']) = 1$ . Hence  $(\mathfrak{G}, \mathfrak{f})$  is a  $p$ -etale group covering of  $\mathfrak{G}'$  in this case.

(5) If  $(\mathfrak{H}, \gamma)$  is an etale group covering of  $\mathfrak{G}'$ , then  $\mathfrak{H}$  is smooth connected semisimple and  $\gamma$  a standard isogeny.

Let  $(\mathfrak{G}, \mathfrak{T})$  be a connected semisimple  $k$ -group scheme having  $(X, \nabla)$  as its root system. Let  $\overline{X}/X$  denote the largest subgroup of  $X^0/X$  whose order is relatively prime to  $p^*$ . Hence  $[X^0: \overline{X}] =$  a power of  $p^*$ .

**2.2. Theorem.** *Let  $k$  be algebraically closed. (i) The standard isogeny  $\mathfrak{G}(\overline{X}, \nabla) \rightarrow \mathfrak{G}(X, \nabla) = \mathfrak{G}$  induced from the inclusion of root systems  $(X, \nabla) \hookrightarrow (\overline{X}, \nabla)$  is a universal group covering, as well as a  $p$ -universal group covering, of  $\mathfrak{G}$ .*

(ii) *In particular the following conditions are equivalent to each other:*

(a)  $\mathfrak{G}$  is (SC); (b)  $\mathfrak{G}$  is (SC) $_p$ ; (c)  $X = \overline{X}$ ; (d)  $[X^0: X] =$  a power of  $p^*$ .

**Proof.** This theorem follows directly from Lemma 2.1, since every etale

group covering of  $\mathfrak{G}$  is (essentially) obtained as the standard isogeny  $\mathfrak{G}(Y, \nabla) \rightarrow \mathfrak{G}(X, \nabla)$  induced from the inclusion  $(X, \nabla) \hookrightarrow (Y, \nabla)$  for some subgroup  $X \subset Y \subset X^0$  with  $(p^*, [Y: X]) = 1$ . Q.E.D.

2.3. Corollary. If  $k$  is algebraically closed, for a connected semisimple  $k$ -group scheme  $\mathfrak{G}$ , the following facts hold:

- (1) The dual Hopf algebra  $\text{hy}(\mathfrak{G})^0$  is finitely generated.
- (2)  $\bar{\mathfrak{G}} = \text{Spec}(\text{hy}(\mathfrak{G})^0)$  is smooth connected semisimple and (SC).
- (3) The canonical epimorphism  $\bar{\mathfrak{G}} \rightarrow \mathfrak{G}$  is a  $p$ -etale group covering.
- (4)  $\text{hy}(\bar{\mathfrak{G}}) \simeq \text{hy}(\mathfrak{G})$ .
- (5)  $\text{hy}(\mathfrak{G})^0 // \mathcal{O}(\mathfrak{G})$  is finite dimensional over  $k$ .
- (6)  $\bar{\mathfrak{G}}$  is a universal group covering, as well as a  $p$ -universal group covering, of  $\mathfrak{G}$ .

This follows directly from Lemma 1.8 and Theorem 2.2.

3. The case of semisimple group schemes. In this section we generalize the results of §2 to the case of perfect ground field. Suppose  $k$  is perfect throughout §3. Let  $\mathfrak{G}$  be a connected smooth semisimple affine algebraic  $k$ -group scheme. This means that the  $\bar{k}$ -group scheme  $\mathfrak{G} \otimes \bar{k}$  is a Chevalley  $\bar{k}$ -group scheme, where  $\bar{k}$  = the algebraic closure of  $k$ . But of course  $\mathfrak{G}$  itself is not necessarily of Chevalley type.

If  $H$  is an arbitrary Hopf algebra over  $k$ , the dual  $\bar{k}$ -Hopf algebra  $(H \otimes \bar{k})^0$  contains  $H^0 \otimes \bar{k}$  as a  $\bar{k}$ -sub-Hopf algebra. If  $H$  is cocommutative and  $(H \otimes \bar{k})^0$  is finitely generated over  $\bar{k}$ , then it follows from 0.1.1(a) that  $H^0$  is finitely generated over  $k$ . In particular put  $H = \text{hy}(\mathfrak{G})$  the hyperalgebra of  $\mathfrak{G}$ . Then  $H \otimes \bar{k} = \text{hy}_{\bar{k}}(\mathfrak{G} \otimes \bar{k})$  is the  $\bar{k}$ -hyperalgebra of the  $\bar{k}$ -group scheme  $\mathfrak{G} \otimes \bar{k}$ . Since  $(H \otimes \bar{k})^0$  is finitely generated over  $\bar{k}$  by Corollary 2.3(1), it follows that  $H^0$  is finitely generated over  $k$ . Let  $\bar{\mathfrak{G}} = \text{Spec}(H^0)$  and  $(\mathfrak{G} \otimes \bar{k})^* = \text{Spec}_{\bar{k}}((H \otimes \bar{k})^0)$ . The canonical inclusions of  $\bar{k}$ -Hopf algebras

$$\mathcal{O}(\mathfrak{G}) \otimes \bar{k} \hookrightarrow H^0 \otimes \bar{k} \hookrightarrow (H \otimes \bar{k})^0$$

induce epimorphisms of affine algebraic  $\bar{k}$ -group schemes

$$\mathfrak{G} \otimes \bar{k} \leftarrow \bar{\mathfrak{G}} \otimes \bar{k} \leftarrow (\mathfrak{G} \otimes \bar{k})^*$$

whose composite is a universal, as well as a  $p$ -universal, group covering of  $\mathfrak{G} \otimes \bar{k}$  by Corollary 2.3(6). Applying the functor  $\text{hy}_{\bar{k}}(?)$ , one can easily conclude that the canonical epimorphism of affine algebraic  $k$ -group schemes  $\bar{\mathfrak{G}} \rightarrow \mathfrak{G}$  (induced from  $\mathcal{O}(\mathfrak{G}) \hookrightarrow H^0$ ) is a  $p$ -etale group covering.

This means in particular that  $\text{hy}(\bar{\mathfrak{G}}) \simeq \text{hy}(\mathfrak{G})$ , so  $\mathcal{O}(\bar{\mathfrak{G}}) \simeq \text{hy}(\bar{\mathfrak{G}})^0$ . Therefore  $\bar{\mathfrak{G}}$  is (SC) by Lemma 1.8. It follows that  $\bar{\mathfrak{G}}$  is a universal, as well as a

$p$ -universal, group covering of  $\mathcal{G}$ . Notice that  $\overline{\mathcal{G}}$  is connected smooth *semi-simple* (since isogenous to  $\mathcal{G}$ ).

In particular if  $\mathcal{G}$  is  $(SC)_p$ , then  $\overline{\mathcal{G}} \simeq \mathcal{G}$ , so  $\mathcal{G}$  is  $(SC)$ . Summarizing we have

**3.1. Theorem.** *Suppose  $k$  is perfect and let  $\mathcal{G}$  be a connected smooth semisimple affine algebraic  $k$ -group scheme. Then the following facts hold.*

- (1) *The dual Hopf algebra  $\text{hy}(\mathcal{G})^0$  is finitely generated.*
- (2)  *$\overline{\mathcal{G}} = \text{Spec}(\text{hy}(\mathcal{G})^0)$  is smooth connected semisimple and  $(SC)$ .*
- (3) *The canonical epimorphism  $\overline{\mathcal{G}} \rightarrow \mathcal{G}$  is a  $p$ -etale group covering.*
- (4)  *$\text{hy}(\overline{\mathcal{G}}) \simeq \text{hy}(\mathcal{G})$ .*
- (5)  *$\text{hy}(\mathcal{G})^0/\mathcal{O}(\mathcal{G})$  is finite dimensional over  $k$ .*
- (6)  *$\mathcal{G}$  is a universal, as well as a  $p$ -universal, group covering of  $\overline{\mathcal{G}}$ .*
- (7)  *$\mathcal{G}$  is  $(SC)_p$  if and only if  $(SC)$ .*

The purpose of the rest of this section is to prove that the canonical epimorphism  $(\mathcal{G} \otimes \overline{k})^* \rightarrow \overline{\mathcal{G}} \otimes \overline{k}$  is isomorphic.

In general if  $\overline{\mathcal{H}}$  is a  $\overline{k}$ -group scheme, we mean by a  $k$ -form of  $\overline{\mathcal{H}}$ , a  $k$ -group scheme  $\mathcal{H}$  with  $\mathcal{H} \otimes \overline{k} \simeq \overline{\mathcal{H}}$ . If  $\overline{f}: \overline{\mathcal{H}} \rightarrow \overline{\mathcal{G}} \otimes \overline{k}$  is a homomorphism of  $\overline{k}$ -group schemes, we say that  $\overline{f}$  is *defined over  $k$*  if there are a  $k$ -form  $\mathcal{H}$  of  $\overline{\mathcal{H}}$  and a homomorphism of  $k$ -group schemes  $f: \mathcal{H} \rightarrow \mathcal{G}$  with

$$\overline{f}: \overline{\mathcal{H}} \simeq \mathcal{H} \otimes \overline{k} \xrightarrow{f \otimes \overline{k}} \mathcal{G} \otimes \overline{k}.$$

**3.2. Lemma.** *The canonical epimorphism of  $\overline{k}$ -group schemes,  $(\mathcal{G} \otimes \overline{k})^* \rightarrow \overline{\mathcal{G}} \otimes \overline{k}$ , is defined over some finite extension field  $l$  of  $k$ .*

**Proof.** Let  $k_0$  be the prime field in  $k$ . Each Chevalley  $\overline{k}$ -group scheme  $\mathcal{G}(X, \nabla)$  has a canonical  $k_0$ -form [5, p. 53] and the standard isogeny  $\mathcal{G}(\overline{X}, \nabla) \rightarrow \mathcal{G}(X, \nabla)$  induced from the inclusion  $X \hookrightarrow \overline{X}$  can be taken to be defined over  $k_0$  [5, p. 60]. Now that  $\mathcal{G}$  is semisimple, there is a unique root system  $(X, \nabla)$  with  $\mathcal{G} \otimes \overline{k} \simeq \mathcal{G}(X, \nabla)$ . By the universal mapping property of universal group coverings, we have a commutative diagram

$$\begin{array}{ccc} (\mathcal{G} \otimes \overline{k})^* & \simeq & \mathcal{G}(\overline{X}, \nabla) \\ \text{cano} \downarrow & & \downarrow \text{cano} \\ \mathcal{G} \otimes \overline{k} & \simeq & \mathcal{G}(X, \nabla) \end{array}$$

where the right vertical arrow is defined over  $k_0$ . Since  $\mathcal{G}$  and  $\mathcal{G}(X, \nabla)$  are both algebraic, there is a finite extension  $l/k$  such that the isomorphism  $\mathcal{G} \otimes \overline{k} \simeq \mathcal{G}(X, \nabla)$  is defined over  $l$ . The lemma follows from this directly. Q.E.D.

If we put  $H = \text{hy}(\mathbb{G})$  as before, the above lemma implies that there is an  $l$ -sub-Hopf algebra  $A$  of  $(H \otimes_k \bar{k})^0$  with

$$\mathcal{O}(\mathbb{G}) \otimes_k l \subset A \quad \text{and} \quad A \otimes_l \bar{k} = (H \otimes_k \bar{k})^0.$$

If we put  $\mathcal{Q} = \text{Spec}_l(A)$ , then the canonical epimorphism of  $l$ -group schemes  $\mathcal{Q} \rightarrow \mathbb{G} \otimes_k l$  is an etale group covering, since so is

$$(\mathbb{G} \otimes_k \bar{k})^* = \mathcal{Q} \otimes_l \bar{k} \rightarrow (\mathbb{G} \otimes_k l) \otimes_l \bar{k} = \mathbb{G} \otimes_k \bar{k}.$$

Hence  $\text{hy}_l(\mathcal{Q}) \simeq H \otimes_k l$ , so  $\text{hy}_l(\mathcal{Q})^0 \simeq H^0 \otimes_k l$ . (In general, if  $l/k$  is finite,  $(B \otimes_k l)^0 = B^0 \otimes_k l$  for all  $k$ -algebras  $B$ .) Thus we get a commutative diagram:

$$\begin{array}{ccc} \text{hy}_l(\mathcal{Q})^0 & \simeq & H^0 \otimes_k l \\ \cup & & \cup \\ A & \supset & \mathcal{O}(\mathbb{G}) \otimes_k l. \end{array}$$

Applying the functor  $?\otimes_l \bar{k}$ , we obtain

$$\begin{array}{ccc} \text{hy}_l(\mathcal{Q})^0 \otimes_l \bar{k} & \simeq & H^0 \otimes_k \bar{k} \\ \cup & & \cup \\ A \otimes_l \bar{k} & \supset & \mathcal{O}(\mathbb{G}) \otimes_k \bar{k}. \end{array}$$

But since  $A \otimes_l \bar{k} = (H \otimes_k \bar{k})^0$ , this means that  $(H \otimes_k \bar{k})^0 = H^0 \otimes_k \bar{k}$  and hence that  $(\mathbb{G} \otimes_k \bar{k})^* \simeq \mathbb{G} \otimes_k \bar{k}$ . Thus we proved

**3.3. Theorem.** *Suppose  $k$  is perfect. Let  $\mathbb{G}$  be a connected smooth semi-simple affine algebraic  $k$ -group scheme. If  $\bar{\mathbb{G}}$  is the universal group covering of  $\mathbb{G}$ , then  $\bar{\mathbb{G}} \otimes_k \bar{k}$  is the universal group covering of the  $\bar{k}$ -scheme  $\mathbb{G} \otimes_k \bar{k}$ . In particular  $\mathbb{G}$  is (SC) (or equivalently  $(SC)_p$ ) if and only if  $\mathbb{G} \otimes_k \bar{k}$  is (SC) (or equivalently  $(SC)_p$ ) as a  $\bar{k}$ -group scheme.*

**4. Semisimple hyperalgebras.** In this section assume  $k$  is perfect. We prove that every semisimple hyperalgebra of finite type is the hyperalgebra of some connected semisimple  $k$ -group scheme. Since connected semisimple  $k$ -group schemes have the universal group covering, it will follow from Theorem 3.1 that the category of (SC) semisimple  $k$ -group schemes is equivalent to the category of semisimple hyperalgebras of finite type.

A hyperalgebra of finite type  $J$  is called *representable* if the dual Hopf algebra  $J^0$  is dense in  $J^*$ , or equivalently if there is an affine algebraic  $k$ -group scheme  $\mathbb{G}$  with  $J \subset \text{hy}(\mathbb{G})$ . Here the  $k$ -group scheme  $\mathbb{G}$  can be taken so that  $J$  is dense in  $\text{hy}(\mathbb{G})$  (i.e.,  $\mathcal{O}(\mathbb{G}) \subset J^0$ ) and hence in particular that  $\mathbb{G}$  is connected. Each finite type hyperalgebra  $J$  contains the *smallest normal sub-hyperalgebra*  $K$  such that the quotient hyperalgebra  $J//K$  is representable;

$K$  is the Hopf kernel of  $J \rightarrow J^{00}$ . We shall denote  $K = K_{\text{rep}}(J)$ . It follows from Dieudonné [1, Proposition 20, p. 365] that  $K_{\text{rep}}(J)$  is contained in the center of  $J$ .

4.1. Lemma. *Let  $J$  be a smooth hyperalgebra of finite type and  $K$  a finite normal subhyperalgebra of  $J$ . If  $J//K$  is representable, then so is  $J$ .*

Proof. We can assume  $p > 0$ . With a sufficiently large positive integer  $r$ , let  $V^r: J \rightarrow J^{(p^r)}$  denote the  $r$ -times iterated *Verschiebung* map of  $J$  (§0.1). Then  $V^r$  is surjective (§0.3) with a finite Hopf kernel containing  $K$  and hence it induces a surjective homomorphism  $J//K \rightarrow J^{(p^r)}$  which has also a finite Hopf kernel. If, in general,  $l|k$  is a field extension and if  $L$  is a representable hyperalgebra over  $k$ , then the  $l$ -hyperalgebra  $l \otimes L$  is representable too. Since now  $k$  is perfect, we have  $J = (J^{(p^r)})^{(p^{-r})}$ . Hence it suffices to show the representability of  $J^{(p^r)}$ , which will follow from the next lemma. Q.E.D.

4.2. Lemma. *If  $J$  is a representable hyperalgebra of finite type and  $K$  a finite normal subhyperalgebra of  $J$ , then the quotient  $J//K$  also is representable.*

Proof. Imbed  $J$  into  $\text{hy}(\mathfrak{G})$  as a dense subhyperalgebra for some connected affine algebraic  $k$ -group scheme  $\mathfrak{G}$ . Since  $[J, K] \subset K$ , it follows from 0.3.4(e) that  $[\text{hy}(\mathfrak{G}), K] \subset K$ . Hence  $K$  is a normal subhyperalgebra of  $\text{hy}(\mathfrak{G})$ . Since any finite dimensional subhyperalgebra of  $\text{hy}(\mathfrak{G})$  is closed (0.3.4(a)), it follows that there is a unique closed normal connected subgroup scheme  $\mathfrak{H}$  of  $\mathfrak{G}$  with  $K = \text{hy}(\mathfrak{H})$ . The induced inclusion

$$J//K \hookrightarrow \text{hy}(\mathfrak{G})//\text{hy}(\mathfrak{H}) = \text{hy}(\mathfrak{G}/\mathfrak{H})$$

proves the representability of  $J//K$ . Q.E.D.

4.3. Proposition. *Let  $J$  be a smooth hyperalgebra of finite type. Then  $K_{\text{rep}}(J)$  is smooth too.*

Proof. Let  $K_{\text{sm}}$  denote the smooth part of  $K = K_{\text{rep}}(J)$  (§0.3). Since  $K$  is central, we can consider the quotient hyperalgebra  $J//K_{\text{sm}}$  which has a finite dimensional normal subhyperalgebra  $K//K_{\text{sm}}$  (§0.3). Since the quotient  $(J//K_{\text{sm}})/(K//K_{\text{sm}}) = J//K$  is representable, it follows from Lemma 4.1 that so is  $J//K_{\text{sm}}$ . By the definition of  $K_{\text{rep}}(J)$ , this proves that  $K = K_{\text{sm}}$ . Q.E.D.

For each hyperalgebra  $J$ , we have defined the derived subhyperalgebra  $[J, J]$  in §0.1. Hence we can define the derived series  $\{J^{(\nu)}\}$  of  $J$  by  $J^{(\nu)} = [J^{(\nu-1)}, J^{(\nu-1)}]$  and  $J^{(0)} = J$ . The hyperalgebra  $J$  is called *solvable* if  $J^{(N)} = k$  for some  $N$ .

Let  $J$  be a smooth hyperalgebra of finite type. The set of normal smooth

*solvable* subhyperalgebras of  $J$  contains clearly the largest element denoted by  $\text{rad}(J)$ , which we shall call the *radical* of  $J$ .  $J$  is called *semisimple* if  $\text{rad}(J) = k$ . In general  $J//\text{rad}(J)$  is semisimple for any smooth hyperalgebra  $J$  of finite type.

**4.4. Proposition.** *Semisimple smooth hyperalgebras of finite type are representable.*

**Proof.** If  $J$  is such a hyperalgebra, then  $K_{\text{rep}}(J)$  is smooth and central by Proposition 4.3 and hence is trivial. Q.E.D.

Let  $J$  be a smooth semisimple hyperalgebra of finite type. Embed as usual  $J$  into the hyperalgebra  $\text{hy}(\mathcal{G})$  as a dense subhyperalgebra for some connected affine algebraic smooth  $k$ -group scheme  $\mathcal{G}$ . Let  $\text{rad}(\mathcal{G})$  denote the radical of  $\mathcal{G}$ .

**4.5. Lemma.** *The hyperalgebra  $J \cap \text{hy}(\text{rad}(\mathcal{G})) = K$  is finite dimensional.*

**Proof.**  $K$  is clearly normal solvable in  $J$ . Since  $[J, K] \subset K$ , it follows from  $J = J_{\text{sm}}$  that  $[J, K_{\text{sm}}] \subset K_{\text{sm}}$ , where  $(\ )_{\text{sm}}$  denotes the smooth part of the hyperalgebra. Thus  $K_{\text{sm}}$  is normal smooth solvable in  $J$  and is trivial by assumption. This means that  $K$  is finite dimensional. Q.E.D.

Notice that  $J//K \subset \text{hy}(\mathcal{G}/\text{rad}(\mathcal{G}))$  is also dense. Hence it follows from 0.3.4(f) that

$$[J//K, J//K] = [\text{hy}(\mathcal{G}/\text{rad}(\mathcal{G})), \text{hy}(\mathcal{G}/\text{rad}(\mathcal{G}))].$$

Since  $\mathcal{G}/\text{rad}(\mathcal{G})$  is a semisimple  $k$ -group scheme, it coincides with the derived group. This means that

$$[J//K, J//K] = \text{hy}(\mathcal{G}/\text{rad}(\mathcal{G}))$$

and hence, in particular, that  $J//K = [J//K, J//K]$ , or equivalently that  $J = [J, J]K$ . Then the quotient hyperalgebra  $J/[J, J]$  is finite and smooth and therefore  $J = [J, J]$ . This proves that  $J$  is a *closed* subhyperalgebra of  $\text{hy}(\mathcal{G})$  (0.3.4(g)). Since  $J$  is dense in  $\text{hy}(\mathcal{G})$ , it follows that  $J = \text{hy}(\mathcal{G})$ . Now  $\text{hy}(\text{rad}(\mathcal{G}))$  is a normal smooth solvable subhyperalgebra of  $J$ . Since  $J$  is semisimple, it follows that  $\text{hy}(\text{rad}(\mathcal{G})) = k$  and hence that  $\text{rad}(\mathcal{G}) = (e)$  or that  $\mathcal{G}$  is *semisimple*. Thus we have proved

**4.6. Theorem.** *If  $k$  is perfect each smooth semisimple hyperalgebra of finite type is the hyperalgebra of some connected semisimple  $k$ -group scheme.*

Conversely we have

**4.7. Proposition.** *Let  $\mathcal{G}$  be a connected smooth affine algebraic  $k$ -group scheme with radical  $\text{rad}(\mathcal{G})$ . Then  $\text{rad}(\text{hy}(\mathcal{G})) = \text{hy}(\text{rad}(\mathcal{G}))$ . In particular*

the  $k$ -group scheme  $\mathcal{G}$  is semisimple if and only if the hyperalgebra  $\text{hy}(\mathcal{G})$  is.

**Proof.** The inclusion  $\text{hy}(\text{rad}(\mathcal{G})) \subset \text{rad}(\text{hy}(\mathcal{G}))$  is clear. Let  $K$  be a normal smooth solvable subhyperalgebra of  $\text{hy}(\mathcal{G})$  and  $A(K)$  the algebraic hull of  $K$  in  $\text{hy}(\mathcal{G})$  (§0.3). Since  $K$  is contained in the smooth part  $A(K)_{\text{sm}}$ , which is also closed, it follows that  $A(K)$  is smooth. Since  $[A(K), A(K)] = [K, K]$ , the solvability of  $A(K)$  follows. Similarly the equality  $[\text{hy}(\mathcal{G}), A(K)] = [\text{hy}(\mathcal{G}), K]$  (0.3.4(d)) implies that  $A(K)$  is normal in  $\text{hy}(\mathcal{G})$ . Let  $\mathcal{H}$  be a unique closed connected normal subgroup scheme of  $\mathcal{G}$  with  $\text{hy}(\mathcal{H}) = A(K)$ . Then  $\mathcal{H}$  is solvable smooth and hence contained in  $\text{rad}(\mathcal{G})$ . Therefore  $K \subset \text{hy}(\text{rad}(\mathcal{G}))$ .

Q.E.D.

By virtue of Theorem 3.1, Theorem 4.6 has the following corollary:

**4.8. Theorem.** *Suppose  $k$  is perfect. If  $J$  is a smooth semisimple hyperalgebra of finite type, then the dual Hopf algebra  $J^0$  is finitely generated and the corresponding affine  $k$ -group scheme  $\text{Spec}(J^0)$  is (SC) semisimple and has  $J$  as its hyperalgebra. The functors  $J \mapsto \text{Spec}(J^0)$  and  $\mathcal{G} \mapsto \text{hy}(\mathcal{G})$  give rise to an equivalence between the categories of smooth semisimple hyperalgebras of finite type and of (SC) semisimple  $k$ -group schemes.*

**5. An example of nonreductive (SC)  $k$ -group schemes.** To conclude this paper we shall provide a simple example of (SC) affine algebraic  $k$ -group schemes which are *not reductive*.

In this section we shall assume  $k = \bar{k}$  with  $p > 0$ . For each left finitely generated  $k[F]$ -module  $M$ , let  $\mathcal{U}(M) = \text{Spec}(U^{[p]}(M))$  denote the corresponding commutative unipotent affine algebraic  $k$ -group scheme killed by the Verschiebung map (see §0.2). Let  $\mathcal{G}$  denote an arbitrary affine algebraic  $k$ -group scheme.

**5.1. Definition.** Let  $M$  be a finitely generated left  $k[F]$ -module. View the affine ring  $\mathcal{O}(\mathcal{G})$  as a left  $k[F]$ -module via  $Fa = a^p$ ,  $a \in \mathcal{O}(\mathcal{G})$ . A right  $\mathcal{O}(\mathcal{G})$ -comodule structure on  $M$ ,  $\rho: M \rightarrow M \otimes \mathcal{O}(\mathcal{G})$ , which is  $k[F]$ -linear, is said to be compatible with the  $k[F]$ -module structure. A left  $k[F]$ - $\mathcal{G}$ -module means a finitely generated left  $k[F]$ -module with a right  $\mathcal{O}(\mathcal{G})$ -comodule structure compatible with the  $k[F]$ -module structure.

**5.2. Lemma.** *The  $k$ -group scheme actions as automorphisms of  $k$ -group schemes  $\mathcal{U}(M) \times \mathcal{G} \rightarrow \mathcal{U}(M)$  correspond bijectively with the right  $\mathcal{O}(\mathcal{G})$ -comodule structures on  $M$  which are compatible with the  $k[F]$ -module structure.*

**Proof.** The  $k$ -group scheme actions of the above mentioned type can be identified with those right comodule structures



$$\rho: U^{[p]}(M) \rightarrow U^{[p]}(M) \otimes \mathcal{O}(\overline{\mathcal{G}})$$

which are compatible with the Hopf algebra structure on  $U^{[p]}(M)$ . This means in particular that  $\rho(M) \subset M \otimes \mathcal{O}(\overline{\mathcal{G}})$  and that the restricted coaction  $\rho|_M$  is compatible with the  $k[F]$ -module structure. Conversely if  $M$  is a left  $k[F]$ - $\overline{\mathcal{G}}$ -module with coaction  $\sigma: M \rightarrow M \otimes \mathcal{O}(\overline{\mathcal{G}})$  then the extended algebra map  $\sigma: U^{[p]}(M) \rightarrow U^{[p]}(M) \otimes \mathcal{O}(\overline{\mathcal{G}})$  is easily seen to be compatible with the Hopf algebra structure of  $U^{[p]}(M)$ . Q.E.D.

In the following let  $M$  be a (finitely generated) left  $k[F]$ - $\overline{\mathcal{G}}$ -module such that  $\mathcal{U}(M)$  is *connected* (see §0.2). The isomorphism classes of exact sequences of homomorphisms of affine  $k$ -group schemes,  $1 \rightarrow \mathcal{U}(M) \rightarrow \mathcal{G} \rightarrow \overline{\mathcal{G}} \rightarrow 1$ , the right action of  $\overline{\mathcal{G}}$  on  $\mathcal{U}(M)$  induced from which coincides with the original action, form an abelian group in a usual manner, which we shall write as  $\text{Ext}(\overline{\mathcal{G}}, \mathcal{U}(M))$ . The neutral element is supplied by the semidirect product  $\overline{\mathcal{G}} \times_s \mathcal{U}(M)$ .

**5.3. Definition.** The  $k[F]$ - $\overline{\mathcal{G}}$ -module  $M$  is of type (#) if the following condition is satisfied:

Let  $k = k[F]/(F - 1)$  be a trivial left  $k[F]$ - $\overline{\mathcal{G}}$ -module. Each exact sequence of  $k[F]$ - $\overline{\mathcal{G}}$ -modules of the form  $0 \rightarrow M \rightarrow N \rightarrow k \rightarrow 0$  splits as  $k[F]$ -modules.

**5.4. Proposition.** If  $\overline{\mathcal{G}}$  is (SC) and the  $k[F]$ - $\overline{\mathcal{G}}$ -module  $M$  is of type (#), then each  $k$ -group scheme  $\mathcal{G}$  in  $\text{Ext}(\overline{\mathcal{G}}, \mathcal{U}(M))$  is (SC) too.

**Proof.** Let  $\mathcal{G} \in \text{Ext}(\overline{\mathcal{G}}, \mathcal{U}(M))$ ,  $(\mathcal{H}, \gamma)$  an etale group covering of  $\mathcal{G}$ , and  $\mathcal{H}_u$  the connected component of  $\gamma^{-1}(\mathcal{U}(M))$ . We have a commutative diagram:

$$\begin{array}{ccccccc} 1 & \rightarrow & \mathcal{H}_u & \rightarrow & \mathcal{H} & \rightarrow & \mathcal{H}/\mathcal{H}_u \rightarrow 1 \\ & & \alpha \downarrow & & \gamma \downarrow & & \beta \downarrow \\ 1 & \rightarrow & \mathcal{U}(M) & \rightarrow & \mathcal{G} & \rightarrow & \overline{\mathcal{G}} \rightarrow 1 \end{array}$$

where the homomorphisms  $\alpha$ ,  $\beta$  and  $\gamma$  are all etale group coverings. Since  $\overline{\mathcal{G}}$  is (SC),  $\beta$  is an isomorphism, via which  $\overline{\mathcal{G}}$  operates on  $\mathcal{H}_u$  which is commutative unipotent and killed by the Verschiebung, because  $\text{hy}(\alpha): \text{hy}(\mathcal{H}_u) \simeq \text{hy}(\mathcal{U}(M))$ . Hence we can write  $\mathcal{H}_u = \mathcal{U}(N)$  for some left  $k[F]$ - $\overline{\mathcal{G}}$ -module  $N$ . The homomorphism  $\alpha$  induces an injection of  $k[F]$ - $\overline{\mathcal{G}}$ -modules  $M \hookrightarrow N$  and the quotient module  $N/M$ , which represents the kernel of  $\alpha$ , which is etale, is therefore a direct sum of some finite number of copies of the trivial  $k[F]$ - $\overline{\mathcal{G}}$ -module  $k = k[F]/(F - 1)$ . Say  $N/M \simeq k^r$ . The exact sequence of  $k[F]$ - $\overline{\mathcal{G}}$ -modules  $0 \rightarrow M \rightarrow N \rightarrow k^r \rightarrow 0$  splits as  $k[F]$ -modules, because  $M$  is of type (#). But since  $\mathcal{U}(N)$  is connected, this means that  $r = 0$  and hence that  $\alpha$  is an isomorphism. Hence  $\gamma$  is an isomorphism and  $\mathcal{G}$  is (SC). Q.E.D.

Suppose that the affine algebraic  $k$ -group scheme  $\overline{\mathfrak{G}}$  is connected smooth and put  $A = \mathcal{O}(\overline{\mathfrak{G}})$ . Let  $V$  be a finite dimensional right  $A$ -comodule with the structure map  $\rho: V \rightarrow V \otimes A$  and put  $M = k[F] \otimes V$ . Extending  $\rho$   $k[F]$ -linearly, we make  $M$  into a  $k[F]$ - $\overline{\mathfrak{G}}$ -module.

**5.5. Proposition.** *Suppose that  $\overline{\mathfrak{G}}$  is connected smooth. Then the  $k[F]$ - $\overline{\mathfrak{G}}$ -module  $M = k[F] \otimes V$  is of type (#) if and only if  $V^{\overline{\mathfrak{G}}} = 0$ .*

**Proof.** The 'only if' part is easy. Suppose that  $V^{\overline{\mathfrak{G}}} = 0$ . Let  $V \otimes_i A$  be the tensor product of  $V$  and  $A$  over  $k$  with the defining relation

$$\lambda v \otimes_i a = v \otimes_i \lambda^{b^i} a \quad \text{for } \lambda \in k, v \in V, a \in A.$$

The space  $M \otimes A$  is the direct sum of  $kF^i \otimes (V \otimes_i A)$ ,  $i \geq 0$ . Since  $A$  is smooth, the  $k$ -linear maps

$$V \otimes_i A \hookrightarrow V \otimes_{i+j} A, \quad v \otimes_i a \mapsto (v \otimes_i a)^{(j)} \stackrel{\text{def}}{=} v \otimes_{i+j} a^{b^j}$$

are injective. Let  $0 \rightarrow M \rightarrow N \rightarrow k[F]/(F - 1) \rightarrow 0$  be an exact sequence of  $k[F]$ - $\overline{\mathfrak{G}}$ -modules. Thus we can write  $N = M \oplus ke$  with  $Fe - e = \sum_i F^i \otimes v_i$  ( $v_i \in V$ ) and  $\rho(e) - e \otimes 1 = \sum_i F^i \otimes x_i$  ( $x_i \in V \otimes_i A$ ). Since  $\rho$  is  $k[F]$ -linear, it follows that  $x_i = \partial(v_i)^{(i)} + x_{i-1}^{(1)}$  for  $i > 0$  and  $x_0 = \partial(v_0)$ , where  $\partial(v) = v \otimes 1 - \rho(v) \in V \otimes A (= V \otimes_0 A)$  for  $v \in V$ . Hence if we put  $u_i = v_i + \dots + v_0 \in V$ , then  $x_i = \partial(u_i)^{(i)}$ . Since the  $k$ -linear map  $\partial: V \rightarrow V \otimes A$  is injective by assumption, it follows that  $u_i$  are equal to zero for almost all  $i$ . Hence we can well define an element of  $M$   $m = -\sum F^i \otimes u_i$ . It is easy to see  $Fm - m = Fe - e$ . This means that the  $k[F]$ - $\overline{\mathfrak{G}}$ -module  $M$  is of type (#). Q.E.D.

If we take  $\overline{\mathfrak{G}}$  to be an (SC) semisimple  $k$ -group scheme and  $V$  a nontrivial irreducible  $k$ - $\overline{\mathfrak{G}}$ -module, then  $V^{\overline{\mathfrak{G}}} = 0$  clearly and hence each element of  $\text{Ext}(\overline{\mathfrak{G}}, \mathfrak{D}_a(V))$  provides an example of nonreductive (SC)  $k$ -group schemes, where  $\mathfrak{D}_a(V) = \text{Spec}(S(V))$  represented by the symmetric algebra  $S(V)$  on  $V$ , which is a Hopf algebra having  $V$  as primitive elements.

**Added in proof.** All subhyperalgebras of a finite type hyperalgebra is of finite type by definition. Each quotient hyperalgebra, which must be of the form  $J//K$  with  $K$  a normal subhyperalgebra by (0.1.2), is also of finite type, when  $J$  is [T<sub>II</sub>, 5.5.2.1]. This is implicitly used in §4.

REFERENCES

[DG] M. Demazure and P. Gabriel, *Groupes algébriques*. Tome I: *Géométrie algébrique, généralités, groupes commutatifs*, North-Holland, Amsterdam, 1970.  
 [T<sub>I</sub>] M. Takeuchi, *Tangent coalgebras and hyperalgebras*. I, Japan. J. Math. 42 (1974), 1-43.  
 [T<sub>II</sub>] ———, *Tangent coalgebras and hyperalgebras*. II, Mem. Amer. Math. Soc (submitted).  
 1. J. Dieudonné, *Lie groups and Lie hyperalgebras over a field of characteristic  $p > 0$* . VI, Amer. J. Math. 79 (1957), 331-388. MR 20 #931.

2. G. Hochschild, *Algebraic groups and Hopf algebras*, Illinois J. Math. 14 (1970), 52–65. MR 41 #1742.
3. N. Iwahori, *The theory of Lie algebras and Chevalley groups*, Seminar Notes 12, 13, University of Tokyo, Tokyo, 1965. (Japanese)
4. M. Miyanishi, *Une caractérisation d'un groupe algébrique simplement connexe*, Illinois J. Math. 16 (1972), 639–650. MR 46 #9052.
5. I. Satake, *Classification theory of semi-simple algebraic groups*, Lecture Notes in Pure and Appl. Math., 3, Dekker, New York, 1971. MR 47 #5135.
6. M. E. Sweedler, *Hopf algebras*, Math. Lecture Note Series, Benjamin, New York, 1969. MR 40 #5705.
7. M. Takeuchi, *A correspondence between Hopf ideals and sub-Hopf algebras*, Manuscripta Math. 7 (1972), 251–270. MR 48 #328.

DEPARTMENT OF THE FOUNDATIONS OF MATHEMATICAL SCIENCES, TOKYO UNIVERSITY OF EDUCATION, OTSUKA, BUNKYO-KU, TOKYO, JAPAN

*Current address:* Department of Mathematics, University of Tsukuba, Ibaraki, 300-31 Japan