

ON DERIVATIVE OF ENERGY FUNCTIONAL
FOR ELASTIC BODIES
WITH CRACKS AND UNILATERAL CONDITIONS

BY

A. M. KHLUDNEV (*Lavrentyev Institute of Hydrodynamics of the Russian Academy of Sciences,
Novosibirsk 630090, Russia*),

K. OHTSUKA (*Department of Computer Science, Hiroshima Kokusai Gakuin University, Aki-ku,
Nakano, Hiroshima, 739-0321, Japan*),

AND

J. SOKOLOWSKI (*Institut Elie Cartan, Laboratoire de Mathématiques, Université Henri Poincaré
Nancy I, B.P. 239, 54506 Vandoeuvre lès Nancy Cedex, France*)

Abstract. In this paper we consider elasticity equations in a domain having a cut (a crack) with unilateral boundary conditions considered at the crack faces. The boundary conditions provide a mutual nonpenetration between the crack faces, and the problem as a whole is nonlinear. Assuming that a general perturbation of the cut is given, we find the derivative of the energy functional with respect to the perturbation parameter. It is known that a calculation of the material derivative for similar problems has the difficulty of finding boundary conditions at the crack faces. We use a variational property of the solution, thus avoiding a direct calculation of the material derivative.

There are many results related to the differentiation of the potential energy functional with respect to variable domains (see, e.g., [9, 4, 5, 16, 18, 17, 3]). The general theory of calculating material and shape derivatives in linear and nonlinear boundary value problems is developed in [6].

Derivatives of energy functionals with respect to the crack length in classical linear elasticity can be found by different ways. It is well known that the classical approach to the crack problem is characterized by the equality-type boundary conditions considered at the crack faces [13, 4, 7, 14, 12, 15]. As for the analysis of solution dependence on the shape domain for a wide class of elastic problems, we refer the reader to [8].

In the works [1, 2] the appropriate technique of finding derivatives of the energy functional with respect to the crack length for unilateral boundary conditions is proposed,

Received December 16, 1999.

2000 *Mathematics Subject Classification.* Primary 49Q12, 74G65; Secondary 65K10, 74G70, 74P10, 90C31.

E-mail address: khlud@hydro.nsc.ru

E-mail address: ohtsuka@barnard.cs.hkg.ac.jp

E-mail address: sokolows@iecn.u-nancy.fr

which can be used for a wide class of the unilateral problems. Qualitative properties of solutions (solution existence, solution regularity, dependence of solutions on parameters, etc.) in the crack problem for plates, shells, two- and three-dimensional bodies with unilateral conditions on the crack faces are analysed in [2] (see also [20, 19, 10, 11]).

1. Problem formulation. Let $D \subset R^3$ be a bounded domain with smooth boundary Γ , and $\Xi \subset D$ be a smooth two-dimensional surface. We assume that this surface can be extended up to the outer boundary Γ in such a way that D is divided into subdomains D_1 and D_2 with Lipschitz boundaries. Assume that this inner surface Ξ is described parametrically by the equations

$$x_i = x_i(y_1, y_2), \quad i = 1, 2, 3, \quad (1)$$

where (y_1, y_2) belong to the closure of an open bounded connected set $\omega \subset R^2$ having a smooth boundary γ . We suppose that the rank of the Jacobi matrix $\partial x_i / \partial y_j$ equals 2 at every point $(y_1, y_2) \in \omega \cup \gamma$, and that the map (1) is one-to-one. Let $\nu = (\nu_1, \nu_2, \nu_3)$ be a unit normal vector to Ξ , for example,

$$\nu = \frac{\frac{\partial x}{\partial y_1} \times \frac{\partial x}{\partial y_2}}{\left| \frac{\partial x}{\partial y_1} \times \frac{\partial x}{\partial y_2} \right|}.$$

Denote $\Omega = D \setminus \Xi$. In the domain Ω , we consider the following boundary value problem for finding a function $u = (u_1, u_2, u_3)$:

$$-\sigma_{ij,j} = f_i, \quad i = 1, 2, 3, \quad (2)$$

$$\sigma_{ij} = a_{ijkl}\varepsilon_{kl}, \quad i, j = 1, 2, 3, \quad (3)$$

$$u = 0 \quad \text{on } \Gamma, \quad (4)$$

$$[u]\nu \geq 0, \quad \sigma_\nu \leq 0, \quad [\sigma_\nu] = 0, \quad \sigma_\tau = 0, \quad \sigma_\nu[u]\nu = 0 \quad \text{on } \Xi. \quad (5)$$

Here $\varepsilon_{kl} = \varepsilon_{kl}(u) = \frac{1}{2}(u_{k,l} + u_{l,k})$ are strain tensor components, $u_{k,l} = \frac{\partial u_k}{\partial x_l}$; $\sigma_{ij} = \sigma_{ij}(u)$ denote the stress tensor components,

$$\{\sigma_{ij}\nu_j\}_{i=1}^3 = \sigma_\tau + \sigma_\nu\nu, \quad \sigma_\nu = \sigma_{ij}\nu_j\nu_i.$$

The brackets $[v] = v^+ - v^-$ mean the jump of v across Ξ , where v^+, v^- stand for the values of v on Ξ^+, Ξ^- , respectively, and where Ξ^+, Ξ^- are defined for a given choice of positive and negative directions of ν on Ξ . Coefficients a_{ijkl} are assumed to be constant, satisfying the usual conditions of symmetry and positive definiteness, i.e.,

$$a_{ijkl} = a_{jikl} = a_{ijlk}, \quad a_{ijkl}\xi_{kl}\xi_{ij} \geq c|\xi|^2, \quad c > 0, \quad \xi_{ij} = \xi_{ji}.$$

The function $f = (f_1, f_2, f_3) \in C^1(R^3)$ is given.

The boundary value problem (2)–(5) describes an equilibrium state of an elastic body occupying the domain Ω in its nondeformable state, and the surface Ξ corresponds to a crack in a body. Conditions (5) provide the mutual nonpenetration between the crack faces without friction [8]. Considering the problem (2)–(5), we have in mind its variational formulation. Denote

$$K_0 = \{u = (u_1, u_2, u_3) \in H^1(\Omega) \mid u = 0 \text{ on } \Gamma; [u]\nu \geq 0 \text{ on } \Xi\}.$$

Then (2)–(5) correspond to the following minimization problem:

$$\min_{u \in K_0} \left\{ \frac{1}{2} \int_{\Omega} a_{ijkl} \varepsilon_{kl}(u) \varepsilon_{ij}(u) - \int_{\Omega} f u \right\}. \quad (6)$$

By the assumptions imposed on Ω, a_{ijkl}, f , the problem (6) has a unique solution u satisfying the following variational inequality:

$$u \in K_0 : \int_{\Omega} a_{ijkl} \varepsilon_{kl}(u) (\varepsilon_{ij}(\bar{u}) - \varepsilon_{ij}(u)) \geq \int_{\Omega} f(\bar{u} - u) \quad \forall \bar{u} \in K_0. \quad (7)$$

In the paper, we consider a general perturbation of the boundary value problem (2)–(5) and find the derivative of the energy functional with respect to the perturbation parameter. Note that the obtained result holds true for other boundary conditions. For example, we may assume that $\Gamma = \Gamma_1 \cup \Gamma_2, \Gamma_1 \cap \Gamma_2 = \emptyset, \text{meas } \Gamma_1 > 0, u = 0$ on $\Gamma_1, \sigma_{ij} n_j = 0$ on Γ_2 . Here $n = (n_1, n_2, n_3)$ is the unit normal vector to Γ .

Let Ω_t be a family of domains such that, for each t , there exists a one-to-one mapping

$$y = \Phi_t(x), \quad x \in \Omega_t, \quad y \in \Omega, \quad (8)$$

with the positive Jacobian $|\frac{\partial \Phi_t}{\partial x}| > c > 0, \Phi_t = (\Phi_t^1, \Phi_t^2, \Phi_t^3)$. We assume that $\Phi_0(x) = x, \Phi \in C^2(0, T; W_{\text{loc}}^{2, \infty}(R^3))$. Let

$$x = x(t, y) = \Phi_t^{-1}(y) \quad (9)$$

be the mapping inverse to Φ_t . By fixing y in (8) and differentiating (8) with respect to t , we have

$$0 = \frac{\partial \Phi_t}{\partial t} + \frac{\partial \Phi_t}{\partial x} \frac{dx(t)}{dt},$$

whence

$$\frac{dx(t)}{dt} = - \left(\frac{\partial \Phi_t}{\partial x} \right)^{-1} \frac{\partial \Phi_t}{\partial t}. \quad (10)$$

It is clear that (10) can be viewed as a system of ordinary differential equations; thus

$$\frac{dx(t)}{dt} = V(t, x(t)), \quad (11)$$

$$x(0) = y, \quad (12)$$

where

$$V(t, x(t)) = - \left(\frac{\partial \Phi_t(x(t))}{\partial x} \right)^{-1} \frac{\partial \Phi_t(x(t))}{\partial t}, \quad (13)$$

and hence for the solution $x(t)$ of (11)–(12) we have $x(t) = x(t, y)$. Note that, by (9),

$$\frac{dx(t, y)}{dt} = \frac{\partial \Phi_t^{-1}(y)}{\partial t},$$

hence

$$V(t, x(t)) = \frac{\partial \Phi_t^{-1}(y)}{\partial t}, \quad x(t) = x(t, y).$$

Let $\Xi_t = \Phi_t^{-1}(\Xi)$ and $\Gamma_t = \Phi_t^{-1}(\Gamma)$. We can assume that Ξ_t has no self-intersections and consider the boundary value problem similar to (2)–(5) for the domain $\Omega_t = \Phi_t^{-1}(\Omega)$. Namely, in the domain Ω_t we want to find a function $u^t = (u_1^t, u_2^t, u_3^t)$ such that

$$-\sigma_{ij,j}^t = f_i, \quad i = 1, 2, 3, \quad (14)$$

$$\sigma_{ij}^t = a_{ijkl}\varepsilon_{kl}^t, \quad i, j = 1, 2, 3, \quad (15)$$

$$u^t = 0 \quad \text{on } \Gamma_t, \quad (16)$$

$$[u^t]\nu^t \geq 0, \quad \sigma_{\nu^t}^t \leq 0, \quad [\sigma_{\nu^t}^t] = 0, \quad \sigma_{\tau^t}^t = 0, \quad \Phi_{\nu^t}^t[u^t]\nu^t = 0 \quad \text{on } \Xi_t. \quad (17)$$

Here ν^t is a unit normal vector to Ξ_t , $\varepsilon_{kl}^t(u^t) = \frac{1}{2}(u_{k,l}^t + u_{l,k}^t)$. All the rest of the notation is similar to that of (2)–(5). In fact, the problem (14)–(17) is written in the variational form

$$u^t \in K_t : \quad \int_{\Omega_t} a_{ijkl}\varepsilon_{kl}(u^t)(\varepsilon_{ij}(\bar{u}^t) - \varepsilon_{ij}(u^t)) \geq \int_{\Omega_t} f(\bar{u}^t - u^t) \quad \forall \bar{u}^t \in K_t, \quad (18)$$

where

$$K_t = \{u = (u_1, u_2, u_3) \in H^1(\Omega) \mid u = 0 \text{ on } \Gamma_t; [u]\nu^t \geq 0 \text{ on } \Xi_t\}.$$

We impose one more condition on the mapping Φ_t . Assume that the condition $v(y) \in K_0$ implies $v^t(x) \in K_t$, $v^t(x) = v(y)$, $x \in \Omega_t$, $y \in \Omega$, $y = \Phi_t(x)$, and conversely, if $v(x) \in K_t$, then $v_t(y) \in K_0$, $v_t(y) = v(x)$, $x = x(t, y)$. Note that this condition is not very restricting, and it holds in many cases [2].

Let u, u^t be the solutions of the problems (7), (18), respectively. Consider the energy functionals

$$J(\Omega) = \frac{1}{2} \int_{\Omega} a_{ijkl}u_{k,l}u_{i,j} - \int_{\Omega} fu,$$

$$J(\Omega_t) = \frac{1}{2} \int_{\Omega_t} a_{ijkl}u_{k,l}^t u_{i,j}^t - \int_{\Omega_t} f u^t.$$

Our purpose is to find the derivative of $J(\Omega_t)$ with respect to the parameter t , namely,

$$\left. \frac{dJ(\Omega_t)}{dt} \right|_{t=0} = \lim_{t \rightarrow 0} \frac{J(\Omega_t) - J(\Omega)}{t}. \quad (19)$$

2. Solution convergence. First of all we prove the convergence of u^t to u in a proper sense. Namely, let $u^t(x) = u_t(y)$, $x \in \Omega_t$, $y \in \Omega$, $x = x(t, y)$. Denote by $\|\cdot\|_{1,\Omega}$ the norm in the space $H^1(\Omega)$.

LEMMA. The following estimate holds:

$$\|u_t - u\|_{1,\Omega} \leq ct,$$

where c is a constant independent of t .

Proof. The functions u, u^t satisfy the following variational inequalities:

$$u \in K_0 : \quad \int_{\Omega} a_{ijkl}u_{k,l}(\bar{u}_{i,j} - u_{i,j}) \geq \int_{\Omega} f(\bar{u} - u) \quad \forall \bar{u} \in K_0, \quad (20)$$

$$u^t \in K_t : \quad \int_{\Omega_t} a_{ijkl}u_{k,l}^t(\bar{u}_{i,j}^t - u_{i,j}^t) \geq \int_{\Omega_t} f(\bar{u}^t - u^t) \quad \forall \bar{u}^t \in K_t. \quad (21)$$

Denote $f_t(y) = f(x(t, y))$, $q_t(y) = \left| \frac{\partial \Phi_t^{-1}(y)}{\partial y} \right|$. We have

$$u_{k,l}^t(x) = u_{tk,p}(y) \Phi_{t,l}^p(x), \quad k, l = 1, 2, 3.$$

Consequently, the inequality (21) can be rewritten in the form

$$u_t \in K_0 : \int_{\Omega} a_{ijkl} u_{tk,p} \Phi_{t,l}^p(\bar{u}_{ti,s} \Phi_{t,j}^s - u_{ti,s} \Phi_{t,j}^s) q_t \geq \int_{\Omega} f_t(\bar{u}_t - u_t) q_t \quad \forall \bar{u}_t \in K_0. \quad (22)$$

It is important to note that $\bar{u}_t \in K_0$ due to the assumption imposed on Φ_t . Here we used the one-to-one mapping between K_t and K_0 . In the inequality (22), we have $\Phi_{t,l}^p = \Phi_{t,l}^p(x(t, y))$. Denote by δ_l^p the Kronecker symbol. The following equalities hold:

$$\Phi_{t,l}^p(x(t, y)) = \Phi_{0,l}^p(x(t, y)) + \frac{\partial \Phi_{\xi,l}^p(x(t, y))}{\partial \xi} t, \quad \xi \in (0, t).$$

Since $\Phi_{0,l}^p = \delta_l^p$, $p, l = 1, 2, 3$, these equalities can be rewritten as follows:

$$\Phi_{t,l}^p(x(t, y)) = \delta_l^p + \Phi_{\xi}^{pl}(x(t, y)) t, \quad (23)$$

where $\xi = \xi(t, p, l)$, and we have denoted $\frac{\partial \Phi_{\xi,l}^p}{\partial \xi}$ by Φ_{ξ}^{pl} ,

$$\|\Phi_{\xi}^{pl}\|_{L^{\infty}(\Omega)} \leq c \quad \text{uniformly in } \xi \in (0, T). \quad (24)$$

Moreover, $q_t(y) = q_0(y) + \frac{\partial q_{\xi}(y)}{\partial \xi} t$, $\xi \in (0, t)$, and $q_0(y) = 1$. Denote $\frac{\partial q_{\xi}(y)}{\partial \xi}$ by $\bar{q}_{\xi}(y)$ which gives

$$q_t(y) = 1 + \bar{q}_{\xi}(y) t, \quad (25)$$

with the uniform in ξ estimate $\|\bar{q}_{\xi}\|_{L^{\infty}(\Omega)} \leq c$. By (23), the inequality (22) can be written in the form

$$\int_{\Omega} a_{ijkl} u_{tk,p} (\delta_l^p + t \Phi_{\xi}^{pl}) [\bar{u}_{ti,s} (\delta_j^s + t \Phi_{\xi}^{sj}) - u_{ti,s} (\delta_j^s + t \Phi_{\xi}^{sj})] q_t \geq \int_{\Omega} f_t(\bar{u}_t - u_t) q_t. \quad (26)$$

Now substitute $\bar{u} = u_t$, $\bar{u} = u$ in (20), (26), respectively. By (25), this yields

$$\int_{\Omega} a_{ijkl} u_{k,l} (u_{ti,j} - u_{i,j}) \geq \int_{\Omega} f(u_t - u), \quad (27)$$

$$\begin{aligned} & \int_{\Omega} a_{ijkl} u_{tk,l} (u_{i,j} - u_{ti,j}) (1 + \bar{q}_{\xi} t) + t \int_{\Omega} a_{ijkl} u_{tk,p} \Phi_{\xi}^{pl} (u_{i,j} - u_{ti,j}) q_t \\ & \quad + t \int_{\Omega} a_{ijkl} u_{tk,l} (u_{i,s} \Phi_{\xi}^{sj} - u_{ti,s} \Phi_{\xi}^{sj}) q_t \\ & \quad + t^2 \int_{\Omega} a_{ijkl} u_{tk,p} \Phi_{\xi}^{pl} (u_{i,s} \Phi_{\xi}^{sj} - u_{ti,s} \Phi_{\xi}^{sj}) q_t \geq \int_{\Omega} f_t(u - u_t) q_t. \quad (28) \end{aligned}$$

Summing (27) and (28) we obtain

$$\begin{aligned} \int_{\Omega} a_{ijkl}(u_{k,l} - u_{tk,l})(u_{i,j} - u_{ti,j}) &\leq t \int_{\Omega} \bar{q}_{\xi} a_{ijkl} u_{tk,l} (u_{i,j} - u_{ti,j}) \\ &+ t \int_{\Omega} [a_{ijkl} u_{tk,p} \Phi_{\xi}^{pl} (u_{i,j} - u_{ti,j}) + a_{ijkl} u_{tk,l} (u_{i,s} \Phi_{\xi}^{sj} - u_{ti,s} \Phi_{\xi}^{sj})] q_t \\ &+ t^2 \int_{\Omega} a_{ijkl} u_{tk,p} \Phi_{\xi}^{pl} (u_{i,s} \Phi_{\xi}^{sj} - u_{ti,s} \Phi_{\xi}^{sj}) q_t \\ &- \int_{\Omega} f(u_t - u) + \int_{\Omega} f_t(u_t - u) q_t. \end{aligned} \quad (29)$$

Taking $\bar{u}_t = 0$ in (26), we derive the uniform in $t \in (0, T)$ estimate

$$\|u_t\|_{1,\Omega} \leq c.$$

Consequently, by (24), (25), from (29) it follows that

$$\int_{\Omega} a_{ijkl}(u_{i,j} - u_{ti,j})(u_{k,l} - u_{tk,l}) \leq ct^2 + \int_{\Omega} |u - u_t| |f - f_t(1 + \bar{q}_{\xi}t)|, \quad (30)$$

where the constant c is independent of $t \in (0, T)$. Since

$$\begin{aligned} f_k(y) - f_{kt}(y)(1 + \bar{q}_{\xi}t) &= f_k(y) - \left[f_k(y) + \frac{\partial f_k}{\partial x_i} \frac{dx_i(\xi_1)}{d\xi_1} (1 + \bar{q}_{\xi}t) \right], \\ &\qquad \qquad \qquad \xi_1 \in (0, t) \quad k = 1, 2, 3, \end{aligned}$$

the inequality (30) implies

$$\|u - u_t\|_{1,\Omega}^2 \leq ct^2,$$

which completes the proof of the lemma. \square

3. Main result. To find the derivative of the energy functional, we shall use the variational property of the solution. Introduce first some notation:

$$\begin{aligned} \Pi(\Omega_t, \varphi) &= \frac{1}{2} \int_{\Omega_t} a_{ijkl} \varphi_{k,l} \varphi_{i,j} dx - \int_{\Omega_t} f \varphi dx, \\ \Pi_t(\Omega, \varphi) &= \frac{1}{2} \int_{\Omega} a_{ijkl} \varphi_{k,p} \Phi_{t,l}^p \varphi_{i,s} \Phi_{t,j}^s q_t dy - \int_{\Omega} f_t \varphi q_t dy. \end{aligned}$$

Since we have the one-to-one mapping between K_t and K_0 , the following equality holds:

$$\min_{\varphi \in K_0} \Pi_t(\Omega; \varphi) = \min_{\varphi \in K_t} \Pi(\Omega_t; \varphi).$$

Note also that

$$J(\Omega) = \Pi(\Omega; u), \quad J(\Omega_t) = \Pi(\Omega_t; u^t),$$

where u, u^t are the solutions of (7) and (18), respectively. Consequently, we conclude

$$\begin{aligned} \frac{J(\Omega_t) - J(\Omega)}{t} &= \frac{\Pi(\Omega_t; u^t) - \Pi(\Omega; u)}{t} \\ &= \frac{\Pi_t(\Omega; u_t) - \Pi(\Omega; u)}{t} \leq \frac{\Pi_t(\Omega; u) - \Pi(\Omega; u)}{t}. \end{aligned}$$

This implies

$$\limsup_{t \rightarrow 0} \frac{J(\Omega_t) - J(\Omega)}{t} \leq \limsup_{t \rightarrow 0} \frac{\Pi_t(\Omega; u) - \Pi(\Omega; u)}{t}. \quad (31)$$

On the other hand,

$$\limsup_{t \rightarrow 0} \frac{J(\Omega_t) - J(\Omega)}{t} = \frac{\Pi(\Omega_t; u^t) - \Pi(\Omega; u)}{t} \geq \frac{\Pi_t(\Omega; u_t) - \Pi(\Omega; u_t)}{t},$$

whence

$$\liminf_{t \rightarrow 0} \frac{J(\Omega_t) - J(\Omega)}{t} \geq \liminf_{t \rightarrow 0} \frac{\Pi_t(\Omega; u_t) - \Pi(\Omega; u_t)}{t}. \quad (32)$$

Now we aim to show that the right-hand sides of (31), (32) coincide, which implies an existence of the limit

$$\lim_{t \rightarrow 0} \frac{J(\Omega_t) - J(\Omega)}{t}.$$

Let us find the right-hand side of (31). It suffices to find the derivative

$$\left. \frac{d}{dt} \Pi_t(\Omega; u) \right|_{t=0} = \left. \frac{d}{dt} \left\{ \frac{1}{2} \int_{\Omega} a_{ijkl} u_{k,p} \Phi_{t,l}^p u_{i,s} \Phi_{t,j}^s q_t - \int_{\Omega} f_t u q_t \right\} \right|_{t=0}. \quad (33)$$

By denoting

$$\Lambda(y) = -V(0, y) = \left. \frac{\partial \Phi_t(x(t, y))}{\partial t} \right|_{t=0}, \quad (34)$$

we have

$$\Phi_{t,l}^p(x(t, y))|_{t=0} = \delta_l^p, \quad \left. \frac{\partial \Phi_{t,l}^p(x(t, y))}{\partial t} \right|_{t=0} = \Lambda_{,l}^p(y), \quad (35)$$

and, moreover, as $t \rightarrow 0$,

$$\Phi_{t,l}^p(x(t, y)) \rightarrow \delta_l^p, \quad \frac{\partial \Phi_{t,l}^p(x(t, y))}{\partial t} \rightarrow \Lambda_{,l}^p(y) \quad \text{in } L^\infty(\Omega). \quad (36)$$

Indeed,

$$\begin{aligned} \Phi_{t,l}^p(x(t, y)) &= \Phi_{t,l}^p(x(t, y)) - \Phi_{0,l}^p(x(t, y)) + \Phi_{0,l}^p(x(t, y)) \\ &= \frac{\partial \Phi_{\xi,l}^p(x(t, y))}{\partial \xi} t + \Phi_{0,l}^p(x(t, y)), \quad \xi \in (0, t). \end{aligned}$$

By (24), we have

$$\frac{\partial \Phi_{\xi,l}^p(x(t, y))}{\partial \xi} \text{ are bounded in } L^\infty(\Omega) \text{ uniformly in } \xi, t \in (0, T),$$

which together with the first equality of (35) implies the first convergence of (36). Similarly, since

$$\frac{\partial^2 \Phi_{\xi,l}^p(x(t, y))}{\partial \xi^2} \text{ are bounded in } L^\infty(\Omega) \text{ for all } \xi, t \in (0, T)$$

and $\frac{\partial \Phi_{0,l}^p(x)}{\partial \xi}$ satisfy the Lipschitz property in x , from the equality

$$\begin{aligned} \frac{\partial \Phi_{t,l}^p(x(t,y))}{\partial t} &= \frac{\partial \Phi_{t,l}^p(x(t,y))}{\partial t} - \frac{\partial \Phi_{0,l}^p(x(t,y))}{\partial t} + \frac{\partial \Phi_{0,l}^p(x(t,y))}{\partial t} \\ &= \frac{\partial^2 \Phi_{\xi,l}^p(x(t,y))}{\partial \xi^2} t + \frac{\partial \Phi_{0,l}^p(x(t,y))}{\partial t}, \quad \xi \in (0,t), \end{aligned}$$

we conclude that the second convergence of (36) takes place.

It is well known that [21]

$$\frac{\partial q_t(y)}{\partial t} = q_t(y) \operatorname{div} V(t, x(t, y)), \quad (37)$$

and by (25), (34),

$$\left. \frac{\partial q_t(y)}{\partial t} \right|_{t=0} = -\operatorname{div} \Lambda(y).$$

We next obtain

$$\begin{aligned} \operatorname{div} V(t, x(t, y)) &= \operatorname{div} V(0, y) + \frac{d}{d\xi} \operatorname{div} V(\xi, x(\xi, y))t, \quad \xi \in (0, t), \\ \left\| \frac{d}{d\xi} \operatorname{div} V(\xi, x(\xi, y)) \right\|_{L^\infty(\Omega)} &\leq c \quad \text{uniformly in } \xi \in (0, T). \end{aligned}$$

Hence, taking into account (25), as $t \rightarrow 0$,

$$q_t(y) \operatorname{div} V(t, x(t, y)) \rightarrow -\operatorname{div} \Lambda(y) \quad \text{in } L^\infty(\Omega).$$

By (37), this gives as $t \rightarrow 0$,

$$\frac{\partial q_t}{\partial t} \rightarrow -\operatorname{div} \Lambda \quad \text{in } L^\infty(\Omega). \quad (38)$$

Also note that (35) implies $\Phi_{t,l}^p(x(t,y))|_{t=0} = 0$, $p, l, s = 1, 2, 3$, so that

$$\left. \frac{d\Phi_{t,l}^p(x(t,y))}{dt} \right|_{t=0} = \left. \frac{\partial \Phi_{t,l}^p(x(t,y))}{\partial t} \right|_{t=0}.$$

Hence, by (36), (38), we can calculate the right-hand side of (33), i.e., the right-hand side of (31):

$$\begin{aligned} \frac{d}{dt} \Pi_t(\Omega; u)|_{t=0} &= \frac{1}{2} \int_{\Omega} (a_{ijkl} u_{k,p} u_{i,s} \Lambda_{,l}^p \delta_j^s + a_{ijkl} u_{k,p} u_{i,s} \Lambda_{,j}^s \delta_l^p) \\ &\quad - \frac{1}{2} \int_{\Omega} a_{ijkl} u_{k,p} u_{i,s} \delta_l^p \delta_j^s (\operatorname{div} \Lambda) + \int_{\Omega} u_k (\nabla f_k \Lambda) + \int_{\Omega} f u \operatorname{div} \Lambda \quad (39) \\ &= \int_{\Omega} \left\{ \sigma_{kl} u_{k,p} \Lambda_{,l}^p - \frac{1}{2} \sigma_{ij} \varepsilon_{ij} \operatorname{div} \Lambda \right\} + \int_{\Omega} u_k \operatorname{div} (f_k \Lambda). \end{aligned}$$

Now we find the right-hand side of (32). To this end we consider the term

$$\Delta_t = \frac{1}{t} \int_{\Omega} (u_{tk,p} u_{ti,s} \Phi_{t,l}^p \Phi_{t,j}^s q_t - u_{tk,l} u_{ti,j})$$

for fixed k, l, i, j . It is possible to write Δ_t in the form

$$\begin{aligned}
\Delta_t = & \frac{1}{t} \int_{\Omega} (u_{tk,p} u_{ti,s} \Phi_{t,l}^p \Phi_{t,j}^s - u_{tk,p} \delta_l^p u_{ti,s} \Phi_{t,j}^s) q_t \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} \delta_l^p u_{ti,s} \Phi_{t,j}^s - u_{tk,l} u_{ti,j}) \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} u_{ti,s} \Phi_{t,l}^p \Phi_{t,j}^s - u_{tk,p} \delta_l^p u_{ti,s} \Phi_{t,j}^s) \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} \delta_l^p u_{ti,s} \Phi_{t,j}^s - u_{tk,p} \delta_l^p u_{ti,s} \delta_j^s) q_t \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} \delta_l^p u_{ti,s} \delta_j^s - u_{tk,p} \Phi_{t,l}^p u_{ti,s} \delta_j^s) q_t \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} \Phi_{t,l}^p u_{ti,s} \delta_j^s q_t - u_{tk,p} \Phi_{t,l}^p u_{ti,s} \delta_j^s) \\
& + \frac{1}{t} \int_{\Omega} (u_{tk,p} \Phi_{t,l}^p u_{ti,s} \delta_j^s - u_{tk,p} \Phi_{t,l}^p u_{ti,s} \Phi_{t,j}^s).
\end{aligned} \tag{40}$$

Recall that, as $t \rightarrow 0$,

$$u_t \rightarrow u \quad \text{in } H^1(\Omega), \quad q_t \rightarrow 1 \quad \text{in } L^\infty(\Omega). \tag{41}$$

Since

$$\begin{aligned}
\frac{q_t(y) - q_0(y)}{t} &= \frac{\partial q_\xi(y)}{\partial \xi}, \quad \xi \in (0, t), \\
\frac{\Phi_{t,l}^p(x(t, y)) - \delta_l^p}{t} &= \frac{\partial \Phi_{\xi,l}^p(x(t, y))}{\partial \xi}, \quad \xi \in (0, t),
\end{aligned}$$

by the convergences (36), (38), we can find the limit of each part of the right-hand side of (40) as $t \rightarrow 0$, which implies

$$\begin{aligned}
\lim_{t \rightarrow 0} \Delta_t &= \int_{\Omega} \{u_{k,p} \Lambda_{,l}^p u_{i,j} + u_{k,l} u_{i,s} \Lambda_{,j}^s + u_{i,j} u_{k,p} \Lambda_{,l}^p + u_{k,l} u_{i,s} \Lambda_{,j}^s \\
&\quad - u_{i,j} u_{k,p} \Lambda_{,l}^p - u_{k,l} u_{i,j} (\operatorname{div} \Lambda) - u_{k,l} u_{i,s} \Lambda_{,j}^s\} \\
&= \int_{\Omega} \{u_{i,j} u_{k,p} \Lambda_{,l}^p + u_{k,l} u_{i,s} \Lambda_{,j}^s - u_{k,l} u_{i,j} (\operatorname{div} \Lambda)\}.
\end{aligned} \tag{42}$$

In addition to this, as $t \rightarrow 0$,

$$\frac{f_{kt} - f_k}{t} \rightarrow -\nabla f_k \cdot \Lambda \quad \text{in } L^\infty(\Omega), \quad k = 1, 2, 3;$$

hence

$$\begin{aligned}
\lim_{t \rightarrow 0} \int_{\Omega} \frac{f_t q_t u_t - f u_t}{t} &= \lim_{t \rightarrow 0} \int_{\Omega} \frac{(f_t - f) q_t u_t}{t} + \lim_{t \rightarrow 0} \int_{\Omega} \frac{f u_t (q_t - 1)}{t} \\
&= - \int_{\Omega} u_k (\nabla f_k \cdot \Lambda) - \int_{\Omega} f u \operatorname{div} \Lambda.
\end{aligned} \tag{43}$$

From (42), (43) it follows that

$$\begin{aligned}
 & \lim_{t \rightarrow 0} \frac{\Pi_t(\Omega; u_t) - \Pi(\Omega; u_t)}{t} \\
 &= \int_{\Omega} \left\{ a_{ijkl} u_{i,j} u_{k,p} \Lambda_{,l}^p - \frac{1}{2} a_{ijkl} u_{i,j} u_{k,l} (\operatorname{div} \Lambda) \right\} \\
 & \quad + \int_{\Omega} u_k (\nabla f_k \Lambda) + \int_{\Omega} f u \operatorname{div} \Lambda \\
 &= \int_{\Omega} \left\{ \sigma_{kl} u_{k,p} \Lambda_{,l}^p - \frac{1}{2} \sigma_{ij} \varepsilon_{ij} \operatorname{div} \Lambda \right\} + \int_{\Omega} u_k \operatorname{div} (f_k \Lambda).
 \end{aligned} \tag{44}$$

By (39), (44), we conclude that the right-hand sides of (31), (32) coincide and we obtain the following statement.

THEOREM. Let the hypotheses concerning Φ_t be fulfilled. Then the derivative of the energy functional is given by the formula

$$\left. \frac{dJ(\Omega_t)}{dt} \right|_{t=0} = \int_{\Omega} \left\{ \sigma_{kl} u_{k,p} \Lambda_{,l}^p - \frac{1}{2} \sigma_{ij} \varepsilon_{ij} \operatorname{div} \Lambda \right\} + \int_{\Omega} u_k \operatorname{div} (f_k \Lambda), \tag{45}$$

where the vector field Λ is defined by (34).

In conclusion, note that the formulae similar to (45) were obtained for isotropic two- and three-dimensional cracked bodies with conditions (5) at the crack faces provided that the perturbation Φ_t of the domain Ω_t describes the crack length change [1, 2].

Acknowledgment. The paper was prepared during a visit of K. Ohtsuka and A. M. Khludnev at the Institute Elie Cartan in Nancy. A. M. Khludnev was supported by the Institute Elie Cartan, of the University Henri Poincaré Nancy I and by the Russian Fund for Basic Research under the grant 00-01-00842.

REFERENCES

- [1] A. M. Khludnev and J. Sokółowski, *The Griffith formula and the Rice-Cherepanov integral for crack problems with unilateral conditions in nonsmooth domains*, European J. Appl. Math. **10**, 379–394 (1999)
- [2] A. M. Khludnev and V. A. Kovtunenکو, *Analysis of Cracks in Solids*, Southampton-Boston, WIT Press, 2000
- [3] V. G. Mazya and S. A. Nazarov, *Asymptotic behavior of energy integrals under small perturbations of the boundary near corner and conic points*, Proceeding of Moscow Math. Society, Moscow State Univ., 1987, pp. 79–129 (in Russian)
- [4] K. Ohtsuka, *Generalized J-integral and three-dimensional fracture mechanics I*, Hiroshima Math. J. **11**, 21–52 (1981)
- [5] K. Ohtsuka, *Generalized J-integral and its applications. I. - Basic theory*, Japan Journal of Applied Mathematics **2**, 329–350 (1985)
- [6] J. Sokółowski and J. P. Zolesio, *Introduction to Shape Optimization—Shape Sensitivity Analysis*, Springer-Verlag, New York, 1992
- [7] V. P. Parton and E. M. Morozov, *Mechanics of Elastoplastic Fracture*, Moscow, Nauka, 1985 (in Russian)
- [8] A. M. Khludnev and J. Sokółowski, *Modelling and Control in Solid Mechanics*, Birkhäuser, Basel-Boston-Berlin, 1997
- [9] P. Destuynder and M. Jaoua, *Sur une Interprétation Mathématique de l'Intégrale de Rice en Théorie de la Rupture Fragile*, Math. Mech. in the Appl. Sci. **3**, 70–87 (1981)

- [10] A. M. Khludnev, *On a Signorini problem for inclusions in shells*, European J. Appl. Math. **7**, 499–510 (1996)
- [11] A. M. Khludnev, *Contact problem for a plate having a crack of minimal opening*, Control and Cybernetics **25**, 605–620 (1996)
- [12] P. Grisvard, *Singularities in Boundary Value Problems*, Masson, Paris; Springer, Berlin, 1992
- [13] G. P. Cherepanov, *Mechanics of Brittle Fracture*, McGraw-Hill, 1979
- [14] M. Bonnet, *Équations intégrales variationnelles pour le problème en vitesse de propagation de fissures en élasticité linéaire*, C. R. Acad. Sci. Paris, série II, vol. 318, 1994, pp. 429–434
- [15] S. A. Nazarov and B. A. Plamenevskii, *Elliptic problems in domains with piecewise smooth boundaries*, Moscow, Nauka, 1991 (in Russian)
- [16] Q. S. Nguen, C. Stolz, and G. Debruyne, *Energy methods in fracture mechanics: Stability, bifurcation and second variations*, European J. of Mech., A/Solids **9**, 157–173 (1990)
- [17] H. Petryk and Z. Mróz, *Time derivatives of integrals and functionals defined on varying volume and surface domains*, Arch. Mech. **38**, 697–724 (1986)
- [18] J. H. Edward, K. C. Kyung, and V. Komkov, *Design Sensitivity Analysis of Structural Systems*, Academic Press, 1986
- [19] A. M. Khludnev, *The contact between two plates, one of which contains a crack*, J. Appl. Math. Mech. **61**, 851–862 (1997)
- [20] A. M. Khludnev, *The contact problem for a shallow shell with a crack*, J. Appl. Math. Mech. **59**, 299–306 (1995)
- [21] L. V. Ovsyannikov, *Lectures on the foundation of gas dynamics*, Moscow, Nauka, 1981 (in Russian)