

Received September 13, 2019, accepted September 24, 2019, date of publication October 1, 2019, date of current version October 16, 2019.

Digital Object Identifier 10.1109/ACCESS.2019.2944860

# On Distance-Based Topological Descriptors of Subdivision Vertex-Edge Join of Three Graphs

HONG YANG<sup>1</sup>, MUHAMMAD IMRAN<sup>2,3</sup>, SHEHNAZ AKHTER<sup>3</sup>,  
ZAHID IQBAL<sup>3</sup>, AND MUHAMMAD KAMRAN SIDDIQUI<sup>4</sup>

<sup>1</sup>School of Information Science and Engineering, Chengdu University, Chengdu 610106, China

<sup>2</sup>Department of Mathematical Sciences, United Arab Emirates University, Al Ain 15551, United Arab Emirates

<sup>3</sup>Department of Mathematics, School of Natural Sciences (SNS), National University of Sciences and Technology (NUST), Islamabad 44000, Pakistan

<sup>4</sup>Department of Mathematics, COMSATS University Islamabad, Lahore Campus, Lahore 54000, Pakistan

Corresponding author: Muhammad Kamran Siddiqui (kamransiddiqui75@gmail.com)

This work was supported in part by the Soft Scientific Research of Sichuan Province under Grant 2018ZR0265, in part by the Sichuan Military and Civilian Integration Strategy Research Center under Grant JMRH-1818, in part by the Sichuan Provincial Department of Education (Key Project) under Grant 18ZA0118, and in part by the Start-Up Research Grant 2016 of United Arab Emirates University (UAEU), Al Ain, United Arab Emirates, under Grant G00002233, and in part by the UPAR Grant of UAEU under Grant G00002590.


**ABSTRACT** The analysis of networks and graphs through topological descriptors carries out a useful role to derive their underlying topologies. This process has been widely used in biomedicine, cheminformatics, and bioinformatics, where assessments based on graph invariants have been made available for effectively communicating with the various challenging schemes. In the studies of quantitative structure-activity relationships (QSARs) and quantitative structure-property relationships (QSPRs), graph invariants are used to approximate the biological activities and properties of chemical compounds. In this paper, we give the results related to the eccentric-connectivity index, connective eccentricity index, total-eccentricity index, average eccentricity index, Zagreb eccentricity indices, eccentric geometric-arithmetic index, eccentric atom-bond connectivity index, eccentric adjacency index, modified eccentric-connectivity index, eccentric distance sum, Wiener index, Harary index, hyper-Wiener index and degree distance index of a new graph operation named as “subdivision vertex-edge join” of three graphs.

**INDEX TERMS** Topological indices, degree, distance, eccentricity, subdivision vertex-edge join.

## I. INTRODUCTION

Graph theory concerned with a lot of applications in a number of domains of chemistry such as Quantitative structure-activity and property relationships, isomer enumeration, prediction of biological activities, topological characterization, graph polynomials for structural analysis, quantum chemistry, NMR spectroscopy, nuclear spin statistics, spectroscopy, proteomics, statistical and other procedures for forecast of toxicity of chemical structures and so on [7], [9], [11], [13], [15], [17], [19], [20], [31], [43], [45], [46]. The QSAR/QSPR studies made use of connection among molecular connectivity and the properties of chemical compounds, therefore a fundamental graph-theoretical characterizations set up the principles for computer-aided predictive toxicology and drug discovery. As a result, successful uses of QSAR/QSPR studies have stimulated the emergence of several topological indices of chemical structures

[10]–[14], [16]–[20], [34], [43]–[45], [47]. The intermolecular interactions rely on the distance and degree criteria and moreover, numerous physico-chemical properties of chemical structures have been proven to correlate with topological characteristics as decent initial points. However, one may require sophisticated bio-descriptors and quantum chemical in addition to quantum molecular dynamics simulations for extra precise forecasts of biological and chemical characteristics, due to computationally extensive nature of such approaches, topological techniques have constructed valuable implementations because of the comparatively easy process with which they can be determined. Many properties such as receptor binding propensity, toxicity, protein-drug interactions, dermal penetrations, drug metabolomics, guest-host interactions, etc., rely on the intermolecular interactions, pore sizes, structural parameters, electrostatic and electronic properties various of which rely on fundamental topological parameter distances and therefore topological descriptors are much more appealing initiating objects to any statistical approximation for securing structure-activity connections.

The associate editor coordinating the review of this manuscript and approving it for publication was Donghyun Kim .




FIGURE 1.  $P_4 \dot{\vee} C_3$ ,  $P_4 \underline{\vee} K_4$  and  $P_4^S \triangleright (C_3^V \cup K_4^I)$ .

Throughout the manuscript, all considered graphs are simple and connected. For a graph  $\mathcal{H}$ ,  $\mathcal{V}(\mathcal{H})$  and  $\mathcal{E}(\mathcal{H})$  appear for vertex and edge sets, respectively, and  $n$  and  $m$  stands for the order and size of  $\mathcal{H}$ , respectively. An edge with end vertices  $h_i$  and  $h_j$  is recognized by  $h_i h_j \in \mathcal{E}(\mathcal{H})$ . For  $h \in \mathcal{V}(\mathcal{H})$ , the number of edges whose an end vertex  $h$  is called the degree of  $h$  in  $\mathcal{H}$  and it is denoted by  $\text{deg}_{\mathcal{H}}(h)$ . A  $(y_1, y_n)$ -path of  $n$ -vertices is described as a graph whose vertex and edge sets are  $\{y_1, \dots, y_n\}$  and  $\{y_i y_{i+1} : 1 \leq i \leq n - 1\}$ , respectively. The notions  $K_n$ ,  $P_n$  and  $C_n$  are commonly used for complete graph, path and cycle, respectively. The distance among two vertices  $a, c \in \mathcal{V}(\mathcal{H})$  is represented by  $d_{\mathcal{H}}(a, c)$  and explained as the length of shortest  $(a, c)$ -path in  $\mathcal{H}$ . For  $a \in \mathcal{V}(\mathcal{H})$ , the eccentricity  $ec_{\mathcal{H}}(a)$  is specified as the largest distance among  $a$  and any other vertex in  $\mathcal{H}$ .

Recently, a new graph operation has been initiated by Wen et al. in [51], they named it as the subdivision vertex-edge join (SVE-join). For a graph  $\mathcal{H}_q$ ,  $\mathcal{S}(\mathcal{H}_q)$  is the subdividing graph of  $\mathcal{H}_q$  whose vertex set has two portions, one the primary vertices  $\mathcal{V}(\mathcal{H}_q)$ , another, represented by  $\mathcal{I}(\mathcal{H}_q)$ , the inserting vertices that are end vertices of the edges of  $\mathcal{H}_q$ . Let  $\mathcal{H}_r$  and  $\mathcal{H}_s$  be the other two disjoint graphs. The SVE-join of  $\mathcal{H}_q$  with  $\mathcal{H}_r$  and  $\mathcal{H}_s$ , expressed by  $\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)$ , is the graph containing of  $\mathcal{S}(\mathcal{H}_q)$ ,  $\mathcal{H}_r$  and  $\mathcal{H}_s$ , all vertex-disjoint, and connecting the  $l$ -th vertex of  $\mathcal{V}(\mathcal{H}_q)$  to every vertex in  $\mathcal{V}(\mathcal{H}_r)$  and  $l$ -th vertex of  $\mathcal{I}(\mathcal{H}_q)$  to each vertex in  $\mathcal{V}(\mathcal{H}_s)$ . It can be observed that  $\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)$  is  $\mathcal{H}_q \dot{\vee} \mathcal{H}_r$  (is attained from  $\mathcal{S}(\mathcal{H}_q)$  and  $\mathcal{H}_r$  by linking each vertex of  $\mathcal{V}(\mathcal{H}_q)$  to every vertex of  $\mathcal{V}(\mathcal{H}_r)$  [38]) if  $\mathcal{H}_s$  is the null graph, and is  $\mathcal{H}_q \underline{\vee} \mathcal{H}_s$  (is attained from  $\mathcal{S}(\mathcal{H}_q)$  and  $\mathcal{H}_s$  by linking each vertex of  $\mathcal{E}(\mathcal{H}_q)$  to every vertex of  $\mathcal{V}(\mathcal{H}_s)$  [38]) if  $\mathcal{H}_r$  is the null graph. The graphs  $P_4 \dot{\vee} C_3$ ,  $P_4 \underline{\vee} K_4$  and  $P_4^S \triangleright (C_3^V \cup K_4^I)$  are illustrated in Figure 1.

Sharma et al. [49] introduced the well-known eccentricity-based index of a graph  $\mathcal{H}$ . They defined it as follows:

$$\xi^c(\mathcal{H}) = \sum_{h \in \mathcal{V}(\mathcal{H})} \text{deg}_{\mathcal{H}}(h) ec_{\mathcal{H}}(h). \quad (1)$$

Gupta et al. [28] gave the concept of connective eccentricity index of  $\mathcal{H}$  as follows:

$$\xi^{ce}(\mathcal{H}) = \sum_{h \in \mathcal{V}(\mathcal{H})} \frac{\text{deg}_{\mathcal{H}}(h)}{ec_{\mathcal{H}}(h)}. \quad (2)$$

If we use only the eccentricities of vertices of  $\mathcal{H}$  in (1), then we can describe the total-eccentricity index as follows [6]:

$$\tau(\mathcal{H}) = \sum_{h \in \mathcal{V}(\mathcal{H})} ec_{\mathcal{H}}(h). \quad (3)$$

The mean value of the eccentricities of elements of  $\mathcal{V}(\mathcal{H})$  is said to be the average eccentricity  $aveg(\mathcal{H})$  of it [50], that is

$$aveg(\mathcal{H}) = \frac{1}{n} \sum_{h \in \mathcal{V}(\mathcal{H})} ec_{\mathcal{H}}(h) = \frac{\tau(\mathcal{H})}{n}. \quad (4)$$

The eccentricity versions of Zagreb indices of  $\mathcal{H}$  were given in [26] as follows:

$$\begin{aligned} \mathcal{M}_1(\mathcal{H}) &= \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} ec_{\mathcal{H}}^2(h_{p_1}), \\ \mathcal{M}_2(\mathcal{H}) &= \sum_{h_{p_1} h_{p_2} \in \mathcal{E}(\mathcal{H})} ec_{\mathcal{H}}(h_{p_1}) ec_{\mathcal{H}}(h_{p_2}). \end{aligned} \quad (5)$$

The eccentric form of geometric-arithmetic index [27] of  $\mathcal{H}$  is as follows:

$$\mathcal{GA}_{ec}(\mathcal{H}) = \sum_{h_{p_1} h_{p_2} \in \mathcal{E}(\mathcal{H})} \frac{2\sqrt{ec_{\mathcal{H}}(h_{p_1}) ec_{\mathcal{H}}(h_{p_2})}}{ec_{\mathcal{H}}(h_{p_1}) + ec_{\mathcal{H}}(h_{p_2})}. \quad (6)$$

The eccentric form of atom-bond connectivity index [22] of  $\mathcal{H}$  is as follows:

$$ABC_{ec}(\mathcal{H}) = \sum_{h_{p_1}h_{p_2} \in \mathcal{E}(\mathcal{H})} \sqrt{\frac{ec_{\mathcal{H}}(h_{p_1}) + ec_{\mathcal{H}}(h_{p_2}) - 2}{ec_{\mathcal{H}}(h_{p_1})ec_{\mathcal{H}}(h_{p_2})}}. \quad (7)$$

The eccentric adjacency index is given by Gupta et al. in [29] as follows:

$$\xi^{ad}(\mathcal{H}) = \sum_{h \in \mathcal{V}(\mathcal{H})} \frac{S_{\mathcal{H}}(h)}{ec_{\mathcal{H}}(h)}. \quad (8)$$

The modified type of eccentric-connectivity index [5] of  $\mathcal{H}$  is given in the following way:

$$\xi_c(\mathcal{H}) = \sum_{h \in \mathcal{V}(\mathcal{H})} S_{\mathcal{H}}(h)ec_{\mathcal{H}}(h). \quad (9)$$

The eccentric distance sum index was first presented in [30] as follows:

$$\begin{aligned} \xi^{ds}(\mathcal{H}) &= \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} (ec_{\mathcal{H}}(h_{p_1}) + ec_{\mathcal{H}}(h_{p_2}))d_{\mathcal{H}}(h_{p_1}, h_{p_2}) \\ &= \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} ec_{\mathcal{H}}(h_{p_1})D_{\mathcal{H}}(h_{p_1}). \end{aligned} \quad (10)$$

where  $D_{\mathcal{H}}(h_{p_1}) = \sum_{h_{p_2} \in \mathcal{V}(\mathcal{H})} d_{\mathcal{H}}(h_{p_1}, h_{p_2})$ .

The Wiener index is a distance-based graph descriptor described by [21] as follows:

$$\begin{aligned} W(\mathcal{H}) &= \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} d_{\mathcal{H}}(h_{p_1}, h_{p_2}) \\ &= \frac{1}{2} \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} D_{\mathcal{H}}(h_{p_1}) \\ &= \frac{1}{2} \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} \sum_{h_{p_2} \in \mathcal{V}(\mathcal{H})} d_{\mathcal{H}}(h_{p_1}, h_{p_2}). \end{aligned} \quad (11)$$

The Harary index of  $\mathcal{H}$  is explained as the sum of reciprocals of distances among all the unordered pairs of its vertices as follows: [42]:

$$\begin{aligned} H(\mathcal{H}) &= \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} \frac{1}{d_{\mathcal{H}}(h_{p_1}, h_{p_2})} \\ &= \frac{1}{2} \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} \sum_{h_{p_2} \in \mathcal{V}(\mathcal{H})} \frac{1}{d_{\mathcal{H}}(h_{p_1}, h_{p_2})}. \end{aligned} \quad (12)$$

As an extension of the Wiener index, Randić put forward the hyper-Wiener index as:

$$WW(\mathcal{H}) = \frac{1}{2} \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} (d_{\mathcal{H}}(h_{p_1}, h_{p_2}) + d_{\mathcal{H}}^2(h_{p_1}, h_{p_2})),$$

where

$$A(\mathcal{H}) = \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} d_{\mathcal{H}}^2(h_{p_1}, h_{p_2})$$

and

$$DD_{\mathcal{H}}(h_{p_1}) = \sum_{h_{p_2} \in \mathcal{V}(\mathcal{H})} d_{\mathcal{H}}^2(h_{p_1}, h_{p_2}),$$

then

$$\begin{aligned} WW(\mathcal{H}) &= \frac{1}{2} (W(\mathcal{H}) + A(\mathcal{H})) \\ &= \frac{1}{4} \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} D_{\mathcal{H}}(h_{p_1}) \\ &\quad + \frac{1}{4} \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} DD_{\mathcal{H}}(h_{p_1}). \end{aligned} \quad (13)$$

Dobrynin and Kochetova [21] introduced the degree-distance index of a graph  $\mathcal{H}$  as follows:

$$\begin{aligned} DD(\mathcal{H}) &= \sum_{\{h_{p_1}, h_{p_2}\} \subseteq \mathcal{V}(\mathcal{H})} (\deg_{\mathcal{H}}(h_{p_1}) + \deg_{\mathcal{H}}(h_{p_2}))d_{\mathcal{H}} \\ &\quad \times (h_{p_1}, h_{p_2}) \\ &= \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} \deg_{\mathcal{H}}(h_{p_1})D_{\mathcal{H}}(h_{p_1}) \\ &= \sum_{h_{p_1} \in \mathcal{V}(\mathcal{H})} \sum_{h_{p_2} \in \mathcal{V}(\mathcal{H})} \deg_{\mathcal{H}}(h_{p_1})d_{\mathcal{H}}(h_{p_1}, h_{p_2}). \end{aligned} \quad (14)$$

For the in depth study of these descriptors and other famous topological descriptors, we recommended the reader to [1]–[4], [23]–[25], [32], [33], [35]–[37], [39]–[41], [52]. Now we state certain properties of the subdivision vertex-edge join of three graphs in the next lemma.

*Lemma 1:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be graphs. Then we have:

$$1) |\mathcal{V}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^Y \cup \mathcal{H}_s^X))| = n_1 + m_1 + n_2 + n_3, \text{ and } |\mathcal{E}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^Y \cup \mathcal{H}_s^X))| = 2m_1 + n_1n_2 + m_1n_3 + m_2 + m_3.$$

$$2) \deg_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^Y \cup \mathcal{H}_s^X)}(h) = \begin{cases} \deg_{\mathcal{H}_q}(h) + n_2, & \text{if } h \in \mathcal{V}(\mathcal{H}_q), \\ n_3 + 2, & \text{if } h \in \mathcal{I}(\mathcal{H}_q), \\ \deg_{\mathcal{H}_r}(z) + n_1, & \text{if } h \in \mathcal{V}(\mathcal{H}_r), \\ \deg_{\mathcal{H}_s}(z) + m_1, & \text{if } h \in \mathcal{V}(\mathcal{H}_s). \end{cases}$$

$$3) d_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^Y \cup \mathcal{H}_s^X)}(h_{p_1}, h_{p_2}) = \begin{cases} 0, & \text{if } h_{p_1} = h_{p_2}, \\ 1, & \text{if } h_{p_1}h_{p_2} \in \mathcal{E}(\mathcal{H}_i), \quad i = 2, 3, \\ & \text{or } h_{p_1}h_{p_2} \in \mathcal{E}(\mathcal{S}(\mathcal{H}_q)), \quad h_{p_1} \in \mathcal{V}(\mathcal{H}_q), \\ & \quad h_{p_2} \in \mathcal{I}(\mathcal{H}_q) \text{ or } h_{p_1} \in \mathcal{V}(\mathcal{H}_q) \\ & \quad \text{and } h_{p_2} \in \mathcal{V}(\mathcal{H}_r), \\ & \text{or } h_{p_1} \in \mathcal{I}(\mathcal{H}_q) \text{ and } h_{p_2} \in \mathcal{V}(\mathcal{H}_s), \\ 2, & h_{p_1}h_{p_2} \notin \mathcal{E}(\mathcal{H}_i), \quad i = 2, 3, \\ & \text{or } h_{p_1}, h_{p_2} \in \mathcal{V}(\mathcal{H}_q), \text{ or } h_{p_1}, h_{p_2} \in \mathcal{I}(\mathcal{H}_q), \\ & \text{or } h_{p_1} \in \mathcal{I}(\mathcal{H}_q) \text{ and } h_{p_2} \in \mathcal{V}(\mathcal{H}_r), \\ & \text{or } h_{p_1} \in \mathcal{V}(\mathcal{H}_q) \text{ and } h_{p_2} \in \mathcal{V}(\mathcal{H}_r). \\ 3, & \text{if } h_{p_1}h_{p_2} \notin \mathcal{E}(\mathcal{S}(\mathcal{H}_q)), \\ & \quad h_{p_1} \in \mathcal{V}(\mathcal{H}_q) \text{ and } h_{p_2} \in \mathcal{I}(\mathcal{H}_q) \\ & \quad \text{or } h_{p_1} \in \mathcal{V}(\mathcal{H}_r) \text{ and } h_{p_2} \in \mathcal{V}(\mathcal{H}_s). \end{cases}$$

$$\begin{aligned}
 & 4) \text{ } ec_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)}(h) \\
 & = \begin{cases} 2, & \text{if } h \in \mathcal{V}(\mathcal{H}_q) \text{ or } h \in \mathcal{I}(\mathcal{H}_q), \\ 3, & \text{if } h \in \mathcal{V}(\mathcal{H}_r) \text{ or } h \in \mathcal{V}(\mathcal{H}_s). \end{cases} \\
 & 5) S_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)}(h) \\
 & = \begin{cases} S_{\mathcal{H}_r}(h) + n_1 \text{deg}_{\mathcal{H}_r}(h) \\ + 2m_1 + n_1 n_2, & \text{if } h \in \mathcal{V}(\mathcal{H}_r), \\ S_{\mathcal{H}_s}(h) + m_1 \text{deg}_{\mathcal{H}_s}(h) \\ + m_1(n_3 + 2), & \text{if } h \in \mathcal{V}(\mathcal{H}_s), \\ \text{deg}_{\mathcal{H}_q}(h)(n_3 + 2) \\ + 2m_2 + n_1 n_2, & \text{if } h \in \mathcal{V}(\mathcal{H}_q), \\ \text{deg}_{\mathcal{H}_q}(h_{p_1}) \\ + \text{deg}_{\mathcal{H}_q}(h_{p_2}) \\ + 2n_2 + 2m_3 + m_1 n_3, & \text{if } h = h_{p_1} h_{p_2} \in \mathcal{I}(\mathcal{H}_q), \end{cases}
 \end{aligned}$$

II. MAIN RESULTS

This section provides the results associated to various distance based indices of the subdivision vertex-edge join of graphs. In next theorem, we present the formulae for eccentric connectivity and connective eccentricity indices of subdivision vertex-edge join for three graphs.

*Theorem 1:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

- 1)  $\xi^c(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = 5m_1 n_3 + 5n_1 n_2 + 8m_1 + 6m_2 + 6m_3.$
- 2)  $\xi^{ce}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = \frac{1}{6}(5m_1 n_3 + 5n_1 n_2 + 12m_1 + 4m_2 + 4m_3).$

*Proof:*

- 1) By using Lemma 1 in Equation (1), we get

$$\begin{aligned}
 & \xi^c(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) \\
 & = \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} 3(\text{deg}_{\mathcal{H}_s}(h_s) + m_1) \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3(\text{deg}_{\mathcal{H}_r}(h_r) + n_1) \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} 2(\text{deg}_{\mathcal{H}_q}(h_q) + n_2) \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} 2(n_3 + 2) \\
 & = 3(2m_3 + m_1 n_3) + 3(2m_2 + n_1 n_2) \\
 & + 2(2m_1 + n_1 n_2) + 2m_1(n_3 + 2) \\
 & = 5m_1 n_3 + 5n_1 n_2 + 8m_1 + 6m_2 \\
 & + 6m_3.
 \end{aligned}$$

- 2) By using Lemma 1 in Equation (2), we get

$$\xi^{ce}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))$$

$$\begin{aligned}
 & = \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \frac{\text{deg}_{\mathcal{H}_s}(h_s) + m_1}{3} \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \frac{\text{deg}_{\mathcal{H}_r}(h_r) + n_1}{3} \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \frac{\text{deg}_{\mathcal{H}_q}(h_q) + n_2}{2} \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \frac{n_3 + 2}{2} \\
 & = \frac{1}{3}(2m_3 + m_1 n_3) + \frac{1}{3}(2m_2 \\
 & + n_1 n_2) + \frac{1}{2}(2m_1 + n_1 n_2) \\
 & + \frac{1}{2}m_1(n_3 + 2) \\
 & = \frac{1}{6}(5m_1 n_3 + 5n_1 n_2 + 12m_1 \\
 & + 4m_2 + 4m_3).
 \end{aligned}$$

This complete the proof. □

Now, we set up the precise values of the total eccentricity and average eccentricity indices of subdivision vertex-edge join for three graphs.

*Theorem 2:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

- 1)  $\tau(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = 3n_3 + 3n_2 + 2n_1 + 2m_1.$
- 2)  $aveg(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = \frac{3n_3 + 3n_2 + 2n_1 + 2m_1}{n_1 + n_2 + n_3 + m_1}.$

*Proof:*

- 1) By using Lemma 1 in Equation (3), we get

$$\begin{aligned}
 \tau(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} 3 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3 \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} 2 + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} 2 \\
 & = 3n_3 + 3n_2 + 2n_1 + 2m_1.
 \end{aligned}$$

- 2) By using Lemma 1 in Equation (4), we obtain

$$\begin{aligned}
 aveg(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = \frac{\tau(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))}{n_1 + n_2 + n_3 + m_1} \\
 & = \frac{3n_3 + 3n_2 + 2n_1 + 2m_1}{n_1 + n_2 + n_3 + m_1}.
 \end{aligned}$$

This complete the proof. □

*Theorem 3:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then

- 1)  $\mathcal{M}_1(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = 9n_3 + 9n_2 + 4n_1 + 4m_1.$
- 2)  $\mathcal{M}_2(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = 9m_3 + 9m_2 + 6n_1 n_2 + 6n_3 m_1 + 8m_1.$

*Proof:*

- 1) By using Lemma 1 in Equation (5), we get

$$\mathcal{M}_1(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))$$

$$\begin{aligned}
 &= \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (3)^2 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (3)^2 \\
 &\quad + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (2)^2 + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (2)^2 \\
 &= 9n_3 + 9n_2 + 4n_1 + 4m_1.
 \end{aligned}$$

2) By using Lemma 1 in Equation (5), we obtain

$$\begin{aligned}
 &\mathcal{M}_2(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) \\
 &= \sum_{h_{s_1} h_{s_2} \in \mathcal{E}(\mathcal{H}_s)} 3 \times 3 \\
 &\quad + \sum_{h_{r_1} h_{r_2} \in \mathcal{E}(\mathcal{H}_r)} 3 \times 3 \\
 &\quad + \sum_{h_{q_1} \in \mathcal{V}(\mathcal{H}_q)} \sum_{h_{r_1} \in \mathcal{V}(\mathcal{H}_r)} 2 \times 3 \\
 &\quad + \sum_{h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_{s_1} \in \mathcal{V}(\mathcal{H}_s)} 2 \times 3 \\
 &\quad + \sum_{\substack{h_{q_1} h'_{q_1} \in \mathcal{E}(\mathcal{S}(\mathcal{H}_q)), \\ h_{q_1} \in \mathcal{V}(\mathcal{H}_q), h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)}} 2 \times 2 \\
 &= 9m_3 + 9m_2 + 6n_1 n_2 + 6n_3 m_1 \\
 &\quad + 8m_1.
 \end{aligned}$$

This complete the proof.  $\square$

*Theorem 4:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

- 1)  $\mathcal{GA}_{ecc}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = m_2 + m_3 + 2m_1 + \frac{2\sqrt{6}}{5}(n_1 n_2 + m_1 n_3)$ .
- 2)  $\mathcal{ABC}_{ecc}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) = \frac{2}{3}(m_2 + m_3) + \frac{1}{\sqrt{2}}(n_1 n_2 + m_1 n_3 + 2m_1)$ .

*Proof:*

1) By using Lemma 1 in Equation (6), we obtain

$$\begin{aligned}
 &\mathcal{GA}_{ecc}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) \\
 &= \sum_{h_{s_1} h_{s_2} \in \mathcal{E}(\mathcal{H}_s)} \frac{2\sqrt{3} \times 3}{3 + 3} + \sum_{h_{r_1} h_{r_2} \in \mathcal{E}(\mathcal{H}_r)} \frac{2\sqrt{3} \times 3}{3 + 3} \\
 &\quad + \sum_{h_{q_1} \in \mathcal{V}(\mathcal{H}_q)} \sum_{h_{r_1} \in \mathcal{V}(\mathcal{H}_r)} \frac{2\sqrt{2} \times 3}{2 + 3} \\
 &\quad + \sum_{h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_{s_1} \in \mathcal{V}(\mathcal{H}_s)} \frac{2\sqrt{2} \times 3}{2 + 3} \\
 &\quad + \sum_{\substack{h_{q_1} h'_{q_1} \in \mathcal{E}(\mathcal{S}(\mathcal{H}_q)), \\ h_{q_1} \in \mathcal{V}(\mathcal{H}_q), h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)}} \frac{2\sqrt{2} \times 2}{2 + 2} \\
 &= m_2 + m_3 + \frac{2\sqrt{6}}{5} n_1 n_2 + \frac{2\sqrt{6}}{5} m_1 n_3 + 2m_1.
 \end{aligned}$$

2) By using Lemma 1 in Equation (7), we obtain

$$\mathcal{ABC}_{ecc}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))$$

$$\begin{aligned}
 &= \sum_{h_{s_1} h_{s_2} \in \mathcal{E}(\mathcal{H}_s)} \sqrt{\frac{3 + 3 - 2}{3 \times 3}} \\
 &\quad + \sum_{h_{r_1} h_{r_2} \in \mathcal{E}(\mathcal{H}_r)} \sqrt{\frac{3 + 3 - 2}{3 \times 3}} \\
 &\quad + \sum_{h_{q_1} \in \mathcal{V}(\mathcal{H}_q)} \sum_{h_{r_1} \in \mathcal{V}(\mathcal{H}_r)} \sqrt{\frac{2 + 3 - 2}{2 \times 3}} \\
 &\quad + \sum_{h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_{s_1} \in \mathcal{V}(\mathcal{H}_s)} \sqrt{\frac{2 + 3 - 2}{2 \times 3}} \\
 &\quad + \sum_{\substack{h_{q_1} h'_{q_1} \in \mathcal{E}(\mathcal{S}(\mathcal{H}_q)), \\ h_{q_1} \in \mathcal{V}(\mathcal{H}_q), h'_{q_1} \in \mathcal{I}(\mathcal{H}_q)}} \sqrt{\frac{2 + 2 - 2}{2 \times 2}} \\
 &= \frac{2}{3} m_2 + \frac{2}{3} m_3 + \frac{1}{\sqrt{2}} n_1 n_2 + \frac{1}{\sqrt{2}} m_1 n_3 \\
 &\quad + \left(\frac{1}{\sqrt{2}}\right) 2m_1.
 \end{aligned}$$

This complete the proof.  $\square$

*Theorem 5:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned}
 &\xi^{ad}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) \\
 &= \frac{1}{6}(3\mathcal{M}_1(\mathcal{H}_q) + 2\mathcal{M}_1(\mathcal{H}_r) + 2\mathcal{M}_1(\mathcal{H}_s)) \\
 &\quad + \frac{5}{3}(n_1 m_2 + m_1 n_2 + m_1 m_3 + n_3 m_1) \\
 &\quad + \frac{1}{6} n_1 n_2 (2n_2 + 3n_1) + \frac{1}{6} n_3 m_1 (2n_3 \\
 &\quad + 3m_1) + 2m_1.
 \end{aligned}$$

*Proof:* By Lemma 1 in Equation (8), we get

$$\begin{aligned}
 &\xi^{ad}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) \\
 &= \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \frac{S_{\mathcal{H}_s}(h_s) + m_1 \deg_{\mathcal{H}_s}(h_s) + m_1(n_3 + 2)}{3} \\
 &\quad + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \frac{S_{\mathcal{H}_r}(h_r) + n_1 \deg_{\mathcal{H}_r}(h_r) + 2m_1 + n_1 n_2}{3} \\
 &\quad + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \frac{\deg_{\mathcal{H}_q}(h_q)(n_3 + 2) + 2m_2 + n_1 n_2}{2} \\
 &\quad + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \frac{\deg_{\mathcal{H}_q}(h_{q_1}) + \deg_{\mathcal{H}_q}(h_{q_2}) + 2n_2 + 2m_3 + m_1 n_3}{2} \\
 &= \frac{1}{3} (\mathcal{M}_1(\mathcal{H}_s) + 2m_1 m_3 + n_3 m_1 (n_3 + 2)) + \frac{1}{3} (\mathcal{M}_1(\mathcal{H}_r) \\
 &\quad + 2n_1 m_2 + 2n_2 m_1 + n_1 n_2^2) \\
 &\quad + \frac{1}{2} (2m_1 (n_3 + 2) + 2m_2 n_1 + n_1^2 n_2) \\
 &\quad + \frac{1}{2} (\mathcal{M}_1(\mathcal{H}_q) + 2n_2 m_1 + 2m_3 m_1 + m_1^2 n_3)
 \end{aligned}$$

After simplification we acquire required result. This complete the proof.  $\square$

**Theorem 6:** Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned} \xi_c(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= 2\mathcal{M}_1(\mathcal{H}_q) + 3\mathcal{M}_1(\mathcal{H}_r) + 3\mathcal{M}_1(\mathcal{H}_s) \\ &\quad + 10(n_1m_2 + m_1n_2 + m_1m_3 + n_3m_1) \\ &\quad + n_1n_2(2n_1 + 3n_2) + n_3m_1(3n_3 + 2m_1) \\ &\quad + 8m_1. \end{aligned}$$

*Proof:* By Lemma 1 in Equation (9), we get

$$\begin{aligned} \xi_c(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} 3(S_{\mathcal{H}_s}(h_s) + m_1 \deg_{\mathcal{H}_s}(h_s) + m_1(n_3 + 2)) \\ &\quad + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3(S_{\mathcal{H}_r}(h_r) + n_1 \deg_{\mathcal{H}_r}(h_r) + 2m_1 + n_1n_2) \\ &\quad + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} 2(\deg_{\mathcal{H}_q}(h_q)(n_3 + 2) + 2m_2 + n_1n_2) \\ &\quad + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} 2(\deg_{\mathcal{H}_q}(h'_q) + \deg_{\mathcal{H}_q} \\ &\quad \times (h_{q_2}) + 2n_2 + 2m_3 + m_1n_3) \\ &= 3(\mathcal{M}_1(\mathcal{H}_s) + 2m_1m_3 + n_3m_1(n_3 + 2)) + 3(\mathcal{M}_1(\mathcal{H}_r) \\ &\quad + 2n_1m_2 + 2n_2m_1 + n_1n_2^2) \\ &\quad + 2(2m_2(n_3 + 2) + 2m_2n_1 + n_1^2n_2) + 2(\mathcal{M}_1(\mathcal{H}_q) \\ &\quad + 2n_2m_1 + 2m_3m_1 + m_1^2n_3) \end{aligned}$$

After simplification we get required result. This complete the proof.  $\square$

**Theorem 7:** Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned} \xi^{ds}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= 6n_3(n_3 - 1) + 6n_2(n_2 - 1) + 4n_1(n_1 - 1) \\ &\quad + 5n_1n_2 + 10n_2m_1 + 4m_1^2 + 18n_2n_3 \\ &\quad + 5m_1n_3 + 10n_1n_3 + 12n_1m_1 + 4m_1^2 - 20m_1 \\ &\quad - 6m_2 - 6m_3. \end{aligned}$$

*Proof:* By Lemma 1 in Equation (10), we get

$$\begin{aligned} \xi^{ds}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in N_{\mathcal{H}_s}(h_s) \cup \mathcal{I}(\mathcal{H}_q)} 3 \times 1 \right. \\ &\quad + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in (\mathcal{V}(\mathcal{H}_s) \setminus N_{\mathcal{H}_s}(h_s)) \cup \mathcal{V}(\mathcal{H}_q)} 3 \times 2 \\ &\quad + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3 \times 3 \left. \right) \\ &\quad + \left( \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in N_{\mathcal{H}_r}(h_r) \cup \mathcal{V}(\mathcal{H}_q)} 3 \times 1 \right. \\ &\quad + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in (\mathcal{V}(\mathcal{H}_r) \setminus N_{\mathcal{H}_r}(h_r)) \cup \mathcal{I}(\mathcal{H}_q)} 3 \times 2 \end{aligned}$$

$$\begin{aligned} &\quad + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} 3 \times 3 \left. \right) \\ &\quad + \left( \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{H}_q}(h_q) \cup \mathcal{V}(\mathcal{H}_r)} 2 \times 1 \right. \\ &\quad + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in (\mathcal{V}(\mathcal{H}_q) \setminus \{h_q\}) \cup \mathcal{V}(\mathcal{H}_s)} 2 \times 2 \\ &\quad + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q) \setminus N_{\mathcal{H}_q}(h'_q)} 2 \times 3 \left. \right) \\ &\quad + \left( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{H}_q}(h'_q) \cup \mathcal{V}(\mathcal{H}_s)} 2 \times 1 \right. \\ &\quad + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in (\mathcal{I}(\mathcal{H}_q) \setminus \{h'_q\}) \cup \mathcal{V}(\mathcal{H}_r)} 2 \times 2 \\ &\quad + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q) \setminus N_{\mathcal{H}_q}(h'_q)} 2 \times 3 \left. \right) \\ &= \left( 3 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}(h_s) + m_1) + 6 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (n_3 - 1 \right. \\ &\quad \left. - \deg_{\mathcal{H}_s}(h_s) + n_1) + 9 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} n_2 \right) \\ &\quad + \left( 3 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}(h_r) + n_1) \right. \\ &\quad + 6 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (n_2 - 1 - \deg_{\mathcal{H}_r}(h_r) + m_1) + 9 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} n_3 \left. \right) \\ &\quad + \left( 2 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) + n_2) + 4 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (n_1 - 1 + n_3) \right. \\ &\quad + 6 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (m_1 - \deg_{\mathcal{H}_q}(h_q)) \left. \right) + \left( 2 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2) \right. \\ &\quad + 4 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (m_1 - 1 + n_2) + 6 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_1 - 2) \left. \right) \\ &= 3(2m_3 + m_1n_3) + 6(n_3^2 - n_3 - 2m_3 + n_1n_3) + 9n_2n_3 \\ &\quad + 3(2m_2 + n_1n_2) + 6(n_2^2 - n_2 - 2m_2 + n_2m_1) + 9n_2n_3 \\ &\quad + 2(2m_1 + n_1n_2) + 4(n_1^2 - n_1 + n_1n_3) + 6(n_1m_1 - 2m_1) \\ &\quad + 2m_1(n_3 + 2) + 4m_1(m_1 - 1 + n_2) + 6m_1(n_1 - 2) \\ &= 6n_3(n_3 - 1) + 6n_2(n_2 - 1) + 4n_1(n_1 - 1) + 5n_1n_2 \\ &\quad + 10n_2m_1 + 4m_1^2 + 18n_2n_3 + 5m_1n_3 + 10n_1n_3 \\ &\quad + 12n_1m_1 + 4m_1^2 - 20m_1 - 6m_2 - 6m_3. \end{aligned}$$

This complete the proof.  $\square$

**Theorem 8:** Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned} W(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= (n_1^2 + n_2^2 + n_3^2 + m_1^2) - (n_1 + n_2 + n_3 \\ &\quad + 5m_1) - (m_2 + m_3) + n_1n_2 + 3n_2n_3 \end{aligned}$$

$$+ 2n_2m_1 + m_1n_3 + 2n_1n_3 + 3n_1m_1.$$

*Proof:* By Lemma 1 in Equation (11), we get

$$\begin{aligned} W(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in N_{\mathcal{H}_s}(h_s) \cup \mathcal{I}(\mathcal{H}_q)} 1 \right. \\ &+ \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in (\mathcal{V}(\mathcal{H}_s) \setminus N_{\mathcal{H}_s}(h_s)) \cup \mathcal{V}(\mathcal{H}_q)} 2 \\ &+ \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3 \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in N_{\mathcal{H}_r}(h) \cup \mathcal{V}(\mathcal{H}_q)} 1 \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in (\mathcal{V}(\mathcal{H}_r) \setminus N_{\mathcal{H}_r}(h_r)) \cup \mathcal{I}(\mathcal{H}_q)} 2 \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in \mathcal{V}(\mathcal{H}_s)} 3 \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{S}(\mathcal{H}_q)}(h_q) \cup \mathcal{V}(\mathcal{H}_r)} 1 \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in (\mathcal{V}(\mathcal{H}_q) \setminus \{h_q\}) \cup \mathcal{V}(\mathcal{H}_s)} 2 \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q) \setminus N_{\mathcal{S}(\mathcal{H}_q)}(h_q)} 3 \\ &+ \left( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{S}(\mathcal{H}_q)}(h'_q) \cup \mathcal{V}(\mathcal{H}_s)} 1 \right. \\ &+ \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in (\mathcal{I}(\mathcal{H}_q) \setminus \{h'_q\}) \cup \mathcal{V}(\mathcal{H}_r)} 2 \\ &+ \left. \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q) \setminus N_{\mathcal{S}(\mathcal{H}_q)}(h'_q)} 3 \right) \\ &= \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}(h_s) + m_1) \right. \\ &+ 2 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (n_3 - 1 - \deg_{\mathcal{H}_s}(h_s) + n_1) \\ &+ 3 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} n_2 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}(h_r) \\ &+ n_1) + 2 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (n_2 - 1 - \deg_{\mathcal{H}_r}(h_r) \\ &+ m_1) + 3 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} n_3 + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) \\ &+ n_2) + 2 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (n_1 - 1 + n_3) \\ &+ 3 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (m_1 - \deg_{\mathcal{H}_q}(h_q)) \\ &+ \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2) + 2 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (m_1 \end{aligned}$$

$$\begin{aligned} &- 1 + n_2) + 3 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_1 - 2) \\ &= \frac{1}{2} \left( 2m_3 + m_1n_3 + 2n_3^2 - 2n_3 - 4m_3 + 2n_1n_3 + 3n_2n_3 \right. \\ &+ 2m_2 + n_1n_2 + 2n_2^2 - 2n_2 - 4m_2 + 2n_2m_1 + 3n_2n_3 \\ &+ 2m_1 + n_1n_2 + 2n_1^2 - 2n_1 + 2n_1n_3 + 3n_1m_1 - 6m_1 \\ &+ m_1(n_3 + 2) + 2m_1(m_1 - 1 + n_2) + 3m_1(n_1 - 2) \left. \right) \\ &= (n_1^2 + n_2^2 + n_3^2 + m_1^2) - (n_1 + n_2 + n_3 + 5m_1) \\ &- (m_2 + m_3) + n_1n_2 + 3n_2n_3 + 2n_2m_1 + m_1n_3 \\ &+ 2n_1n_3 + 3n_1m_1. \end{aligned}$$

This complete the proof.  $\square$

*Theorem 9:* Let  $\mathcal{H}_q$ ,  $\mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned} H(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= \frac{1}{2} \left( \frac{13}{6}m_1 + m_2 + m_3 + 2n_1n_2 + \frac{1}{2}n_1(n_1 \right. \\ &- 1) + \frac{1}{2}n_2(n_2 - 1) + \frac{1}{2}n_3(n_3 - 1) \\ &+ m_1 \left( n_2 + 2n_3 + \frac{2}{3}n_1 \right) + n_3 \left( \frac{2}{3}n_2 + \frac{1}{2}n_1 \right) \\ &\left. + \frac{1}{2}m_1^2 \right). \end{aligned}$$

*Proof:* By Lemma 1 in Equation (12), we get

$$\begin{aligned} H(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) &= \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in N_{\mathcal{H}_s}(h_s) \cup \mathcal{I}(\mathcal{H}_q)} \frac{1}{1} \right. \\ &+ \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in (\mathcal{V}(\mathcal{H}_s) \setminus N_{\mathcal{H}_s}(h_s)) \cup \mathcal{V}(\mathcal{H}_q)} \frac{1}{2} \\ &+ \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \frac{1}{3} \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in N_{\mathcal{H}_r}(h_r) \cup \mathcal{V}(\mathcal{H}_q)} \frac{1}{1} \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in (\mathcal{V}(\mathcal{H}_r) \setminus N_{\mathcal{H}_r}(h_r)) \cup \mathcal{I}(\mathcal{H}_q)} \frac{1}{2} \\ &+ \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \frac{1}{3} \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{S}(\mathcal{H}_q)}(h_q) \cup \mathcal{V}(\mathcal{H}_r)} \frac{1}{1} \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in (\mathcal{V}(\mathcal{H}_q) \setminus \{h_q\}) \cup \mathcal{V}(\mathcal{H}_s)} \frac{1}{2} \\ &+ \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q) \setminus N_{\mathcal{S}(\mathcal{H}_q)}(h_q)} \frac{1}{3} \end{aligned}$$

$$\begin{aligned}
 & + \left( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in N_{S(\mathcal{H}_q)}(h'_q) \cup \mathcal{V}(\mathcal{H}_s)} \frac{1}{1} \right. \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in (\mathcal{I}(\mathcal{H}_q) \setminus \{h'_q\}) \cup \mathcal{V}(\mathcal{H}_r)} \frac{1}{2} \\
 & \left. + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q) \setminus N_{S(\mathcal{H}_q)}(h'_q)} \frac{1}{3} \right) \\
 = & \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\text{deg}_{\mathcal{H}_s}(h_s) + m_1) \right. \\
 & + \frac{1}{2} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (n_3 - 1 - \text{deg}_{\mathcal{H}_s}(h_s) + n_1) \\
 & + \frac{1}{3} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} n_2 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\text{deg}_{\mathcal{H}_r}(h_r) + n_1) \\
 & + \frac{1}{2} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (n_2 - 1 - \text{deg}_{\mathcal{H}_r}(h_r) + m_1) \\
 & + \frac{1}{3} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} n_3 + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\text{deg}_{\mathcal{H}_q}(h_q) + n_2) \\
 & + \frac{1}{2} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (n_1 - 1 + n_3) + \frac{1}{3} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (m_1 - \text{deg}_{\mathcal{H}_q}(h_q)) \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2) + \frac{1}{2} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (m_1 - 1 + n_2) \\
 & \left. + \frac{1}{3} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_1 - 2) \right) \\
 = & \frac{1}{2} \left( 2m_3 + m_1n_3 + \frac{1}{2}(n_3^2 - n_3 - 2m_3 + n_1n_3) + \frac{1}{3}n_2n_3 \right. \\
 & + 2m_2 + n_1n_2 + \frac{1}{2}(n_2^2 - n_2 - 2m_2 + n_2m_1) + \frac{1}{3}n_2n_3 \\
 & + 2m_1 + n_1n_2 + \frac{1}{2}n_1(n_1 - 1 + n_3) + \frac{1}{3}n_1m_1 - \frac{2}{3}m_1 \\
 & \left. + m_1(n_3 + 2) + \frac{1}{2}m_1(m_1 - 1 + n_2) + \frac{1}{3}m_1(n_1 - 2) \right) \\
 = & \frac{1}{2} \left( \frac{13}{6}m_1 + m_2 + m_3 + 2n_1n_2 + \frac{1}{2}n_1(n_1 - 1) \right. \\
 & + \frac{1}{2}n_2(n_2 - 1) + \frac{1}{2}n_3(n_3 - 1) + m_1 \left( n_2 + 2n_3 + \frac{2}{3}n_1 \right) \\
 & \left. + n_3 \left( \frac{2}{3}n_2 + \frac{1}{2}n_1 \right) + \frac{1}{2}m_1^2 \right).
 \end{aligned}$$

This complete the proof.  $\square$   
**Theorem 10:** Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned}
 WW(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = \frac{3}{2}(n_1^2 + n_2^2 + n_3^2 + m_1^2) - \frac{3}{2}(n_1 + n_2 \\
 & + n_3) + \frac{23}{2}m_1 - 2(m_2 + m_3) + n_1n_2 \\
 & + 6n_2n_3 + 3n_2m_1 + m_1n_3 + 3n_1n_3
 \end{aligned}$$

$$+ 6n_1m_1.$$

*Proof:* By Lemma 1 in Equation (13), we need to find only  $A(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))$ .

$$\begin{aligned}
 A(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = \frac{1}{2} \left( \sum_{h_1 \in \mathcal{V}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))} DD_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)}(h_1) \right) \\
 & = \frac{1}{2} \left( \sum_{h_1 \in \mathcal{V}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))} \sum_{h_2 \in \mathcal{V}(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I))} d_{\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)}^2(h_1, h_2) \right) \\
 & = \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in N_{\mathcal{H}_s}(h_s) \cup \mathcal{I}(\mathcal{H}_q)} (1)^2 \right. \\
 & + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in (\mathcal{V}(\mathcal{H}_s) \setminus N_{\mathcal{H}_s}(h_s)) \cup \mathcal{V}(\mathcal{H}_q)} (2)^2 \\
 & + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (3)^2 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h_r \in N_{\mathcal{H}_r}(h_r) \cup \mathcal{V}(\mathcal{H}_q)} (1)^2 \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in (\mathcal{V}(\mathcal{H}_r) \setminus N_{\mathcal{H}_r}(h_r)) \cup \mathcal{I}(\mathcal{H}_q)} (2)^2 \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (3)^2 \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in N_{S(\mathcal{H}_q)}(h_q) \cup \mathcal{V}(\mathcal{H}_r)} (1)^2 \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in (\mathcal{V}(\mathcal{H}_q) \setminus \{h_q\}) \cup \mathcal{V}(\mathcal{H}_s)} (2)^2 \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q) \setminus N_{S(\mathcal{H}_q)}(h_q)} (3)^2 \\
 & + \left( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in N_{S(\mathcal{H}_q)}(h'_q) \cup \mathcal{V}(\mathcal{H}_s)} (1)^2 \right. \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in (\mathcal{I}(\mathcal{H}_q) \setminus \{h'_q\}) \cup \mathcal{V}(\mathcal{H}_r)} (2)^2 \\
 & \left. + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q) \setminus N_{S(\mathcal{H}_q)}(h'_q)} (3)^2 \right) \\
 & = \frac{1}{2} \left( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\text{deg}_{\mathcal{H}_s}(h_s) + m_1) + 4 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (n_3 - 1 \right. \\
 & - \text{deg}_{\mathcal{H}_s}(h_s) + n_1) + 9 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} n_2 + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\text{deg}_{\mathcal{H}_r}(h_r) \\
 & + n_1) + 4 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (n_2 - 1 - \text{deg}_{\mathcal{H}_r}(h_r) + m_1) \\
 & + 9 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} n_3 + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\text{deg}_{\mathcal{H}_q}(h_q) + n_2) \\
 & + 4 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (n_1 - 1 + n_3) + 9 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (m_1 - \text{deg}_{\mathcal{H}_q}(h_q))
 \end{aligned}$$



$$\begin{aligned}
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2) + 4 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (m_1 - 1 + n_2) \\
 & + 9 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_1 - 2) \Big) \\
 = & \frac{1}{2} \Big( 2m_3 + m_1n_3 + 4n_3^2 - 4n_3 - 8m_3 + 4n_1n_3 \\
 & + 9n_2n_3 + 2m_2 + n_1n_2 + 4n_2^2 - 4n_2 - 8m_2 + 4n_2m_1 \\
 & + 9n_2n_3 + 2m_1 + n_1n_2 + 4n_1^2 - 4n_1 + 4n_1n_3 + 9n_1m_1 \\
 & - 18m_1 + m_1(n_3 + 2) + 4m_1(m_1 - 1 + n_2) \\
 & + 9m_1(n_1 - 2) \Big) \\
 = & 2(n_1^2 + n_2^2 + n_3^2 + m_1^2) - 2(n_1 + n_2 + n_3 + 5m_1) \\
 & - 3(m_2 + m_3) + n_1n_2 + 9n_2n_3 + 4n_2m_1 + m_1n_3 \\
 & + 4n_1n_3 + 9n_1m_1. \tag{15}
 \end{aligned}$$

Therefore by using Theorem 8 and (15) in (13), we get the required result. This complete the proof.  $\square$

*Theorem 11:* Let  $\mathcal{H}_q, \mathcal{H}_r$  and  $\mathcal{H}_s$  be three graphs. Then we have

$$\begin{aligned}
 DD(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = -\mathcal{M}_1(\mathcal{H}_s) - \mathcal{M}(\mathcal{H}_r) - 2\mathcal{M}_1(\mathcal{H}_q) \\
 & + n_1n_2(n_1 + n_2 - 4) + 4(n_1 - 1)(m_1 \\
 & + m_2 + m_3) + 3m_1n_3(m_1 + n_3 - 2) \\
 & + 6(m_1n_2 - n_2m_1 + n_3m_2 + n_2m_3) \\
 & + 4n_3m_3 + 3n_1n_2(n_3 + m_1) + 5n_3m_1(n_1 \\
 & + n_2) + 2m_1(5m_1 - 6).
 \end{aligned}$$

*Proof:* By Lemma 1 in Equation (14), we get

$$\begin{aligned}
 DD(\mathcal{H}_q^S \triangleright (\mathcal{H}_r^V \cup \mathcal{H}_s^I)) & = \Big( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in N_{\mathcal{H}_s}(h_s) \cup \mathcal{I}(\mathcal{H}_q)} 1(\deg_{\mathcal{H}_s}(h_s) + m_1) \\
 & + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h \in (\mathcal{V}(\mathcal{H}_s) \setminus N_{\mathcal{H}_s}(h_s)) \cup \mathcal{V}(\mathcal{H}_q)} 2(\deg_{\mathcal{H}_s}(h_s) + m_1) \\
 & + \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} 3(\deg_{\mathcal{H}_s}(h_s) + m_1) \Big) \\
 & + \Big( \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in N_{\mathcal{H}_r}(h_r) \cup \mathcal{V}(\mathcal{H}_q)} 1(\deg_{\mathcal{H}_r}(h_r) + n_1) \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h \in (\mathcal{V}(\mathcal{H}_r) \setminus N_{\mathcal{H}_r}(h_r)) \cup \mathcal{I}(\mathcal{H}_q)} 2(\deg_{\mathcal{H}_r}(h_r) + n_1) \\
 & + \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} 3(\deg_{\mathcal{H}_r}(h_r) + n_1) \Big) \\
 & + \Big( \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{H}_q}(h_q) \cup \mathcal{V}(\mathcal{H}_r)} 1(\deg_{\mathcal{H}_q}(h_q) + n_2) \\
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h \in (\mathcal{V}(\mathcal{H}_q) \setminus \{h_q\}) \cup \mathcal{V}(\mathcal{H}_s)} 2(\deg_{\mathcal{H}_q}(h_q) + n_2)
 \end{aligned}$$

$$\begin{aligned}
 & + \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q) \setminus N_{\mathcal{H}_q}(h_q)} 2(\deg_{\mathcal{H}_q}(h_q) + n_2) \Big) \\
 & + \Big( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in N_{\mathcal{H}_q}(h'_q) \cup \mathcal{V}(\mathcal{H}_s)} 1(n_3 + 2) \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h \in (\mathcal{I}(\mathcal{H}_q) \setminus \{h'_q\}) \cup \mathcal{V}(\mathcal{H}_r)} 2(n_3 + 2) \\
 & + \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} \sum_{h_q \in \mathcal{V}(\mathcal{H}_q) \setminus N_{\mathcal{H}_q}(h'_q)} 3(n_3 + 2) \Big) \\
 = & \Big( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}(h_s) + m_1)^2 + 2 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}(h_s) + m_1) \\
 & (n_3 - 1 - \deg_{\mathcal{H}_s}(h_s) + n_1) + 3 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} n_2(\deg_{\mathcal{H}_s}(h_s) + m_1) \Big) \\
 & + \Big( \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}(h_r) + n_1)^2 \\
 & + 2 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}(h_r) + n_1)(n_2 - 1 - \deg_{\mathcal{H}_r}(h_r) + m_1) \\
 & + 3 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} n_3(\deg_{\mathcal{H}_r}(h_r) + n_1) \Big) \\
 & + \Big( \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) + n_2)^2 \\
 & + 2 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) + n_2)(n_1 - 1 + n_3) \\
 & + 3 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) + n_2)(m_1 - \deg_{\mathcal{H}_q}(h_q)) \Big) \\
 & + \Big( \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2)^2 + 2 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2)(m_1 - 1 + n_2) \\
 & + 3 \sum_{h'_q \in \mathcal{I}(\mathcal{H}_q)} (n_3 + 2)(n_1 - 2) \Big) \\
 = & \Big( \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}^2(h_s) + m_1^2 + 2m_1 \deg_{\mathcal{H}_s}(h_s)) \\
 & + 2 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} ((n_3 - 1 + n_1)(\deg_{\mathcal{H}_s}(h_s) + m_1) \\
 & - \deg_{\mathcal{H}_s}^2(h_s) - m_1 \deg_{\mathcal{H}_s}(h_s)) \\
 & + 3n_2 \sum_{h_s \in \mathcal{V}(\mathcal{H}_s)} (\deg_{\mathcal{H}_s}(h_s) + m_1) \Big) \\
 & + \Big( \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}^2(h_r) + n_1^2 + 2n_1 \deg_{\mathcal{H}_r}(h_r)) \\
 & + 2 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} ((n_2 - 1 + m_1)(\deg_{\mathcal{H}_r}(h_r) + n_1) - \deg_{\mathcal{H}_r}^2(h_r) \\
 & - n_1 \deg_{\mathcal{H}_r}(h_r)) + 3n_3 \sum_{h_r \in \mathcal{V}(\mathcal{H}_r)} (\deg_{\mathcal{H}_r}(h_r) + n_1) \Big) \\
 & + \Big( \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}^2(h_q) + n_2^2 + 2n_2 \deg_{\mathcal{H}_q}(h_q))
 \end{aligned}$$

$$\begin{aligned}
& + 2(n_1 - 1 + n_3) \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (\deg_{\mathcal{H}_q}(h_q) + n_2) \\
& + 3 \sum_{h_q \in \mathcal{V}(\mathcal{H}_q)} (m_1 \deg_{\mathcal{H}_q}(h_q) \\
& - \deg_{\mathcal{H}_q}^2(h_q) + n_2 m_1 - n_2 \deg_{\mathcal{H}_q}(h_q)) \\
& + \left( m_1(n_3 + 2)^2 + 2m_1(n_3 + 2)(m_1 - 1 + n_2) \right. \\
& \left. + 3m_1(n_3 + 2)(n_1 - 2) \right) \\
= & \mathcal{M}_1(\mathcal{H}_s) + m_1^2 n_3 + 4m_1 m_3 + 2(n_3 + n_1 - 1) \\
& \times (2m_3 + m_1 n_3) - 2\mathcal{M}_1(\mathcal{H}_s) - 4m_1 m_3 + 6n_2 m_3 \\
& + 3n_2 m_1 n_3 + \mathcal{M}_1(\mathcal{H}_r) + n_1^2 n_2 + 4n_1 m_2 \\
& + 2(n_2 - 1 + n_1) - 2\mathcal{M}_1(\mathcal{H}_r) - 4n_1 m_2 + 6n_3 m_2 \\
& + 3n_1 n_2 n_3 + \mathcal{M}(\mathcal{H}_q) + n_1 n_2^2 + 4n_2 m_1 + 2(n_1 + n_2 - 1) \\
& \times (2m_1 + n_1 n_2) + 3(2m_1^2 - \mathcal{M}_1(\mathcal{H}_q) + n_1 n_2 m_1 - 2n_2 m_1) \\
& + m_1(n_3 + 2)(n_3 + 2 + 2m_1 - 2 + 2n_2 + 3n_1 + 6) \\
= & -\mathcal{M}_1(\mathcal{H}_s) - \mathcal{M}(\mathcal{H}_r) - 2\mathcal{M}_1(\mathcal{H}_q) + n_1 n_2 (n_1 + n_2 - 4) \\
& + 4(n_1 - 1)(m_1 + m_2 + m_3) + 3m_1 n_3 (m_1 + n_3 - 2) \\
& + 6(m_1 n_2 - n_2 m_1 + n_3 m_2 + n_2 m_3) + 4n_3 m_3 \\
& + 3n_1 n_2 (n_3 + m_1) + 5n_3 m_1 (n_1 + n_2) + 2m_1 (5m_1 - 6).
\end{aligned}$$

This complete the proof.  $\square$

### III. CONCLUSION

The analysis of graphs and networks through structural properties is a research topic with growing significance. One of the approaches in investigating structural characteristics is discussing the quantitative measure that encodes structural statistics of the entire network by a real number. In this article, we have provided the results related to the eccentric-connectivity, connective eccentricity, total-eccentricity, average eccentricity, Zagreb eccentricity, eccentric geometric-arithmetic, eccentric atom-bond connectivity, eccentric adjacency, modified eccentric-connectivity, eccentric distance sum, Wiener, Harary, hyper-Wiener and degree distance indices of subdivision vertex-edge join of graphs.

### REFERENCES

- [1] S. Akhter and M. Imran, "The sharp bounds on general sum-connectivity index of four operations on graphs," *J. Inequalities Appl.*, vol. 2016, pp. 241–251, Dec. 2016.
- [2] S. Akhter and M. Imran, "Computing the forgotten topological index of four operations on graphs," *AKCE Int. J. Graphs Combinatorics*, vol. 14, no. 1, pp. 70–79, 2017.
- [3] S. Akhter, M. Imran, W. Gao, and M. R. Farahani, "On topological indices of honeycomb networks and graphene networks," *Hacettepe J. Math. Statist.*, vol. 47, no. 1, pp. 19–35, 2018.
- [4] S. Akhter, M. Imran, and Z. Raza, "Bounds for the general sum-connectivity index of composite graphs," *J. Inequalities Appl.*, vol. 2017, Dec. 2017, Art. no. 76.
- [5] A. R. Ashrafi and M. Ghorbani, "A study of fullerenes by MEC polynomials," *Electron. Mater. Lett.*, vol. 6, no. 2, pp. 87–90, 2010.
- [6] A. R. Ashrafi, M. Ghorbani, and M. A. Hossein-Zadeh, "The eccentric-connectivity polynomial of some graph operations," *Serdica J. Comput.*, vol. 5, no. 2, pp. 101–116, 2011.
- [7] A. T. Balaban, "Applications of graph theory in chemistry," *J. Chem. Inf. Comput. Sci.*, vol. 25, no. 3, pp. 334–343, 1985.
- [8] K. Balasubramanian, "A method for nuclear spin statistics in molecular spectroscopy," *J. Chem. Phys.*, vol. 74, no. 12, pp. 6824–6829, 1981.
- [9] K. Balasubramanian, "Operator and algebraic methods for NMR spectroscopy. I. Generation of NMR spin species," *J. Chem. Phys.*, vol. 78, no. 11, pp. 6358–6368, 1983.
- [10] K. Balasubramanian, "Applications of combinatorics and graph theory to spectroscopy and quantum chemistry," *Chem. Rev.*, vol. 85, no. 6, pp. 599–618, 1985.
- [11] K. Balasubramanian, "Characteristic polynomials of organic polymers and periodic structures," *J. Comput. Chem.*, vol. 6, no. 6, pp. 656–661, 1985.
- [12] K. Balasubramanian, "Nuclear-spin statistics of  $C_{60}$ ,  $C_{60}H_{60}$  and  $C_{60}D_{60}$ ," *Chem. Phys. Lett.*, vol. 183, nos. 3–4, pp. 292–296, Aug. 1991.
- [13] K. Balasubramanian, "Exhaustive generation and analytical expressions of matching polynomials of fullerenes  $C_{20}$ - $C_{50}$ ," *J. Chem. Inf. Comput. Sci.*, vol. 34, no. 2, pp. 421–427, 1994.
- [14] K. Balasubramanian, "Matching polynomials of fullerene clusters," *Chem. Phys. Lett.*, vol. 201, nos. 1–4, pp. 306–314, 1994.
- [15] A. T. Balaban, I. Motoc, D. Bonchev, and O. Mekenyan, "Topological indices for structure-activity correlations," *Steric Effects in Drug Design (Topics in Current Chemistry)*, vol. 114. New York, NY, USA: Academic, 1983, pp. 21–55.
- [16] K. Balasubramanian, K. Khokhani, and S. C. Basak, "Complex graph matrix representations and characterizations of proteomic maps and chemically induced changes to proteomes," *J. Proteome Res.*, vol. 5, no. 5, pp. 1133–1142, 2006.
- [17] K. Balasubramanian and M. Randić, "The characteristic polynomials of structures with pending bonds," *Theoretica Chimica Acta*, vol. 61, no. 4, pp. 307–323, 1982.
- [18] S. C. Basak, G. D. Grunwald, B. D. Gute, K. Balasubramanian, and D. Opitz, "Use of statistical and neural net approaches in predicting toxicity of chemicals," *J. Chem. Inf. Comput. Sci.*, vol. 40, no. 4, pp. 885–890, 2000.
- [19] S. C. Basak, D. Mills, M. M. Mumtaz, and K. Balasubramanian, "Use of topological indices in predicting aryl hydrocarbon receptor binding potency of dibenzofurans: A hierarchical QSAR approach," *Indian J. Chem.*, vol. 42A, no. 6, pp. 1385–1391, 2003.
- [20] C. Cao and Y. Hua, "Topological indices based on vertex, distance, and ring: On the boiling points of paraffins and cycloalkanes," *J. Chem. Inf. Comput. Sci.*, vol. 41, no. 4, pp. 867–877, 2001.
- [21] A. A. Dobrynin and A. A. Kochetova, "Degree distance of a graph: A degree analog of the wiener index," *J. Chem. Inf. Comput.*, vol. 34, pp. 1082–1086, Sep. 1994.
- [22] M. R. Farahani, "Eccentricity version of atom-bond connectivity index of benzenoid family  $ABC_5(H_k)$ ," *World Appl. Sci. J. Chem.*, vol. 21, no. 9, pp. 1260–1265, 2013.
- [23] W. Gao, M. Aamir, Z. Iqbal, M. Ishaq, and A. Aslam, "On irregularity measures of some dendrimers structures," *Mathematics*, vol. 7, no. 3, pp. 271–281, 2019.
- [24] W. Gao, Z. Iqbal, M. Ishaq, A. Aslam, M. Aamir, and M. A. Binyamin, "Bounds on topological descriptors of the corona product of  $F$ -sum of connected graphs," *IEEE Access*, vol. 7, pp. 26788–26796, 2019.
- [25] W. Gao, Z. Iqbal, M. Ishaq, A. Aslam, and R. Sarfraz, "Topological aspects of dendrimers via distance based descriptors," *IEEE Access*, vol. 7, pp. 35619–35630, 2019.
- [26] M. Ghorbani and M. A. Hosseinzadeh, "A new version of Zagreb indices," *Filomat*, vol. 26, pp. 93–100, Jan. 2012.
- [27] M. Ghorbani and A. Khaki, "A note on the fourth version of geometric-arithmetic index," *Optoelectron. Adv. Mat.*, vol. 4, no. 12, pp. 2212–2215, 2010.
- [28] S. Gupta, M. Singh, and A. K. Madan, "Connective eccentricity index: A novel topological descriptor for predicting biological activity," *J. Mol. Graph. Model.*, vol. 18, no. 1, pp. 18–25, 2000.
- [29] S. Gupta, M. Singh, and A. K. Madan, "Predicting anti-HIV activity: Computational approach using a novel topological descriptor," *J. Comput. Aided. Mol. Des.*, vol. 15, no. 7, pp. 671–678, 2001.
- [30] S. Gupta, M. Singh, and A. K. Madan, "Eccentric distance sum: A novel graph invariant for predicting biological and physical properties," *J. Math. Anal. Appl.*, vol. 275, no. 1, pp. 386–401, 2002.
- [31] I. Gutman and N. Trinajstić, "Graph theory and molecular orbitals. Total  $\varphi$ -electron energy of alternant hydrocarbons," *Chem. Phys. Lett.*, vol. 17, no. 4, pp. 535–538, Dec. 1972.
- [32] I. Gutman, "Degree-based topological indices," *Croatica Chem. Acta*, vol. 86, pp. 351–361, Dec. 2013.

- [33] S. Hayat and M. Imran, "Computation of topological indices of certain networks," *Appl. Math. Comput.*, vol. 240, pp. 213–228, Aug. 2014.
- [34] B. Hemmateenejad and A. Mohajeri, "Application of quantum topological molecular similarity descriptors in QSPR study of the O-methylation of substituted phenols," *J. Comput. Chem.*, vol. 29, no. 2, pp. 266–274, 2008.
- [35] M. Imran and S. Akhter, "Degree-based topological indices of double graphs and strong double graphs," *Discrete Math. Algorithms Appl.*, vol. 9, no. 5, 2017, Art. no. 1750066.
- [36] M. Imran, M. K. Siddiqui, A. A. E. Abunamous, D. Adi, S. H. Rafique, and A. Q. Baig, "Eccentricity based topological indices of an oxide network," *Mathematics*, vol. 6, no. 7, pp. 126–136, 2018.
- [37] S. Imran, M. K. Siddiqui, M. Imran, and M. F. Nadeem, "Computing topological indices and polynomials for line graphs," *Mathematics*, vol. 6, no. 8, pp. 137–148, 2018.
- [38] G. Indulal, "Spectrum of two new joins of graphs and infinite families of integral graphs," *Kragujevac J. Math.*, vol. 36, no. 38, pp. 133–139, 2012.
- [39] Z. Iqbal, M. Ishaq, and M. Aamir, "On eccentricity-based topological descriptors of dendrimers," *Iranian J. Sci. Technol., Trans. A, Sci.*, vol. 43, no. 4, pp. 1523–1533, 2019.
- [40] Z. Iqbal, A. Aslam, M. Ishaq, and M. Aamir, "Characteristic study of irregularity measures of some nanotubes," *Can. J. Phys.*, vol. 89, pp. 123–133, 2019.
- [41] Z. Iqbal, M. Ishaq, A. Aslam, and W. Gao, "On eccentricity-based topological descriptors of water-soluble dendrimers," *Zeitschrift Naturforschung*, vol. 74, nos. 1–2, pp. 25–33, 2018.
- [42] O. Ivanciuc, A. T. Balaban, and T. S. Balaban, "Design of topological indices. Part 4. Reciprocal distance matrix, related local vertex invariants and topological indices," *J. Math. Chem.*, vol. 12, no. 1, pp. 309–320, Dec. 1993.
- [43] M. Manoharan, M. M. Balakrishnarajan, P. Venuvanalingam, and K. Balasubramanian, "Topological resonance energy predictions of the stability of fullerene clusters," *Chem. Phys. Lett.*, vol. 222, nos. 1–2, pp. 95–100, May 1994.
- [44] A. Mohajeri and M. H. Dinpajooh, "Structure–toxicity relationship for aliphatic compounds using quantum topological descriptors," *J. Mol. Struct., THEOCHEM*, vol. 855, nos. 1–3, pp. 1–5, Apr. 2008.
- [45] A. Mohajeri, P. Manshour, and M. Mousaee, "A novel topological descriptor based on the expanded Wiener index: Applications to QSPR/QSAR studies," *Iran. J. Math. Chem.*, vol. 8, no. 2, pp. 107–135, 2017.
- [46] T. Parsons-Moss, L. K. Schwaiger, A. Hubaud, Y. J. Hu, H. Tuysuz, P. Yang, K. Balasubramanian, and H. Nitsche, "Plutonium complexation by phosphonate-functionalized mesoporous silica," in *Proc. ACS Nat. Meeting*, Anaheim, CA, USA, Mar. 2011, p. 241.
- [47] R. Ramaraj and K. Balasubramanian, "Computer generation of matching polynomials of chemical graphs and lattices," *J. Comput. Chem.*, vol. 6, no. 2, pp. 122–141, 1985.
- [48] M. Randić, "Novel molecular descriptor for structure—property studies," *Chem. Phys. Lett.*, vol. 211, pp. 478–483, Aug. 1993.
- [49] V. Sharma, R. Goswami, and A. K. Madan, "Eccentric-connectivity index: A novel highly discriminating topological descriptor for structure-property and structure-activity studies," *J. Chem. Inf. Comput. Sci.*, vol. 37, no. 2, pp. 273–282, 1997.
- [50] V. A. Skorobogatov and A. A. Dobrynin, "Metric analysis of graphs," *MATCH Commun. Math. Comput. Chem.*, vol. 23, no. 1, pp. 105–151, 1988.
- [51] F. Wen, Y. Zhang, and M. Li, "Spectra of subdivision vertex-edge join of three graphs," *Mathematics*, vol. 7, no. 2, pp. 171–181, 2019.
- [52] J. Zheng, Z. Iqbal, A. Fahad, A. Zafar, A. Aslam, R. Irfan, and M. I. Qureshi, "Some eccentricity-based topological indices and polynomials of poly(ETHyleneAmidoAmine) (PETAA) dendrimers," *Processes*, vol. 7, no. 7, pp. 433–443, 2019.



**MUHAMMAD IMRAN** received the Ph.D. degree in mathematics with a specialization in graph theory from the Abdus Salam School of Mathematical Sciences, Government College University, Lahore, Pakistan, in 2011, under the supervision of Prof. I. Tomescu of the Faculty of Mathematics and Informatics, University of Bucharest, Romania. He served as an Assistant Professor with the National University of Sciences and Technology (NUST), Islamabad, Pakistan, from May 2011 to July 2016. He has been an Assistant Professor with the Department of Mathematical Sciences, United Arab Emirates University, Al Ain, Abu Dhabi, United Arab Emirates, since August 2016. He successfully supervised one Ph.D. and six M.S. students of mathematics at NUST. He has published research articles in reputed international journals of mathematics and informatics. His research interests include metric graph theory, graph labeling, and spectral graph theory. He is a Referee for several international mathematical journals.



**SHEHNAZ AKHTER** received the M.Sc. degree in mathematics from the University of Sargodha, Pakistan, and the M.Phil. degree in mathematics from GC University Faisalabad, Pakistan. She is currently pursuing the Ph.D. degree in mathematics with the School of Natural Sciences, National University of Sciences and Technology, Islamabad, Pakistan. Her current research interests include chemical graph theory and extremal graph theory.



**ZAHID IQBAL** received the M.Sc. degree in mathematics from the University of Gujrat, Pakistan, and the M.Phil. degree in mathematics from Quaid-i-Azam University. He is currently pursuing the Ph.D. degree in mathematics with the School of Natural Sciences, National University of Sciences and Technology, Islamabad, Pakistan. His current research interests include algebra, combinatorics, and chemical graph theory.



**MUHAMMAD KAMRAN SIDDIQUI** received the M.Sc. degree in applied mathematics from the University of the Punjab, Pakistan, in 2005, the M.Phil. degree in applied mathematics from the Government College University, Lahore, Pakistan, in 2009, and the Ph.D. degree in discrete mathematics with a specialization in graph theory from the Abdus Salam School of Mathematical Sciences, Government College University, Lahore, in 2014. Since 2014, he has been an Assistant Professor with the Department of Mathematics, COMSATS University Islamabad, Lahore Campus, Pakistan. Since 2018, he has been a Postdoctoral Fellow with the Department of Mathematical Sciences, United Arab Emirates University, Al Ain, United Arab Emirates. He successfully supervised 12 M.Sc. students of mathematics at COMSATS University Islamabad, Sahiwal Campus, Pakistan. His current research interests include discrete mathematics, graph theory and its applications, chemical graph theory, combinatorics, neural networks, and complex dynamical networks. He is a Reviewer of *Ars Combinatoria*, *Utilitas Mathematica*, *Math Reports*, *Symmetry*, *IET Control Theory and Application*, *IEEE Access*, *Mathematics*, and *Discrete Applied Mathematics*.



**HONG YANG** received the M.S. degree from Sichuan University, in 1997, and the Ph.D. degree from the Southwestern Institute of Physics of Nuclear Industry, in 2003. He is currently an Associate Professor with the School of Information Science and Engineering, Chengdu University, Chengdu, China. His research interests include applied mathematics, graph theory, and algorithm design.