

On doubly transitive permutation groups

Cheryl E. Praeger

Suppose that G is a doubly transitive permutation group on a finite set Ω , and that for α in Ω the stabilizer G_α of α has a set $\Sigma = \{B_1, \dots, B_t\}$ of nontrivial blocks of imprimitivity in $\Omega - \{\alpha\}$. If G_α is 3-transitive on Σ it is shown that either G is a collineation group of a desarguesian projective or affine plane or no nonidentity element of G_α fixes B_1 pointwise.

Introduction

Suppose that G is a doubly transitive but not doubly primitive permutation group on a finite set Ω . Let $\Sigma = \{B_1, \dots, B_t\}$ be a set of nontrivial blocks of imprimitivity in $\Omega - \{\alpha\}$ for the stabilizer G_α of a point $\alpha \in \Omega$.

This paper completes an investigation which began with [11]. In that paper it was shown that if G_α^Σ is the alternating or symmetric group or one of the Mathieu groups in its usual representation then either G is a collineation group of a projective or affine plane or no nonidentity element of G_α fixes B_1 pointwise. In subsequent papers, [10, 12], it was shown that the same conclusions are valid if we assume only that G_α is 3-transitive and not faithful on Σ . The assumption that G_α is not

Received 1 May 1978.

faithful on Σ is unattractive but unfortunately it was crucial in the proofs given in those papers. Our aim in this paper is to show that the assumption that G_α is not faithful on Σ is unnecessary. We prove

THEOREM. *Suppose that G is 2-transitive on Ω of degree n , and that for $\alpha \in \Omega$, G_α has a set $\Sigma = \{B_1, \dots, B_t\}$ of blocks of imprimitivity in $\Omega - \{\alpha\}$, where $t = |\Sigma| \geq 3$, $|B_i| = b > 1$, $bt = n - 1$. Suppose that G_α is 3-transitive on Σ and that G_α contains a nonidentity element fixing B_1 pointwise. Then G is a collineation group of a desarguesian projective or affine plane of order $t - 1$ such that the lines containing α are precisely the sets $B_i \cup \{\alpha\}$, for $i = 1, \dots, t$.*

We remark that the affine planes arising in the conclusions of [11] and [12] are desarguesian. This was shown in [13]. Most of the notation used here follows the conventions of Wielandt's book [14]. If a group G has a permutation representation on a set Ω then $\text{fix}_\Omega G$ and $\text{supp}_\Omega G$ will denote the subset of Ω fixed by G and the subset of Ω permuted nontrivially by G respectively. By a block design we shall mean a set of v points and a set of b blocks with a relation called incidence between points and blocks, such that any block is incident with k points, where $2 \leq k < v-1$, and any pair of points is incident with λ blocks, where $\lambda > 0$. The number r of blocks incident with a given point is also constant and a counting argument shows $vr = bk$. It is well known that $b \geq v$ and hence that $r \geq k$.

Proof of the theorem

Let G satisfy the hypothesis of the theorem and let K_i, \bar{K}_i denote the setwise and pointwise stabilizers respectively of $B_i \in \Sigma$, for $i = 1, \dots, t$. By [12] and [13], Proposition D, the theorem is true if either G_α is not faithful on Σ or if \bar{K}_1 is 2-transitive on $\Sigma - \{B_1\}$. Thus we may assume that G_α is faithful on Σ and \bar{K}_1 is not 2-transitive on $\Sigma - \{B_1\}$. By [13], Theorem A, and since G_α is

3-transitive on Σ , the translates under G of $B_1 \cup \{\alpha\}$ are the blocks of a block design with $\lambda = 1$ preserved by G .

LEMMA 1. *The theorem is true if K_1 has a normal subgroup which acts regularly on $\Sigma - \{B_1\}$.*

Proof. Suppose that K_1 has a normal subgroup N which is regular on $\Sigma - \{B_1\}$. By [5] and since G_α is 3-transitive on Σ , it follows that either G_α^Σ is a normal extension of $\text{PSL}(2, q)$ of degree $q + 1$ or G_α has a normal subgroup M which is regular on Σ . In the latter case, by [14], 11.3, either $t = |M| = 3$ or M is elementary abelian of order 2^a for some $a \geq 2$. If $t = 3$, then $G_\alpha \simeq S_3$ and hence $K_1 \simeq Z_2$. Since K_1 is transitive on B_1 , $b = 2$ and $n = 7$. However there is no 2-transitive group G of degree 7 with $G_\alpha \simeq S_3$. If $|M| = 2^a$ then by [4] and since G_α is 3-transitive on Σ , it follows that G has a regular normal subgroup L , say. Then ML is a Frobenius group with Frobenius complement M . However a Frobenius complement contains at most one involution and so we have a contradiction (see [3], 10.3.1). Thus G_α^Σ is a normal extension of $\text{PSL}(2, q)$ with $t = q + 1$.

Now K_1 has a normal subgroup V of order q which is regular on $\Sigma - \{B_1\}$, and since \bar{K}_1 is nontrivial, $V \subseteq \bar{K}_1$. Then $K_1 = (K_1 \cap K_2)\bar{K}_1$, so that $K_1^{B_1} = (K_1 \cap K_2)^{B_1}$, which is metacyclic. If Z is the largest cyclic normal subgroup of $K_1 \cap K_2$, then Z has order $q - 1$ or $(q-1)/2$. Let $Y = Z \cap \bar{K}_1$ and suppose that Y is nontrivial. Then Y is semi-regular on $\Sigma - \{B_1, B_2\}$, so that $\text{fix}_\Omega Y \subseteq \{\alpha\} \cup B_1 \cup B_2$. Moreover Y, Z are characteristic subgroups of $Z, K_1 \cap K_2$ respectively.

Now there is an element g in G_α which interchanges B_1 and B_2 and hence normalises $K_1 \cap K_2$. It follows that g normalises Y and Z ,

and hence that $Y = Z \cap \bar{K}_2$. Thus $Y \leq \bar{K}_1 \cap \bar{K}_2$ and so $\text{fix}_\Omega Y = \{\alpha\} \cup B_1 \cup B_2$.

If $\Sigma(\beta)$ is the set of blocks of imprimitivity of G_β corresponding to Σ , then $C = (B_1 - \{\beta\}) \cup \{\alpha\}$ is the block in $\Sigma(\beta)$ containing α and \bar{K}_1 is the pointwise stabilizer of $\{\beta\} \cup C$. Now V must be regular on both $\Sigma(\beta) - \{C\}$ and $\Sigma - \{B_1\}$, so that the representations of \bar{K}_1 on $\Sigma(\beta) - \{C\}$ and $\Sigma - \{B_1\}$ are equivalent. Thus $\bar{K}_1 \cap K_2$ is the stabilizer in \bar{K}_1 of an element C' , say, of $\Sigma(\beta) - \{C\}$. By a similar argument Y fixes C' pointwise, so that $C' = B_2$, which is a contradiction, since $\{\alpha\} \cup B_2$ and $\{\beta\} \cup C'$ are distinct blocks of the block design. Thus $Y = 1$, and Z is faithful and hence semi-regular on B_1 . Therefore b is divisible by $|Z|$.

Now the size $1 + b$ of a block of the block design is at most the number $t = q + 1$ of blocks containing α , and hence b is $q - 1$ or $(q-1)/2$. If b is $q - 1$, the design is an affine plane, which can be shown to be desarguesian as in [13]. So assume that $b = (q-1)/2 = |Z|$. Then q is odd and $G_\alpha \not\leq \text{PGL}(2, q)$. In this case V has odd order q , and if $\beta \in B_1$, $G_{\alpha\beta}/V$ is cyclic. Thus a Sylow 2-subgroup of $G_{\alpha\beta}$ is cyclic or trivial. By [1] and [2], respectively, it follows that G has a regular normal subgroup L , say. Now $\text{PSL}(2, q) \leq G_\alpha \leq \text{P}\Gamma\text{L}(2, q)$. Let M be a subgroup of $\text{PSL}(2, q)$, which is elementary abelian of order 4, $M \leq G_\alpha$. An involution g in M fixes either 0 or 2 elements of Σ . In the former case g fixes only the point α of Ω . In the latter case we may assume that $g \in K_1 \cap K_2$ and hence that $g \in Z$. We showed above that Z is semi-regular on B_1 and similarly Z is semi-regular on B_2 . Thus in this case, also, $\text{fix}_\Omega g = \{\alpha\}$. Thus M acts fixed point freely on L and so is a Frobenius complement, whereas a Frobenius complement of even order contains only one involution (see [3], 10.3.1). Thus the lemma is proved.

We may assume therefore that K_1 has no normal subgroup acting

regularly on $\Sigma - \{B_1\}$; in particular \bar{K}_1 is not regular on $\Sigma - \{B_1\}$.

LEMMA 2. *The theorem is true if $K_1 \cap K_2$ has a nontrivial abelian normal subgroup.*

Proof. Suppose that $K_1 \cap K_2$ has a nontrivial abelian normal subgroup. By [9], Lemma 1, and [10], the theorem is true unless G_α^Σ has a regular normal elementary abelian 2-subgroup. Thus we may assume that G_α has an elementary abelian normal 2-subgroup M of order $t \geq 4$ which is semiregular on $\Omega - \{\alpha\}$. By [4], G has a regular normal subgroup L , and so M is a Frobenius complement. However, as Frobenius complements contain at most one involution, this is a contradiction (see [3], 10.3.1), and Lemma 2 is proved.

Thus we may assume that $K_1 \cap K_2$ has no nontrivial abelian normal subgroups. By [8], Theorem A, $K_1 \cap K_2$ has a unique minimal normal subgroup S which is a nonabelian simple group and hence has even order. Since \bar{K}_1 is not regular on $\Sigma - \{B_1\}$, $\bar{K}_1 \cap K_2$ and similarly $K_1 \cap \bar{K}_2$ are nontrivial normal subgroups of $K_1 \cap K_2$. Thus

$$S \subseteq (\bar{K}_1 \cap K_2) \cap (K_1 \cap \bar{K}_2) = \bar{K}_1 \cap \bar{K}_2 = X,$$

say. Since $K_1 \cap K_2$ is transitive on $\Sigma - \{B_1, B_2\}$, X is $\frac{1}{2}$ -transitive on $\Sigma - \{B_1, B_2\}$ with orbits of length, say, x , where $x > 1$, and x divides $t - 2$.

LEMMA 3. *$X = \bar{K}_1 \cap \bar{K}_2$ is not semi-regular on $\Sigma - \{B_1, B_2\}$.*

Proof. If X is semi-regular on $\Sigma - \{B_1, B_2\}$, then, since $|X|$ is even, it follows from [3] and Lemma 1 that $K_1^{\Sigma - \{B_1\}}$ has a unique normal subgroup, say X^1 , which is 2-transitive and simple. Then $X^1 \leq \bar{K}_1^{\Sigma - \{B_1\}}$ which is a contradiction, since \bar{K}_1 is not 2-transitive on $\Sigma - \{B_1\}$.

Let $\Sigma(\beta)$ be the set of blocks of imprimitivity for G_β

corresponding to Σ , where $\beta \in B_1$. Then the element of $\Sigma(\beta)$ containing α is $C = (B_1 - \{\beta\}) \cup \{\alpha\}$. Let $C^1 \in \Sigma(\beta) - \{C\}$ be chosen so that $C^1 \cap B_2$ is non-empty. Then as $\{\beta\} \cup C^1$ and $\{\alpha\} \cup B_2$ are distinct blocks of the block design, $C^1 \cap B_2 = \{\gamma\}$, say. Let Y be the pointwise stabilizer of C^1 in \bar{K}_1 . Then Y is conjugate to X in G (for \bar{K}_1 is normal in $G_{\{\alpha, \beta\}}$, so if $g \in G_{\{\alpha, \beta\}} - G_{\alpha\beta}$, then $Y^g = \bar{K}_1 \cap \bar{K}_i$, where $(C^1)^g = B_i$, and there is an h in \bar{K}_1 such that $B_i^h = B_2$ and hence $Y^{gh} = \bar{K}_1 \cap \bar{K}_2 = X$). It follows that $\text{fix}_\Omega Y = \{\alpha\} \cup B_1 \cup C^1$, and all Y -orbits in $\text{supp}_\Omega Y$ have length a multiple of x . Since Y fixes γ , Y fixes B_2 setwise, and clearly $B_2 - \{\gamma\} \subseteq \text{supp}_\Omega Y$. Therefore $b - 1 = |B_2 - \{\gamma\}|$ is a multiple of x .

LEMMA 4. *The subset of Σ fixed by Y is precisely*

$$\text{fix}_\Sigma Y = \{B_1\} \cup \{B \in \Sigma; B \cap C^1 \neq \emptyset\},$$

and $\text{fix}_\Sigma Y$ is a union of X -orbits in Σ .

Proof. Clearly $\{B_1\} \cup \{B \in \Sigma; B \cap C^1 \neq \emptyset\} \subseteq \text{fix}_\Sigma Y$, and is the subset of $\text{fix}_\Sigma Y$ of those elements of Σ which contain a point of $\text{fix}_\Omega Y$. If $\text{fix}_\Sigma Y$ contains an additional element B , then $B \subseteq \text{supp}_\Omega Y$, and so B is a union of nontrivial Y -orbits; that is, b is divisible by x . However $b - 1$ is divisible by x and $x > 1$, so we conclude that $\text{fix}_\Sigma Y = \{B_1\} \cup \{B \in \Sigma; B \cap C^1 \neq \emptyset\}$.

Now let $B \in \text{fix}_\Sigma Y - \{B_1, B_2\}$ and let Δ be the X -orbit in Σ containing B . It is sufficient to show that $\Delta \subseteq \text{fix}_\Sigma Y$. Let $B \cap C^1 = \{\delta\}$ and let Δ' be the X -orbit in Ω containing δ . Then $|\Delta| = x$ and $|\Delta'| = xy$, where $y = |\Delta' \cap B| \geq 1$. Now $Y \subseteq K_1 \cap K_2$, and so Y normalises X . Therefore Y fixes $\Delta' = \{\delta^x; x \in X\}$ setwise (for

if $y \in Y$, then $\delta^{xy} = \delta^{y^{-1}xy} \in \Delta'$. Thus Y fixes $B \cap \Delta'$ setwise, and so $(B \cap \Delta') - \{\delta\}$ is a union of nontrivial Y -orbits, that is $y - 1 = |(B \cap \Delta') - \{\delta\}| \equiv 0 \pmod{x}$.

Let $a = |\text{supp}_\Sigma Y \cap \Delta|$. Then $0 \leq a < x$. If $a > 0$, then $U\{B' \cap \Delta'; B' \in \text{supp}_\Sigma Y \cap \Delta\}$ is a union of nontrivial Y -orbits and is a set of ay points. Thus $ay \equiv 0 \pmod{x}$, and so $a \equiv 0 \pmod{x}$, which is a contradiction, since $0 < a < x$. Thus $a = 0$ and $\Delta \subseteq \text{fix}_\Sigma Y$.

LEMMA 5. $X \cap Y = 1$.

Proof. If $X \cap Y$ is nontrivial, then it follows from Lemma 4 that X does not act faithfully on its orbits in $\Sigma - \{B_1, B_2\}$, and so by [8],

Proposition 4, $K_1^{\Sigma - \{B_1\}}$ is a normal extension of $L_r(q)$, for some $r \geq 3$ and prime power q , in its natural representation. Then $\bar{K}_1 \supseteq L_r(q)$ and hence is 2-transitive on $\Sigma - \{B_1\}$, which is a contradiction.

Thus $X \cap Y = 1$ and Y normalises X . Similarly since X fixes C and C^1 setwise, X normalises Y . It follows that X and Y centralise each other. Thus, by Lemma 3, the centraliser of X in K_1 is not semiregular on $\Sigma - \{B_1, B_2\}$, and it follows from [6], Corollary B3, and Lemma 2.8 that X is a T.I. set in K_1 (that is distinct conjugates of X by elements of K_1 intersect only in the identity). Thus by [7], Theorem A, and since X is not semiregular on $\Sigma - \{B_1, B_2\}$, $K_1^{\Sigma - \{B_1\}}$ is a normal extension of $L_r(q)$ in its natural representation for some $r \geq 3$ and prime power q . This is impossible, as \bar{K}_1 is not 2-transitive on $\Sigma - \{B_1\}$.

Thus the theorem is proved.

References

- [1] Michael Aschbacher, "2-transitive groups whose 2-point stabilizer has 2-rank 1", *J. Algebra* 36 (1975), 98-127.
- [2] Helmut Bender, "Transitive Gruppen gerader Ordnung, in denen jede Involution genau einen Punkt festläßt", *J. Algebra* 17 (1971), 527-554.
- [3] Daniel Gorenstein, *Finite groups* (Harper and Row, New York, Evanston, London, 1968).
- [4] Christoph Hering, "On subgroups with trivial normalizer intersection", *J. Algebra* 20 (1972), 622-629.
- [5] Christoph Hering and William M. Kantor and Gary M. Seitz, "Finite groups with a split BN -pair of rank 1. I", *J. Algebra* 20 (1972), 435-475.
- [6] Michael O'Nan, "A characterization of $L_n(q)$ as a permutation groups", *Math. Z.* 127 (1972), 301-314.
- [7] Michael E. O'Nan, "Normal structure of the one-point stabilizer of a doubly-transitive permutation group. I", *Trans. Amer. Math. Soc.* 214 (1975), 1-42.
- [8] Michael E. O'Nan, "Normal structure of the one-point stabilizer of a doubly-transitive permutation group. II", *Trans. Amer. Math. Soc.* 214 (1975), 43-74.
- [9] Michael E. O'Nan, "Triply-transitive permutation groups whose two-point stabilizer is local", submitted.
- [10] Cheryl E. Praeger, "Doubly transitive permutation groups involving the one-dimensional projective special linear group", *Bull. Austral. Math. Soc.* 14 (1976), 349-358.
- [11] Cheryl E. Praeger, "Doubly transitive permutation groups which are not doubly primitive", *J. Algebra* 44 (1977), 389-395.
- [12] Cheryl E. Praeger, "Doubly transitive permutation groups in which the one-point stabilizer is triply transitive on a set of blocks", *J. Algebra* 47 (1977), 433-440.

- [13] Cheryl E. Praeger, "Doubly transitive automorphism groups of block designs", *J. Combinatorial Theory Ser. A* (to appear).
- [14] Helmut Wielandt, *Finite permutation groups* (translated by R. Bercov. Academic Press, New York, London, 1964).

Department of Mathematics,
University of Western Australia,
Nedlands,
Western Australia.