

 Open access • Journal Article • DOI:10.2298/FIL0802065E

## On $ds$ / -sets and decompositions of continuous functions — [Source link](#)

Erdal Ekici, Saeid Jafari

**Institutions:** Çanakkale Onsekiz Mart University

**Published on:** 01 Jan 2008 - Filomat (National Library of Serbia)

Related papers:

- [Algorithmic characterization results for the Kerr-NUT-\(A\)dS space-time. I. A space-time approach](#)
- [Three variable exponential functions of the alternating group](#)
- [Parter, Periodic and Coperiodic Functions on Groups and their Characterization](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/on-ds-sets-and-decompositions-of-continuous-functions-3h00622455>

## ON $\mathcal{DS}^*$ -SETS AND DECOMPOSITIONS OF CONTINUOUS FUNCTIONS

Erdal Ekici and Saeid Jafari

### Abstract

In this paper, the notions of  $\mathcal{DS}^*$ -sets and  $\mathcal{DS}^*$ -continuous functions are introduced and their properties and their relationships with some other types of sets are investigated. Moreover, some new decompositions of continuous functions are obtained by using  $\mathcal{DS}^*$ -continuous functions,  $\mathcal{DS}$ -continuous functions and  $\mathcal{D}$ -continuous functions.

## 1 Introduction

In a recent paper, Ekici and Jafari [12] have studied  $\mathcal{DS}$ -sets and  $\mathcal{D}$ -sets and obtained some decompositions of continuous functions via  $\mathcal{DS}$ -continuous functions and  $\mathcal{D}$ -continuous functions. In this paper, we introduce a new class of sets called  $\mathcal{DS}^*$ -sets. Properties of this class are investigated. Furthermore, the notion of  $\mathcal{DS}^*$ -continuous functions is introduced via  $\mathcal{DS}^*$ -sets to establish some new decompositions of continuous functions. On the other hand, by using  $\mathcal{DS}$ -sets and  $\mathcal{D}$ -sets, other new decompositions of continuous functions are obtained.

In this paper  $(X, \tau)$  and  $(Y, \sigma)$  represent topological spaces. For a subset  $A$  of a space  $X$ ,  $cl(A)$  and  $int(A)$  denote the closure of  $A$  and the interior of  $A$ , respectively. A subset  $A$  of a space  $X$  is called regular open (resp regular closed) [22] if  $A = int(cl(A))$  (resp.  $A = cl(int(A))$ ).  $A$  is called  $\delta$ -open [24] if for each  $x \in A$ , there exists a regular open set  $U$  such that  $x \in U \subset A$ .  $A$  is called  $\delta$ -closed if its complement is  $\delta$ -open. A point  $x \in X$  is called a  $\delta$ -cluster point of  $A$  if  $A \cap int(cl(U)) \neq \emptyset$  for each open set  $U$  containing  $x$ . The set of all  $\delta$ -cluster points of  $A$  is called the  $\delta$ -closure of  $A$  and is denoted by  $\delta-cl(A)$ . The union of all regular open sets, each contained in  $A$  called the  $\delta$ -interior of  $A$  and is denoted by  $\delta-int(A)$ . A subset  $A$  of a space  $(X, \tau)$  is called semiopen [15] (resp. semi-regular [7],  $\alpha$ -open [19], preopen [16] or locally dense [6],  $b$ -open [4] or  $\gamma$ -open [13] or sp-open [8],  $\beta$ -open [1] or

---

2000 *Mathematics Subject Classifications.* 54C08.

*Key words and Phrases.*  $\mathcal{DS}^*$ -sets,  $\mathcal{DS}^*$ -continuous function, decomposition.

Received: November 5, 2007

Communicated by Dragan Djordjević

semi-preopen [3],  $\delta$ -semiopen [20],  $\delta$ -preopen [21]) if  $A \subset cl(int(A))$  (resp. semiopen and semiclosed,  $A \subset int(cl(int(A)))$ ,  $A \subset int(cl(A))$ ,  $A \subset int(cl(A)) \cup cl(int(A))$ ,  $A \subset cl(int(cl(A)))$ ,  $A \subset cl(\delta-int(A))$ ,  $A \subset int(\delta-cl(A))$ ). The complement of a  $\delta$ -semiopen (resp. semiopen) set is called a  $\delta$ -semiclosed (resp. semiclosed) set. The union (resp. intersection) of all  $\delta$ -preopen (resp.  $\delta$ -semiclosed) sets, each contained in (resp. containing) a set  $A$  in a topological space  $X$  is called the  $\delta$ -preinterior (resp.  $\delta$ -semiclosure) of  $A$  and it is denoted by  $\delta-pint(A)$  (resp.  $\delta-scl(A)$ ).

**Definition 1** A subset  $A$  of a space  $(X, \tau)$  is called

- (1) a  $\mathcal{D}$ -set [12] if  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -closed,
- (2) a  $\mathcal{DS}$ -set [12] if  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -semiclosed,
- (3) a  $\mathcal{B}$ -set [23] if  $A \in \mathcal{B}(X) = \{U \cap V : U \in \tau, int(cl(V)) \subset V\}$ ,
- (4) an  $\mathcal{AB}$ -set [9] if  $A \in \mathcal{AB}(X) = \{U \cap V : U \in \tau \text{ and } V \text{ is semi-regular}\}$ .

The family of all  $\mathcal{DS}$ -sets (resp.  $\mathcal{D}$ -sets) of a topological space  $X$  will be denoted by  $\mathcal{DS}(X)$  (resp.  $\mathcal{D}(X)$ ). A topological space  $X$  is called a locally indiscrete [10] if every open subset of  $X$  is closed and called submaximal [5] if every dense subset of  $X$  is open.

**Definition 2** A function  $f : X \rightarrow Y$  is called


- (1)  $\beta$ -continuous [1] if  $f^{-1}(A)$  is  $\beta$ -open for each  $A \in \sigma$ .
- (2)  $\alpha$ -continuous [17] if  $f^{-1}(A)$  is  $\alpha$ -open for each  $A \in \sigma$ .
- (3)  $\gamma$ -continuous [13] if  $f^{-1}(A)$  is  $\gamma$ -open for each  $A \in \sigma$ .
- (4) quasi-continuous [14] if  $f^{-1}(A)$  is semiopen for each  $A \in \sigma$ .
- (5) precontinuous [16] if  $f^{-1}(A)$  is preopen for each  $A \in \sigma$ .
- (6)  $\delta$ -almost continuous [21] if  $f^{-1}(A)$  is  $\delta$ -preopen for each  $A \in \sigma$ .
- (7)  $\delta$ -semicontinuous [11] if  $f^{-1}(A)$  is  $\delta$ -semiopen for each  $A \in \sigma$ .
- (8) super-continuous [18] if  $f^{-1}(A)$  is  $\delta$ -open for each  $A \in \sigma$ .

## 2 $\mathcal{DS}^*$ -sets in topological spaces

**Definition 3** A subset  $A$  of a topological space  $X$  is called a  $\mathcal{DS}^*$ -set if  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -semiclosed and  $int(\delta-cl(V)) = cl(\delta-int(V))$ .

The family of all  $\mathcal{DS}^*$ -sets of a topological space  $X$  will be denoted by  $\mathcal{DS}^*(X)$ .

**Remark 4** The following diagram holds for a subset of a space  $X$ :



The following examples show that these implications are not reversible.

**Example 5** Let  $X = \{a, b, c, d\}$  and let  $\tau = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ . The set  $\{a, c, d\}$  is a  $\mathcal{DS}$ -set but it is not a  $\mathcal{DS}^*$ -set.

**Example 6** ([12]) Let  $X = \{a, b, c, d\}$  and let  $\tau = \{\emptyset, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}, X\}$ . The set  $\{b, c, d\}$  is a  $\mathcal{B}$ -set but it is not a  $\mathcal{DS}$ -set.

**Remark 7** Every open set is a  $\mathcal{DS}^*$ -set. The converse is not true.

**Example 8** Let  $X = \{a, b, c, d\}$  and let  $\tau = \{\emptyset, \{a\}, \{b\}, \{a, b\}, X\}$ . The set  $\{a, c\}$  is a  $\mathcal{DS}^*$ -set but it is not open.

**Theorem 9** The following are equivalent for a subset  $A$  of a space  $X$ :

- (1)  $A$  is open,
- (2)  $A$  is  $\alpha$ -open and a  $\mathcal{DS}^*$ -set,
- (3)  $A$  is semiopen and a  $\mathcal{DS}^*$ -set,
- (4)  $A$  is preopen and a  $\mathcal{DS}^*$ -set,
- (5)  $A$  is  $\gamma$ -open and a  $\mathcal{DS}^*$ -set.
- (6)  $A$  is  $\beta$ -open and a  $\mathcal{DS}^*$ -set.

**Proof.** (1)  $\Rightarrow$  (2) : It follows from the fact that every open set is  $\alpha$ -open and a  $\mathcal{DS}^*$ -set.

(2)  $\Rightarrow$  (3)  $\Rightarrow$  (5) : Obvious.

(2)  $\Rightarrow$  (4)  $\Rightarrow$  (5) : Obvious.

(5)  $\Rightarrow$  (6) : Obvious.

(6)  $\Rightarrow$  (1) : Let  $A$  be  $\beta$ -open and a  $\mathcal{DS}^*$ -set. Since  $A$  is  $\beta$ -open,  $A \subset cl(int(cl(A)))$ . Since  $A$  is a  $\mathcal{DS}^*$ -set, then  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -semiclosed and  $int(\delta-cl(V)) = cl(\delta-int(V))$ . Also, by  $\delta$ -semiclosedness of  $V$ , we have  $\delta-int(V) = \delta-int(\delta-cl(V))$ . Furthermore, we obtain

$$\begin{aligned}
 A = A \cap U \subset cl(int(cl(A))) \cap U &= cl(int(cl(U \cap V))) \cap U \\
 &\subset cl(int(cl(U))) \cap cl(int(cl(V))) \cap U \\
 &= cl(int(cl(V))) \cap U \\
 &\subset cl(int(\delta-cl(V))) \cap U \\
 &= cl(\delta-int(V)) \cap U \\
 &= int(\delta-cl(V)) \cap U \\
 &= \delta-int(V) \cap U.
 \end{aligned}$$

Thus,  $A = \delta-int(V) \cap U$  and hence  $A$  is open. ■

**Theorem 10** ([12]) For a space  $X$ , the following are equivalent:

- (1)  $X$  is indiscrete,
- (2) the  $\mathcal{DS}$ -sets in  $X$  are only the trivial ones,
- (3) the  $\mathcal{D}$ -sets in  $X$  are only the trivial ones.

**Theorem 11** *The following are equivalent for a space  $X$ :*

- (1)  $X$  is indiscrete,
- (2) the  $\mathcal{DS}^*$ -sets in  $X$  are the trivial ones.

**Proof.** (1)  $\Rightarrow$  (2) : Let  $A$  be a  $\mathcal{DS}^*$ -set in  $X$ . Then there exist an open set  $U$  and a  $\delta$ -semiclosed set  $V$  such that  $A = U \cap V$  and  $\delta\text{-int}(\delta\text{-cl}(V)) = \delta\text{-cl}(\delta\text{-int}(V))$ . Let  $A \neq \emptyset$ . We have  $U \neq \emptyset$ . By (1),  $U = X$  and  $A = V$ . Hence,  $X = \delta\text{-scl}(A) \subset A$  and so  $A = X$ .

(2)  $\Rightarrow$  (1) : Since every  $\mathcal{DS}^*$ -set is  $\mathcal{DS}$ -set, it follows from Theorem 10. ■

**Theorem 12** *Let  $X$  be a topological space and  $A \subset X$ . If  $A \in \mathcal{DS}(X)$ , then  $\delta\text{-pint}(A) = \text{int}(A)$ .*

**Proof.** Let  $A \in \mathcal{DS}(X)$ . Then,  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -semiclosed. Since  $V$  is  $\delta$ -semiclosed, then we have  $\delta\text{-int}(V) = \delta\text{-int}(\delta\text{-cl}(V))$ . Moreover, we obtain

$$\begin{aligned} \delta\text{-pint}(A) = A \cap \delta\text{-int}(\delta\text{-cl}(A)) &\subset U \cap \delta\text{-int}(\delta\text{-cl}(V)) \\ &= U \cap \delta\text{-int}(V) \\ &\subset U \cap \text{int}(V) \\ &= \text{int}(A). \end{aligned}$$

Thus,  $\delta\text{-pint}(A) = \text{int}(A)$ . ■

**Theorem 13** *The following are equivalent for a subset  $A$  of a space  $X$ :*

- (1)  $A$  is open,
- (2)  $A$  is  $\delta$ -preopen and a  $\mathcal{D}$ -set,
- (3)  $A$  is  $\delta$ -preopen and a  $\mathcal{DS}$ -set.

**Proof.** (1)  $\Rightarrow$  (2) : Since every open set is  $\delta$ -preopen and a  $\mathcal{D}$ -set, it is completed.

(2)  $\Rightarrow$  (3) : Obvious.

(3)  $\Rightarrow$  (1) : Let  $A$  be  $\delta$ -preopen and a  $\mathcal{DS}$ -set. By Theorem 12,  $\delta\text{-pint}(A) = \text{int}(A)$ . Also, since  $A$  is  $\delta$ -preopen,  $A = \delta\text{-pint}(A) = \text{int}(A)$ . Thus,  $A$  is open. ■

**Theorem 14** *Let  $X$  be a topological space and  $A \subset X$ . If  $A \in \mathcal{DS}^*(X)$ , then  $A = U \cap \delta\text{-scl}(A)$  for some open set  $U$ .*

**Proof.** Let  $A \in \mathcal{DS}^*(X)$ . This implies that  $A = U \cap V$ , where  $U$  is open and  $V$  is  $\delta$ -semiclosed and  $\text{int}(\delta\text{-cl}(V)) = \text{cl}(\delta\text{-int}(V))$ . Since  $A \subset V$ ,  $\delta\text{-scl}(A) \subset \delta\text{-scl}(V) = V$ . Moreover,  $U \cap \delta\text{-scl}(A) \subset U \cap V = A \subset U \cap \delta\text{-scl}(A)$  and hence  $A = U \cap \delta\text{-scl}(A)$ . ■

**Theorem 15** ([12]) *The following are equivalent for a subset  $P$  of a space  $X$ :*

- (1)  $P$  is an  $\mathcal{AB}$ -set,
- (2)  $P$  is semiopen and a  $\mathcal{DS}$ -set,
- (3)  $P$  is  $b$ -open and a  $\mathcal{DS}$ -set,
- (4)  $P$  is  $\beta$ -open and a  $\mathcal{DS}$ -set.

**Theorem 16** *Let  $X$  be a topological space and  $A \subset X$ . If  $A$  is  $\beta$ -open and a  $\mathcal{DS}^*$ -set, then it is an  $\mathcal{AB}$ -set.*

**Proof.** Let  $A$  be  $\beta$ -open and a  $\mathcal{DS}^*$ -set. Since  $A$  is a  $\mathcal{DS}$ -set, by Theorem 15,  $A$  is an  $\mathcal{AB}$ -set. ■

**Definition 17** *Let  $X$  be a topological space and  $A \subset X$ . Then  $A$  is called a  $\delta^*$ -set if  $\delta\text{-int}(A)$  is  $\delta$ -closed.*

**Theorem 18** *Let  $X$  be a topological space and  $A \subset X$ . If  $A$  is a  $\delta^*$ -set and  $\delta$ -semiopen, then it is  $\delta$ -open.*

**Proof.** Let  $A$  be a  $\delta^*$ -set and  $\delta$ -semiopen. Then  $A \subset cl(\delta\text{-int}(A)) = \delta\text{-int}(A)$ . Thus,  $A$  is  $\delta$ -open. ■

**Theorem 19** *Let  $X$  be a topological space and  $A \subset X$ . Then  $A$  is open if  $A$  is a  $\delta$ -semiopen  $\mathcal{DS}^*$ -set and  $A$  is preopen or a  $\delta^*$ -set.*

**Proof.** Let  $A$  be a  $\delta$ -semiopen  $\mathcal{DS}^*$ -set. Suppose that  $A$  is preopen or a  $\delta^*$ -set. If  $A$  is a preopen  $\mathcal{DS}^*$ -set, then it is a preopen  $\mathcal{B}$ -set. So, by Proposition 9 [23],  $A$  is open. Also, if  $A$  is a  $\delta^*$ -set and  $\delta$ -semiopen, by Theorem 18,  $A$  is open. Thus, the proof is completed. ■

**Theorem 20** *The following are equivalent for a space  $X$  :*

- (1)  $X$  is a locally indiscrete space,
- (2) every  $\mathcal{DS}^*$ -set is clopen,
- (3) every  $\mathcal{DS}^*$ -set is closed.

**Proof.** (1)  $\Rightarrow$  (2) : Let  $A$  be a  $\mathcal{DS}^*$ -set. Then there exist an open set  $U$  and a  $\delta$ -semiclosed set  $V$  such that  $A = U \cap V$  and  $int(\delta\text{-cl}(V)) = cl(\delta\text{-int}(V))$ . Since  $U$  is clopen, then  $A$  is semiclosed. By [2], since  $X$  is a locally indiscrete space, then  $A$  is clopen.

(2)  $\Rightarrow$  (3) : Obvious.

(3)  $\Rightarrow$  (1) : Let  $A \subset X$  be an open set. Since  $A$  is a  $\mathcal{DS}^*$ -set, then  $A$  is closed. Hence,  $X$  is a locally indiscrete space. ■

**Theorem 21** ([12]) *For a space  $X$  the following are equivalent:*

- (1)  $X$  is submaximal,
- (2) every dense subset of  $X$  is a  $\mathcal{D}$ -set,
- (3) every dense subset of  $X$  is a  $\mathcal{DS}$ -set.

**Theorem 22** *Let  $X$  be a topological space. Then  $X$  is submaximal if and only if every dense subset of  $X$  is a  $\mathcal{DS}^*$ -set.*

**Proof.** ( $\Rightarrow$ ) : Let  $A$  be a dense subset of  $X$ . Since  $X$  submaximal, then  $A$  is open and so  $A$  is a  $\mathcal{DS}^*$ -set.

( $\Leftarrow$ ) : Since every dense subset is a  $\mathcal{DS}^*$ -set and every  $\mathcal{DS}^*$ -set is a  $\mathcal{DS}$ -set, then by Theorem 21,  $X$  is submaximal. ■

### 3 Some new decompositions of continuity

**Definition 23** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called

- (1)  $\mathcal{DS}^*$ -continuous if  $f^{-1}(V) \in \mathcal{DS}^*(X)$  for each  $V \in \sigma$ .
- (2)  $\delta^*$ -continuous if  $f^{-1}(V)$  is a  $\delta^*$ -set for each  $V \in \sigma$ .

**Definition 24** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called

- (1)  $\mathcal{D}$ -continuous [12] if  $f^{-1}(V) \in \mathcal{D}(X)$  for each  $V \in \sigma$ .
- (2)  $\mathcal{DS}$ -continuous [12] if  $f^{-1}(V) \in \mathcal{DS}(X)$  for each  $V \in \sigma$ .
- (3)  $\mathcal{AB}$ -continuous [9] if  $f^{-1}(V) \in \mathcal{AB}(X)$  for each  $V \in \sigma$ .
- (4)  $\mathcal{B}$ -continuous [23] if  $f^{-1}(V) \in \mathcal{B}(X)$  for each  $V \in \sigma$ .

**Remark 25** The following diagram holds for a function  $f : X \rightarrow Y$ :

$$\begin{array}{c}
 \mathcal{B}\text{-continuous} \\
 \uparrow \\
 \mathcal{DS}\text{-continuous} \\
 \uparrow \\
 \mathcal{DS}^*\text{-continuous}
 \end{array}$$

None of these implications is reversible as shown in the following examples.

**Example 26** Let  $X = Y = \{a, b, c, d\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ ,  $\sigma = \{\emptyset, X, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}\}$ . Then the function  $f : (X, \tau) \rightarrow (Y, \sigma)$ , defined as:  $f(a) = c$ ,  $f(b) = b$ ,  $f(c) = c$ ,  $f(d) = d$ , is  $\mathcal{DS}$ -continuous but it is not  $\mathcal{DS}^*$ -continuous.

**Example 27** ([12]) Let  $X = Y = \{a, b, c, d\}$  and  $\tau = \sigma = \{\emptyset, X, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}\}$ . Then the function  $f : (X, \tau) \rightarrow (Y, \sigma)$ , defined as:  $f(a) = b$ ,  $f(b) = a$ ,  $f(c) = c$ ,  $f(d) = d$ , is  $\mathcal{B}$ -continuous but it is not  $\mathcal{DS}$ -continuous.

**Remark 28** Every continuous function is  $\mathcal{DS}^*$ -continuous but not conversely.

**Example 29** Let  $X = Y = \{a, b, c, d\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}\}$ ,  $\sigma = \{\emptyset, X, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}\}$ . Then the function  $f : (X, \tau) \rightarrow (Y, \sigma)$ , defined as:  $f(a) = c$ ,  $f(b) = b$ ,  $f(c) = c$ ,  $f(d) = b$ , is  $\mathcal{DS}^*$ -continuous but it is not continuous.

**Theorem 30** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function. If  $f$  is  $\beta$ -continuous and  $\mathcal{DS}^*$ -continuous, then it is  $\mathcal{AB}$ -continuous.

**Proof.** It follows from Theorem 16. ■

**Theorem 31** *The following are equivalent for a function  $f : X \rightarrow Y$ :*

- (1)  $f$  is continuous,
- (2)  $f$  is  $\alpha$ -continuous and  $\mathcal{DS}^*$ -continuous,
- (3)  $f$  is quasi-continuous and  $\mathcal{DS}^*$ -continuous,
- (4)  $f$  is precontinuous and  $\mathcal{DS}^*$ -continuous,
- (5)  $f$  is  $\gamma$ -continuous and  $\mathcal{DS}^*$ -continuous,
- (6)  $f$  is  $\beta$ -continuous and  $\mathcal{DS}^*$ -continuous.

**Proof.** It is an immediate consequence of Theorem 9. ■

**Theorem 32** *The following are equivalent for a function  $f : X \rightarrow Y$ :*

- (1)  $f$  is continuous,
- (2)  $f$  is  $\delta$ -almost continuous and  $\mathcal{D}$ -continuous,
- (3)  $f$  is  $\delta$ -almost continuous and  $\mathcal{DS}$ -continuous.

**Proof.** It follows from Theorem 13. ■

**Theorem 33** *Let  $f : X \rightarrow Y$  be a function. Then  $f$  is continuous if  $f$  is  $\delta$ -semicontinuous,  $\mathcal{DS}^*$ -continuous and precontinuous or  $\delta^*$ -continuous.*

**Proof.** It is an immediate consequence of Theorem 19. ■

**Theorem 34** *Let  $f : X \rightarrow Y$  be a function. Then  $f$  is super-continuous if  $f$  is  $\delta^*$ -continuous and  $\delta$ -semicontinuous.*

**Proof.** It is an immediate consequence of Theorem 18. ■

## References

- [1] M. E. Abd El-Monsef, S. N. El-Deeb and R. A. Mahmoud,  $\beta$ -open sets and  $\beta$ -continuous mappings, Bull. Fac. Sci. Assiut Univ. 12 (1983) 77-90.
- [2] T. Aho and T. Nieminen, Spaces in which preopen subsets are semiopen, Rich. di Mat., vol. XLIII (1) (1994), 45-59.
- [3] D. Andrijević, Semi-preopen sets, Mat. Vesnik, 38 (1986), 24-32.
- [4] D. Andrijević, On  $b$ -open sets, Mat. Bech., 48 (1996), 59-64.
- [5] N. Bourbaki, General Topology, Part I, Addison Wesley, Reading, Mass 1996.
- [6] H. Carson and E. Michael, Metrizable of certain countable unions, Illinois J. Math., 8 (1964), 351-360.
- [7] Di Maio and T. Noiri, On  $s$ -closed spaces, Indian J. Pure Appl. Math., 18 (3) (1987), 226-233.



- [8] J. Dontchev and M. Przemski, On the various decompositions of continuous and some weakly continuous functions, *Acta Math. Hungar.*, 71 (1-2) (1996), 109–120.
- [9] J. Dontchev, Between  $\mathcal{A}$ - and  $\mathcal{B}$ -sets, *Math. Balkanica (N.S.)*, 12 (3-4) (1998), 295-302.
- [10] W. Dunham, Weakly Hausdorff spaces, *Kyungpook Math. J.*, 15 (1975), 41-50.
- [11] E. Ekici and G. B. Navalagi,  $\delta$ -semicontinuous functions, *Mathematical Forum*, Vol. XVII (2004-2005), 29-42.
- [12] E. Ekici and S. Jafari, On  $\mathcal{D}$ -sets,  $\mathcal{DS}$ -sets and decompositions of continuous,  $\mathcal{A}$ -continuous and  $\mathcal{AB}$ -continuous functions, *Italian J. Pure and Appl. Math.*, in press.
- [13] A. A. El-Atik, A study of some types of mappings on topological spaces, Master's Thesis, Faculty of Science, Tanta Univ., Egypt, 1997.
- [14] S. Kempisty, Sur les fonctions quasicontinues, *Fund. Math.*, 19 (1932), 184-197.
- [15] N. Levine, Semi-open sets and semi-continuity in topological spaces, *Amer. Math. Monthly* 70 (1963) 36-41.
- [16] A. S. Mashhour, M. E. Abd El-Monsef and S. N. El-Deeb, On precontinuous and weak precontinuous mappings, *Proc. Math. Phys. Soc. Egypt*, 53 (1982), 47-53.
- [17] A. S. Mashhour, I. A. Hasanein and S. N. El-Deeb,  $\alpha$ -continuous and  $\alpha$ -open mappings, *Acta Math. Hungar.*, 41 (1983), 213-218.
- [18] B. M. Munshi and D. S. Bassan, Super continuous mappings, *Indian J. Pure and Appl. Math.*, 13 (1982), 229-236.
- [19] O. Njåstad, On some classes of nearly open sets, *Pacific J. Math.*, 15 (1965), 961-970.
- [20] J. H. Park, B. Y. Lee and M. J. Son, On  $\delta$ -semiopen sets in topological space, *J. Indian Acad. Math.*, 19 (1) (1997), 59-67.
- [21] S. Raychaudhuri and M. N. Mukherjee, On  $\delta$ -almost continuity and  $\delta$ -preopen sets, *Bull. Inst. Math. Acad. Sinica*, 21 (1993), 357-366.
- [22] M. Stone, Application of the theory of Boolean rings to general topology, *Trans. Amer. Math. Soc.*, 41 (1937), 374-481.
- [23] J. Tong, On decomposition of continuity in topological spaces, *Acta Math. Hungar.*, 54 (1-2) (1989), 51–55.
- [24] N. V. Veličko, H-closed topological spaces, *Amer. Math. Soc. Transl.*, (2) 78 (1968), 103-118.

Address

Erdal Ekici (Corresponding Author): Department of Mathematics, Canakkale Onsekiz Mart University, Terzioğlu Campus, 17020 Canakkale, Turkey

*E-mail:* eekici@comu.edu.tr

Saeid Jafari: College of Vestsjaelland South, Herrestraede 11, 4200 Slagelse, Denmark

*E-mail:* jafari@stofanet.dk