

Title	On equivariant J-homomorphism for involutions
Author(s)	Minami, Haruo
Citation	Osaka Journal of Mathematics. 20(1) P.109-P.122
Issue Date	1983
Text Version	publisher
URL	https://doi.org/10.18910/8855
DOI	10.18910/8855
rights	
Note	

Osaka University Knowledge Archive : OUKA

<https://ir.library.osaka-u.ac.jp/>

Osaka University

ON EQUIVARIANT J -HOMOMORPHISM FOR INVOLUTIONS

HARUO MINAMI

(Received April 17, 1981)

Introduction. Let G be the cyclic group of order 2.

We denote by $\pi_S^{*,*}$ the equivariant stable cohomotopy theory [2, 3] and by KO_G^* the K -theory of real G -vector bundles on G -spaces. For a finite pointed G -complex we then have an equivariant J_G -map $J: \widetilde{KO}_G^{-1}(X) \rightarrow \pi_S^{0,0}(X)$ [14], which becomes a homomorphism if X is a suspension in the usual sense.

Let $R^{p,q}$ be the euclidean space R^{p+q} with non trivial G -action on the first p coordinates and $\Sigma^{p,q}$ be the one point compactification of $R^{p,q}$, with ∞ as base point. We have the canonical isomorphism $\pi_S^{0,0}(\Sigma^{p,q}) \approx \pi_{p,q}^S$, the (p, q) -th equivariant stable homotopy group of Landweber [9, 3] (which is $\pi_{p+q,p}$ of Bredon [5]), and therefore we get an induced map

$$\widetilde{KO}_G^{-1}(\Sigma^{p,q}) \rightarrow \pi_S^{0,0}(\Sigma^{p,q}) \approx \pi_{p,q}^S$$

which we also denote by J_G . P. Löffler [10] showed that if $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ is a torsion group then J_G is a split injection. In this paper we shall study the image of J_G when $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ is torsion free. And then we shall give a supplement to [11] on $\text{Im } J_R$. The J_G is also studied by M.C. Crabb [6].

We denote by Z/n a cyclic group of order n , by $R \cdot x$ the free module over a ring R generated by x . If $p \equiv i \pmod 8$ and $q \equiv j \pmod 8$, then we write $(p, q) \equiv (i, j) \pmod 8$.

The author would like to express his gratitude to Professor S. Araki for his kindly advice.

1. The J -homomorphism J_G

In this section we shall give the relations between various homomorphisms and collect some basic tools. Let X be a finite pointed G -complex.

Let KR denote the K -functor of [4]. By regarding a Real vector bundle on X as a real G -vector bundle on X we get a homomorphism $\sigma: \widetilde{KR}^{-1}(X) \rightarrow \widetilde{KO}_G^{-1}(X)$. We define a map $J_R: \widetilde{KR}^{-1}(X) \rightarrow \pi_S^{0,0}(X)$ by

$$(1.1) \quad J_R = J_G \sigma$$

which is the same as in [11].

Let ρ , ψ , \mathcal{X} and δ be the homomorphisms defined in [2], §§1-5 and J_o be the usual real J -map. From the definitions of these maps it follows that

$$(1.2) \quad \psi J_G = J_o \psi, \mathcal{X} J_G = J_G \mathcal{X}, \delta J_o = J_G \delta \quad \text{and} \quad \rho J_G = J_G \rho$$

when each map is valid.

Let $RO(G)$ denote the real representation ring of G . If we put $1=R^{0,1}$ and $H=R^{1,0}$ then $RO(G)=Z \cdot 1 \oplus Z \cdot H$. Define a homomorphism $\varepsilon: RO(G) \rightarrow Z$ by the assignment $s+tH \mapsto s$ and define a homomorphism $\phi: \widetilde{KO}_G^{-1}(X) \rightarrow \widetilde{KO}^{-1}(X^G)$ to be the composite of the homomorphisms

$$\begin{aligned} \widetilde{KO}_G^{-1}(X) &\xrightarrow{i^*} \widetilde{KO}_G^{-1}(X^G) \approx \widetilde{KO}^{-1}(X^G) \otimes RO(G) \\ &\xrightarrow{1 \otimes \varepsilon} \widetilde{KO}^{-1}(X^G) \otimes Z = \widetilde{KO}^{-1}(X^G) \end{aligned}$$

where i denotes the inclusion of the fixed point set X^G of X and the isomorphism is the same as in Remark of [13], p. 133. Then we also have

$$(1.3) \quad \phi J_G = J_o \phi$$

where ϕ on the left hand side denotes the homomorphism as in [2], §4.

Let $\hat{\eta}: \Sigma^{2,1} \rightarrow \Sigma^{1,1}$ be the G -map defined in [3], §8 such that $\psi(\hat{\eta}) = \eta: S^3 \rightarrow S^2$ is the Hopf map. We see that $\hat{\eta}$ and η yield natural homomorphisms of our cohomology theories. Denote these homomorphisms by the same letters, i.e.,

$$\begin{aligned} \hat{\eta}: \widetilde{KO}_G^{-1}(\Sigma^{1,1} \wedge X) &\rightarrow \widetilde{KO}_G^{-1}(\Sigma^{2,1} \wedge X), \quad \hat{\eta}: \pi_s^{0,0}(\Sigma^{1,1} \wedge X) \rightarrow \pi_s^{0,0}(\Sigma^{2,1} \wedge X), \\ \eta: \widetilde{KO}^{-1}(S^2 \wedge X) &\rightarrow \widetilde{KO}^{-1}(S^3 \wedge X), \quad \eta: \pi_s^0(S^2 \wedge X) \rightarrow \pi_s^0(S^3 \wedge X). \end{aligned}$$

It is clear that

$$(1.4) \quad \text{i) } \mathcal{X}\hat{\eta} = 1 + \rho, \quad \text{ii) } \psi\hat{\eta} = \eta\psi \quad \text{and} \quad \text{iii) } \hat{\eta}J_G = J_G\hat{\eta}.$$

Note that the first formula follows from [3], (8.1).

Let $\widetilde{Sph}_G^{-1}(X)$ be the group of stable fibre homotopy equivalence classes of spherical G -fibrations on $\Sigma^{0,1}X$ ([7], §7). As in the non equivariant case J_G can be factorized as the composition of maps $\widetilde{KO}_G^{-1}(X) \rightarrow \widetilde{Sph}_G^{-1}(X) \rightarrow \pi_s^{0,0}(X)$. By [7], Theorem 0.4 we therefore have

Theorem 1.5 ([7], Theo. 0.4). *If X is a suspension then, for any $x \in \widetilde{KO}_G^{-1}(X)$ there exists $e > 0$ such that $3^e J_G(\psi^3 - 1)x = 0$.*

Finally we recall the following theorems.

Theorem 1.6 ([1, 12]).

$$\text{Im } \{J_0: \widetilde{KO}^{-1}(S^{4p-1}) \rightarrow \pi_{4p-1}^S\} \approx Z/m(2p)$$

where $m(t)$ denotes the numerical function as in [1, II].

Theorem 1.7 ([5], p. 272).

$$\psi \oplus \phi: \pi_{p,q}^S \left[\frac{1}{2} \right] \rightarrow (\pi_{p+q}^S \oplus \pi_q^S) \left[\frac{1}{2} \right] \quad \text{for } p \text{ even}$$

and

$$\phi: \pi_{p,q}^S \left[\frac{1}{2} \right] \rightarrow \pi_q^S \left[\frac{1}{2} \right] \quad \text{for } p \text{ odd}$$

are isomorphisms.

2. The equivariant K-theory of spheres

According to [10] $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ is an abelian 2-group or a free abelian group (with a single generator as a module over $RO(G)$). We refer the reader to Table in [10] for these groups. We extract only the cases when $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ is free.

- (2.1) a) If $(p, q) \equiv (2, 1), (6, 1), (2, 5)$ or $(6, 5) \pmod 8$ then $\widetilde{KO}_G^{-1}(\Sigma^{p,q}) \approx Z$,
 b) If $(p, q) \equiv (0, 3), (4, 3), (0, 7)$ or $(4, 7) \pmod 8$ then $\widetilde{KO}_G^{-1}(\Sigma^{p,q}) \approx RO(G)$ (cf. [9]),
 c) If $(p, q) \equiv (i, 3)$ or $(i, 7) \pmod 8$ for $i=1, 2, 3, 5, 6$ or 7 then $\widetilde{KO}_G^{-1}(\Sigma^{p,q}) \approx Z$.

In this section we provide preliminary lemmas. By [9], Lemma 4.1 and [13], $\widetilde{KO}_G^{-1}(\Sigma^{p,q}/\Sigma^{0,q}) \approx KO^{-q-2}(P^{p-1})$, where P^n is the real projective n -space. This isomorphism and the exact sequence of the pair $(\Sigma^{p,q}, \Sigma^{0,q})$ induce the exact sequence:

$$(2.2) \quad \rightarrow KO^{-q-2}(P^{p-1}) \rightarrow \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \xrightarrow{i^*} \widetilde{KO}_G^{-1}(\Sigma^{0,q}) \rightarrow KO^{-q-1}(P^{p-1}) \rightarrow$$

where i is the inclusion $\Sigma^{0,q} \subset \Sigma^{p,q}$.

Lemma 2.3. *Let t be a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$, $p \geq 1$. Then*

- i) $\rho(t) = -t$ and $Ht = t$ in case of (2.1), a),
- ii) $\rho(t) = -Ht$ in case of (2.1), b),
- iii) $\rho(t) = t$ and $Ht = -t$ in case of (2.1), c).

Proof. i) Using [2], (5.1) of KO_G^* we see that $\psi: \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \rightarrow \widetilde{KO}^{-1}(S^{p+q})$ is monic. From this and the equalities $\psi\rho = -\psi$ ([2], §3) and $\psi H = \psi$ the claim follows immediately.

ii) From the equality $\rho^2 = 1$ it follows that $\rho(t) = \pm t$ or $\pm Ht$. Since $\psi(t)$ is a generator of $\widetilde{KO}^{-1}(S^{p+q})$, the relation $\psi\rho = -\psi$ implies that $\rho(t) = -t$ or

— Ht . Using [8], by (2.2) we get $i^*(t) = 2^s(1-H)t_0$ for some $s > 0$ up to sign, where t_0 denotes a generator of $\widetilde{KO}_G^{-1}(\Sigma^{0,q})$. This and the relation $i^*\rho = i^*$ (cf. [2]) show that $\rho(t) = -Ht$.

iii) Analogous to ii). To prove the second we have to use that i^* is a homomorphism of $RO(G)$ -modules. q.e.d.

Lemma 2.4. i) In case of (2.1), a), if $(p, q) \equiv (2, 1)$ or $(6, 5) \pmod{8}$ then $\psi: \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \approx \widetilde{KO}^{-1}(S^{p+q})$, and if $(p, q) \equiv (2, 5)$ or $(6, 1) \pmod{8}$ then ψ carries a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ to twice a generator of $\widetilde{KO}^{-1}(S^{p+q})$. And $\phi = 0$ on $\widetilde{KO}^{-1}(S^q) \left[\frac{1}{2} \right]$,

ii) In case of (2.1), b), $\psi(t)$ and $\phi(t)$ generate $\widetilde{KO}^{-1}(S^{p+q})$ and $\widetilde{KO}^{-1}(S^q) \left[\frac{1}{2} \right]$,

iii) In case of (2.1), c), $\psi = 0$ on $\widetilde{KO}_G^{-1}(\Sigma^{p,q}) \left[\frac{1}{2} \right]$ and $\phi(t)$ generates $\widetilde{KO}^{-1}(S^q) \left[\frac{1}{2} \right]$, where t denotes a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$.

Proof. i) Making use of [2], (5.1) of KO_G^* we get $\text{Im } \psi = \widetilde{KO}^{-1}(S^{p+q})$ or $2\widetilde{KO}^{-1}(S^{p+q})$ according as $p+q \equiv 3$ or $7 \pmod{8}$. The second is immediate.

ii) The claim for ψ is obvious by the definition of t . As in the proof of Lemma 2.3, ii) we obtain $i^*(t) = 2^s(1-H)t_0$, with the above notation, when $p > 0$. So $\phi(t)$ is 2^s times of a generator of $\widetilde{KO}^{-1}(S^q)$ and hence the claim for $p > 0$ follows immediately. Clear when $p = 0$.

iii) Since $\widetilde{KO}^{-1}(S^{p+q})$ is a 2-group, the claim for ψ is obvious. For ϕ , same as in ii). q.e.d.

Lemma 2.5. For $\chi: \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \rightarrow \widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$, $p \geq 1$, $q \geq 0$, and $\hat{\eta}: \widetilde{KO}_G^{-1}(\Sigma^{p-1,q}) \rightarrow \widetilde{KO}_G^{-1}(\Sigma^{p,q})$, $p \geq 2$, $q \geq 1$, we have

i) If $(p, q) \equiv (2, 3), (3, 3), (6, 7)$ or $(7, 7) \pmod{8}$ then $\chi: \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \approx \widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$, and $\hat{\eta}$ carries a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$ to twice a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$,

ii) If $(p, q) \equiv (2, 7), (3, 7), (6, 3)$ or $(7, 3) \pmod{8}$ then χ carries a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ to twice a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$, and $\hat{\eta}: \widetilde{KO}_G^{-1}(\Sigma^{p-1,q}) \approx \widetilde{KO}_G^{-1}(\Sigma^{p,q})$,

iii) If $(p, q) \equiv (1, 3), (5, 3), (1, 7)$ or $(5, 7) \pmod{8}$ then χ carries a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ to $(1+\rho)$ or $(1-H)$ times a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$ according as $p > 1$ or $p = 1$, and $\hat{\eta}$ is epic,

iv) If $(p, q) \equiv (0, 3), (4, 3), (0, 7)$ or $(4, 7) \pmod{8}$ then χ is epic, and $\hat{\eta}$ carries

a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$ to $(1+\rho)$ times a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$.

Proof. For the proof we use Lemma 2.3 freely. The claims for χ follow immediately from [2], (5.1) of KO_G^* . Hence i)–iii) for $\hat{\eta}$ follow easily from the results for χ and (1.4), i). We now prove iv) for $\hat{\eta}$. Let t be a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ and write $\chi(t)=t'$. Then t' becomes a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p-1,q})$ by iv) for χ . Put $\hat{\eta}(t')=(a+bH)t$, then, by (1.4), i) we see that $a-b=2$. By (1.4), ii) $\psi\hat{\eta}(t')=\eta\psi(t')=0$ because $\widetilde{KO}^{-1}(S^{p+q-1})=0$. This implies that $a+b=0$ and therefore $a=-b=1$. Hence $\hat{\eta}(t')=(1-H)t=(1+\rho)t$. q.e.d.

Finally we consider

$$\sigma: \widetilde{KR}^{-1}(\Sigma^{p,q}) \rightarrow \widetilde{KO}_G^{-1}(\Sigma^{p,q})$$

when $\widetilde{KR}^{-1}(\Sigma^{p,q}) \approx Z$.

Lemma 2.6. i) If $(p, q) \equiv (2, 1)$ or $(6, 5) \pmod 8$ then $\sigma: \widetilde{KR}^{-1}(\Sigma^{p,q}) \approx \widetilde{KO}_G^{-1}(\Sigma^{p,q})$, and if $(p, q) \equiv (2, 5)$ or $(6, 1) \pmod 8$ then σ carries a generator of $\widetilde{KR}^{-1}(\Sigma^{p,q})$ to twice a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$,

ii) If $(p, q) \equiv (0, 3), (4, 3), (0, 7)$ or $(4, 7) \pmod 8$ then σ carries a generator of $\widetilde{KR}^{-1}(\Sigma^{p,q})$ to $(1-\rho)$ or $(1+H)$ times a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ according as $p > 0$ or $p = 0$.

Proof. By r and c we denote the realification homomorphism and the complexification homomorphism. Then it is clear that $\psi\sigma=rc$. Let \bar{t} , t and t_0 be generators of $\widetilde{KR}^{-1}(\Sigma^{p,q})$, $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$ and $\widetilde{KO}^{-1}(S^{p+q})$ respectively. From the relations between c , r and the coefficients of K -theories it follows that

$$rc(\bar{t}) = \begin{cases} t_0 & \text{for } (p, q) \equiv (2, 1), (6, 5) \pmod 8 \\ 4t_0 & \text{for } (p, q) \equiv (2, 5), (6, 1) \pmod 8 \\ 2t_0 & \text{otherwise} \end{cases}$$

up to sign.

i) Immediate from this and Lemma 2.4, i).

ii) Since $\widetilde{KR}^{-1}(\Sigma^{0,q}) \approx KO^{-1}(S^q)$ the assertion for $p=0$ is obvious. So we consider the case $p > 0$. Put $\sigma(\bar{t})=(a+bH)t$. Since $\psi(t)=t_0$ (up to sign) and $rc(\bar{t})=2t_0$, we get $a+b=\pm 2$. ρ acts on $\widetilde{KR}^{-1}(\Sigma^{p,q})$ as -1 because c is monic. Using this and Lemma 2.3, ii), from the equality $\sigma\rho=\rho\sigma$ it follows that $a=b$ and hence $a=b=\pm 1$. This and Lemma 2.3, ii) complete the proof. q.e.d.

3. $\text{Im } J_G$ and $\text{Im } J_R$

Let

$$I_{p,q} = \text{Im} \{J_G: \widetilde{KO}_G^{-1}(\Sigma^{p,q}) \rightarrow \pi_{p,q}^S\} \quad \text{for } q \geq 1.$$

Denote by t a generator of $\widetilde{KO}_G^{-1}(\Sigma^{p,q})$, so that $\widetilde{KO}_G^{-1}(\Sigma^{p,q}) = Z \cdot t$ or $RO(G) \cdot t$ in our cases. Put

$$\alpha = J_G(t).$$

Let $m(s)$ be as in Theorem 1.6 and $v_p(s)$, p prime, be the power in s . (We write $\rho(x) = \rho x$.)

Theorem 3.1. *Suppose that $(p, q) \equiv (2, 1), (6, 1), (2, 5)$ or $(6, 5) \pmod{8}$. Then*

$$I_{p,q} = Z/m\left(\frac{p+q+1}{2}\right) \cdot \alpha.$$

Theorem 3.2. *Suppose that $(p, q) \equiv (i, 3)$ or $(i, 7) \pmod{8}$ for $i=1, 2, 3, 5, 6$ or 7 . Then*

$$I_{p,q} = Z/m\left(\frac{q+1}{2}\right) \cdot \alpha.$$

Theorem 3.3. i) *Suppose that $q \equiv 3$ or $7 \pmod{8}$. Then*

$$I_{0,q} = Z/m\left(\frac{q+1}{2}\right) \cdot \alpha \oplus Z/m\left(\frac{q+1}{2}\right) \cdot \beta$$

where $\beta = J_G(Ht)$,

ii) *Suppose that $\varepsilon = 0$ or 1 , $v_2(2q+\varepsilon) \leq v_2(p+q+\varepsilon)$ and*

$$\left(\frac{m(4p+4q+4\varepsilon)}{2^{v_2(p+q+\varepsilon)+4}d}, d\right) = 1$$

where

$$d = \left(\frac{m(4p+4q+4\varepsilon)}{2^{v_2(p+q+\varepsilon)+4}}, \frac{m(4q+2\varepsilon)}{2^{v_2(2q+\varepsilon)+3}}\right).$$

Then, for $8p+4\varepsilon > 0$

$$I_{8p+4\varepsilon, 8q+4\varepsilon-1} = Z/\frac{m(4p+4q+4\varepsilon)m(4q+2\varepsilon)}{2^{v_2(2q+\varepsilon)+2}d} \cdot \alpha \oplus Z/2^{v_2(2q+\varepsilon)+2}d \cdot \gamma$$

where

$$\gamma = \left(\frac{m(4p+4q+4\varepsilon)}{2^{v_2(2p+\varepsilon)+3}d} + \frac{m(4q+2\varepsilon)}{2^{v_2(2q+\varepsilon)+3}}\right)(1+\rho)\alpha - \frac{m(4p+4q+4\varepsilon)}{2^{v_2(2p+\varepsilon)+2}d} \alpha.$$

Corollary 3.4.

i) $I_{4p, 4p-1} = Z/2m(4p) \cdot \alpha \oplus Z/\frac{m(2p)}{2} \cdot \gamma$

where

$$\gamma = \left(\frac{m(4p)}{m(2p)} + \frac{m(2p)}{2^{\nu_2(p)+3}} \right) (1+\rho)\alpha - \frac{2m(4p)}{m(2p)} \alpha,$$

ii) Suppose that $\nu_3(p) \leq \nu_3(2p-1)$. Then

$$I_{4,8p-5} = Z/\frac{m(4p)m(4p-2)}{4 \cdot 3^{\nu_3(p)+1}} \cdot \alpha \oplus Z/4 \cdot 3^{\nu_3(p)+1} \cdot \gamma$$

where

$$\gamma = \left(\frac{m(4p)}{8 \cdot 3^{\nu_3(p)+1}} + \frac{m(4p-2)}{8} \right) (1+\rho)\alpha - \frac{m(4p)}{4 \cdot 3^{\nu_3(p)+1}} \alpha.$$

REMARK. We can take γ to be the following simpler elements respectively in Corollary 3.4:

i) $\frac{m(4p)}{2m(2p)} ((1+\rho)\alpha - 4\alpha),$

ii) $(1+\rho)\alpha - \frac{3^{4p}-1}{4}\alpha$ for $\nu_3(2p-1)-2 \leq \nu_3(p) \leq \nu_3(2p-1)$. (ii) was sug-

gested by the computation of $\pi_{4,3}^S$ by Araki-Iriye [3] and also this suggested (5.4) of §5.)

Let $o(x)$ denote the order of an element x of a finite group. Then we obtain

Proposition 3.5. Let $\alpha \in \pi_{8p+4\epsilon, 8q+4\delta-1}^S$, $8p+4\epsilon > 0$, where $\epsilon, \delta = 0$ or 1 . Then

i) $o((1-\rho)\alpha) = m(4p+4q+2\epsilon+2\delta),$

$o((1+\rho)\alpha) = m(4q+2\delta),$

ii) $o(\alpha) = \frac{m(4p+4q+2\epsilon+2\delta)m(4q+2\delta)}{2^{\zeta}d}$

where

$$d = \left(\frac{m(4p+4q+2\epsilon+2\delta)}{2^{\nu_2(2p+2q+\epsilon+\delta)+3}}, \frac{m(4q+2\delta)}{2^{\nu_2(2q+\delta)+3}} \right)$$

and ζ is the following integer:

$\nu_2(2q+\epsilon)+2$ if $\epsilon = \delta$ and $\nu_2(2q+\epsilon) \leq \nu_2(p+q+\epsilon),$

$\nu_2(2q+\epsilon)+2$ or $\nu_2(2q+\epsilon)+3$

if $\epsilon = \delta$ and $\nu_2(2q+\epsilon) = \nu_2(p+q+\epsilon)+1,$

$\nu_2(p+q+\epsilon)+3$ if $\epsilon = \delta$ and $\nu_2(2q+\epsilon) \geq \nu_2(p+q+\epsilon)+2,$

2 or 3 if $\epsilon = 0$ and $\delta = 1,$

2 if $\epsilon = 1$ and $\delta = 0.$

For a finite abelian group A , we denote by A_p the p -component of A and by $\tilde{x} \in A_p$ the p -component of $x \in A$.

Proposition 3.6. i) Suppose that $v_2(2q+\varepsilon) \leq v_2(p+q+\varepsilon)$, $\varepsilon=0$ or 1 . Then

$$(I_{8p+4\varepsilon, 8q+4\varepsilon-1})_2 = Z/2^{v_2(p+q+\varepsilon)+5} \cdot \bar{\alpha} \oplus Z/2^{v_2(2p+\varepsilon)+2} \cdot \bar{\gamma}.$$

for $8p+4\varepsilon > 0$ where

$$\gamma = (1+\rho)\alpha - 2^{v_2(p+q+\varepsilon)-v_2(2p+\varepsilon)+2}\alpha,$$

ii) Suppose that $v_2(2q+\varepsilon) \geq v_2(p+q+\varepsilon)+2$, $\varepsilon=0$ or 1 . Then

$$(I_{8p+4\varepsilon, 8p+4\varepsilon-1})_2 = Z/2^{v_2(2q+\varepsilon)+4} \cdot \bar{\alpha} \oplus Z/2^{v_2(2p+\varepsilon)+2} \cdot \bar{\gamma}$$

for $8p+4\varepsilon > 0$ where

$$\gamma = \frac{3^{4p+2\varepsilon}-1}{2^{v_2(2p+\varepsilon)+3}}(1+\rho)\alpha - \frac{3^{4p+4q+4\varepsilon}-1}{2^{v_2(2p+\varepsilon)+2}}\alpha,$$

iii) $(I_{8p+4, 8q-1})_2 = Z/2^{v_2(q)+5} \cdot \bar{\alpha} \oplus Z/4 \cdot \bar{\gamma}$

for $p \geq 0$ where

$$\gamma = \frac{3^{4p+2}-1}{8}(1+\rho)\alpha - \frac{3^{4p+4q+2}-1}{4}\alpha.$$

Let

$$\bar{I}_{p,q} = \text{Im} \{J_R: \widetilde{KR}^{-1}(\Sigma^{p,q}) \rightarrow \pi_{p,q}^S\} \quad \text{for } q \geq 1.$$

Suppose that $\widetilde{KR}^{-1}(\Sigma^{p,q}) \approx Z$, generated by \bar{t} , and set

$$\bar{\alpha} = J_R(\bar{t}).$$

Then we have

Theorem 3.7.

$$\bar{I}_{p,q} = \begin{cases} Z/m(2s) \cdot \bar{\alpha} & \text{if } (p, q) \equiv (2, 5) \text{ or } (6, 1) \pmod{8} \\ Z/m(2s) \cdot \bar{\alpha} & \text{if } (p, q) \equiv (2, 1), (6, 5), (0, 3), (4, 3) \\ & (0, 7) \text{ or } (4, 7) \pmod{8} \end{cases}$$

where $s = \frac{p+q+1}{2}$.

4. Proofs of Theos. 3.1, 3.2 and 3.3, i)

Proof of Theorem 3.1. Put $m = m\left(\frac{p+q+1}{2}\right)$ for brevity.

We divide the proof into two cases: (i) $(p, q) \equiv (2, 1)$ or $(6, 5) \pmod{8}$ and (ii) $(p, q) \equiv (2, 5)$ or $(6, 1) \pmod{8}$.

Let t_0 be a generator of $\widetilde{KO}^{-1}(S^{p+q})$. By Lemma 2.4, i), (1.2) and (1.3) we have

$$\psi(\alpha) = \begin{cases} J_o(t_0) & \text{for (i)} \\ 2J_o(t_0) & \text{for (ii)} \end{cases}$$

up to sign and $\phi=0$ on $\pi_{p,q}^S\left[\frac{1}{2}\right]$ in any case. Therefore, by Theorems 1.6 and 1.7 we see that the odd components of $o(\alpha)$ are equal to those of m , and also $m|o(\alpha)$ for (i) and $\frac{m}{2}|o(\alpha)$ for (ii). Since ψ of KO_G^* is monic, $\psi^3(t)=3^{(p+q+1)/2}t$. From Theorem 1.5 and the equality $\nu_2(3^{(p+q+1)/2}-1)=\nu_2(m)$ ([1, II], Lemma (2.12)) it follows that $\nu_2(m)\geq\nu_2(o(\alpha))$. Hence $o(\alpha)=m$ for (i).

For (ii) it suffices to prove that $\frac{m}{2}\alpha\neq 0$. Observe [2], (5.1) of KO_G^* . Then, using (1.2) we have $\delta(J_o(t_0))=\alpha$ (up to sign). So if $\frac{m}{2}\alpha=0$, then, by [2], (10.5) there exists an element β of $\pi_{p-1,q+1}^S$ such that $\psi(\beta)=\frac{m}{2}J_o(t_0)$. Using the equalities $\widetilde{KO}_G(\Sigma^{8n+p-1,8n+q+1})=\widetilde{KO}_G(\Sigma^{8n+p-1,8n+q+2})=0$ and the formula for ψ^k on $\widetilde{KO}_G(\Sigma^{8n,8n})$ of [9] we see that β gives rise to a contradiction such that $e_k\left(\frac{m}{2}J_o(t_0)\right)=0$ [1, IV]. Therefore also we have $o(\alpha)=m$ for (ii). Thus we obtain $\text{Im } J_G=Z/m\cdot\alpha$ because $\widetilde{KO}_G^{-1}(\Sigma^{p,q})=Z\cdot t$.

Proof of Theorem 3.2. Let $t\in\widetilde{KO}_G^{-1}(\Sigma^{8p+4\varepsilon+i,8q+4\delta-1})$, $0\leq\varepsilon, \delta\leq 1$ and $1\leq i\leq 3$. By Lemma 2.5 we have

$$\mathcal{X}^i(t) = (1+\rho)t_0 \text{ or } (1-H)t_0$$

according as $8p+4\varepsilon>0$ or $8p+4\varepsilon=0$ for $(\varepsilon, \delta)=(0, 1)$ or $(1, 0)$ and

$$\hat{\eta}^{4-i}(t) = (1+\rho)t_1$$

for $(\varepsilon, \delta)=(0, 0)$ or $(1, 1)$, where $t_s, s=0, 1$, is a generator of $\widetilde{KO}_G^{-1}(\Sigma^{8p+4\varepsilon+4s,8q+4\delta-1})$. Therefore, using (1.2) and (1.4), iii), we get

$$(*) \quad m(4q+2\delta)|o(\alpha)$$

by Theorem 3.3, i) and Proposition 3.5, i), which will be proved in the following.

The proof of the inverse of (*) is analogous to that of Theorem 3.1 for (i). Then we have need of the equality $\psi^3(t)=3^{4q+2\delta}t$, which is obtained by the injectivity of \mathcal{X}^i or $\hat{\eta}^{4-i}$ and the equality $\psi^3((1-H)t_s)=3^{4q+2\delta}(1-H)t_s, 0\leq s\leq 1$ [9].

Proof of Theorem 3.3, i). Let t_0 be a generator of $\widetilde{KO}^{-1}(S^q)$. Then, $\psi(\alpha)$

$=\psi(\beta)=J_o(t_0)$, $\phi(\alpha)=J_o(t_0)$ (up to sign) and $\phi(\beta)=0$ clearly. We get $o(\alpha)=o(\beta)=m\left(\frac{q+1}{2}\right)$ as in the above, so that it follows that α and β have no relation.

5. Proofs of Theos. 3.3, ii), 3.7, Cor. 3.4, Props. 3.5 and 3.6

As we saw in §4, by Theorems 1.6, 1.7, Lemma 2.4, ii), (1.2) and (1.3) we can determine the odd components of the order of elements of $\text{Im } J_G$. So we observe only the 2-component. Let $o_2(\)$ denote the 2-component of $o(\)$. We use freely the equalities

$$v_2(3^{2s}-1) = v_2(m(2s)) = v_2(s)+3$$

([1, II], p. 139 and Lemma (2.12)) in the following.

Let α and $\bar{\alpha}$ be as in §3. By Lemma 2.3, ii) and (1.2)

$$\rho\alpha = -J_G(Ht), \quad t \in \widetilde{KO}_G^{-1}(\Sigma^{8p+4\varepsilon, 8q+4\delta-1}).$$

We begin by considering $o((1-\rho)\alpha)$. Write $t_0=\psi(t)$. Then t_0 is a generator of $\widetilde{KO}_G^{-1}(S^{8p+8q+4\varepsilon+4\delta-1})$. Put $\delta(t_0)=(a+bH)t$. By the definition of δ we get $\psi\delta(t_0)=2t_0$, hence $a+b=2$. Using [9], Lemma 3.3 and Proposition 3.5, from the equality $\delta\psi\psi^2=\psi^2\delta\psi$ it follows that $a=b$. Therefore $a=b=1$, so that $\delta(t_0)=(1+H)t$. This and (1.2) show $\delta(J_o(t_0))=(1-\rho)\alpha$. Since $o(J_o(t_0))=m(4p+4q+2\varepsilon+2\delta)$ by Theorem 1.6, we therefore have

$$o((1-\rho)\alpha) \mid m(4p+4q+2\varepsilon+2\delta).$$

Suppose that $o((1-\rho)\alpha)=\frac{m(4p+4q+2\varepsilon+2\delta)}{2}$. Then, by [2], (10.5) we have an element $\beta \in \pi_{8p+4\varepsilon-1, 8q+4\delta}^S$ such that $\psi(\beta)=\frac{m(4p+4q+2\varepsilon+2\delta)}{2}J_o(t_0)$. As in the proof of Theorem 3.1, β yields a contradiction: $e'_R(\frac{m(4p+4q+2\varepsilon+2\delta)}{2}J_o(t_0))=0$. This proves

$$(5.1) \quad o((1-\rho)\alpha) = m(4p+4q+2\varepsilon+2\delta).$$

By Theorem 1.5, [9], Lemma 3.3 and Proposition 3.5, we obtain

$$(5.2) \quad \frac{3^{4p+4q+2\varepsilon+2\delta}-1}{2}(\rho-1)\bar{\alpha} = \frac{3^{4q+2\delta}-1}{2}(\rho+1)\bar{\alpha}.$$

From (5.1) and (5.2) it follows that

$$(5.3) \quad o((1+\rho)\alpha) = m(4q+2\delta).$$

Using (5.3), the equality (5.2) can be transformed into the following

$$(5.4) \quad (3^{4p+4q+2\varepsilon+2\delta}-1)\tilde{\alpha}-\frac{3^{4p+2\varepsilon}-1}{2}(1+\rho)\tilde{\alpha}=0.$$

Proof of Proposition 3.5. i) Immediate from (5.1) and (5.3).

ii) The proof breaks up into the 4 cases.

Case 1) $\varepsilon=\delta$ and $\nu_2(2q+\varepsilon)\leq\nu_2(p+q+\varepsilon)$. We have

$$(*) \quad 2^{\nu_2(p+q+\varepsilon)+3}(1+\rho)\tilde{\alpha}=0$$

by (5.3). By (5.1), $2^{\nu_2(p+q+\varepsilon)+4}(1-\rho)\tilde{\alpha}=0$. Hence we get

$$2^{\nu_2(p+q+\varepsilon)+5}\tilde{\alpha}=0.$$

If $2^{\nu_2(p+q+\varepsilon)+4}\tilde{\alpha}=0$, then

$$\begin{aligned} 2^{\nu_2(p+q+\varepsilon)+3}\tilde{\alpha} &= -2^{\nu_2(p+q+\varepsilon)+3}\tilde{\alpha} \\ &= 2^{\nu_2(p+q+\varepsilon)+3}\rho\tilde{\alpha} \quad \text{by } (*). \end{aligned}$$

This shows $2^{\nu_2(p+q+\varepsilon)+3}(1-\rho)\tilde{\alpha}=0$, which contradicts to (5.1). Therefore we have

$$(5.5) \quad o_2(\alpha) = 2^{\nu_2(p+q+\varepsilon)+5}.$$

By (5.4) and (5.5) we have $o_2((1+\rho)\alpha)=2^{\nu_2(2p+\varepsilon)+3}$. Comparing this with (5.3) we get

$$(5.6) \quad \nu_2(2p+\varepsilon) = \nu_2(2q+\varepsilon).$$

Case 2) $\varepsilon=\delta$ and $\nu_2(2q+\varepsilon)=\nu_2(p+q+\varepsilon)+1$. As in Case 1) we can prove $\nu_2(o_2(\alpha))\leq\nu_2(2q+\varepsilon)+4$. To show the inverse we have to use the fact that $(\text{Im } J_o)_2 = Z/2^{\nu_2(2q+\varepsilon)+3} \cdot \nu(\alpha)$ (Lem. 2.4, ii) and Theo. 1.6). Hence we have

$$(5.7) \quad o_2(\alpha) = 2^{\nu_2(2q+\varepsilon)+3} \text{ or } 2^{\nu_2(2q+\varepsilon)+4}.$$

Case 3) $\varepsilon=\delta$ and $\nu_2(2q+\varepsilon)\geq\nu_2(p+q+\varepsilon)+2$. The same argument as in Case 1) shows

$$(5.8) \quad o_2(\alpha) = 2^{\nu_2(2q+\varepsilon)+4}$$

and

$$(5.9) \quad \nu_2(2p+\varepsilon) = \nu_2(p+q+\varepsilon)+1.$$

Case 4) $(\varepsilon, \delta)=(0, 1)$ or $(1, 0)$. As in Cases 1) and 2) we can prove

$$(5.10) \quad o_2(\alpha) = 2^3 \text{ or } 2^4 \text{ for } (\varepsilon, \delta) = (0, 1) \text{ and } 2^{\nu_2(q)+5} \text{ for } (\varepsilon, \delta) = (1, 0).$$

(5.5), (5.7), (5.8) and (5.10) complete the proof of ii).

Proof of Theorem 3.3, ii). For the simplicity we put

$$m(4p+4q+4\varepsilon)=2^{\nu_2(p+q+\varepsilon)+4}ad \text{ and } m(4q+2\varepsilon)=2^{\nu_2(2q+\varepsilon)+3}bd$$

where d is as in §3. Then a , b and d are odd and $(a, b)=1$.

Since $\nu_2(2p+\varepsilon)=\nu_2(2q+\varepsilon)$ by (5.6), we have

$$(5.11) \quad 2^{\nu_2(2p+\varepsilon)+2}(1+\rho)\tilde{\alpha} = 2^{\nu_2(p+q+\varepsilon)+4}\tilde{\alpha}$$

by (5.4), (5.3) and (5.5).

Define

$$\gamma = (2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+1}a+bd)(1+\rho)\alpha - 2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}a\alpha.$$

We observe $o(\gamma)$. Since $\nu_2(p+q+\varepsilon) \geq \nu_2(2p+\varepsilon) = \nu_2(2q+\varepsilon)$, $o_2((1+\rho)\alpha) = 2^{\nu_2(2q+\varepsilon)+3}$ and $o_2(\alpha) = 2^{\nu_2(p+q+\varepsilon)+5}$ we have

$$2^{\nu_2(2p+\varepsilon)+2}\tilde{\gamma} = 0$$

by (5.11). Note that

$$\psi(\gamma) = -2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}a\psi(\alpha), \quad \phi(\gamma) = 2bd\phi(\alpha)$$

and

$$o(\psi(\alpha)) = 2^{\nu_2(p+q+\varepsilon)+4}ad, \text{ the odd component of } o(\phi(\alpha)) = bd.$$

Since $\psi(2^{\nu_2(2q+\varepsilon)+1}\tilde{\gamma}) = -2^{\nu_2(p+q+\varepsilon)+3}a\psi(\tilde{\alpha}) \neq 0$,

$$o(\tilde{\gamma}) = 2^{\nu_2(2q+\varepsilon)+2}$$

and so

$$(5.12) \quad o(\gamma) = 2^{\nu_2(2q+\varepsilon)+2}d.$$

From the assumption such that $(a, d)=1$ it follows that

$$(*) \quad (2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+1}a+bd, 2^{\nu_2(2p+\varepsilon)+3}bd) = 1.$$

Using (5.3), we then have

$$(1+\rho)\alpha = x(\gamma + 2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}a\alpha)$$

for some integer x . This shows that α and γ generate $\text{Im } J_G$.

Next we prove that there is no relation between α and γ . Suppose that $x\alpha = y\gamma$. Then

$$(x + 2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}ay)\alpha = y(2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+1}a+bd)(1+\rho)\alpha$$

and

$$(x + 2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}ay)\psi(\alpha) = 0.$$

Hence

$$2^{\nu_2(p+q+\varepsilon)+4}ad \mid x + 2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+2}ay .$$

Since $o(\alpha) = 2^{\nu_2(p+q+\varepsilon)+5}abd$,

$$2by(2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+1}a + bd) (1 + \rho)\alpha = 0 .$$

So

$$2^{\nu_2(2q+\varepsilon)+3}bd \mid 2by(2^{\nu_2(p+q+\varepsilon)-\nu_2(2p+\varepsilon)+1}a + bd)$$

since $o((1 + \rho)\alpha) = 2^{\nu_2(2q+\varepsilon)-3}bd$. From this and (*), it follows that

$$2^{\nu_2(2q+\varepsilon)+2}d \mid y .$$

Therefore $x\alpha = y\gamma = 0$ by (5.12).

Proof of Corollary 3.4. i) Let $p = q$ and replace $2p + \varepsilon$ by p in Theorem 3.3, ii). If $\nu_q(m(2p)) \geq 1$ for odd prime q then $\nu_q(m(2p)) = \nu_q(m(4p))$ ([1, II], p. 139). Hence, $d = \frac{m(2p)}{2^{\nu_2(p)+3}}$ and so the assumption holds. The remaining follows easily.

ii) Take $p = 0$ and $\varepsilon = 1$, and replace q by $p - 1$ in Theorem 3.3, ii). By the definition of $m(t)$ [1, II] we have

$$\nu_3(m(4p - 2)) = 1 + \nu_3(2p - 1), \nu_3(m(4p)) = 1 + \nu_3(p)$$

and if $\nu_q(m(4p - 2)) \geq 1$ for a prime $q \geq 5$, then $\nu_q(m(4p)) = 0$. Therefore,

$$d = 3^{1 + \min\{\nu_3(p), \nu_3(2p-1)\}} = 3^{1 + \nu_3(p)}$$

using the condition. This shows that the assumption holds. The rest is immediate.

Proof of Proposition 3.6. Similar to the proof of Theorem 3.3, ii). We make only a remark. In order to determine $o_2(\gamma)$ we have to use (5.4) and $o_2(\psi(\alpha))$. From $o_2(\gamma)$, $o_2(\alpha)$ and $o_2((1 + \rho)\alpha)$ it follows that there is no relation between α and γ .

Proof of Theorem 3.7. This theorem follows immediately from Lemma 2.6, Theorem 3.1 and Proposition 3.5, i).

ADDENDUM TO THE REMARK OF §3. This is proved by the analogous argument to Theorem 3.3, ii) and Proposition 3.6. We note that in order to determine $o(\gamma)$ in ii) we have need of the inequality of [1, II], Lemma (2.12): $\nu_q(3^{2s} - 1) \geq \nu_q(m(2s))$, q prime ≥ 5 .

References

- [1] J.F. Adams: *On the groups $J(X)$* — II, IV, *Topology* **3** (1965), 137–171, **5** (1966), 21–71.
- [2] S. Araki and M. Murayama: *τ -Cohomology theories*, *Japan. J. Math.* **4** (1978), 363–416.
- [3] S. Araki and K. Iriye: *Equivariant stable homotopy groups of spheres with involutions*, I, *Osaka J. Math.* **19** (1982), 1–55.
- [4] M.F. Atiyah: *K-theory and reality*, *Quart. J. Math.*, Oxford, **17** (1966), 367–386.
- [5] G.E. Bredon: *Equivariant stable stems*, *Bull. Amer. Math. Soc.* **73** (1967), 269–273.
- [6] M.C. Crabb: *$Z/2$ -homotopy theory*, *London Math. Soc. Lecture Note Series*, **44** (1981).
- [7] Z. Fiedorowicz, H. Hauschild and J.P. May: *Equivariant algebraic K-theory*, Preprint (1981).
- [8] M. Fujii: *K_o -groups of projective spaces*, *Osaka J. Math.* **4** (1967), 141–149.
- [9] P.S. Landweber: *On equivariant maps between spheres with involutions*, *Ann. of Math.* **89** (1969), 125–137.
- [10] P. Löffler: *Equivariant framability of involutions on homotopy spheres*, *Manuscripta Math.* **23** (1978), 161–171.
- [11] H. Minami: *On Real J -homomorphisms*, *Osaka J. Math.* **16** (1979), 529–537.
- [12] D. Quillen: *The Adams conjecture*, *Topology* **10** (1971), 67–80.
- [13] G.B. Segal: *Equivariant K-theory*, *Publ. Math. I.H.E.S.* **34** (1968), 129–151.
- [14] G.B. Segal: *Equivariant stable homotopy theory*, *Actes Congrès intern. math.*, 1970, t.2, 59–63.

Department of Mathematics
Osaka City University
Sumiyoshi-ku, Osaka 558
Japan