

RESEARCH

Open Access

On fixed points of α - ψ -contractive multifunctions

J Hasanzade Asl¹, S Rezapour^{1,2} and N Shahzad^{3*}

*Correspondence:

nshahzad@kau.edu.sa

³Department of Mathematics, King Abdulaziz University, P.O. Box 80203, Jeddah, 21859, Saudi Arabia
Full list of author information is available at the end of the article

Abstract

Recently Samet, Vetro and Vetro introduced the notion of α - ψ -contractive type mappings and established some fixed point theorems in complete metric spaces. In this paper, we introduce the notion of α_* - ψ -contractive multifunctions and give a fixed point result for these multifunctions. We also obtain a fixed point result for self-maps in complete metric spaces satisfying a contractive condition.

Keywords: α_* - ψ -contractive multifunction; fixed point; partial metric

1 Introduction

Fixed point theory has many applications in different branches of science. During the last few decades, there has been a lot of activity in this area and several well-known fixed point theorems have been extended by a number of authors in different directions (see, for example, [1–38]). Recently Samet, Vetro and Vetro introduced the notion of α - ψ -contractive type mappings [33]. Denote with Ψ the family of nondecreasing functions $\psi : [0, \infty) \rightarrow [0, \infty)$ such that $\sum_{n=1}^{\infty} \psi^n(t) < \infty$ for all $t > 0$, where ψ^n is the n th iterate of ψ . It is known that $\psi(t) < t$ for all $t > 0$ and $\psi \in \Psi$ [33]. Let (X, d) be a metric space, T be a self-map on X , $\psi \in \Psi$ and $\alpha : X \times X \rightarrow [0, \infty)$ be a function. Then T is called an α - ψ -contraction mapping whenever $\alpha(x, y)d(Tx, Ty) \leq \psi(d(x, y))$ for all $x, y \in X$. Also, we say that T is α -admissible whenever $\alpha(x, y) \geq 1$ implies $\alpha(Tx, Ty) \geq 1$ [33]. Also, we say that α has the property (B) if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all $n \geq 1$ and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all $n \geq 1$. Let (X, d) be a complete metric space and T be an α -admissible α - ψ -contractive mapping on X . Suppose that there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$. If T is continuous or T has the property (B), then T has a fixed point (see [33]; Theorems 2.1 and 2.2). Finally, we say that X has the property (H) whenever for each $x, y \in X$ there exists $z \in X$ such that $\alpha(x, z) \geq 1$ and $\alpha(y, z) \geq 1$. If X has the property (H) in Theorems 2.1 and 2.2, then X has a unique fixed point ([33]; Theorem 2.3). It is considerable that the results of Samet *et al.* generalize similar ordered results in the literature (see the results of the third section in [33]). The aim of this paper is to introduce the notion of α_* - ψ -contractive multifunctions and give a fixed point result about the multifunctions. Let (X, d) be a metric space, $T : X \rightarrow 2^X$ be a closed-valued multifunction, $\psi \in \Psi$ and $\alpha : X \times X \rightarrow [0, \infty)$ be a function. In this case, we say that T is an α_* - ψ -contractive multifunction whenever $\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(d(x, y))$ for $x, y \in X$, where H is the Hausdorff generalized metric, $\alpha_*(A, B) = \inf\{\alpha(a, b) : a \in A, b \in B\}$ and 2^X denotes the family of all nonempty subsets of X . Also, we say that T is α_* -admissible whenever $\alpha(x, y) \geq 1$ implies $\alpha_*(Tx, Ty) \geq 1$.

Example 1.1 Let $X = [0, \infty)$, $d(x, y) = |x - y|$ and $\delta \in (0, 1)$ be a fixed number. Define $T : X \rightarrow 2^X$ by $Tx = [0, \delta x]$ for all $x \in X$ and $\alpha : X \times X \rightarrow [0, \infty)$ by $\alpha(x, y) = 1$ whenever $x, y \in [0, 1]$ and $\alpha(x, y) = 0$ whenever $x \notin [0, 1]$ or $y \notin [0, 1]$. Now, we show that T is α_* -admissible. If $\alpha(x, y) \geq 1$, then $x, y \in [0, 1]$ and so Tx and Ty are subsets of $[0, 1]$. Thus, $a, b \in [0, 1]$ for all $a \in Tx$ and $b \in Ty$. Hence, $\alpha(a, b) = 1$ for all $a \in Tx$ and $b \in Ty$. This implies that

$$\alpha_*(Tx, Ty) = \inf\{\alpha(a, b) : a \in Tx, b \in Ty\} = 1.$$

Therefore, T is α_* -admissible. Now, we show that T is an α_* - ψ -contractive multifunction, where $\psi(t) = \delta t$ for all $t \geq 0$. If $x \notin [0, \frac{1}{\delta}]$ or $y \notin [0, \frac{1}{\delta}]$, then an easy calculation shows us that $\alpha_*(Tx, Ty) = 0$. If $0 \leq x, y \leq \frac{1}{\delta}$, then $\alpha_*(Tx, Ty) = 1$. By using the definition of the Hausdorff metric, it is easy to see that $H(Tx, Ty) \leq \delta d(x, y)$ for $x, y \in [0, \frac{1}{\delta}]$. Thus, $\alpha_*(Tx, Ty)H(Tx, Ty) \leq \psi(d(x, y))$ for $x, y \in X$. Therefore, T is an α_* - ψ -contractive multifunction.

Let (X, \leq, d) be an ordered metric space and $A, B \subseteq X$. We say that $A \leq B$ whenever for each $a \in A$ there exists $b \in B$ such that $a \leq b$. Also, we say that $A \leq_r B$ whenever for each $a \in A$ and $b \in B$ we have $a \leq b$.

2 Main results

Now, we are ready to state and prove our main results. In the following result, we use the argument similar to that in the proof of Theorem 3.1 in [22].

Theorem 2.1 *Let (X, d) be a complete metric space, $\alpha : X \times X \rightarrow [0, \infty)$ be a function, $\psi \in \Psi$ be a strictly increasing map and T be a closed-valued, α_* -admissible and α_* - ψ -contractive multifunction on X . Suppose that there exist $x_0 \in X$ and $x_1 \in Tx_0$ such that $\alpha(x_0, x_1) \geq 1$. Assume that if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all n and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all n . Then T has a fixed point.*

Proof If $x_1 = x_0$, then we have nothing to prove. Let $x_1 \neq x_0$. If $x_1 \in Tx_1$, then x_1 is a fixed point of T . Let $x_1 \notin Tx_1$ and $q > 1$ be given. Then

$$0 < d(x_1, Tx_1) \leq \alpha_*(Tx_0, Tx_1)H(Tx_0, Tx_1) < q\alpha_*(Tx_0, Tx_1)H(Tx_0, Tx_1).$$

Hence, there exists $x_2 \in Tx_1$ such that

$$0 < d(x_1, x_2) < q\alpha_*(Tx_0, Tx_1)H(Tx_0, Tx_1) \leq q\psi(d(x_0, x_1)).$$

It is clear that $x_2 \neq x_1$ and $\alpha(x_1, x_2) \geq 1$. Thus, $\alpha_*(Tx_1, Tx_2) \geq 1$. Now, put $t_0 = d(x_0, x_1)$. Then, $t_0 > 0$ and $d(x_1, x_2) < q\psi(t_0)$. Since ψ is strictly increasing, $\psi(d(x_1, x_2)) < \psi(q\psi(t_0))$. Put $q_1 = \frac{\psi(q\psi(t_0))}{\psi(d(x_1, x_2))}$. Then $q_1 > 1$. If $x_2 \in Tx_2$, then x_2 is a fixed point of T . Assume that $x_2 \notin Tx_2$. Then

$$0 < d(x_2, Tx_2) \leq \alpha_*(Tx_1, Tx_2)H(Tx_1, Tx_2) < q_1\alpha_*(Tx_1, Tx_2)H(Tx_1, Tx_2).$$

Hence, there exists $x_3 \in Tx_2$ such that

$$0 < d(x_2, x_3) < q_1\alpha_*(Tx_1, Tx_2)H(Tx_1, Tx_2) \leq q_1\psi(d(x_1, x_2)) = \psi(q\psi(t_0)).$$

It is clear that $x_3 \neq x_2$, $\alpha(x_2, x_3) \geq 1$ and $\psi(d(x_2, x_3)) < \psi^2(q\psi(t_0))$. Now, put $q_2 = \frac{\psi^2(q\psi(t_0))}{\psi(d(x_2, x_3))}$. Then $q_2 > 1$. If $x_3 \in Tx_3$, then x_3 is a fixed point of T . Assume that $x_3 \notin Tx_3$. Then

$$0 < d(x_3, Tx_3) \leq \alpha_*(Tx_2, Tx_3)H(Tx_2, Tx_3) < q_2\alpha_*(Tx_2, Tx_3)H(Tx_2, Tx_3).$$

Thus, there exists $x_4 \in Tx_3$ such that

$$0 < d(x_3, x_4) < q_1\alpha_*(Tx_2, Tx_3)H(Tx_2, Tx_3) \leq q_2\psi(d(x_2, x_3)) = \psi^2(q\psi(t_0)).$$

By continuing this process, we obtain a sequence $\{x_n\}$ in X such that $x_n \in Tx_{n-1}$, $x_n \neq x_{n-1}$, $\alpha(x_n, x_{n+1}) \geq 1$ and $d(x_n, x_{n+1}) \leq \psi^{n-1}(q\psi(t_0))$ for all n . Now, for each $m > n$, we have

$$d(x_n, x_m) \leq \sum_{i=n}^{m-1} d(x_i, x_{i+1}) \leq \sum_{i=n}^{m-1} \psi^{i-1}(q\psi(t_0)).$$

Hence, $\{x_n\}$ is a Cauchy sequence in X . Choose $x^* \in X$ such that $x_n \rightarrow x^*$. Since $\alpha(x_n, x^*) \geq 1$ for all n and T is α_* -admissible, $\alpha_*(Tx_n, Tx^*) \geq 1$ for all n , thus

$$\begin{aligned} d(x^*, Tx^*) &\leq H(Tx^*, Tx_n) + d(x_{n+1}, x^*) \leq \alpha_*(Tx_n, Tx^*)H(Tx_n, Tx^*) + d(x_{n+1}, x^*) \\ &\leq \psi(d(x_n, x^*)) + d(x_{n+1}, x^*) \end{aligned}$$

for all n . Therefore, $d(x^*, Tx^*) = 0$ and so $x^* \in Tx^*$. □

Example 2.1 Let $X = [0, \infty)$ and $d(x, y) = |x - y|$. Define $T : X \rightarrow 2^X$ by $Tx = [2x - \frac{3}{2}, \infty)$ for all $x > 1$, $Tx = [0, \frac{x}{2}]$ for all $0 \leq x \leq 1$ and $\alpha : X \times X \rightarrow [0, \infty)$ by $\alpha(x, y) = 1$ whenever $x, y \in [0, 1]$ and $\alpha(x, y) = 0$ whenever $x \notin [0, 1]$ or $y \notin [0, 1]$. Then it is easy to check that T is an α_* -admissible and α_* - ψ -contractive multifunction, where $\psi(t) = \frac{t}{2}$ for all $t \geq 0$. Put $x_0 = 1$ and $x_1 = \frac{1}{2}$. Then $\alpha(x_0, x_1) \geq 1$. Also, if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all n and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all n . Note that T has infinitely many fixed points.

Corollary 2.2 Let (X, \leq, d) be a complete ordered metric space, $\psi \in \Psi$ be a strictly increasing map and T be a closed-valued multifunction on X such that

$$H(Tx, Ty) \leq \psi(d(x, y))$$

for all $x, y \in X$ with $x \leq y$. Suppose that there exists $x_0 \in X$ and $x_1 \in Tx_0$ such that $x_0 \leq x_1$. Assume that if $\{x_n\}$ is a sequence in X such that $x_n \leq x_{n+1}$ for all n and $x_n \rightarrow x$, then $x_n \leq x$ for all n . If $x \leq y$ implies $Tx \leq_r Ty$, then T has a fixed point.

Proof Define $\alpha : X \times X \rightarrow [0, \infty)$ by $\alpha(x, y) = 1$ whenever $x \leq y$ and $\alpha(x, y) = 0$ whenever $x \not\leq y$. Since $x \leq y$ implies $Tx \leq_r Ty$, $\alpha(x, y) = 1$ implies $\alpha_*(Tx, Ty) = 1$. Thus, it is easy to check that T is an α_* -admissible and α_* - ψ -contractive multifunction on X . Now, by using Theorem 2.1, T has a fixed point. □

Now, we prove the following result for self-maps.

Theorem 2.3 Let (X, d) be a complete metric space, $\alpha : X \times X \rightarrow [0, \infty)$ be a function, $\psi \in \Psi$ and T be a self-map on X such that $\alpha(x, y)d(Tx, Ty) \leq \psi(m(x, y))$ for all $x, y \in X$, where $m(x, y) = \max\{d(x, y), d(x, Tx), d(y, Ty), \frac{1}{2}[d(x, Ty) + d(y, Tx)]\}$. Suppose that T is α -admissible and there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$. Assume that if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all n and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all n . Then T has a fixed point.

Proof Take $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$ and define the sequence $\{x_n\}$ in X by $x_{n+1} = Tx_n$ for all $n \geq 0$. If $x_n = x_{n+1}$ for some n , then $x^* = x_n$ is a fixed point of T . Assume that $x_n \neq x_{n+1}$ for all n . Since T is α -admissible, it is easy to check that $\alpha(x_n, x_{n+1}) \geq 1$ for all natural numbers n . Thus, for each natural number n , we have

$$\begin{aligned} d(x_n, x_{n+1}) &= d(Tx_{n-1}, Tx_n) \leq \alpha(x_{n-1}, x_n)d(Tx_{n-1}, Tx_n) \\ &\leq \psi \left(\max \left\{ d(x_n, x_{n-1}), d(x_n, x_{n+1}), d(x_{n-1}, x_n), \frac{1}{2}[d(x_n, x_n) + d(x_{n-1}, x_{n+1})] \right\} \right) \\ &\leq \psi \left(\max \left\{ d(x_n, x_{n-1}), d(x_n, x_{n+1}), \frac{1}{2}[d(x_n, x_{n-1}) + d(x_n, x_{n+1})] \right\} \right) \\ &= \psi(\max\{d(x_n, x_{n-1}), d(x_n, x_{n+1})\}). \end{aligned}$$

If $\max\{d(x_n, x_{n-1}), d(x_n, x_{n+1})\} = d(x_n, x_{n+1})$, then

$$d(x_{n+1}, x_n) \leq \psi(d(x_n, x_{n+1})) < d(x_{n+1}, x_n)$$

which is contradiction. Thus, $\max\{d(x_n, x_{n-1}), d(x_n, x_{n+1})\} = d(x_n, x_{n-1})$ for all n . Hence, $d(x_{n+1}, x_n) \leq \psi(d(x_n, x_{n-1}))$ and so $d(x_{n+1}, x_n) \leq \psi^n(d(x_1, x_0))$ for all n . It is easy to check that $\{x_n\}$ is a Cauchy sequence. Thus, there exists $x^* \in X$ such that $x_n \rightarrow x^*$. By using the assumption, we have $\alpha(x_n, x^*) \geq 1$ for all n . Thus,

$$\begin{aligned} d(Tx^*, x^*) &\leq d(Tx_n, Tx_n) + d(x_{n+1}, x^*) \leq \alpha(x_n, x^*)d(Tx_n, Tx_n) + d(x_{n+1}, x^*) \\ &\leq \psi \left(\max \left\{ d(x_n, x^*), d(x_n, x_{n+1}), d(x^*, Tx_n), \right. \right. \\ &\quad \left. \left. \frac{1}{2}[d(x_n, Tx_n) + d(x^*, x_{n+1})] \right\} \right) + d(x_{n+1}, x^*) \\ &\leq \psi(d(x^*, Tx_n)) + d(x_{n+1}, x^*) \end{aligned}$$

for sufficiently large n . Hence, $d(Tx^*, x^*) = 0$ and so $Tx^* = x^*$. □

Example 2.2 Let $X = [0, \infty)$ and $d(x, y) = |x - y|$. Define the self-map T on X by $Tx = 2x - \frac{5}{3}$ for $x > 1$, $Tx = \frac{x}{3}$ for $0 \leq x \leq 1$ and $\alpha : X \times X \rightarrow [0, \infty)$ by $\alpha(x, y) = 1$ whenever $x, y \in [0, 1]$ and $\alpha(x, y) = 0$ whenever $x \notin [0, 1]$ or $y \notin [0, 1]$. Then it is easy to check that T is α -admissible and $\alpha(x, y)d(Tx, Ty) \leq \psi(m(x, y))$ for all $x, y \in X$, where $\psi(t) = \frac{t}{3}$ for all $t \geq 0$. Also, $\alpha(1, T1) = 1$ and if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all n and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all n . Note that, T has two fixed points.

Corollary 2.4 Let (X, \leq, d) be a complete ordered metric space, $\psi \in \Psi$ and T be a self-map on X such that $d(Tx, Ty) \leq \psi(m(x, y))$ for all $x, y \in X$ with $x \leq y$. Suppose that there

exists $x_0 \in X$ such that $x_0 \leq Tx_0$. If $\{x_n\}$ is a sequence in X such that $x_n \leq x_{n+1}$ for all n and $x_n \rightarrow x$, then $x_n \leq x$ for all n . If $x \leq y$ implies $Tx \leq Ty$, then T has a fixed point.

If we substitute a partial metric ρ for the metric d in Theorem 2.3, it is easy to check that a similar result holds for the partial metric case as follows.

Theorem 2.5 *Let (X, ρ) be a complete partial metric space, $\alpha : X \times X \rightarrow [0, \infty)$ be a function, $\psi \in \Psi$ and T be a self-map on X such that $\alpha(x, y)\rho(Tx, Ty) \leq \psi(m(x, y))$ for all $x, y \in X$, where $m(x, y) = \max\{\rho(x, y), \rho(x, Tx), \rho(y, Ty), \frac{1}{2}[\rho(x, Ty) + \rho(y, Tx)]\}$. Suppose that T is α -admissible and there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$. Assume that if $\{x_n\}$ is a sequence in X such that $\alpha(x_n, x_{n+1}) \geq 1$ for all n and $x_n \rightarrow x$, then $\alpha(x_n, x) \geq 1$ for all n . Then T has a fixed point.*

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Science and Research Branch, Islamic Azad University, Tehran, Iran. ²Department of Mathematics, Azarbaijan Shahid Madani University, Tabriz, Azarshahr, Iran. ³Department of Mathematics, King Abdulaziz University, P.O. Box 80203, Jeddah, 21859, Saudi Arabia.

Acknowledgements

The authors are grateful to the reviewers for their useful comments.

Received: 5 May 2012 Accepted: 26 October 2012 Published: 26 November 2012

References

1. Abbas, M, Nazir, T, Radenovic, S: Common fixed points of four maps in partially ordered metric spaces. *Appl. Math. Lett.* **24**, 1520-1526 (2011)
2. Aleomraninejad, SMA, Rezapour, S, Shahzad, N: On generalizations of the Suzuki's method. *Appl. Math. Lett.* **24**, 1037-1040 (2011)
3. Aleomraninejad, SMA, Rezapour, S, Shahzad, N: Fixed points of hemi-convex multifunctions. *Topol. Methods Nonlinear Anal.* **37**(2), 383-389 (2011)
4. Aleomraninejad, SMA, Rezapour, S, Shahzad, N: Some fixed point results on a metric space with a graph. *Topol. Appl.* **159**, 659-663 (2012)
5. Alghamdi, MA, Alnafe'i, SH, Radenovic, S, Shahzad, N: Fixed point theorems for convex contraction mappings on cone metric spaces. *Math. Comput. Model.* **54**, 2020-2026 (2011)
6. Altun, I, Damjanovic, B, Djoric, D: Fixed point and common fixed point theorems on ordered cone metric spaces. *Appl. Math. Lett.* **23**, 310-316 (2010)
7. Aydi, H, Damjanovic, B, Samet, B, Shatanawi, W: Coupled fixed point theorems for nonlinear contractions in partially ordered G -metric spaces. *Math. Comput. Model.* **54**, 2443-2450 (2011)
8. Aydi, H, Nashine, HK, Samet, B, Yazidi, H: Coincidence and common fixed point results in partially ordered cone metric spaces and applications to integral equations. *Nonlinear Anal.* **74**, 6814-6825 (2011)
9. Berinde, V: Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces. *Nonlinear Anal.* **74**, 7347-7355 (2011)
10. Berinde, V, Borcut, M: Tripled fixed point theorems for contractive type mappings in partially ordered metric spaces. *Nonlinear Anal.* **74**, 4889-4897 (2011)
11. Berinde, V, Vetro, F: Common fixed points of mappings satisfying implicit contractive conditions. *Fixed Point Theory Appl.* **2012**, 105 (2012)
12. Borcut, M, Berinde, V: Tripled coincidence theorems for contractive type mappings in partially ordered metric spaces. *Appl. Math. Comput.* (2012). doi:10.1016/j.amc.2011.11.049
13. Derafshpour, M, Rezapour, S, Shahzad, N: Best proximity points of cyclic φ -contractions in ordered metric spaces. *Topol. Methods Nonlinear Anal.* **37**, 193-202 (2011)
14. Di Bari, C, Vetro, P: Fixed points for weak φ -contractions on partial metric spaces. *Int. J. Eng. Contemp. Math. Sci.* **1**, 4-9 (2011)
15. Di Bari, C, Vetro, P: φ -pairs and common fixed points in cone metric spaces. *Rend. Circ. Mat. Palermo* **57**, 279-285 (2008). doi:10.1007/s12215-008-0020-9
16. Di Bari, C, Vetro, P: Weakly φ -pairs and common fixed points in cone metric spaces. *Rend. Circ. Mat. Palermo* **58**, 125-132 (2009). doi:10.1007/s12215-009-0012-4
17. Ding, HS, Lu, L: Coupled fixed point theorems in partially ordered cone metric spaces. *Filomat* **25**(2), 137-149 (2011)

18. Du, W-S: Coupled fixed point theorems for nonlinear contractions satisfied Mizoguchi-Takahashi's condition in quasi-ordered metric spaces. *Fixed Point Theory Appl.* **2010**, Article ID 876372 (2010)
19. Haghi, RH, Rezapour, S, Shahzad, N: Some fixed point generalizations are not real generalizations. *Nonlinear Anal.* **74**, 1799-1803 (2011)
20. Hu, X-Q, Ma, X-Y: Coupled coincidence point theorems under contractive conditions in partially ordered probabilistic metric spaces. *Nonlinear Anal.* **74**, 6451-6458 (2011)
21. Jachymski, J: Equivalent conditions for generalized contractions on (ordered) metric spaces. *Nonlinear Anal.* **74**, 768-774 (2011)
22. Lazar, VL: Fixed point theory for multivalued ϕ -contractions. *Fixed Point Theory Appl.* **2011**, 50 (2011)
23. Luong, NV, Thuan, NX: Coupled fixed points in partially ordered metric spaces and application. *Nonlinear Anal.* **74**, 983-992 (2011)
24. Nashine, HK, Samet, B, Vetro, C: Monotone generalized nonlinear contractions and fixed point theorems in ordered metric spaces. *Math. Comput. Model.* **54**, 712-720 (2011)
25. Nashine, HK, Samet, B: Fixed point results for mappings satisfying (ψ, φ) -weakly contractive condition in partially ordered metric spaces. *Nonlinear Anal.* **74**, 2201-2209 (2011)
26. Nashine, HK, Shatanawi, W: Coupled common fixed point theorems for a pair of commuting mappings in partially ordered complete metric spaces. *Comput. Math. Appl.* **62**, 1984-1993 (2011)
27. Rezapour, S, Amiri, P: Some fixed point results for multivalued operators in generalized metric spaces. *Comput. Math. Appl.* **61**, 2661-2666 (2011)
28. Rezapour, S, Haghi, RH: Some notes on the paper 'Cone metric spaces and fixed point theorems of contractive mappings'. *J. Math. Anal. Appl.* **345**, 719-724 (2008)
29. Rezapour, S, Haghi, RH, Rhoades, BE: Some results about T -stability and almost T -stability. *Fixed Point Theory* **12**(1), 179-186 (2011)
30. Rezapour, S, Haghi, RH, Shahzad, N: Some notes on fixed points of quasi-contraction maps. *Appl. Math. Lett.* **23**, 498-502 (2010)
31. Rus, MD: Fixed point theorems for generalized contractions in partially ordered metric spaces with semi-monotone metric. *Nonlinear Anal.* **74**, 1804-1813 (2011)
32. Samet, B, Vetro, C: Coupled fixed point theorems for multi-valued nonlinear contraction mappings in partially ordered metric spaces. *Nonlinear Anal.* **74**, 4260-4268 (2011)
33. Samet, B, Vetro, C, Vetro, P: Fixed point theorems for α - ψ -contractive type mappings. *Nonlinear Anal.* **75**, 2154-2165 (2012)
34. Shatanawi, W, Samet, B: On (ψ, φ) -weakly contractive condition in partially ordered metric spaces. *Comput. Math. Appl.* **62**, 3204-3214 (2011)
35. Vetro, F: On approximating curves associated with nonexpansive mappings. *Carpath. J. Math.* **27**, 142-147 (2011)
36. Vetro, P: Common fixed points in cone metric spaces. *Rend. Circ. Mat. Palermo* **56**, 464-468 (2007). doi:10.1007/BF03032097
37. Zhang, X: Fixed point theorems of multivalued monotone mappings in ordered metric spaces. *Appl. Math. Lett.* **23**, 235-240 (2010)
38. Zhilong, L: Fixed point theorems in partially ordered complete metric spaces. *Math. Comput. Model.* **54**, 69-72 (2011)

doi:10.1186/1687-1812-2012-212

Cite this article as: Asl et al.: On fixed points of α - ψ -contractive multifunctions. *Fixed Point Theory and Applications* 2012 **2012**:212.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
