

ON GO -COMPACT SPACES

M. Caldas, S. Jafari, S. P. Moshokoa and T. Noiri

Abstract

The purpose of this paper is to offer some more properties of GO -compact spaces and to introduce and investigate some properties of g -continuous multifunctions. We also investigate GO -compact spaces in the context of multifunctions.

1 Introduction and preliminaries

Levine [11] introduced the concept of generalized closed sets of a topological space. Since the advent of these notions, several research papers with interesting results in different respects came to existence (see, [1], [3], [4], [5], [6], [8], [9], [10], [12]). Recently Caldas and Jafari [4] introduced and investigated the concepts of g - US spaces, g -convergency, sequential GO -compactness, sequential g -continuity and sequential g -sub-continuity.

Throughout the present paper (X, τ) and (Y, σ) (or simply X and Y) denote topological spaces. Let A be a subset of X . We denote the interior and the closure of a set A by $Int(A)$ and $Cl(A)$, respectively. $A \subset X$ is called a generalized closed set (briefly g -closed set) of X [11] if $Cl(A) \subset G$ holds whenever $A \subset G$ and G is open in X . The union of two g -closed sets is a g -closed set. A subset A of X is called a g -open set of X , if its complement A^c is g -closed in X . The intersection of all g -closed sets containing a set A is called the g -closure of A [10] and is denoted by $gCl(A)$. If $A \subset X$, then $A \subset gCl(A) \subset Cl(A)$. The collection of all g -closed (resp. g -open) subsets of X will be denoted by $GC(X)$ (resp. $GO(X)$). We set $GC(X, x) = \{V \in GC(X) / x \in V\}$ for $x \in X$. We define similarly $GO(X, x)$. Let p be a point of X and N be a subset of X . N is called a g -neighborhood of p in X [3] if there exists a g -open set O of X such that $p \in O \subset N$.

A space X is GO -compact if every g -open cover of X has a finite subcover. Since every open sets is a g -open set, it follows that every GO -compact space is compact.

2000 *Mathematics Subject Classifications*. 54B05, 54C08; Secondary: 54D05.

Key words and Phrases. Ag -open sets, g -closed sets, g -complete accumulation point, GO -compact spaces, multifunctions.

Received: June 27, 2007

Communicated by Dragan S. Djordjević

But, the converse may be false. Let $X = \{x\} \cup \{x_i : i \in I\}$ where the indexed set I is uncountable. Let $\tau = \{\emptyset, \{x\}, X\}$ be the topology on X . Evidently, X is a compact space. However, it is not a GO -compact space because $\{\{x, x_i\} : i \in I\}$ is a g -open covering of X but it has no finite subcover. A subset A of a space X is said to be GO -compact if A is GO -compact as a subspace of X . If the product space of two non-empty spaces is GO -compact, then each factor space is GO -compact [1]. If A is g -open in X and B is g -open in Y , then $A \times B$ is g -open in $X \times Y$ [11]. A function $f : X \rightarrow Y$ is said to be g -continuous [1] if the inverse image of every closed set in Y is g -closed in X .

It is the purpose of this paper to offer some more characterizations of GO -compact spaces. We also introduce the notion of g -complete accumulation points by which we give some characterizations of GO -compact spaces. By introducing the notion of 1-lower (resp. 1-upper) g -continuous functions and considering the known notion of 1-lower (resp. 1-upper) compatible partial orders we investigate some more properties of GO -compactness. We also investigate GO -compact spaces in the context of multifunctions by introducing 1-lower (resp. 1-upper) g -continuous multifunctions. Lastly we also obtain some characterizations of GO -compact spaces by using lower (resp. upper) g -continuous multifunctions. In this paper we are working in ZFC.

Recall that a function $f : X \rightarrow Y$ is said to be g -continuous [3] if the inverse image of each open set in Y is g -open in X .

Let Λ be a directed set. Now we introduce the following notions which will be used in this paper. A net $\xi = \{x_\alpha \mid \alpha \in \Lambda\}$ g -accumulates at a point $x \in X$ if the net is frequently in every $U \in GO(X, x)$, i.e. for each $U \in GO(X, x)$ and for each $\alpha_0 \in \Lambda$, there is some $\alpha \geq \alpha_0$ such that $x_\alpha \in U$. The net ξ g -converges to a point x of X if it is eventually in every $U \in GO(X, x)$. We say that a filterbase $\Theta = \{F_\alpha \mid \alpha \in \Gamma\}$ g -accumulates at a point $x \in X$ if $x \in \bigcap_{\alpha \in \Gamma} gCl(F_\alpha)$. Given a set S with $S \subset X$, a g -cover of S is a family of g -open subsets U_α of X for each $\alpha \in I$ such that $S \subset \bigcup_{\alpha \in I} U_\alpha$. A filterbase $\Theta = \{F_\alpha \mid \alpha \in \Gamma\}$ g -converges to a point x in X if for each $U \in GO(X, x)$, there exists an F_α in Θ such that $F_\alpha \subset U$.

Recall that a multifunction (also called multivalued function [2]) F on a set X into a set Y , denoted by $F : X \rightarrow Y$, is a relation on X into Y , i.e. $F \subset X \times Y$. Let $F : X \rightarrow Y$ be a multifunction. The upper and lower inverse of a set V of Y are denoted by $F^+(V)$ and $F^-(V)$:

$$F^+(V) = \{x \in X \mid F(x) \subset V\} \text{ and } F^-(V) = \{x \in X \mid F(x) \cap V \neq \emptyset\}.$$

2 GO -compact spaces

We begin with the following notions:

Definition 1 A point x in a space X is said to be a g -complete accumulation point of a subset S of X if $Card(S \cap U) = Card(S)$ for each $U \in GO(X, x)$, where

$Card(S)$ denotes the cardinality of S .

Example 2.1 Let $X = \{a, b, c, d\}$ with topology $\tau = \{X, \emptyset, \{a\}, \{a, b, c\}\}$. Observe that both b and c are g -complete accumulation points of $\{a, b\}$. Notice that a is not a g -complete accumulation point of $\{a, b\}$

Definition 2 In a topological space X , a point x is said to be a g -adherent point of a filterbase Θ on X if it lies in the g -closure of all sets of Θ .

Observe that the Frechet filter does not satisfy Definition 2. But take a topological space X such that $A \subset X$. Then any point of the g -closure of A is a g -adherent point of $\Omega = \{U \subset X \mid A \subset U\}$.

Theorem 2.2 A space X is GO -compact if and only if each infinite subset of X has a g -complete accumulation point.

Proof. Let the space X be GO -compact and S an infinite subset of X . Let K be the set of points x in X which are not g -complete accumulation points of S . Now it is obvious that for each point x in K , we are able to find $U(x) \in GO(X, x)$ such that $Card(S \cap U(x)) \neq Card(S)$. If K is the whole space X , then $\Theta = \{U(x) \mid x \in X\}$ is a g -cover of X . By the hypothesis X is GO -compact, so there exists a finite subcover $\Psi = \{U(x_i)\}$, where $i = 1, 2, \dots, n$ such that $S \subset \bigcup\{U(x_i) \cap S \mid i = 1, 2, \dots, n\}$. Then $Card(S) = \max\{Card(U(x_i) \cap S) \mid i = 1, 2, \dots, n\}$ which does not agree with what we assumed. This implies that S has a g -complete accumulation point. Now assume that X is not GO -compact and that every infinite subset $S \subset X$ has a g -complete accumulation point in X . It follows that there exists a g -cover Ξ with no finite subcover. Set $\delta = \min\{Card(\Phi) \mid \Phi \subset \Xi, \text{ where } \Phi \text{ is a } g\text{-cover of } X\}$. Fix $\Psi \subset \Xi$ for which $Card(\Psi) = \delta$ and $\bigcup\{U \mid U \in \Psi\} = X$. Let N denote the set of natural numbers. Then by hypothesis $\delta \geq Card(N)$. By well-ordering of Ψ by some minimal well-ordering " \sim ", suppose that U is any member of Ψ . By minimal well-ordering " \sim " we have $Card(\{V \mid V \in \Psi, V \sim U\}) < Card(\{V \mid V \in \Psi\})$. Since Ψ can not have any subcover with cardinality less than δ , then for each $U \in \Psi$ we have $X \neq \bigcup\{V \mid V \in \Psi, V \sim U\}$. For each $U \in \Psi$, choose a point $x(U) \in X \setminus \bigcup\{V \cup \{x(V)\} \mid V \in \Psi, V \sim U\}$. We are always able to do this if not one can choose a cover of smaller cardinality from Ψ . If $H = \{x(U) \mid U \in \Psi\}$, then to finish the proof we will show that H has no g -complete accumulation point in X . Suppose that z is a point of the space X . Since Ψ is a g -cover of X then z is a point of some set W in Ψ . By the fact that $U \sim W$, we have $x(U) \in W$. It follows that $T = \{U \mid U \in \Psi \text{ and } x(U) \in W\} \subset \{V \mid V \in \Psi, V \sim W\}$. But $Card(T) < \delta$. Therefore $Card(H \cap W) < \delta$. But $Card(H) = \delta \geq Card(N)$ since for two distinct points U and W in Ψ , we have $x(U) \neq x(W)$. This means that H has no g -complete accumulation point in X which contradicts our assumptions. Therefore X is GO -compact.

Theorem 2.3 For a space X the following statements are equivalent:

- (1) X is GO -compact;
- (2) Every net in X with a well-ordered directed set as its domain g -accumulates to some point of X .

Proof. (1) \Rightarrow (2): Suppose that X is GO -compact and $\xi = \{x_\alpha \mid \alpha \in \Lambda\}$ a net with a well-ordered directed set Λ as domain. Assume that ξ has no g -adherent point in X . Then for each point x in X , there exist $V(x) \in GO(X, x)$ and an $\alpha(x) \in \Lambda$ such that $V(x) \cap \{x_\alpha \mid \alpha \geq \alpha(x)\} = \emptyset$. This implies that $\{x_\alpha \mid \alpha \geq \alpha(x)\}$ is a subset of $X \setminus V(x)$. Then the collection $C = \{V(x) \mid x \in X\}$ is a g -cover of X . By hypothesis of the theorem, X is GO -compact and so C has a finite subfamily $\{V(x_i)\}$, where $i = 1, 2, \dots, n$ such that $X = \bigcup\{V(x_i)\}$. Suppose that the corresponding elements of Λ be $\{\alpha(x_i)\}$, where $i = 1, 2, \dots, n$. Since Λ is well-ordered and $\{\alpha(x_i)\}$, where $i = 1, 2, \dots, n$ is finite, the largest element of $\{\alpha(x_i)\}$ exists. Suppose it is $\{\alpha(x_l)\}$. Then for $\gamma \geq \{\alpha(x_l)\}$, we have $\{x_\delta \mid \delta \geq \gamma\} \subset \bigcap_{i=1}^n (X \setminus V(x_i)) = X \setminus \bigcup_{i=1}^n V(x_i) = \emptyset$ which is impossible. This shows that ξ has at least one g -adherent point in X .

(2) \Rightarrow (1): Now it is enough to prove that each infinite subset has a g -complete accumulation point by utilizing Theorem 3.1. Suppose that $S \subset X$ is an infinite subset of X . According to Zorn's Lemma, the infinite set S can be well-ordered. This means that we can assume S to be a net with a domain which is a well-ordered index set. It follows that S has a g -adherent point z . Therefore z is a g -complete accumulation point of S . This shows that X is GO -compact.

Theorem 2.4 *A space X is GO -compact if and only if each family of g -closed subsets of X with the finite intersection property has a nonempty intersection.*

Proof. Straightforward.

Question 2.5 *Which condition or conditions should be imposed on a topological space X such that the following statements are equivalent:*

- (1) X is GO -compact;
- (2) Each filterbase on X with at most one g -adherent point is g -convergent.

3 GO -compactness and 1-lower and 1-upper g -continuous functions

In this section we further investigate properties of GO -compactness by 1-lower and 1-upper g -continuous functions. We begin with the following notions and in what follows R denotes the set of real numbers.

Definition 3 *A function $f : X \rightarrow R$ is said to be 1-lower (resp. 1-upper) g -continuous at the point y in X if for each $\lambda > 0$, there exists a g -open set $U(y) \in GO(X, y)$ such that $f(x) > f(y) - \lambda$ (resp. $f(x) > f(y) + \lambda$) for every point x in $U(y)$. The function f is 1-lower (resp. 1-upper) g -continuous in X if it has these properties for every point x of X .*

Example 3.1 *Take $f : (R, \tau_u) \rightarrow (R, \tau)$, where τ_u is the usual topology and τ is the family of sets $\tau = \{(\eta, \infty) \mid \eta \in R\} \cup R$. Such a function is 1-lower g -continuous. Now take $f : (R, \tau_u) \rightarrow (R, \sigma)$, where τ_u is the usual topology and σ is the family of sets $\sigma = \{(-\infty, \eta) \mid \eta \in R\} \cup R$. Such a function is 1-upper g -continuous.*

Theorem 3.2 *A function $f : X \rightarrow R$ is 1-lower g -continuous if and only if for each $\eta \in R$, the set of all x such that $f(x) \leq \eta$ is g -closed.*

Proof. It is obvious that the family of sets $\tau = \{(\eta, \infty) \mid \eta \in R\} \cup R$ establishes a topology on R . Then the function f is 1-lower g -continuous if and only if $f : X \rightarrow (R, \tau)$ is g -continuous. The interval $(-\infty, \eta]$ is closed in (R, τ) . It follows that $f^{-1}((-\infty, \eta])$ is g -closed. Therefore the set of all x such that $f(x) \leq \eta$ is equal to $f^{-1}((-\infty, \eta])$ and thus is g -closed.

Corollary 3.3 *A subset S of X is GO-compact if and only if the characteristic function X_S is 1-lower g -continuous.*

Theorem 3.4 *A function $f : X \rightarrow R$ is 1-upper g -continuous if and only if for each $\eta \in R$, the set of all x such that $f(x) \geq \eta$ is g -closed.*

Corollary 3.5 *A subset S of X is GO-compact if and only if the characteristic function X_S is 1-upper g -continuous.*

Question 3.6 *Is it true that if the function $G(x) = \inf_{i \in I} f_i(x)$ exists, where f_i , are 1-upper g -continuous functions from X into R , then $G(x)$ is 1-upper g -continuous?*

Theorem 3.7 *Let $f : X \rightarrow R$ be a 1-lower g -continuous function, where X is GO-compact. Then f assumes the value $m = \inf_{x \in X} f(x)$.*

Proof. Suppose $\eta > m$. Since f is 1-lower g -continuous, then the set $K(\eta) = \{x \in X \mid f(x) \leq \eta\}$ is a non-empty g -closed set in X by infimum property. Hence the family $\{K(\eta) \mid \eta > m\}$ is a collection of non-empty g -closed sets with finite intersection property in X . By Theorem 2.4 this family has non-empty intersection. Suppose $z \in \bigcap_{\eta > m} K(\eta)$. Therefore $f(z) = m$ as we wished to prove.

4 GO-compactness and g -continuous multifunctions

In this section, we give some characterizations of GO-compact spaces by using lower (resp. upper) g -continuous multifunctions.

Definition 4 *A multifunction $F : X \rightarrow Y$ is said to be lower (resp. upper) g -continuous if $X \setminus F^-(S)$ (resp. $F^-(S)$) is g -closed in X for each open (resp. closed) set S in Y .*

For the following two lemmas we shall assume that if $gCl(A) = A$, then A is g -closed”.

Lemma 4.1 *For a multifunction $F : X \rightarrow Y$, the following statements are equivalent:*

(1) F is lower g -continuous;

- (2) If $x \in F^-(U)$ for a point x in X and an open set $U \subset Y$, then $V \subset F^-(U)$ for some $V \in GO(X)$;
- (3) If $x \notin F^+(D)$ for a point x in X and a closed set $D \subset Y$, then $F^+(D) \subset K$ for some g -closed set K with $x \notin K$;
- (4) $F^-(U) \in GO(X)$ for each open set $U \subset Y$.

Proof. (1) \Rightarrow (4): Let U be any open set in Y . By (1), $X - F^-(U)$ is g -closed in X and hence $F^-(U) \in GO(X)$

(4) \Rightarrow (2): Let U be any open set of Y and $x \in F^-(U)$. By (4), $F^-(U) \in GO(X)$. Put $V = F^-(U)$. Then $V \in GO(X)$ and $V \subset F^-(U)$.

(2) \Rightarrow (3): Let D be closed in Y and $x \notin F^+(D)$. Then $Y - D$ is open in Y and $x \in X - F^+(D) = F^-(X - D)$. Therefore, There exists $V \in GO(x)$ such that $V \subset F^-(U)$. Now, put $K = X - V$, then $x \notin K$, K is g -closed and $K = X - V \supset X - F^-(Y - D) = F^+(D)$.

(3) \Rightarrow (1): We show that $F^+(H)$ is g -closed for any closed set H of Y . Let H be any closed set and $x \notin F^+(H)$. By (3) there exists a g -closed set K such that $x \notin K$ and $F^+(H) \subset K$; hence $F^+(H) \subset gCl(F^+(H)) \subset K$. Since $x \notin K$, we have $x \notin gCl(F^+(H))$. This implies that $gCl(F^+(H)) \subset F^+(H)$. In general, we have $F^+(H) \subset gCl(F^+(H))$ and hence $F^+(H) = gCl(F^+(H))$. Hence $F^+(H)$ is g -closed for any closed set H of Y .

Lemma 4.2 For a multifunction $F : X \rightarrow Y$, the following statements are equivalent:

- (1) F is upper g -continuous;
- (2) If $x \in F^+(V)$ for a point x in X and an open set $V \subset Y$, then $F(U) \subset V$ for some $U \in GO(X)$;
- (3) If $x \notin F^-(D)$ for a point x in X and a closed set $D \subset Y$, then $F^-(D) \subset K$ for some g -closed set K with $x \notin K$;
- (4) $F^+(U) \in GO(X)$ for each open set $U \subset Y$.

Proof. (1) \Rightarrow (4): Let U be any open set in Y . Then $Y - U$ is closed. By (1), $F^-(Y - U) = X - F^+(U)$ is g -closed in X and hence $F^+(U) \in GO(X)$.

(4) \Rightarrow (2): Let V be any open set of Y and $x \in F^+(V)$. By (4), $F^+(V) \in GO(X)$. Put $U = F^+(V)$. Then $U \in GO(X)$ and $F(U) \subset V$.

(2) \Rightarrow (3): Let D be closed in Y and $x \notin F^-(D)$. Then $Y - D$ is open and $x \in X - F^-(D) = F^+(Y - D)$. By (2), there exists $U \in GO(X)$ such that $F(U) \subset Y - D$. Now, put $K = X - U$, then $x \notin K$, K is g -closed and $K = X - U \supset X - F^+(Y - D) = F^-(D)$.

(3) \Rightarrow (1): We show that $F^-(H)$ is g -closed for any closed set H of Y . Let H be any closed set and $x \notin F^-(H)$. By (3), there exists a g -closed set K such that $x \notin K$ and $F^-(H) \subset K$; hence $F^-(H) \subset gCl(F^-(H)) \subset K$. Since $x \notin K$, we have $x \notin gCl(F^-(H))$. This implies that $gCl(F^-(H)) \subset F^-(H)$. In general, we have $F^-(H) \subset gCl(F^-(H))$ and hence $F^-(H) = gCl(F^-(H))$. Hence $F^-(H)$ is g -closed for any closed set H of Y .

Theorem 4.3 The following two statements are equivalent for a space X :

- (1) X is GO -compact.

(2) Every lower g -continuous multifunction from X into the closed sets of a space assumes a minimal value with respect to set inclusion relation.

Proof. (1) \Rightarrow (2): Suppose that F is a lower g -continuous multifunction from X into the closed subsets of a space Y . We denote the poset of all closed subsets of Y with the set inclusion relation " \subseteq " by Λ . Now we show that $F : X \rightarrow \Lambda$ is a lower g -continuous function. We will show that $N = F^{-}(\{S \subset Y \mid S \in \Lambda \text{ and } S \subseteq C\})$ is g -closed in X for each closed set C of Y . Let $z \notin N$, then $F(z) \neq S$ for every closed set S of Y . It is obvious that $z \in F^{-}(Y \setminus C)$, where $Y \setminus C$ is open in Y . By Lemma 4.1 (2), we have $W \subset F^{-}(Y \setminus C)$ for some $W \in GO(z)$. Hence $F(w) \cap (Y \setminus C) \neq \emptyset$ for each w in W . So for each w in W , $F(w) \setminus C \neq \emptyset$. Consequently, $F(w) \setminus S \neq \emptyset$ for every closed subset S of Y for which $S \subseteq C$. We consider that $W \cap N = \emptyset$. This means that N is g -closed. Thus we observe that F assumes a minimal value.

(2) \Rightarrow (1): Suppose that X is not GO -compact. It follows that we have a net $\{x_i \mid i \in \Lambda\}$, where Λ is a well-ordered set with no g -accumulation point by ([8], Theorem 3.2). We give Λ the order topology. Let $M_j = gCl\{x_i \mid i \geq j\}$ for every j in Λ . We establish a multifunction $F : X \rightarrow \Lambda$ where $F(x) = \{i \in \Lambda \mid i \geq j_x\}$, j_x is the first element of all those j 's for which $x \notin M_j$. Since Λ has the order topology, $F(x)$ is closed. By the fact that $\{j_x \mid x \in X\}$ has no greatest element in Λ , then F does not assume any minimal value with respect to set inclusion. We now show that $F^{-}(U) \in GO(X)$ for every open set U in Λ . If $U = \Lambda$, then $F^{-}(U) = X$ which is g -open. Suppose that $U \subset \Lambda$ and $z \in F^{-}(U)$. It follows that $F(z) \cap U \neq \emptyset$. Suppose $j \in F(z) \cap U$. This means that $j \in U$ and $j \in F(z) = \{i \in \Lambda \mid i \geq j_x\}$. Therefore $M_j \geq M_{j_x}$. Since $z \notin M_{j_x}$, then $z \notin M_j$. There exists $W \in GO(z)$ such that $W \cap \{x_i \mid i \in \Lambda\} = \emptyset$. This means that $W \cap M_j = \emptyset$. Let $w \in W$. Since $W \cap M_j = \emptyset$, it follows that $w \notin M_j$ and since j_w is the first element for which $w \notin M_j$, then $j_w \leq j$. Therefore $j \in \{i \in \Lambda \mid i \geq j_w\} = F(w)$. By the fact that $j \in U$, then $j \in F(w) \cap U$. It follows that $F(w) \cap U \neq \emptyset$ and therefore $w \in F^{-}(U)$. So we have $W \subset F^{-}(U)$ and thus $z \in W \subset F^{-}(U)$. Therefore $F^{-}(U)$ is g -open. This shows that F is lower g -continuous which contradicts the hypothesis of the theorem. So the space X is GO -compact.

Theorem 4.4 *The following two statements are equivalent for a space X :*

- (1) X is GO -compact.
- (2) Every upper g -continuous multifunction from X into the subsets of a T_1 -space attains a maximal value with respect to set inclusion relation.

Proof. Its proof is similar to that of Theorem 4.3.

The following result concerns the existence of a fixed point for multifunctions on GO -compact spaces.

Theorem 4.5 *Suppose that $F : X \rightarrow Y$ is a multifunction from a GO -compact domain X into itself. Let $F(S)$ be g -closed for S being a g -closed set in X . If $F(x) \neq \emptyset$ for every point $x \in X$, then there exists a nonempty, g -closed set C of X such that $F(C) = C$.*

Proof. Let $\Lambda = \{S \subset X \mid S \neq \emptyset, S \in GC(X) \text{ and } F(S) \subset S\}$. It is evident that x belongs to Λ . Therefore $\Lambda \neq \emptyset$ and also it is partially ordered by set inclusion. Suppose that $\{S_\gamma\}$ is a chain in Λ . Then $F(S_\gamma) \subset S_\gamma$ for each γ . By the fact that the domain is GO -compact and by ([8], Theorem 3.3), $S = \bigcap_\gamma S_\gamma \neq \emptyset$ and also $S \in GC(X)$. Moreover, $F(S) \subset F(S_\gamma) \subset S_\gamma$ for each γ . It follows that $F(S) \subset S_\gamma$. Hence $S \in \Lambda$ and $S = \inf\{S_\gamma\}$. It follows from Zorn's lemma that Λ has a minimal element C . Therefore $C \in GC(X)$ and $F(C) \subset C$. Since C is the minimal element of Λ , we have $F(C) = C$.

We close with the following open question:

Question 4.6 Give a nontrivial example of a GO -compact space?

Acknowledgment. S. P. Moshokoa acknowledges the support by the South African National Research Foundation under Grant number 2053847. Parts of this paper were written while S. Jafari was a visitor to the Department of Mathematical Sciences of the University of South Africa during october 2005.

References

- [1] K. Balachandran, P. Sundaram and H. Maki, On generalized continuous maps in topological spaces, *Mem. Fac. Sci. Kochi Univ. (Math.)*, **12** (1991), 5-13.
- [2] C. Berge, Topological spaces, Macmillan, New York, 1963.
- [3] M. Caldas, On g -closed sets and g -continuous mappings, *Kyungpook Math. J.* **33**(1993),205-209.
- [4] M. Caldas and S. Jafari, On g -US spaces, *Univ. Bacau. Stud. Cerc. St. Ser. Mat. Romania*, **14** (2005), 13-20.
- [5] M. Caldas and S. Jafari, On rarely g -continuous functions, *Glasnik Mat.*, **40** (2) (2005), 317-322.
- [6] M. Caldas, S. Jafari and T. Noiri, Notions via g -open sets, *Kochi J. Math.* **2**(2007), 43-50.
- [7] M. Caldas, S. Jafari, T. Noiri and M. Simões, A new generalization of contra-continuity via Levine's g -closed sets, *Chaos, Solitons and Fractals*, **32** (4)(2007), 1597-1603.
- [8] W. Dunham and N. Levine, Further results on generalized closed sets in topology, *Kyungpook Math. J.*, **20** (1980), 169-175.
- [9] W. Dunham, $T_{1/2}$ -spaces, *Kyungpook Math. J.*, **17**(1977), 161-169.
- [10] W. Dunham, A new closure operator for non- T_1 topologies, *Kyungpook Math. J.*, **22**(1982), 55-60.

- [11] N. Levine, Generalized closed sets in topology, *Rend. Circ. Mat. Palermo* (2), **19** (1970), 89-96.
- [12] S. R. Malghan, Generalized closed sets maps, *J. Karnatak Univ. Sci.*, **27**(1982), 82-88.

Addresses :

M. Caldas
Departamento de Matemática Aplicada,
Universidade Federal Fluminense,
Rua Mário Santos Braga, s/n^o
24020-140, Niterói, RJ BRAZIL.
E-mail: gmamccs@vm.uff.br

S. Jafari
College of Vestsjaelland South,
Herrestraede 11
4200 Slagelse, DENMARK.
E-mail: jafari@stofanet.dk

S. P. Moshokoa
Department of Mathematical Sciences,
University of South Africa,
P.O. Box 392, Pretoria 0003, SOUTH AFRICA.
E-mail: moshosp@unisa.ac.za

T. Noiri
2949-1 Shiokita-cho, Hinagu,
Yatsushiro-shi, Kumamoto-ken,
869-5142 JAPAN
E-mail: t.noiri@nifty.com