

## On harmonic Hardy and Bergman spaces

By Stevo STEVIĆ

(Received Nov. 6, 2000)

(Revised Apr. 16, 2001)

**Abstract.** In this paper we use an identity of Hardy-Stein type in investigations of the harmonic  $\mathcal{H}^p(B)$  and Bergman  $b^p(B)$  spaces.

### 1. Introduction and auxiliary results.

Throughout this paper  $n$  is an integer greater than 1,  $B(a, r) = \{x \in \mathbf{R}^n \mid |x - a| < r\}$  denotes the open ball centered at  $a$  of radius  $r$ , where  $|x|$  denotes the norm of  $x \in \mathbf{R}^n$  and  $B$  is the open unit ball in  $\mathbf{R}^n$ .  $S = \partial B = \{x \in \mathbf{R}^n \mid |x| = 1\}$  is the boundary of  $B$ .

Let  $dV$  denote the Lebesgue measure on  $\mathbf{R}^n$ ,  $d\sigma$  the surface measure on  $S$ ,  $\sigma_n$  the surface area of a  $S$ ,  $dV_N$  the normalized Lebesgue measure on  $B$ ,  $d\sigma_N$  the normalized surface measure on  $S$ .

Let  $\mathcal{H}(B)$  denote the set of complex-valued harmonic functions on  $B$ ,  $\mathcal{H}^p(B)$  denote the set of harmonic functions on  $B$  such that:

$$\|u\|_{\mathcal{H}^p(B)} = \sup_{0 < r < 1} \left( \int_S |u(r\zeta)|^p d\sigma_N(\zeta) \right)^{1/p} < +\infty$$

and let  $b^p(B)$  denote the Bergman space i.e., the set of harmonic functions  $u$  on  $B$  such that:

$$\|u\|_p = \left( \int_B |u(x)|^p dV(x) \right)^{1/p} < +\infty.$$

A function  $f \in C^1(B)$  is said to be a Bloch function if

$$\|f\|_{\mathcal{B}} = \sup_{x \in B} (1 - |x|) |\nabla f(x)| < +\infty$$

where  $|\nabla f(x)| = (\sum_1^n |\partial f(x)/\partial x_i|^2)^{1/2}$ . The space of Bloch functions is denoted by  $\mathcal{B}(B)$ .

Let  $p > 0$ . A Borel function  $f$ , locally integrable on  $B$ , is said to be a  $BMO_p(B)$  function if

$$\|f\|_{BMO_p} = \sup_{B(a,r) \subset B} \left( \frac{1}{V(B(a,r))} \int_{B(a,r)} |f(x) - f_{B(a,r)}|^p dV(x) \right)^{1/p} < +\infty$$

where the supremum is taken over all balls  $B(a,r)$  in  $B$ , and  $f_{B(a,r)}$  is the mean value of  $f$  over  $B(a,r)$ .

In [6] for  $p \geq 1$ , Muramoto proved that  $\mathcal{B}(B) \cap \mathcal{H}(B)$  is isomorphic to  $BMO_p \cap \mathcal{H}(B)$  as Banach spaces. That paper inspired us to calculate exactly  $BMO$  norm which is the theme of [8]. We proved the following:

**THEOREM A.** *Let  $u \in \mathcal{H}(B)$ ,  $p > 1$ . Then*

a)

$$\|u\|_{BMO_p}^p = \sup_{a \in B, 0 < r < 1 - |a|} \frac{p(p-1)}{2n(n-2)} \times \left( \int_B |u_{a,r}(x) - u_{a,r}(0)|^{p-2} |\nabla u_{a,r}(x)|^2 (2|x|^{2-n} + (n-2)|x|^2 - n) dV_N(x) \right)$$

for  $n \geq 3$

b)

$$\|u\|_{BMO_p}^p = \sup_{a \in B, 0 < r < 1 - |a|} p(p-1) \times \left( \int_B |u_{a,r}(x) - u_{a,r}(0)|^{p-2} |\nabla u_{a,r}(x)|^2 (\ln(1/|x|) - 1 + |x|) dV_N(x) \right)$$

for  $n = 2$ .

In the proof of this theorem we essentially also proved a generalization of Hardy-Stein identity, [2]. This identity is included in the following lemma.

**LEMMA 1.** *Let  $1 < p < +\infty$ ,  $u \in \mathcal{H}(B)$ , then*

$$\int_S |u(r\zeta)|^p d\sigma_N(\zeta) = |u(0)|^p + \frac{p(p-1)}{n(n-2)} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (|x|^{2-n} - r^{2-n}) dV_N(x), \quad n \geq 3. \quad (1)$$

It is interesting that Lemma 1 is a consequence of the Riesz decomposition theorem, see, for example [4], of the subharmonic function  $|u|^p$  on  $rB$  into a difference between a harmonic function and a potential.

COROLLARY 1. Let  $1 < p < +\infty$ ,  $u \in \mathcal{H}(B)$ ,  $r \in (0, 1)$  then

$$\frac{d}{dr} \int_S |u(r\zeta)|^p d\sigma_N(\zeta) = \frac{p(p-1)}{n} r^{1-n} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x). \tag{2}$$

Note that this identity is of Hardy-Stein type.

This lemma will be the main tool in this paper. We will use this lemma in investigations of spaces  $\mathcal{H}^p(B)$  and  $b^p(B)$ . Similar identity exists in the case of holomorphic functions in [9], but the author does not use it. We shall keep our attention to the case  $n \geq 3$ . Analogous results hold in the case  $n = 2$ . Formulations and proofs of this results we leave to the reader.

We will need another lemma in our consideration.

LEMMA 2. Let  $I(r)$  be a nonnegative nondecreasing function on the interval  $[0, 1)$ ,  $n \in \mathbb{N}$ ,  $M > 0$ , and let

$$r^n I(r) \leq n \int_0^r I(\rho) \rho^{n-1} d\rho + M, \tag{3}$$

then for each  $\varepsilon \in (0, r)$  the following inequality

$$I(r) \leq \frac{M}{\varepsilon^n} + I(\varepsilon)$$

holds.

PROOF. From (3) we have

$$\frac{nt^{n-1}I(t)}{(M + n \int_0^t I(\rho)\rho^{n-1} d\rho)} \leq n/t. \tag{4}$$

Integrating (4) from  $\varepsilon$  to  $r$  in the variable  $t$ , we obtain

$$\ln \frac{(M + n \int_0^r I(\rho)\rho^{n-1} d\rho)}{(M + n \int_0^\varepsilon I(\rho)\rho^{n-1} d\rho)} \leq n \ln \frac{r}{\varepsilon}$$

i.e.

$$M + n \int_0^r I(\rho)\rho^{n-1} d\rho \leq \left( M + n \int_0^\varepsilon I(\rho)\rho^{n-1} d\rho \right) \left( \frac{r}{\varepsilon} \right)^n.$$

Since  $I(r)$  is nondecreasing we have

$$\left( M + n \int_0^\varepsilon I(\rho)\rho^{n-1} d\rho \right) \left( \frac{r}{\varepsilon} \right)^n \leq (M + I(\varepsilon)\varepsilon^n) \frac{r^n}{\varepsilon^n} = M \frac{r^n}{\varepsilon^n} + I(\varepsilon)r^n.$$

From which the result follows. □

In section 2 we prove several necessary and sufficient conditions for a harmonic function in the unit ball to be in  $\mathcal{H}^p(B)$  (Theorems 1–4). In Theorem 5 we investigate the rate of growth of

$$E(r) = \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV(x), \quad u \in \mathcal{H}^p(B),$$

as  $r \rightarrow 1$ . In Theorem 6 we give a local estimate of derivatives of such functions.

In section 3, we prove necessary and sufficient condition for a harmonic function in the unit ball to be in harmonic Bergman space  $b^p(B)$  (Theorem 7). We also investigate the rate of growth of  $E(r)$  as  $r \rightarrow 1$ , for  $u \in b^p(B)$  (Theorem 8).

### 2. Hardy harmonic $\mathcal{H}^p(B)$ space.

In this section we consider Hardy harmonic  $\mathcal{H}^p(B)$  space.

**THEOREM 1.** *Let  $1 < p < +\infty$ . Function  $u \in \mathcal{H}(B)$  belongs to  $\mathcal{H}^p(B)$  if and only if*

$$\int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|^2) dV_N(x) < +\infty.$$

If  $u \in \mathcal{H}^p(B)$ ,  $1 < p < +\infty$ , then

$$\|u\|_{\mathcal{H}^p}^p = \int_B |u(x)|^p dV_N(x) + \frac{p(p-1)}{2n} \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|^2) dV_N(x).$$

**PROOF.** Multiplying the formula (2) by  $r^n$  and integrating from 0 to  $r$ , we obtain

$$\int_0^r \rho^n \frac{d}{d\rho} \int_S |u(\rho\zeta)|^p d\sigma_N(\zeta) d\rho = \frac{p(p-1)}{n} \int_0^r \rho \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x) d\rho.$$

Applying integration by parts on the left hand side of previous equality and using Fubini's theorem on the right hand side we obtain

$$\begin{aligned} & r^n \int_S |u(r\zeta)|^p d\sigma_N(\zeta) \\ &= \int_{rB} |u(x)|^p dV_N(x) + \frac{p(p-1)}{2n} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (r^2 - |x|^2) dV_N(x). \end{aligned} \tag{5}$$

If  $u \in \mathcal{H}^p(B)$ , then by polar coordinates easily follows  $u \in b^p(B)$ . From that by the monotone convergence theorem we have

$$\lim_{r \rightarrow 1-0} \int_{rB} |u(x)|^p dV_N(x) = \int_B |u(x)|^p dV_N(x).$$

Since functions  $r^n$  and  $\int_S |u(r\zeta)|^p d\sigma_N(\zeta)$  are nondecreasing then function  $r^n \int_S |u(r\zeta)|^p d\sigma_N(\zeta)$  is also such a function. From  $u \in \mathcal{H}^p(B)$  also follows

$$\lim_{r \rightarrow 1-0} r^n \int_S |u(r\zeta)|^p d\sigma_N(\zeta) = \int_S |u^*(\zeta)|^p d\sigma_N(\zeta),$$

where  $u^*(\zeta)$  is a usual radial limit for a function in  $\mathcal{H}^p$ .

From all of the above we have there exists

$$\lim_{r \rightarrow 1-0} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (r^2 - |x|^2) dV_N(x).$$

By the monotone convergence theorem we have

$$\begin{aligned} & \lim_{r \rightarrow 1-0} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (r^2 - |x|^2) dV_N(x) \\ &= \int_B \lim_{r \rightarrow 1-0} |u(x)|^{p-2} |\nabla u(x)|^2 (r^2 - |x|^2) \chi_{rB}(x) dV_N(x) \\ &= \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|^2) dV_N(x) \end{aligned}$$

as required.

Conversely, if  $M = \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|^2) dV_N(x) < +\infty$ , then by (5) we have

$$r^n \int_S |u(r\zeta)|^p d\sigma_N(\zeta) \leq \int_{rB} |u(x)|^p dV_N(x) + M_1, \quad M_1 \in \mathbf{R}_+, \quad r \in (0, 1). \quad (6)$$

Let  $I(r) = \int_S |u(r\zeta)|^p d\sigma_N(\zeta)$ , then we may simply write (6) in the following form

$$r^n I(r) \leq n \int_0^r \rho^{n-1} I(\rho) d\rho + M_1.$$

Taking  $\varepsilon = 1/2$  in Lemma 2 we have  $I(r) \leq M_1 2^n + I(1/2)$ , for  $r \in [1/2, 1)$ . Since  $I(r)$  is nondecreasing function we have  $I(r) \leq I(1/2)$ , for  $r \in [0, 1/2]$ . Thus the result follows. □

**THEOREM 2.** *Let  $u \in \mathcal{H}^p(B)$ ,  $1 < p < +\infty$ , then*

$$\|u\|_{\mathcal{H}^p}^p = |u(0)|^p + \frac{p(p-1)}{n(n-2)} \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (|x|^{2-n} - 1) dV_N(x).$$

**PROOF.** Since  $u \in \mathcal{H}^p(B)$ , letting  $r \rightarrow 1$  in formula (1), and noting that the integrand is positive and increasing in  $r$  we obtain the above formula. □

Next theorem is a generalization of Lemma 1 and Lemma 2 in [2]. We use the following notation:

$$I_s(r, u) = \left( \int_S |u(r\zeta)|^s d\sigma_N(\zeta) \right)^{1/s}.$$

**THEOREM 3.** *Let  $s > 0$ . If  $u \in \mathcal{H}^s(B)$ ,  $1 < p \leq 2$ , then*

$$\int_0^1 (1 - \rho) I_{2s/(s-p+2)}^2(\rho, \nabla u) d\rho < +\infty. \quad (7)$$

*If  $2 \leq p < s + 2$ ,  $s \leq p$ , then (7) implies  $u \in \mathcal{H}^s(B)$ .*

**PROOF.** By Theorem 1 condition  $u \in \mathcal{H}^p(B)$  is equivalent with

$$\begin{aligned} & \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|^2) dV(x) \\ &= \int_0^1 r^{n-1} (1 - r^2) \int_S |u(r\zeta)|^{p-2} |\nabla u(r\zeta)|^2 d\sigma(\zeta) dr < +\infty. \end{aligned} \quad (8)$$

Let  $J_p(r) = \int_S |u(r\zeta)|^{p-2} |\nabla u(r\zeta)|^2 d\sigma_N(\zeta)$ .

If  $p \leq 2$ , we can use Jensen's inequality to the integral

$$\frac{J_p(r)}{I_s^s(r, u)} = \int_S \frac{|\nabla u(r\zeta)|^2}{|u(r\zeta)|^{s-p+2}} \frac{|u(r\zeta)|^s}{I_s^s(r, u)} d\sigma_N(\zeta).$$

Since in that case  $0 < s/(s-p+2) \leq 1$ , we have

$$\begin{aligned} \left( \frac{J_p(r)}{I_s^s(r, u)} \right)^{s/(s-p+2)} &\geq \int_S \frac{|\nabla u(r\zeta)|^{2s/(s-p+2)}}{|u(r\zeta)|^s} \frac{|u(r\zeta)|^s}{I_s^s(r, u)} d\sigma_N(\zeta) \\ &= \int_S |\nabla u(r\zeta)|^{2s/(s-p+2)} d\sigma_N(\zeta) I_s^{-s}(r, u), \end{aligned} \quad (9)$$

i.e.

$$J_p(r) \geq I_s^{p-2}(r, u) I_{2s/(s-p+2)}^2(r, \nabla u). \quad (10)$$

From that we have

$$J_p(r) \geq \|u\|_{\mathcal{H}^s}^{p-2} I_{2s/(s-p+2)}^2(r, \nabla u). \quad (11)$$

By (8) and (11) we get

$$\int_0^1 r^{n-1} (1 - r^2) I_{2s/(s-p+2)}^2(r, \nabla u) dr < +\infty.$$

It is easy to see that the last integral is equiconvergent with integral in (7).

If  $2 \leq p < s + 2$ ,  $s \leq p$ , then in (10) the converse inequality holds. Therefore from (1) we obtain, using polar coordinates,

$$\begin{aligned} I_p^p(r, u) &\leq |u(0)|^p + \frac{p(p-1)}{(n-2)} \int_0^r J_p(\rho)(\rho^{2-n} - r^{2-n})\rho^{n-1} d\rho \\ &\leq |u(0)|^p + \frac{p(p-1)}{(n-2)} \int_0^r I_s^{p-2}(\rho, u) I_{2s/(s-p+2)}^2(\rho, \nabla u)(\rho - r^{2-n}\rho^{n-1}) d\rho. \end{aligned}$$

Since  $u \in \mathcal{H}(B)$ , function  $I_p(r, u)$  is nondecreasing in  $r$ ; see, for example [4]. By Jensen's inequality we get  $I_s(r, u) \leq I_p(r, u)$  for  $s \leq p$ . Thus we have

$$\begin{aligned} I_p^2(r, u) &\leq |u(0)|^2 + \frac{p(p-1)}{(n-2)} \int_0^r I_{2s/(s-p+2)}^2(\rho, \nabla u)(\rho - r^{2-n}\rho^{n-1}) d\rho \\ &\leq |u(0)|^2 + \frac{p(p-1)}{(n-2)} \int_0^r I_{2s/(s-p+2)}^2(\rho, \nabla u)\rho(1 - \rho^{n-2}) d\rho \\ &\leq |u(0)|^2 + p(p-1) \int_0^r I_{2s/(s-p+2)}^2(\rho, \nabla u)(1 - \rho) d\rho. \end{aligned}$$

From that follows second part of this theorem. □

**COROLLARY 2.** *Let  $p \geq 2$ ,  $u \in \mathcal{H}(B)$ . If  $\int_B(1 - |x|)|\nabla u(x)|^p dV(x) < +\infty$ , then  $u \in \mathcal{H}^p(B)$ .*

**PROOF.** Since  $u \in \mathcal{H}(B)$ , function  $|\nabla u|^p$  is subharmonic, see [7], and therefore  $I_p(r, \nabla u)$  is nondecreasing in  $r$ . If  $\sup_{0 < r < 1} I_p^p(r, \nabla u)$  is finite, then  $\sup_{0 < r < 1} I_p^2(r, \nabla u)$  is also finite. Thus we have

$$\int_0^1 (1 - \rho)I_p^2(\rho, \nabla u) d\rho < +\infty.$$

By Theorem 3, we obtain  $u \in \mathcal{H}^p(B)$ .

If  $\sup_{0 < r < 1} I_p^p(r, \nabla u) = +\infty$  we have there exists  $r_0$  such that for  $r \in [r_0, 1)$ ,  $I_p^p(r, \nabla u) > 1$ . Since  $p \geq 2$  we have  $I_p^2(r, \nabla u) \leq I_p^p(r, \nabla u)$  for  $r \in [r_0, 1)$ . Therefore

$$\int_{r_0}^1 (1 - \rho)I_p^2(\rho, \nabla u) d\rho \leq \int_{r_0}^1 (1 - \rho)I_p^p(\rho, \nabla u) d\rho < +\infty.$$

Hence

$$\int_0^1 (1 - \rho)I_p^2(\rho, \nabla u) d\rho < +\infty.$$

Applying Theorem 3 we obtain our result. □

By Theorem 3, as in Corollary 2 we can prove this result. Details we leave to the reader.

**COROLLARY 3.** *Let  $1 < p \leq 2$ ,  $u \in \mathcal{H}^p(B)$ . Then*

$$\int_B (1 - |x|) |\nabla u(x)|^p dV(x) < +\infty.$$

We shall give an elementary proof of the following theorem (i.e. without using maximal theorem).

**THEOREM 4.** *Let  $u \in \mathcal{H}(B)$  and  $\sup_{0 < r < 1} \int_S |\nabla u(r\zeta)|^p d\sigma_N(\zeta) < +\infty$ ,  $p \geq 2$ , then  $u \in \mathcal{H}^p(B)$ .*

**PROOF.** Let  $I(r) = \int_S |u(r\zeta)|^p d\sigma_N(\zeta)$  and

$$J_p(r) = J(r) = \int_S |u(r\zeta)|^{p-2} |\nabla u(r\zeta)|^2 d\sigma_N(\zeta)$$

in polar coordinates (2) becomes

$$I'(r) = p(p - 1)r^{1-n} \int_0^r \rho^{n-1} J(\rho) d\rho. \tag{12}$$

Let us estimate  $J(r)$  by  $I(r)$ . For that purpose we write  $J(r)$  as follows

$$J(r) = \int_S \frac{|u(r\zeta)|^{p-2}}{|\nabla u(r\zeta)|^{p-2}} \frac{|\nabla u(r\zeta)|^p}{\|\nabla u(r \cdot)\|_p^p} d\sigma_N(\zeta) \cdot \|\nabla u(r \cdot)\|_p^p,$$

where

$$\|\nabla u(r \cdot)\|_p^p = \int_S |\nabla u(r\zeta)|^p d\sigma_N(\zeta).$$

For  $p > 2$  we have  $p/(p - 2) > 1$ . Applying Jensen's inequality we obtain

$$\begin{aligned} J(r)^{p/(p-2)} &\leq \int_S \frac{|u(r\zeta)|^p}{|\nabla u(r\zeta)|^p} \frac{|\nabla u(r\zeta)|^p}{\|\nabla u(r \cdot)\|_p^p} d\sigma_N(\zeta) \cdot \|\nabla u(r \cdot)\|_p^{p^2/(p-2)} \\ &= I(r) \left( \int_S |\nabla u(r\zeta)|^p d\sigma_N(\zeta) \right)^{2p/(p-2)}. \end{aligned}$$

Thus

$$J(r) \leq I(r)^{(p-2)/p} \left( \sup_{0 < r < 1} \int_S |\nabla u(r\zeta)|^p d\sigma_N(\zeta) \right)^2 = C_{\nabla u} I(r)^{(p-2)/p}. \tag{13}$$

Combining (12) and (13) we obtain



$$I'(r) \leq C_{\nabla u} p(p-1)r^{1-n} \int_0^r \rho^{n-1} I(\rho)^{(p-2)/p} d\rho.$$

Integrating from 0 to  $r$  we get

$$\begin{aligned} I(r) &\leq I(0) + C_{\nabla u} p(p-1) \int_0^r \rho^{1-n} \int_0^\rho s^{n-1} I(s)^{(p-2)/p} ds d\rho \\ &= I(0) + C_{\nabla u} p(p-1) \cdot \int_0^r s^{n-1} I(s)^{(p-2)/p} \int_s^r \rho^{1-n} d\rho ds \end{aligned}$$

i.e.

$$\begin{aligned} I(r) &\leq I(0) + \frac{C_{\nabla u} p(p-1)}{n-2} \int_0^r s^{n-1} I(s)^{(p-2)/p} (s^{2-n} - r^{2-n}) ds \\ &\leq I(0) + \frac{C_{\nabla u} p(p-1)}{n-2} \int_0^r s I(s)^{(p-2)/p} ds. \end{aligned} \tag{14}$$

Let  $\alpha = (p-2)/p$  then (14) can be written

$$I(r)^\alpha \leq \left( I(0) + C \int_0^r s I(s)^\alpha ds \right)^\alpha, \quad C = \frac{C_{\nabla u} p(p-1)}{n-2}. \tag{15}$$

From that we have

$$\frac{CrI(r)^\alpha}{\left( I(0) + C \int_0^r s I(s)^\alpha ds \right)^\alpha} \leq Cr, \quad 0 < r < 1.$$

Integrating from 0 to  $r$  we get

$$\left( I(0) + C \int_0^r s I(s)^\alpha ds \right)^{1-\alpha} - I(0)^{1-\alpha} \leq (1-\alpha) \frac{Cr^2}{2}. \tag{16}$$

By (15) and (16) we have

$$I(r) \leq \left( (1-\alpha) \frac{Cr^2}{2} + I(0)^{1-\alpha} \right)^{1/(1-\alpha)} \leq \left( \frac{(1-\alpha)C}{2} + I(0)^{1-\alpha} \right)^{1/(1-\alpha)} < +\infty$$

thus  $u \in \mathcal{H}^p(B)$ .

REMARK 1. Theorem 4 is a consequence of Minkowski's inequality and maximal theorem, also in the case  $p \in (1, 2)$ .

THEOREM 5. Let  $1 < p < +\infty$ ,  $u \in \mathcal{H}^p(B)$ , then

$$\lim_{r \rightarrow 1-0} (1-r) \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV(x) = 0.$$

PROOF. Integrating the following formula

$$r^{n-1} \frac{d}{dr} \int_S |u(r\xi)|^p d\sigma_N(\xi) = \frac{p(p-1)}{n} \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x)$$

from 0 to  $r$  we get

$$\begin{aligned} r^{n-1} \int_S |u(r\xi)|^p d\sigma_N(\xi) - (n-1) \int_0^r \rho^{n-2} \int_S |u(\rho\xi)|^p d\sigma_N(\xi) d\rho \\ = \frac{p(p-1)}{n} \int_0^r \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x) d\rho. \end{aligned}$$

Since  $u \in \mathcal{H}^p(B)$  we have  $u \in b^p(B)$ . On the other hand

$$\begin{aligned} \int_0^r \rho^{n-2} \int_S |u(\rho\xi)|^p d\sigma_N(\xi) d\rho \\ = \int_0^{1/2} \rho^{n-2} \int_S |u(\rho\xi)|^p d\sigma_N(\xi) d\rho + \int_{1/2}^r \rho^{n-1} \int_S |u(\rho\xi)|^p d\sigma_N(\xi) \frac{d\rho}{\rho} \\ \leq c_1 + c_2 \int_{rB} |u(x)|^p dV(x) < +\infty. \end{aligned}$$

Therefore, there exists

$$\lim_{r \rightarrow 1-0} \int_0^r \rho^{n-2} \int_S |u(\rho\xi)|^p d\sigma_N(\xi) d\rho,$$

from which it follows that

$$\lim_{r \rightarrow 1-0} \int_0^r \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x) d\rho$$

exists.

Thus by Cauchy's criterion we have

$$\lim_{r \rightarrow 1-0} \int_r^1 \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 dV_N(x) d\rho = 0.$$

Since function  $J(\rho) = \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 dV(x)$  is nondecreasing we obtain

$$(1-r)J(r) \leq \int_r^1 J(\rho) d\rho \rightarrow 0, \quad \text{as } r \rightarrow 1-0,$$

which is what we needed to prove. □

In the following theorem we generalize Lemma 2 in [5]. In that paper Luecking proved theorem in the case of  $n = 2$ . Also Luecking reproved Hardy-Stein identity in that case and applied it in proving Littlewood-Paley inequality.

**THEOREM 6.** *Let  $p \geq 2$  and  $u \in \mathcal{H}^p(B)$ , then*

$$|\nabla u(0)|^p \leq \frac{n^{p/2} p(p-1)}{(n-2)n} \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (|x|^{2-n} - 1) dV_N(x), \quad n \geq 3.$$

**PROOF.** It is well-known that if  $u \in \mathcal{H}(B)$  then  $u(x) = \sum_{m=0}^{+\infty} p_m(x)$ , where  $p_m(x)$  is harmonic homogeneous polynomial of degree  $m$  on  $B$ . By Hölder inequality we have  $\|u\|_{\mathcal{H}^2} \leq \|u\|_{\mathcal{H}^p}$ . For  $u \in \mathcal{H}^2(B)$ , the following formula

$$\|u\|_{\mathcal{H}^2}^2 = \sum_{m=0}^{+\infty} \int_S |p_m(\zeta)|^2 d\sigma_N(\zeta) \tag{17}$$

holds; see, for example [1, p. 122].

On the other hand, for homogeneous polynomial of degree  $m$  the following formula  $\langle \nabla p_m(x), x \rangle = mp_m(x)$ ,  $x \in \mathbf{R}^n$ , holds. From (17) we have

$$\begin{aligned} \|u\|_{\mathcal{H}^2}^2 - |u(0)|^2 &= \sum_{m=1}^{+\infty} \int_S |p_m(\zeta)|^2 d\sigma_N(\zeta) \geq \int_S |p_1(\zeta)|^2 d\sigma_N(\zeta) \\ &= \int_S |\langle \nabla p_1(\zeta), \zeta \rangle|^2 d\sigma_N(\zeta) \end{aligned}$$

since  $u(0) = p_0(0)$ . Let  $p_1(x) = a_1x_1 + \dots + a_nx_n$ ,  $a_1, \dots, a_n \in \mathbf{C}$ , then

$$\begin{aligned} \int_S |\langle \nabla p_1(\zeta), \zeta \rangle|^2 d\sigma_N(\zeta) &= \int_S |a_1\zeta_1 + \dots + a_n\zeta_n|^2 d\sigma_N(\zeta) \\ &= \int_S \left( \sum_{i=1}^n |a_i|^2 \zeta_i^2 + 2 \sum_{i \neq j} a_i \bar{a}_j \zeta_i \zeta_j \right) d\sigma_N(\zeta). \end{aligned}$$

By symmetry of the sphere and subintegral functions we have  $\int_S \zeta_i \zeta_j d\sigma_N(\zeta) = 0$ ,  $i \neq j$ , and  $\int_S \zeta_i^2 d\sigma_N(\zeta) = 1/n$ ,  $i = 1, \dots, n$ .

Thus

$$\int_S |\langle \nabla p_1(\zeta), \zeta \rangle|^2 d\sigma_N(\zeta) = \frac{1}{n} \sum_{i=1}^n |a_i|^2.$$

Since  $\sum_{i=1}^n |a_i|^2 = |\nabla p_1(x)|^2 = |\nabla p_1(0)|^2 = |\nabla u(0)|^2$  we have

$$\frac{1}{n} |\nabla u(0)|^2 \leq \|u\|_{\mathcal{H}^2}^2 - |u(0)|^2 \leq \|u\|_{\mathcal{H}^p}^2 - |u(0)|^2.$$

By the inequality  $(a - b)^q + b^q \leq a^q$ ,  $a \geq b > 0$ ,  $q \geq 1$ , we obtain

$$|\nabla u(0)|^p \leq n^{p/2} (\|u\|_{\mathcal{H}^p}^2 - |u(0)|^2)^{p/2} \leq n^{p/2} (\|u\|_{\mathcal{H}^p}^p - |u(0)|^p). \tag{18}$$

By the following formula

$$\|u\|_{\mathcal{H}^p}^p = |u(0)|^p + \frac{p(p-1)}{n(n-2)} \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (|x|^{2-n} - 1) dV_N(x), \tag{19}$$

(18) and (19) the result follows. □

### 3. Bergman $b^p(B)$ space.

In this section we consider harmonic Bergman space  $b^p(B)$ .

**THEOREM 7.** *Let  $u \in \mathcal{H}(B)$ ,  $p > 1$ , then  $u \in b^p(B)$  if and only if*

$$I_u = \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (2|x|^{2-n} + (n-2)|x|^2 - n) dV_N(x) < +\infty.$$

**PROOF.** Multiplying the formula (1) by  $nr^{n-1}$ , and integrating from 0 to 1, applying Fubini's theorem, for  $u \in \mathcal{H}(B)$ , we obtain the following formula

$$\|u\|_{b^p}^p = |u(0)|^p + \frac{p(p-1)}{2n(n-2)} \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (2|x|^{2-n} + (n-2)|x|^2 - n) dV_N(x),$$

thus the result follows. □

**COROLLARY 4.**  *$u \in b^p(B)$ ,  $p > 1$  if and only if*

$$\int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|)^2 dV(x) < +\infty.$$

**PROOF.** Let

$$I_u = \int_B |u(x)|^{p-2} |\nabla u(x)|^2 ((n-2)|x|^n - n|x|^{n-2} + 2)/|x|^{n-2} dV_N(x).$$

We leave to the reader that  $I_u$  is equiconvergent with  $\int_B |u(x)|^{p-2} |\nabla u(x)|^2 \cdot (1 - |x|)^2 / |x|^{n-2} dV(x)$ .

Using polar coordinates it is easy to see that the last integral is equiconvergent with

$$\int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|)^2 dV(x).$$

If  $n \geq 3$ , the Green function for the ball  $B$  with pole in the origin is given by  $G(|x|, 1) = |x|^{2-n} - 1$ . □

**COROLLARY 5.** *Let  $1 < p < +\infty$ , then  $u \in b^p(B)$  if and only if*

$$L_u = \int_B |u(x)|^{p-2} |\nabla u(x)|^2 G(|x|, 1)(1 - |x|) dV(x) < +\infty.$$

PROOF. By Theorem 7 it is enough to show that integrals  $L_u$  and  $I_u$  are equiconvergent. We know that

$$\begin{aligned} I_u &\sim \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|)^2 / |x|^{n-2} dV(x) \\ &= \int_B |u(x)|^{p-2} |\nabla u(x)|^2 G(|x|, 1)(1 - |x|)^2 / (1 - |x|^{n-2}) dV(x). \end{aligned}$$

It is clear that expression

$$\frac{1 - |x|}{1 - |x|^{n-2}} = \frac{1}{1 + |x| + \dots + |x|^{n-3}}$$

for  $x \in B$  takes values from the interval  $(1/(n - 2), 1]$ . Thus we have the required result. □

REMARK 2. Note that from previous estimates we have

$$|u(0)|^p + \frac{p(p - 1)}{2n(n - 2)^2} c_1 I_u \leq \|u\|_{b_p}^p \leq |u(0)|^p + \frac{p(p - 1)}{2n(n - 2)} c_2 I_u,$$

for positive  $c_1 = c_1(n)$  and  $c_2 = c_2(n)$ .

THEOREM 8. Let  $u \in b^p(B)$ ,  $p > 1$ , then

$$\lim_{r \rightarrow 1-0} (1 - r)^2 \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV(x) = 0.$$

PROOF. By Fubini's theorem we get

$$\int_0^1 \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|) dV(x) d\rho = \int_B |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|)^2 dV(x).$$

Thus, by Corollary 4 we have

$$\lim_{r \rightarrow 1-0} \int_r^1 \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|) dV(x) d\rho = 0.$$

It is obvious that

$$\begin{aligned} &(1 - r) \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|) dV(x) \\ &\leq \int_r^1 \int_{\rho B} |u(x)|^{p-2} |\nabla u(x)|^2 (1 - |x|) dV(x) d\rho. \end{aligned}$$

Therefore

$$\lim_{r \rightarrow 1-0} (1-r) \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (1-|x|) dV(x) = 0.$$

For  $|x| < r$  we have  $1 - |x| > 1 - r$ , so we have

$$(1-r)^2 \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 dV(x) \leq (1-r) \int_{rB} |u(x)|^{p-2} |\nabla u(x)|^2 (1-|x|) dV(x),$$

which finish the proof of the theorem.  $\square$

### References

- [ 1 ] S. Axler, P. Bourdon and W. Ramey, Harmonic function theory, Springer-Verlag, New York, 1992.
- [ 2 ] G. H. Hardy and J. E. Littlewood, Notes on the theory of series (XX) Generalization of a theorem of Paley, Quart. J. Math., Oxford Ser., **8** (1937), 161–171.
- [ 3 ] W. Hayman, Multivalent functions, Cambridge Univ. Press, London, 1958.
- [ 4 ] W. Hayman and P. B. Kennedy, Subharmonic functions, volume I, Academic Press, London, New York, San Francisco, 1976.
- [ 5 ] D. Luecking, A new proof of an inequality of Littlewood and Paley, Proc. Amer. Math. Soc., **103** (1988), 887–893.
- [ 6 ] K. Muramoto, Harmonic Bloch and BMO functions on the unit ball in several variables, Tokyo J. Math., **11**, no. 2, (1988), 381–386.
- [ 7 ] E. Stein, Singular integrals and differentiability properties of functions, Princeton Univ. Press, 1970.
- [ 8 ] S. Stević, An equivalent norm on BMO spaces, Acta Sci. Math., **66** (2000), 553–563.
- [ 9 ] M. Stoll, A characterization of Hardy spaces on the unit ball of  $\mathbf{C}^n$ , J. London Math. Soc., **48** (1993), 126–136.

Stevo STEVIĆ

Matematički Fakultet  
 Studentski Trg 16  
 11000 Beograd, Serbia  
 Yugoslavia  
 E-mail: sstevo@matf.bg.ac.yu