

## On Hermite-Hadamard Type Inequalities for $(\alpha, m)$ -Convex Functions

Shu-Hong Wang<sup>\*1</sup>, Bo-Yan Xi<sup>\*2</sup>, Feng Qi<sup>†</sup>

<sup>\*</sup>College of Mathematics, Inner Mongolia University for Nationalities,  
Tongliao City, Inner Mongolia Autonomous Region, 028043, China.

E-mail: shuhong7682@163.com (Wang), baoyintu78@qq.com (Xi).

<sup>†</sup>Department of Mathematics, School of Science, Tianjin Polytechnic  
University, Tianjin City, 300387, China.

E-mail: qifeng618@gmail.com, qifeng618@hotmail.com.

URL: <http://qifeng618.wordpress.com>.

### Abstract

*In the paper, some new inequalities of Hermite-Hadamard type for  $(\alpha, m)$ -convex functions are obtained.*

**Keywords:** *Hermite-Hadamard's integral inequality;  $(\alpha, m)$ -convex function; Hölder's inequality.*

## 1 Introduction

Throughout this paper, we use  $b^*$  to represent a positive number and adopt the following notations:

$$\begin{aligned} \mathbb{R} &= (-\infty, \infty), & \mathbb{R}_0 &= [0, +\infty), & \mathbb{R}_+ &= (0, \infty), \\ \bar{a} &= \min\{a, mb\}, & \bar{b} &= \max\{a, mb\}, & \|g\|_\infty &= \sup_{t \in [\bar{a}, \bar{b}]} |g(t)|, \end{aligned}$$

where  $a, b \in \mathbb{R}$  and  $m \in (0, 1]$ .

---

<sup>1</sup>The first author was supported by the Science Research Fund of Inner Mongolia University for Nationalities under Grant No. NMD1103.

<sup>2</sup>The second author was supported in part by the National Natural Science Foundation of China under Grant No. 10962004.

Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a convex function on an interval  $I$  of real numbers and  $a, b \in I$  with  $a < b$ . Then

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}. \quad (1)$$

This inequality is well known in the literature as Hermite-Hadamard's integral inequality for convex functions. See [9] and closely related references therein.

We now recite two definitions.

**Definition 1.1** ([10]). Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a function and  $m \in (0, 1]$ . If

$$f(\lambda x + m(1-\lambda)y) \leq \lambda f(x) + m(1-\lambda)f(y) \quad (2)$$

holds for all  $x, y \in [0, b^*]$  and  $\lambda \in [0, 1]$ , then we say that the function  $f(x)$  is  $m$ -convex on  $[0, b^*]$ .

**Definition 1.2** ([8]). Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a function and  $(\alpha, m) \in (0, 1]^2$ . If

$$f(\lambda x + m(1-\lambda)y) \leq \lambda^\alpha f(x) + m(1-\lambda^\alpha)f(y) \quad (3)$$

is valid for all  $x, y \in [0, b^*]$  and  $\lambda \in [0, 1]$ , then we say that  $f(x)$  is an  $(\alpha, m)$ -convex function on  $[0, b^*]$ .

The following theorems are some known results obtained in recent years.

**Theorem 1.1** ([6, Theorem 2.2]). Let  $f : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I^\circ$  with  $a < b$ . If  $|f'(x)|$  is convex on  $[a, b]$ , then

$$\left| \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{(b-a)(|f'(a)| + |f'(b)|)}{8}. \quad (4)$$

**Theorem 1.2** ([5, Theorem 2]). Let  $f : \mathbb{R}_0 \rightarrow \mathbb{R}$  be  $m$ -convex and  $m \in (0, 1]$ . If  $f \in L[a, b]$  for  $0 \leq a < b < \infty$ , then

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \min \left\{ \frac{f(a) + mf(b/m)}{2}, \frac{mf(a/m) + f(b)}{2} \right\}. \quad (5)$$

**Theorem 1.3** ([3, Theorem 2.2]). Let  $I \supseteq \mathbb{R}_0$  be an open real interval and let  $f : I \rightarrow \mathbb{R}$  be a differentiable function such that  $f'(x) \in L[a, b]$  for  $0 \leq a < b < \infty$ . If  $|f'(x)|^q$  is  $m$ -convex on  $[a, b]$  for some given numbers  $m \in (0, 1]$  and  $q \geq 1$ , then

$$\left| f\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{b-a}{4} \times \min \left\{ \left( \frac{|f'(a)|^q + m|f'(b/m)|^q}{2} \right)^{1/q}, \left( \frac{m|f'(a/m)|^q + |f'(b)|^q}{2} \right)^{1/q} \right\}. \quad (6)$$

**Theorem 1.4** ([11, Theorem 1]). *Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I$  with  $a < b$ . If  $|f'(x)|^q$  is convex on  $[a, b]$  and  $g : [a, b] \rightarrow \mathbb{R}$  is a continuous function, then for all  $x \in [a, b]$  we have*

$$\left| f(a) \int_a^x g(s) ds + f(b) \int_x^b g(s) ds - \int_a^b f(s)g(s) ds \right| \leq \|g\|_\infty \times \left[ \frac{(x-a)^2 + (b-x)^2}{2} \right]^{1-1/q} \left[ \frac{(3b-x-2a)(x-a)^2 + (b-x)^3}{6(b-a)} |f'(a)|^q + \frac{(x-a)^3 + (2b+x-3a)(b-x)^2}{6(b-a)} |f'(b)|^q \right]^{1/q}. \quad (7)$$

**Theorem 1.5** ([11, Theorem 2]). *Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function and  $a, b \in I$  with  $a < b$ . If  $|f'(x)|^q$  is convex on  $[a, b]$  and  $g : [a, b] \rightarrow \mathbb{R}$  is a continuous function, then for all  $x \in [a, b]$  we have*

$$\left| f(x) \int_a^b g(s) ds - \int_a^b f(s)g(s) ds \right| \leq \|g\|_\infty \left[ \frac{(x-a)^2 + (b-x)^2}{2} \right]^{1-1/q} \times \left[ \frac{(3b-2x-a)(x-a)^2 + 2(b-x)^3}{6(b-a)} |f'(a)|^q + \frac{2(x-a)^3 + (b+2x-3a)(b-x)^2}{6(b-a)} |f'(b)|^q \right]^{1/q}. \quad (8)$$

For more information on this topic, please refer to [1, 2, 4, 7, 9, 12, 13, 14, 15, 16] and plenty of references cited therein.

Our goal of this paper is to establish some new Hermite-Hadamard type inequalities for  $(\alpha, m)$ -convex functions.

## 2 Lemmas

For establishing new integral inequalities of Hermite-Hadamard type for  $(\alpha, m)$ -convex functions, we need the following lemmas.

**Lemma 2.1.** *Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in I$  with  $a < b$ ,  $m \in (0, 1]$ ,  $a \neq mb$ , and  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$ . If  $f', g \in L[\bar{a}, \bar{b}]$ , then, for all  $x \in [\bar{a}, \bar{b}]$ , we have*

$$f(a) \int_a^x g(s) ds + f(mb) \int_x^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds = \int_a^{mb} \int_x^t g(s)f'(t) ds dt. \quad (9)$$

**Lemma 2.2.** *Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in I$  with  $a < b$ ,  $m \in (0, 1]$ ,  $a \neq mb$ , and  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$ . If  $f', g \in L[\bar{a}, \bar{b}]$ , then for all  $x \in [\bar{a}, \bar{b}]$ , we have*

$$f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds = \int_a^{mb} S_g(t)f'(t) dt, \quad (10)$$

where

$$S_g(t) = \begin{cases} \int_{\bar{a}}^t g(s) ds, & t \in [\bar{a}, x), \\ -\int_t^{\bar{b}} g(s) ds, & t \in [x, \bar{b}] \end{cases} \quad \text{and} \quad S_1(t) = \begin{cases} t - \bar{a}, & t \in [\bar{a}, x), \\ \bar{b} - t, & t \in [x, \bar{b}]. \end{cases}$$

*Proof of Lemmas 2.1 and 2.2.* These lemmas can be deduced directly from integrating by part the right-hand sides of (9) and (10) respectively.  $\square$

### 3 Some new integral inequalities of Hermite-Hadamard type for $(\alpha, m)$ -convex functions

Now we are in a position to establish some new integral inequalities of Hermite-Hadamard type for functions whose derivatives are  $(\alpha, m)$ -convex.

**Theorem 3.1.** *Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in [0, b^*]$  with  $a < b$ ,  $(\alpha, m) \in (0, 1]^2$ ,  $a \neq mb$ , and  $f' \in L[\bar{a}, \bar{b}]$ . If  $|f'(x)|^q$  for  $q \geq 1$  is  $(\alpha, m)$ -convex on  $[0, b]$ ,  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$  is a continuous function, and  $x \in [\bar{a}, \bar{b}]$ , then*

$$\begin{aligned} & \left| f(a) \int_a^x g(s) ds + f(mb) \int_x^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| \\ & \leq \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \left\{ m \frac{(x-a)^2 + (mb-x)^2}{2} |f'(b)|^q \right. \\ & \quad \left. + \frac{|f'(a)|^q - m|f'(b)|^q}{(\alpha+1)(\alpha+2)} \left[ 2(mb-x)^2 \left( \frac{mb-x}{mb-a} \right)^\alpha \right. \right. \\ & \quad \left. \left. + (mb-a)[(\alpha+1)(x-a) - (mb-x)] \right] \right\}^{1/q}. \quad (11) \end{aligned}$$

*Proof.* Using Lemma 2.1, Hölder inequality, and the  $(\alpha, m)$ -convexity of  $|f'|^q$ , it follows that

$$\left| f(a) \int_a^x g(s) ds + f(mb) \int_x^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right|$$

$$\begin{aligned}
 &\leq \int_{\bar{a}}^{\bar{b}} \left| \int_x^t g(s) \, ds \right| |f'(t)| \, dt \leq \left( \int_{\bar{a}}^{\bar{b}} \left| \int_x^t g(s) \, ds \right| dt \right)^{1-1/q} \\
 &\quad \times \left( \int_{\bar{a}}^{\bar{b}} \left| \int_x^t g(s) \, ds \right| |f'(t)|^q \, dt \right)^{1/q} \leq \|g\|_\infty \left[ \int_{\bar{a}}^{\bar{b}} |t-x| \, dt \right]^{1-1/q} \\
 &\quad \times \left[ \int_{\bar{a}}^{\bar{b}} |t-x| |f'(t)|^q \, dt \right]^{1/q} = \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \\
 &\quad \times \left[ \int_{\bar{a}}^{\bar{b}} |t-x| \left| f' \left( \frac{\bar{b}-t}{\bar{b}-\bar{a}} \bar{a} + \frac{t-\bar{a}}{\bar{b}-\bar{a}} \bar{b} \right) \right|^q \, dt \right]^{1/q} \\
 &\leq \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \left\{ \int_{\bar{a}}^{\bar{b}} |t-x| \left[ \left( \frac{mb-t}{mb-a} \right)^\alpha |f'(a)|^q \right. \right. \\
 &\quad \left. \left. + m \left( 1 - \left( \frac{mb-t}{mb-a} \right)^\alpha \right) |f'(b)|^q \right] dt \right\}^{1/q} \\
 &= \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \left\{ m |f'(b)|^q \int_{\bar{a}}^{\bar{b}} |t-x| \, dt \right. \\
 &\quad \left. + (|f'(a)|^q - m |f'(b)|^q) \int_{\bar{a}}^{\bar{b}} |t-x| \left( \frac{mb-t}{mb-a} \right)^\alpha \, dt \right\}^{1/q}
 \end{aligned}$$

for  $x \in [\bar{a}, \bar{b}]$ . Substituting  $\int_{\bar{a}}^{\bar{b}} |t-x| \, dt = \frac{1}{2}[(x-a)^2 + (mb-x)^2]$  and

$$\begin{aligned}
 &\int_{\bar{a}}^{\bar{b}} |t-x| \left( \frac{mb-t}{mb-a} \right)^\alpha \, dt = \frac{1}{(\alpha+1)(\alpha+2)} \\
 &\quad \times \left\{ 2(mb-x)^2 \left( \frac{mb-x}{mb-a} \right)^\alpha + (mb-a)[(\alpha+1)(x-a) - (mb-x)] \right\}
 \end{aligned}$$

into the above inequality leads to (11). Theorem 3.1 is thus proved.  $\square$

**Remark 3.2.** The inequality (7) is a special case of (11) applied to  $m = \alpha = 1$ .

**Corollary 3.1.** Under the conditions of Theorem 3.1, if  $q = 1$ , we have

$$\begin{aligned}
 &\left| f(a) \int_a^x g(s) \, ds + f(mb) \int_x^{mb} g(s) \, ds - \int_a^{mb} f(s)g(s) \, ds \right| \leq \|g\|_\infty \\
 &\quad \times \left\{ m \frac{(x-a)^2 + (mb-x)^2}{2} |f'(b)| + \left[ 2(mb-x)^2 \left( \frac{mb-x}{mb-a} \right)^\alpha \right. \right. \\
 &\quad \left. \left. + (mb-a)[(\alpha+1)(x-a) - (mb-x)] \right] \frac{|f'(a)| - m|f'(b)|}{(\alpha+1)(\alpha+2)} \right\}. \quad (12)
 \end{aligned}$$

**Theorem 3.3.** Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in [0, b^*]$  with  $a < b$ ,  $(\alpha, m) \in (0, 1]^2$ ,  $a \neq mb$ , and  $f' \in L[\bar{a}, \bar{b}]$ . If  $|f'(x)|^q$  for  $q > 1$

is  $(\alpha, m)$ -convex on  $[0, b]$ ,  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$  is a continuous function, then, for all  $x \in [\bar{a}, \bar{b}]$ , we have

$$\begin{aligned} & \left| f(a) \int_a^x g(s) ds + f(mb) \int_x^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| \\ & \leq \|g\|_\infty \left\{ \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} + |mb-x|^{(2q-1)/(q-1)}] \right\}^{1-1/q} \\ & \quad \times \left\{ \frac{|mb-a|}{\alpha+1} [|f'(a)|^q + \alpha m |f'(b)|^q] \right\}^{1/q}. \quad (13) \end{aligned}$$

*Proof.* By Lemma 2.1, Hölder inequality, and the  $(\alpha, m)$ -convexity of  $|f'|^q$ , for  $x \in [\bar{a}, \bar{b}]$ , it follows that

$$\begin{aligned} & \left| f(a) \int_a^x g(s) ds + f(mb) \int_x^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| \\ & \leq \int_{\bar{a}}^{\bar{b}} \left| \int_x^t g(s) ds \right| |f'(t)| dt \leq \left( \int_{\bar{a}}^{\bar{b}} \left| \int_x^t g(s) ds \right|^{q/(q-1)} dt \right)^{1-1/q} \\ & \quad \times \left( \int_{\bar{a}}^{\bar{b}} |f'(t)|^q dt \right)^{1/q} \leq \|g\|_\infty \left[ \int_{\bar{a}}^{\bar{b}} |t-x|^{q/(q-1)} dt \right]^{1-1/q} \left[ \int_{\bar{a}}^{\bar{b}} |f'(t)|^q dt \right]^{1/q} \\ & = \|g\|_\infty \left( \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} + |mb-x|^{(2q-1)/(q-1)}] \right)^{1-1/q} \\ & \quad \times \left[ \int_{\bar{a}}^{\bar{b}} \left| f' \left( \frac{\bar{b}-t}{\bar{b}-\bar{a}} \bar{a} + \frac{t-\bar{a}}{\bar{b}-\bar{a}} \bar{b} \right) \right|^q dt \right]^{1/q} \\ & \leq \|g\|_\infty \left( \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} + |mb-x|^{(2q-1)/(q-1)}] \right)^{1-1/q} \\ & \quad \times \left\{ \int_{\bar{a}}^{\bar{b}} \left[ \left( \frac{mb-t}{mb-a} \right)^\alpha |f'(a)|^q + m \left( 1 - \left( \frac{mb-t}{mb-a} \right)^\alpha \right) |f'(b)|^q \right] dt \right\}^{1/q} \\ & = \|g\|_\infty \left\{ \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} + |mb-x|^{(2q-1)/(q-1)}] \right\}^{1-1/q} \\ & \quad \times \left\{ \frac{|mb-a|}{\alpha+1} [|f'(a)|^q + \alpha m |f'(b)|^q] \right\}^{1/q}. \end{aligned}$$

The proof of Theorem 3.3 is complete.  $\square$

**Corollary 3.2.** Under the conditions of Theorem 3.3, if  $m = \alpha = 1$ , we have

$$\left| f(a) \int_a^x g(s) ds + f(b) \int_x^b g(s) ds - \int_a^b f(s)g(s) ds \right|$$

$$\begin{aligned} &\leq \|g\|_\infty \left\{ \frac{q-1}{2q-1} [(x-a)^{(2q-1)/(q-1)} + (b-x)^{(2q-1)/(q-1)}] \right\}^{1-1/q} \\ &\quad \times \left\{ \frac{b-a}{2} [|f'(a)|^q + |f'(b)|^q] \right\}^{1/q}. \quad (14) \end{aligned}$$

**Theorem 3.4.** Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in [0, b^*]$  with  $a < b$ ,  $(\alpha, m) \in (0, 1]^2$ ,  $a \neq mb$ , and  $f' \in L[\bar{a}, \bar{b}]$ . If  $|f'(x)|^q$  for  $q \geq 1$  is  $(\alpha, m)$ -convex on  $[0, b]$ ,  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$  is a continuous function, and  $x \in [\bar{a}, \bar{b}]$ , then

$$\begin{aligned} &\left| f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| \leq \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \\ &\quad \times \left\{ m \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right] |f'(b)|^q + \left[ \alpha(mb-x)^2 - (\alpha+2) \right. \right. \\ &\quad \left. \left. \times (mb-x)(x-a) \right] \left( \frac{mb-x}{mb-a} \right)^\alpha + (mb-a)^2 \right\} \frac{|f'(a)|^q - m|f'(b)|^q}{(\alpha+1)(\alpha+2)} \Bigg\}^{1/q}. \quad (15) \end{aligned}$$

*Proof.* Using Lemma 2.2 yields  $|S_g(t)| \leq \|g\|_\infty S_1(t)$  for  $t \in [\bar{a}, \bar{b}]$ . By Hölder inequality and the  $(\alpha, m)$ -convexity of  $|f'|^q$ , it follows that

$$\begin{aligned} &\left| f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| \leq \int_{\bar{a}}^{\bar{b}} |S_g(t)| |f'(t)| dt \\ &\leq \|g\|_\infty \left[ \int_{\bar{a}}^{\bar{b}} S_1(t) dt \right]^{1-1/q} \left[ \int_{\bar{a}}^{\bar{b}} S_1(t) |f'(t)|^q dt \right]^{1/q} \\ &\leq \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \left\{ m|f'(b)|^q \int_{\bar{a}}^{\bar{b}} S_1(t) dt \right. \\ &\quad \left. + (|f'(a)|^q - m|f'(b)|^q) \int_{\bar{a}}^{\bar{b}} S_1(t) \left( \frac{mb-t}{mb-a} \right)^\alpha dt \right\}^{1/q} \\ &\leq \|g\|_\infty \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right]^{1-1/q} \left\{ m \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right] |f'(b)|^q \right. \\ &\quad \left. + \frac{1}{(\alpha+1)(\alpha+2)} \left[ \left( \frac{mb-x}{mb-a} \right)^\alpha [\alpha(mb-x)^2 \right. \right. \right. \\ &\quad \left. \left. \left. - (\alpha+2)(mb-x)(x-a) \right] + (mb-a)^2 \right] (|f'(a)|^q - m|f'(b)|^q) \right\}^{1/q}. \end{aligned}$$

The proof of Theorem 3.4 is complete. □

**Remark 3.5.** The inequality (8) is a special case of (15) applied to  $m = \alpha = 1$ .

**Corollary 3.3.** Under the conditions of Theorem 3.4, if  $q = 1$ , we have

$$\begin{aligned} \left| f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| &\leq \|g\|_\infty \left\{ m \left[ \frac{(x-a)^2 + (mb-x)^2}{2} \right] \right. \\ &\times |f'(b)| + \frac{1}{(\alpha+1)(\alpha+2)} \left[ \alpha(mb-x)^2 - (\alpha+2)(mb-a)(x-a) \right. \\ &\left. \left. \times \left( \frac{mb-x}{mb-a} \right)^\alpha + (mb-a)^2 \right] (|f'(a)| - m|f'(b)|) \right\}. \quad (16) \end{aligned}$$

**Theorem 3.6.** Let  $f : [0, b^*] \rightarrow \mathbb{R}$  be a differentiable function,  $a, b \in [0, b^*]$  with  $a < b$ ,  $(\alpha, m) \in (0, 1]^2$ ,  $a \neq mb$ , and  $f' \in L[\bar{a}, \bar{b}]$ . If  $|f'(x)|^q$  for  $q > 1$  is  $(\alpha, m)$ -convex on  $[0, b]$ ,  $g : [\bar{a}, \bar{b}] \rightarrow \mathbb{R}$  is a continuous function, and  $x \in [\bar{a}, \bar{b}]$ , then

$$\begin{aligned} \left| f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| &\leq \|g\|_\infty \left\{ \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} \right. \\ &\left. + |mb-x|^{(2q-1)/(q-1)}] \right\}^{1-1/q} \left\{ \frac{|mb-a|}{\alpha+1} [|f'(a)|^q + \alpha m |f'(b)|^q] \right\}^{1/q}. \quad (17) \end{aligned}$$

*Proof.* By Lemma 2.2, Hölder inequality, and the  $(\alpha, m)$ -convexity of  $|f'|^q$ , for  $x \in [\bar{a}, \bar{b}]$ , it follows that

$$\begin{aligned} \left| f(x) \int_a^{mb} g(s) ds - \int_a^{mb} f(s)g(s) ds \right| &\leq \|g\|_\infty \left\{ \int_{\bar{a}}^{\bar{b}} [S_1(t)]^{q/(q-1)} dt \right\}^{1-1/q} \\ &\times \left\{ \int_{\bar{a}}^{\bar{b}} |f'(t)|^q dt \right\}^{1/q} \leq \|g\|_\infty \left\{ \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} \right. \\ &\left. + |mb-x|^{(2q-1)/(q-1)}] \right\}^{1-1/q} \left[ \int_{\bar{a}}^{\bar{b}} \left| f' \left( \frac{\bar{b}-t}{\bar{b}-\bar{a}} \bar{a} + \frac{t-\bar{a}}{\bar{b}-\bar{a}} \bar{b} \right) \right|^q dt \right]^{1/q} \\ &= \|g\|_\infty \left\{ \frac{q-1}{2q-1} [|x-a|^{(2q-1)/(q-1)} + |mb-x|^{(2q-1)/(q-1)}] \right\}^{1-1/q} \\ &\times \left\{ \frac{|mb-a|}{\alpha+1} [|f'(a)|^q + \alpha m |f'(b)|^q] \right\}^{1/q}. \end{aligned}$$

The proof of Theorem 3.6 is complete.  $\square$

**Corollary 3.4.** Under the conditions of Theorem 3.6, if  $m = \alpha = 1$ , we have

$$\begin{aligned} \left| f(x) \int_a^b g(s) ds - \int_a^b f(s)g(s) ds \right| &\leq \|g\|_\infty \left\{ \frac{q-1}{2q-1} [(x-a)^{(2q-1)/(q-1)} \right. \\ &\left. + (b-x)^{(2q-1)/(q-1)}] \right\}^{1-1/q} \left\{ \frac{b-a}{2} [|f'(a)|^q + |f'(b)|^q] \right\}^{1/q}. \quad (18) \end{aligned}$$



## 4 Two open problems

Finally, we would like to pose two open problems as follows.

1. When the function  $g$  is symmetric with respect to the midpoint  $\frac{\bar{a}+\bar{b}}{2}$  of the interval  $[\bar{a}, \bar{b}]$ , can one simplify the inequalities (11), (13), (15), and (17) in Theorems 3.1, 3.3, 3.4, and 3.6 respectively?
2. Can these inequalities established in this paper be applied to construct inequalities of special means of two positive numbers, as done in [6, 14] and other papers?

## References

- [1] R.-F. Bai, F. Qi, and B.-Y. Xi, *Hermite-Hadamard type inequalities for the  $m$ - and  $(\alpha, m)$ -logarithmically convex functions*, *Filomat* **27** (2013), no. 1, 1–7.
- [2] S.-P. Bai, S.-H. Wang, and F. Qi, *Some Hermite-Hadamard type inequalities for  $n$ -time differentiable  $(\alpha, m)$ -convex functions*, *J. Inequal. Appl.* 2012, **2012**:267, 11 pages; Available online at <http://dx.doi.org/10.1186/1029-242X-2012-267>.
- [3] M. K. Bakula, M. E. Özdemir, and J. Pečarić, *Hadamard type inequalities for  $m$ -convex and  $(\alpha, m)$ -convex functions*, *J. Inequal. Pure Appl. Math.* **9** (2008), no. 4, Art. 96, 12 pages; Available online at <http://www.emis.de/journals/JIPAM/article1032.html>.
- [4] L. Chun and F. Qi, *Integral inequalities of Hermite-Hadamard type for functions whose 3rd derivatives are  $s$ -convex*, *Appl. Math.* **3** (2012), no. 11, 1680–1685; Available online at <http://dx.doi.org/10.4236/am.2012.311232>.
- [5] S. S. Dragomir, *On some new inequalities of Hermite-Hadamard type for  $m$ -convex functions*, *Tamkang J. Math.* **33** (2002), no. 1, 45–55; Available online at <http://dx.doi.org/10.5556/j.tkjm.33.2002.45-55>.
- [6] S. S. Dragomir and R. P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, *Appl. Math. Lett.* **11** (1998), no. 5, 91–95; Available online at [http://dx.doi.org/10.1016/S0893-9659\(98\)00086-X](http://dx.doi.org/10.1016/S0893-9659(98)00086-X).
- [7] W.-D. Jiang, D.-W. Niu, Y. Hua, and F. Qi, *Generalizations of Hermite-Hadamard inequality to  $n$ -time differentiable functions which are  $s$ -convex*

- in the second sense*, Analysis (Munich) **32** (2012), no. 3, 209–220; Available online at <http://dx.doi.org/10.1524/anly.2012.1161>.
- [8] V. G. Miheşan, *A generalization of the convexity*, Seminar on Functional Equations, Approx. Convex, Cluj-Napoca, 1993. (Romania)
- [9] F. Qi, Z.-L. Wei, and Q. Yang, *Generalizations and refinements of Hermite-Hadamard's inequality*, Rocky Mountain J. Math. **35** (2005), no. 1, 235–251; Available online at <http://dx.doi.org/10.1216/rmj/1181069779>.
- [10] G. Toader, *Some generalizations of the convexity*, Univ. Cluj-Napoca, Cluj-Napoc. 1985, 329–338.
- [11] K.-L. Tseng, S.-R. Hwang, G.-S. Yang, and J.-C. Lo, *Two inequalities for differentiable mappings and applications to weighted trapezoidal formula, weighted midpoint formula and random variable*, Math. Comput. Modelling **53** (2011), no. 1-2, 179–188; Available online at <http://dx.doi.org/10.1016/j.mcm.2010.08.002>.
- [12] S.-H. Wang, B.-Y. Xi, and F. Qi, *Some new inequalities of Hermite-Hadamard type for  $n$ -time differentiable functions which are  $m$ -convex*, Analysis (Munich) **32** (2012), no. 3, 247–262; Available online at <http://dx.doi.org/10.1524/anly.2012.1167>.
- [13] B.-Y. Xi, R.-F. Bai, and F. Qi, *Hermite-Hadamard type inequalities for the  $m$ - and  $(\alpha, m)$ -geometrically convex functions*, Aequationes Math. **84** (2012), no. 3, 261–269; Available online at <http://dx.doi.org/10.1007/s00010-011-0114-x>.
- [14] B.-Y. Xi and F. Qi, *Some integral inequalities of Hermite-Hadamard type for convex functions with applications to means*, J. Funct. Spaces Appl. **2012** (2012), Article ID 980438, 14 pages; Available online at <http://dx.doi.org/10.1155/2012/980438>.
- [15] B.-Y. Xi, S.-H. Wang, and F. Qi, *Some inequalities of Hermite-Hadamard type for functions whose 3rd derivatives are  $P$ -convex*, Appl. Math. **3** (2012), no. 12, 1898–1902; Available online at <http://dx.doi.org/10.4236/am.2012.312260>.
- [16] T.-Y. Zhang, A.-P. Ji, and F. Qi, *On integral inequalities of Hermite-Hadamard type for  $s$ -geometrically convex functions*, Abstr. Appl. Anal. **2012** (2012), Article ID 560586, 14 pages; Available online at <http://dx.doi.org/10.1155/2012/560586>.