

# ON INCOMPLETE INFINITE DIRECT PRODUCT OF $W^*$ -ALGEBRAS

TEISHIRÔ SAITÔ

(Received March 5, 1958)

The purpose of this paper is to show some results on incomplete infinite direct products of  $W^*$ -algebras. As we can see from [7: Part IV, Chap. 7], the behaviour of complete infinite direct product of  $W^*$ -algebras is not caught. Hence we shall do not pursue the complete infinite direct product of  $W^*$ -algebras.<sup>1)</sup>

**1. Preliminaries.** Throughout this paper, let  $\mathbf{I}$  be a set of indices of arbitrary size and let for each  $i \in \mathbf{I}$  a  $W^*$ -algebra  $\mathbf{M}_i$  on a Hilbert space  $\mathfrak{H}_i$  (or, sometimes  $\mathfrak{H}$ ) be given. By  $\mathbf{B}(\mathfrak{H})$  we denote the full operator algebra on a Hilbert space  $\mathfrak{H}$ . We owe other notations and terminology used here to [5] and [7]. J. von Neumann introduced the concept of infinite direct product of full operator algebras, in which the infinite direct product of underlying Hilbert spaces has important meaning. For convenience, we shall sketch the outlines of general theory in [7]. For any pair of  $C$ -sequences

$$X = (x_i : i \in \mathbf{I}), \quad Y = (y_i : i \in \mathbf{I}) \quad (x_i, y_i \in \mathfrak{H}_i \text{ for each } i \in \mathbf{I})$$

we associate

$$(X, Y) = \prod_{i \in \mathbf{I}} (x_i, y_i).$$

Then, for linear combinations  $\sum_{j=1}^m \lambda_j X_j, \sum_{k=1}^n \mu_k Y_k$  we define the inner product

$$\left( \sum_{j=1}^m \lambda_j X_j, \sum_{k=1}^n \mu_k Y_k \right) = \sum_{j=1}^m \sum_{k=1}^n \lambda_j \overline{\mu_k} (X_j, Y_k),$$

and get a prehilbert space. Its completion is called the complete infinite direct product of  $\mathfrak{H}_i$  and denoted by  $\prod_{i \in \mathbf{I}} \bigotimes \mathfrak{H}_i$ . An element in  $\prod_{i \in \mathbf{I}} \bigotimes \mathfrak{H}_i$  determined by  $(x_i : i \in \mathbf{I})$  is denoted by  $\prod_{i \in \mathbf{I}} \bigotimes x_i$ . Next, the family of all  $C_0$ -sequences is divided into equivalence-classes by the relation " $\approx$ " ([7: Chap. 3]). We denote equivalence-class of a given  $C_0$ -sequence  $(x_i : i \in \mathbf{I})$  by  $\mathfrak{C}(x_i : i \in \mathbf{I})$  (or, simply by  $\mathfrak{C}$  if there is no confusion). Then  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{C}} \mathfrak{H}_i$  means the closed linear

---

1) In preparing this paper, the author has received valuable advice and suggestions from Dr. Z. Takeda and Dr. Y. Misonou which he gratefully acknowledges.

set determined by all  $\prod_{i \in \mathbf{I}} \otimes x_i$ , where  $(x_i : i \in \mathbf{I}) \in \mathfrak{G}$  and is called the  $\mathfrak{G}$ -adic incomplete infinite direct product. Then the following results are obtained.

**THEOREM A** ([7: LEMMA 4.1.4]). *Let  $(x_i^0 : i \in \mathbf{I})$  be a  $C_0$ -sequence with  $\|x_i^0\| = 1$  for all  $i \in \mathbf{I}$  and  $d_i$  be the dimension of  $\mathfrak{H}_i$  for each  $i \in \mathbf{I}$ . Choosing a complete orthonormal set  $\{x_{i,j} : j \in \mathbf{J}_i\}$  of  $\mathfrak{H}_i$  such as  $0 \in \mathbf{J}_i$  and  $x_{i,0} = x_i^0$ , where the cardinal of the set of indices  $\mathbf{J}_i$  is equal to  $d_i$  for each  $i \in \mathbf{I}$ . If we construct  $\prod_{i \in \mathbf{I}} \otimes x_{i,j}$  where  $j \in \mathbf{J}_i = 0$  except for a finite number of  $i$ 's, then the totality of such  $\prod_{i \in \mathbf{I}} \otimes x_{i,j}$  is a complete orthonormal set in  $\prod_{i \in \mathbf{I}} \otimes^{\mathfrak{G}} \mathfrak{H}_i$  for  $\mathfrak{G} = \mathfrak{G}(x_i^0 : i \in \mathbf{I})$ .*

**THEOREM B** ([7: THEOREM VI]). *Let  $\mathbf{J}$  be a set of indices. If  $\mathbf{I}$  be divided into mutually disjoint sets  $\mathbf{I}_j (j \in \mathbf{J})$ , and if  $(x_i^0 : i \in \mathbf{I})$  be a  $C_0$ -sequence, we can form the equivalence classes  $\mathfrak{G} = \mathfrak{G}(x_i^0 : i \in \mathbf{I})$  (in  $\prod_{i \in \mathbf{I}} \otimes \mathfrak{H}_i$ ),  $\mathfrak{G}_j = \mathfrak{G}(x_i : i \in \mathbf{I}_j)$  (in  $\prod_{i \in \mathbf{I}_j} \otimes \mathfrak{H}_i$ ) and  $\mathfrak{G}_0 = \mathfrak{G}(\prod_{i \in \mathbf{I}_j} \otimes x_i^0 : j \in \mathbf{J})$  (in  $\prod_{j \in \mathbf{J}} (\prod_{i \in \mathbf{I}_j} \otimes \mathfrak{H}_i)$ ). Then the classes  $\mathfrak{G}_j, \mathfrak{G}_0$  depend on  $\mathfrak{G}$  only and there exists a unique isomorphism of  $\prod_{i \in \mathbf{I}} \otimes^{\mathfrak{G}} \mathfrak{H}_i$  and  $\prod_{j \in \mathbf{J}} \otimes^{\mathfrak{G}_0} (\prod_{i \in \mathbf{I}_j} \otimes^{\mathfrak{G}_j} \mathfrak{H}_i)$  such that  $\prod_{i \in \mathbf{I}} \otimes x_i$  corresponds to  $\prod_{j \in \mathbf{J}} (\prod_{i \in \mathbf{I}_j} \otimes x_i)$  for all  $C_0$ -sequences  $(x_i : i \in \mathbf{I}) \in \mathfrak{G}$ .*

**2. Infinite direct product of  $W^*$ -algebras.** First we define the infinite direct product of  $W^*$ -algebras  $\mathbf{M}_i (i \in \mathbf{I})$  as J. von Neumann has done for full operator algebras. For any fixed  $i_0 \in \mathbf{I}$  every  $A_{i_0} \in \mathbf{M}_{i_0}$  is extended to an operator  $\bar{A}_{i_0}$  on  $\prod_{i \in \mathbf{I}} \otimes \mathfrak{H}_i$  such that for each  $\prod_{i \in \mathbf{I}} \otimes x_i \in \prod_{i \in \mathbf{I}} \otimes \mathfrak{H}_i$ ,

$$\bar{A}_{i_0} (\prod_{i \in \mathbf{I}} \otimes x_i) = A_{i_0} x_{i_0} \otimes (\prod_{i \in \mathbf{I}, i \neq i_0} \otimes x_i).$$

The set of all extended operators  $\bar{A}_{i_0}$  of elements of  $\mathbf{M}_{i_0}$  is denoted by  $\bar{\mathbf{M}}_{i_0}$ . Then we have the following

**DEFINITION 1.** Let  $\mathbf{I}, \mathbf{M}_i$  and  $\mathfrak{H}_i (i \in \mathbf{I})$  be as above. Then the  $W^*$ -algebra on  $\prod_{i \in \mathbf{I}} \otimes \mathfrak{H}_i$  generated by all  $\bar{A}_i \in \bar{\mathbf{M}}_i$  with  $i \in \mathbf{I}$  is called *the complete infinite direct product of  $\mathbf{M}_i$  on  $\mathfrak{H}_i$*  and denoted by  $\prod_{i \in \mathbf{I}} \otimes \mathbf{M}_i$ .

Since every  $\bar{A}_{i_0} (i_0 \in \mathbf{I})$  commutes with the projection on  $\prod_{i \in \mathbf{I}} \otimes^{\mathfrak{G}} \mathfrak{H}_i$ ,  $\prod_{i \in \mathbf{I}} \otimes^{\mathfrak{G}} \mathfrak{H}_i$  is stable under  $\prod_{i \in \mathbf{I}} \otimes \mathbf{M}_i$ ; so any operator  $\mathbf{A} \in \prod_{i \in \mathbf{I}} \otimes \mathbf{M}_i$  may be considered as an operator in each  $\prod_{i \in \mathbf{I}} \otimes^{\mathfrak{G}} \mathfrak{H}_i$  for each equivalence-class  $\mathfrak{G}$  (cf. [7]). Then we have:

DEFINITION 2. For any equivalence-class  $\mathcal{C}$  the  $W^*$ -algebra  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}} \mathbf{M}_i$  restricted in  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}} \mathfrak{H}_i$  is denoted by  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathcal{C}} \mathbf{M}_i$  and we call it  $\mathcal{C}$ -adic incomplete infinite direct product of  $\mathbf{M}_i$ .

Now we have the following

THEOREM 1. If all  $\mathbf{M}_i$  are factors, then for each equivalence-class  $\mathcal{C}$  the  $\mathcal{C}$ -adic incomplete infinite direct product  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathcal{C}} \mathbf{M}_i$  is a factor.

To prove this theorem we provide following lemmas.

LEMMA 1. If  $\mathbf{M}_i$  ( $i \in \mathbf{I}$ ) are  $W^*$ -algebras on a Hilbert space  $\mathfrak{H}$  satisfying the following conditions:

- (i)  $\mathbf{M}_i \subset \mathbf{M}_j'$  for  $i \neq j, i, j \in \mathbf{I}$ ;
- (ii)  $\mathbf{R}(\mathbf{M}_i : i \in \mathbf{I}) = \mathbf{B}(\mathfrak{H})$ .

Then every  $\mathbf{M}_i$  is a factor.

PROOF. 
$$\mathbf{M}_i \cap \mathbf{M}_i' \subset \left[ \bigcap_{j \in \mathbf{I}, j \neq i} \mathbf{M}_j' \right] \cap \mathbf{M}_i' = \bigcap_{j \in \mathbf{I}} \mathbf{M}_j'$$

$$= \mathbf{R}(\mathbf{M}_j : j \in \mathbf{I})' = \mathbf{B}(\mathfrak{H})' = (\alpha \mathbf{I}) \text{ for each } i \in \mathbf{I}.$$

Hence we have  $\mathbf{M}_i \cap \mathbf{M}_i' = (\alpha \mathbf{I})$  for each  $i \in \mathbf{I}$ .

LEMMA 2. Let  $\mathbf{M}_i$  be a factor on a Hilbert space  $\mathfrak{H}$  for each  $i \in \mathbf{I}$  satisfying the following conditions: For each  $i \in \mathbf{I}$  there exists  $\mathbf{P}_i$ , a  $W^*$ -algebra on  $\mathfrak{H}$ , such that

- (i)  $\mathbf{M}_i \subset \mathbf{P}_i, \mathbf{M}_j \subset \mathbf{P}_i'$  for any  $j \neq i, j \in \mathbf{I}$ ,
- (ii)  $\mathbf{P}_i$  and  $\mathbf{P}_i'$  are both normal, and
- (iii)  $\mathbf{R}(\mathbf{P}_i : i \in \mathbf{I}) = \mathbf{B}(\mathfrak{H})$ .

Then  $\mathbf{R}(\mathbf{M}_i : i \in \mathbf{I})$  is a factor.

PROOF. From (i), for each  $i, j \in \mathbf{I}, i \neq j, \mathbf{M}_i \subset \mathbf{M}_j'$ . We have, by (ii),  $\mathbf{R}(\mathbf{M}_i, \mathbf{M}_i' \cap \mathbf{P}_i)' \cap \mathbf{P}_i = \mathbf{M}_i' \cap [(\mathbf{M}_i' \cap \mathbf{P}_i)' \cap \mathbf{P}_i] = \mathbf{M}_i' \cap \mathbf{M}_i = (\alpha \mathbf{I})$ , for each  $i \in \mathbf{I}$ . Therefore, considering  $\mathbf{R}(\mathbf{M}_i, \mathbf{M}_i' \cap \mathbf{P}_i) \subset \mathbf{P}_i$ , we get

$$\mathbf{R}(\mathbf{M}_i, \mathbf{M}_i' \cap \mathbf{P}_i) = [\mathbf{R}(\mathbf{M}_i, \mathbf{M}_i' \cap \mathbf{P}_i)' \cap \mathbf{P}_i] \cap \mathbf{P}_i = (\alpha \mathbf{I})' \cap \mathbf{P}_i = \mathbf{P}_i,$$
 for each  $i \in \mathbf{I}$ . Hence

$$\begin{aligned} & \mathbf{R}(\mathbf{R}(\mathbf{M}_i : i \in \mathbf{I}), \mathbf{R}(\mathbf{M}_i' \cap \mathbf{P}_i : i \in \mathbf{I})) \\ &= \mathbf{R}(\mathbf{M}_i, \mathbf{M}_i' \cap \mathbf{P}_i : i \in \mathbf{I}) = \mathbf{R}(\mathbf{R}(\mathbf{M}_i, \mathbf{M}_i \cap \mathbf{P}_i) : i \in \mathbf{I}) \\ &= \mathbf{R}(\mathbf{P}_i : i \in \mathbf{I}) = \mathbf{B}(\mathfrak{H}). \end{aligned}$$

Now, for any  $i \in \mathbf{I}, \mathbf{M}_i$  commutes with all  $\mathbf{M}_j' \cap \mathbf{P}_j, (j \in \mathbf{I})$  by (i). Thus  $\mathbf{R}(\mathbf{M}_i : i \in \mathbf{I})$  commutes with  $\mathbf{R}(\mathbf{M}_i' \cap \mathbf{P}_i : i \in \mathbf{I})$ . Hence, by Lemma 1,  $\mathbf{R}(\mathbf{M}_i : i \in \mathbf{I})$  is a factor.

From [7] we quote the following result ([7: Theorems VIII and X]).

LEMMA 3. Let  $\mathbf{B}_i$  be a full operator algebra on a Hilbert space  $\mathfrak{H}_i$  for each  $i \in \mathbf{I}$ . Then  $\overline{\mathbf{B}}_i$  is isomorphic to  $\mathbf{B}_i$  for each  $i \in \mathbf{I}$ , and moreover, each  $\mathfrak{C}$ -adic incomplete infinite direct product  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathbf{B}_i$  coincides with  $\mathbf{B}(\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathfrak{H}_i)$ .

PROOF OF THEOREM 1. Let each  $\mathbf{B}_i$  be the full operator algebra on each  $\mathfrak{H}_i$ . Put  $\overline{\mathbf{B}}_i = \mathbf{P}_i$ . By Lemma 3 and [5: Lemma 11.2.2] all  $\mathbf{P}_i, \mathbf{P}_i'$  are normal. It is clear that if  $i, j \in \mathbf{I}, i \neq j, \overline{\mathbf{M}}_i \subset \mathbf{P}_i$  and  $\overline{\mathbf{M}}_j \subset \mathbf{P}_j'$ . Thus, by Lemmas 2 and 3, we see that  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathbf{M}_i$  is a factor for each  $\mathfrak{C}$ .

Now, let  $\mathbf{M}_i$  be a  $W^*$ -algebra on a Hilbert space  $\mathfrak{H}_i$  with normal state  $\sigma_i$  for each  $i \in \mathbf{I}$ . Z. Takeda [10] defined the restricted infinite direct product of  $\mathbf{M}_i$  with  $\sigma_i$  denoted by  $\bigotimes_{i \in \mathbf{I}} (\mathbf{M}_i, \sigma_i)$  as follows: We construct a positive functional  $\sigma_0$  on the algebraical direct product of  $\mathbf{M}_i$  such that

$$\sigma_0(\dots \otimes I_{i_p} \otimes A_{i_q} \otimes \dots \otimes A_{i_r} \otimes I_{i_s} \otimes \dots) = \sigma_{i_q}(A_{i_q}) \dots \sigma_{i_r}(A_{i_r}).$$

The weak closure of the representation of the algebraical direct product of  $\mathbf{M}_i$  by  $\sigma_0$  is called the restricted infinite direct product of  $\mathbf{M}_i$  with  $\sigma_i$ . Then the next theorem states the relation between the restricted infinite direct product and the incomplete infinite direct product of  $W^*$ -algebras.

THEOREM 2. Let  $\mathbf{M}_i$  be a  $W^*$ -algebra on a Hilbert space  $\mathfrak{H}_i$  having a generating and separating vector  $x_i$  with unity norm for each  $i \in \mathbf{I}$ . Define a normal state  $\sigma_i$  by  $\sigma_i(A_i) = (A_i x_i, x_i)$  for every  $A_i \in \mathbf{M}_i$  for each  $i \in \mathbf{I}$ . Then  $\bigotimes_{i \in \mathbf{I}} (\mathbf{M}_i, \sigma_i)$  and  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathbf{M}_i$  are spatially isomorphic, where  $\mathfrak{C} = \mathfrak{C}(x_i : i \in \mathbf{I})$ .

PROOF. If we denote the representative space of  $\bigotimes_{i \in \mathbf{I}} (\mathbf{M}_i, \sigma_i)$  by  $\mathfrak{H}$ , the image of the algebraical direct product  $\mathbf{M}_0$  of  $\mathbf{M}_i (i \in \mathbf{I})$  is dense in  $\mathfrak{H}$ . If we denote the natural mapping from  $\bigotimes_{i \in \mathbf{I}} (\mathbf{M}_i, \sigma_i)$  into  $\mathfrak{H}$  by  $V$ , we have

$$\begin{aligned} & \|V(\dots \otimes I_{i_p} \otimes A_{i_q} \otimes \dots \otimes A_{i_r} \otimes I_{i_s} \otimes \dots)\| \\ &= \|A_{i_q} x_{i_q}\| \dots \|A_{i_r} x_{i_r}\| \end{aligned}$$

for any  $(\dots \otimes I_{i_p} \otimes A_{i_q} \otimes \dots \otimes A_{i_r} \otimes I_{i_s} \otimes \dots) \in \mathbf{M}_0$ . Now, consider the mapping  $W$  from  $V(\mathbf{M}_0)$  into  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathfrak{H}_i$  determined by

$$\begin{aligned} & W\{V(\dots \otimes I_{i_p} \otimes A_{i_q} \otimes \dots \otimes A_{i_r} \otimes I_{i_s} \otimes \dots)\} \\ &= (\dots x_{i_p} \otimes A_{i_q} x_{i_q} \otimes \dots \otimes A_{i_r} x_{i_r} \otimes x_{i_s} \otimes \dots) \end{aligned}$$

Then we see that this mapping is linear and isometric, for we have

$$\begin{aligned} & \|(\dots \otimes x_{i_p} \otimes A_{i_q} x_{i_q} \otimes \dots \otimes A_{i_r} x_{i_r} \otimes x_{i_s} \otimes \dots)\| \\ &= \|A_{i_q} x_{i_q}\| \dots \|A_{i_r} x_{i_r}\|. \end{aligned}$$

As each  $x_i$  is a generating vector for  $\mathbf{M}_i$ , we know that  $W$ -image of  $V(\mathbf{M}_0)$  is dense in  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\mathfrak{C}} \mathfrak{H}_i$  by [7: Lemma 4.1.2.]. Hence the mapping  $W$  can be

extended to a linear isometric mapping  $U$  from  $\mathfrak{H}$  onto  $\prod_{i \in I} \bigotimes_{\mathfrak{G}} \mathfrak{H}_i$ .

For any  $A = (\dots \otimes I_{i_p} \otimes A_{i_q} \otimes \dots \otimes A_{i_r} \otimes I_i, \dots) \in \mathbf{M}_0$  and  $y = \prod_{i \in I} \bigotimes_{\mathfrak{G}} y_i \in \prod_{i \in I} \bigotimes_{\mathfrak{G}} \mathfrak{H}_i$ , we have

$$(UAU^{-1})y = Ay.$$

Therefore it suffices to show that if  $A_\lambda \in \mathbf{M}_0$  is a directed set which converges to  $A \in \prod_{i \in I} (\mathbf{M}_i, \sigma_i)$  in the weak topology,  $UA_\lambda U^{-1}$  converges to  $UAU^{-1}$  in the weak topology. For any  $\varepsilon > 0$  and  $f \in \mathfrak{H}$  there exists a  $\lambda_0$  such that

$$|(Af, f) - (A_\lambda f, f)| < \varepsilon \text{ for } \lambda \geq \lambda_0.$$

On the other hand, for any  $x \in \prod_{i \in I} \bigotimes_{\mathfrak{G}} \mathfrak{H}_i$  there exists an  $f \in \mathfrak{H}$  such that  $x = Uf$ . Thus we have

$$|(UAU^{-1}x, x) - (UA_\lambda U^{-1}x, x)| = |(AU^{-1}x, U^{-1}x) - (A_\lambda U^{-1}x, U^{-1}x)| < \varepsilon$$

for  $\lambda \geq \lambda_0(x, \varepsilon)$ . This shows that  $UA_\lambda U^{-1}$  converges to  $UAU^{-1}$  in the weak topology.

**3. The type of the incomplete infinite direct product.** The type of the incomplete infinite direct product of  $W^*$ -algebras of the same type varies according as the choice of equivalence-class ([7: Part IV, Chap. 7]). The most favourable result on this type problem is to make the equivalence-class correspond to the type of the incomplete infinite direct product, and *vice versa*, but we have only a partial results. Namely, we have merely the following two results.

**THEOREM 3.** *If  $\mathbf{M}_i$  be a factor of type I on a Hilbert space  $\mathfrak{H}_i$  for each  $i \in \mathbf{I}$ , there exists an equivalence-class  $\mathfrak{G}$  of  $C_0$ -sequences such that  $\prod_{i \in I} \bigotimes_{\mathfrak{G}} \mathbf{M}_i$  is a factor of type I.*

**THEOREM 4.** *If  $\mathbf{M}_i$  be a factor which is not of type I on a Hilbert space  $\mathfrak{H}_i$  for each  $i \in \mathbf{I}$ , there exists an equivalence-class  $\mathfrak{G}$  of  $C_0$ -sequences such that  $\prod_{i \in I} \bigotimes_{\mathfrak{G}} \mathbf{M}_i$  is a factor of type III.*

To prove these theorems we need the following lemma.

**LEMMA 4.** *Let  $\mathbf{M}_{1i}, \mathbf{M}_{2i}$  be  $W^*$ -algebras on Hilbert spaces  $\mathfrak{H}_{1i}, \mathfrak{H}_{2i}$ , respectively. Consider three equivalence-classes of  $C_0$ -sequences  $\mathfrak{G}_1 = \mathfrak{G}(x_{1i} : i \in \mathbf{I})$ ,  $\mathfrak{G}_2 = \mathfrak{G}(x_{2i} : i \in \mathbf{I})$  and  $\mathfrak{G}_0 = \mathfrak{G}(x_{1i} \otimes x_{2i} : i \in \mathbf{I})$  where  $x_{1i} \in \mathfrak{H}_{1i}$ ,  $x_{2i} \in \mathfrak{H}_{2i}$ , then  $\prod_{i \in I} \bigotimes_{\mathfrak{G}_0} (\mathbf{M}_{1i} \otimes \mathbf{M}_{2i})$  is spatially isomorphic to  $(\prod_{i \in I} \bigotimes_{\mathfrak{G}_1} \mathbf{M}_{1i}) \otimes (\prod_{i \in I} \bigotimes_{\mathfrak{G}_2} \mathbf{M}_{2i})$ .*

**PROOF.** Considering  $\prod_{i \in I, j=1,2} \bigotimes_{\mathfrak{G}} \mathfrak{H}_{ji}$  for  $\mathfrak{G} = \mathfrak{G}(x_{ji} : i \in \mathbf{I}, j = 1, 2)$ , we can

find a linear isometric mapping  $U$  of  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{1i} \otimes \mathfrak{H}_{2i}}^{\mathfrak{G}_0}$  onto  $(\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{1i}}^{\mathfrak{G}_1} \mathfrak{H}_{1i}) \otimes (\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{2i}}^{\mathfrak{G}_2} \mathfrak{H}_{2i})$  such that

$$U \prod_{i \in \mathbf{I}} \otimes (y_{1i} \otimes y_{2i}) = (\prod_{i \in \mathbf{I}} \otimes y_{1i}) \otimes (\prod_{i \in \mathbf{I}} \otimes y_{2i})$$

for all  $C_0$ -sequences  $(y_{1i} \otimes y_{2i} : i \in \mathbf{I}) \in \mathfrak{C}_0$  (cf. Theorem A).

For any  $A = \prod_{i \in \mathbf{I}} \otimes (A_{1i} \otimes A_{2i}) \in \prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i}}^{\mathfrak{G}_0}(\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i})$  and  $z = (\prod_{i \in \mathbf{I}} \otimes z_{1i}) \otimes (\prod_{i \in \mathbf{I}} \otimes z_{2i}) \in (\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{1i}}^{\mathfrak{G}_1} \mathfrak{H}_{1i}) \otimes (\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{2i}}^{\mathfrak{G}_2} \mathfrak{H}_{2i})$  we have  $UAU^{-1}z = U(\prod_{i \in \mathbf{I}} \otimes (A_{1i} \otimes A_{2i}))U^{-1}z = [\prod_{i \in \mathbf{I}} \otimes A_{1i}] \otimes (\prod_{i \in \mathbf{I}} \otimes A_{2i})z$ .

Hence it is easily seen that  $U$  provides an isomorphism between

$$\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i}}^{\mathfrak{G}_0}(\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i}) \text{ and } (\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{1i}}^{\mathfrak{G}_1} \mathfrak{M}_{1i}) \otimes (\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{2i}}^{\mathfrak{G}_2} \mathfrak{M}_{2i}).$$

PROOF OF THEOREM 3. By the assumption, there exist Hilbert spaces  $\mathfrak{H}_{1i}$ ,  $\mathfrak{H}_{2i}$  such that  $\mathfrak{M}_i$  is spatially isomorphic to  $\mathbf{B}(\mathfrak{H}_{1i}) \otimes I_{\mathfrak{H}_{2i}}$  for each  $i \in \mathbf{I}$ . So it suffices to consider  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{B}(\mathfrak{H}_{1i}) \otimes I_{\mathfrak{H}_{2i}}}$  for  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_i}$ . Choosing a  $C_0$ -sequence  $(x_{ij}, i \in \mathbf{I}, j = 1, 2)$  in  $\prod_{i \in \mathbf{I}, j=1,2} \bigotimes_{\mathfrak{H}_{j,i}} \mathfrak{H}_{j,i}$ , we put  $\mathfrak{C}_1 = \mathfrak{C}(x_{1i} : i \in \mathbf{I})$ ,  $\mathfrak{C}_2 = \mathfrak{C}(x_{2i} : i \in \mathbf{I})$  and  $\mathfrak{C} = \mathfrak{C}(x_{1i} \otimes x_{2i} : i \in \mathbf{I})$ . By Lemma 4  $[\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{B}(\mathfrak{H}_{1i})}^{\mathfrak{G}_1} \mathfrak{B}(\mathfrak{H}_{1i})] \otimes I_{\mathfrak{C}_2}$  is spatially isomorphic to  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{B}(\mathfrak{H}_{1i}) \otimes I_{\mathfrak{H}_{2i}}}$  and the assertion of the theorem is clear by Lemma 3.

PROOF OF THEOREM 4. It is known that a  $W^*$ -algebra not of type  $I$  is the direct product of two  $W^*$ -algebras, one of which is of type  $I_p$  for any positive integer  $p$ . Accordingly we represent each  $\mathfrak{M}_i$  by the direct product of factors  $\mathfrak{M}_{1i}$ ,  $\mathfrak{M}_{2i}$  acting on  $\mathfrak{H}_{1i}$ ,  $\mathfrak{H}_{2i}$  respectively, the latter being of type  $I_2$ . Now there exists an equivalence-class  $\mathfrak{C}$  for which  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{2i}}^{\mathfrak{G}} \mathfrak{M}_{2i}$  is a factor of type  $III$  [8]. Hence if we choose an equivalence-class  $\mathfrak{C} = \mathfrak{C}(x_{1i} \otimes x_{2i} : i \in \mathbf{I})$  (in  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{1i} \otimes \mathfrak{H}_{2i}} (\mathfrak{H}_{1i} \otimes \mathfrak{H}_{2i})$ ) such that  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{2i}}^{\mathfrak{G}_2} \mathfrak{M}_{2i}$  is of type  $III$  for  $\mathfrak{C}_2 = \mathfrak{C}(x_{2i} : i \in \mathbf{I})$ , we get a factor  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i}}^{\mathfrak{G}} (\mathfrak{M}_{1i} \otimes \mathfrak{M}_{2i})$  of type  $III$  by Lemma 4 and [9].

By Lemma 4 and [6: Lemma 5.2.1] we can easily show the following:

COROLLARY ([1]. [2]. [3]). *The direct product of two approximately finite factors is also an approximately finite factor.*

2)  $I_{\mathfrak{C}_2}$  means the identity operator on  $\prod_{i \in \mathbf{I}} \bigotimes_{\mathfrak{H}_{2i}}^{\mathfrak{G}_2} \mathfrak{H}_{2i}$

**4. Normalcy of the infinite direct product.** Concerning normalcy of  $W^*$ -algebra we have the following

**THEOREM 5** (cf. [4]). *Let  $\mathbf{M}_i$  be factors on Hilbert spaces  $\mathfrak{H}_i$  for all  $i \in \mathbf{I}$ . If a certain factor  $\mathbf{M}_{i_0}$  is not normal, then  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i$  is not normal for any equivalence-class  $\mathfrak{C}$ .*

**PROOF.** As  $\mathbf{M}_{i_0}$  is not normal, then there exists a  $W^*$ -subalgebra  $\mathbf{N}_{i_0}$  of  $\mathbf{M}_{i_0}$  such that

$$(\mathbf{N}'_{i_0} \cap \mathbf{M}_{i_0}) \cap \mathbf{M}_{i_0} \not\equiv \mathbf{N}_{i_0}.$$

Now  $\bar{\mathbf{N}}_{i_0} \subset \prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i$  on  $\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathfrak{H}_i$  for any  $\mathfrak{C}$  and

$$\begin{aligned} & [ \bar{\mathbf{N}}'_{i_0} \cap \prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i ] \cap \prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i = \mathbf{R}(\bar{\mathbf{N}}_{i_0}, (\prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i)') \cap \prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathbf{M}_i \\ & \supset \mathbf{R}(\bar{\mathbf{N}}_{i_0}, \bar{\mathbf{M}}'_{i_0}) \cap \bar{\mathbf{M}}_{i_0} = \mathbf{R}(\mathbf{N}_{i_0}, \mathbf{M}'_{i_0}) \cap \bar{\mathbf{M}}_{i_0} \\ & \supset \mathbf{R}(\mathbf{N}_{i_0}, \mathbf{M}'_{i_0}) \cap \mathbf{M}_{i_0} = (\mathbf{N}'_{i_0} \cap \mathbf{M}_{i_0})' \cap \mathbf{M}_{i_0} \not\equiv \mathbf{N}_{i_0} \text{ in } \prod_{i \in \mathbf{I}} \bigotimes_{i \in \mathbf{I}}^{\otimes} \mathfrak{H}_i \end{aligned}$$

for any  $\mathfrak{C}$ , which proves the theorem.

REFERENCES

- [1] J. DIXMIER, Les algèbres d'opérateurs dans l'espace hilbertien, Paris, (1957).
- [2] Y. MISONOU, On the direct product of  $W^*$ -algebras, Tôhoku Math. Journ., 6 (1954), 189-204.
- [3] Y. MISONOU, Generalized approximately finite  $W^*$ -algebras, Tôhoku Math. Journ., 7(1955), 192-205.
- [4] Y. MISONOU, On divisors of factors, Tôhoku Math. Journ., 8(1956), 63-69.
- [5] F. J. MURRAY and J. VON NEUMANN, On rings of operators, Ann. of Math., 37(1936), 116-229.
- [6] F. J. MURRAY and J. VON NEUMANN, On rings of operators IV, Ann. of Math., 44(1943), 716-808.
- [7] J. VON NEUMANN, On infinite direct products, Compositio Mathematica 6(1938), 1-77.
- [8] J. VON NEUMANN, On rings of operators III, Ann. of Math., 41(1940), 94-161.
- [9] S. SAKAI, On topological properties of  $W^*$ -algebras, Proc. Japan Acad., 33(1957), 439-444.
- [10] Z. TAKEDA, Inductive limit and infinite direct product of operator algebras. Tôhoku Math. Journ., 7(1955), 67-86.

DEPARTMENT OF MATHEMATICS, TÔHOKU UNIVERSITY.