



# On Initial Chebyshev Polynomial Coefficient Problem for Certain Subclass of Bi-Univalent Functions

F. Müge Sakar<sup>1,\*</sup>, and Ertuğrul Doğan<sup>2</sup>,

<sup>1</sup> Department of Business Administration, Faculty of Management and Economics, Dicle University, Diyarbakır, Turkey

<sup>2</sup> Institute of Science, Batman, Batman University, Turkey

\*Corresponding author: [mugesakar@hotmail.com](mailto:mugesakar@hotmail.com)

**Abstract.** In this paper, we firstly, introduced the subclass  $R_{\Sigma}(\tau, \alpha, \gamma; t)$  satisfying subordinate conditions. Subsequently, considering this defined subclass, initial coefficient estimations are established using by Chebyshev polynomials expansions, and Fekete-Szegő inequalities are also derived for functions belonging to the said subclass. Furthermore, Some relevant consequences of these results are also discussed.

**Keywords.** Initial coefficients problem; Bi-univalent function; Chebyshev polynomials; Fekete-Szegő problem

**MSC.** 30C45; 30C50

**Received:** December 2, 2019

**Accepted:** December 18, 2019

Copyright © 2020 F. Müge Sakar and Ertuğrul Doğan. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

## 1. Introduction

Let  $\mathcal{A}$  denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

which are analytic in the open unit disc  $\Delta = \{z : z \in \mathbb{C} \text{ and } |z| < 1\}$ . In addition, we indicate by  $\mathcal{S}$  the class of all univalent functions in  $\Delta$ .

For  $f$  and  $g$ , analytic functions in  $\Delta$ , the function  $f(z)$  is subordinate to  $g(z)$  in  $\Delta$ , and it can be represented by

$$f(z) < g(z) \quad (z \in \Delta)$$

if there exists an analytic Schwarz function  $w(z)$  being as

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \Delta)$$

such that

$$f(z) = g(w(z)) \quad (z \in \Delta).$$

Especially, if the function  $g$  is univalent in  $\Delta$ , then the above subordination is equivalent to  $f(0) = g(0)$  and  $f(\Delta) \subset g(\Delta)$ .

Every function  $f \in \mathcal{S}$  is known to have an inverse  $f^{-1}$ , given by

$$f^{-1}(f(z)) = z \quad (z \in \Delta)$$

and

$$f^{-1}(f(w)) = w \left( |w| < r_0(f); r_0(f) \geq \frac{1}{4} \right),$$

where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots \quad (2)$$

In order to have bi-univalent  $f \in \mathcal{A}$  in  $\Delta$ , both  $f(z)$  and  $f^{-1}(z)$  must be univalent in  $\Delta$ .

Let represent the class of bi-univalent functions in  $\Delta$ , given by the Taylor-Maclaurin series expansion (1) by  $\Sigma$ .

Some well-known subclasses of the  $\mathcal{S}$ , which are denoted by class  $S^*(\alpha)$  of starlike functions of order  $\alpha$  in  $\Delta$  and the class  $K(\alpha)$  of convex functions of order  $\alpha$  in  $\Delta$ , are respectively shown as follows:

$$S^*(\alpha) := \left\{ f : f \in \mathcal{A} \text{ and } \Re \left( \frac{zf'(z)}{f(z)} \right) > \alpha; z \in \Delta; 0 \leq \alpha < 1 \right\} \quad (3)$$

and

$$K(\alpha) := \left\{ f : f \in \mathcal{A} \text{ and } \Re \left( 1 + \frac{zf''(z)}{f'(z)} \right) > \alpha; z \in \Delta; 0 \leq \alpha < 1 \right\}. \quad (4)$$

For  $0 \leq \alpha < 1$ , if both  $f$  and  $f^{-1}$  are respectively starlike or convex functions of order  $\alpha$ ,  $f \in \Sigma$  is in the class  $S_{\Sigma}^*(\alpha)$  of bi-starlike function of order  $\alpha$ , or  $K_{\Sigma}(\alpha)$  of bi-convex function of order  $\alpha$ .

The significance of Chebyshev polynomial in numerical analysis is increased in terms of both theoretical and practical points of view. On the other hand, many researchers have also been dealing with orthogonal Chebyshev polynomials. One can see the details of Chebyshev polynomials of first kind  $T_n(t)$ , the second kind  $U_n(t)$  and their numerous uses in different applications in the references [4, 5, 7]. The well known first and second kinds Chebyshev polynomials are defined as follows:

$$T_n(t) = \cos(n\theta) \quad \text{and} \quad U_n(t) = \frac{\sin(n+1)\theta}{\sin\theta} \quad (-1 < t < 1)$$

where the subscript  $n$  shows the polynomial degree and  $t$  is equal to  $\cos\theta$ .

In geometric function theory, the Fekete-Szegő functional  $|\alpha_3 - \eta\alpha_2^2|$  for normalized univalent functions of the form given by (1) is well known in this field. Historically, its origin is based on Fekete and Szegő of the 1933 conjecture of Littlewood and Paley. In their study the coefficients of odd univalent functions are bounded by unity (see [6]). Since Fekete-Szegő functional has received great attention, especially in many subclasses of the family of univalent functions, this topic had become of interest among researchers (see, e.g., [1, 2, 8, 9, 11]).

**Definition 1.1.** A function  $f \in \Sigma$  given by the equation (1) is said to be in the class  $R_\Sigma(\tau, \alpha, \gamma; t)$ , for  $\alpha \geq 1, \gamma \geq 0, \tau \in \mathbb{C} \setminus \{0\}, t \in (\frac{1}{2}, 1]$  and all  $z, w \in \Delta$  if the following subordination conditions hold:

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{f(z)}{z} + \alpha f'(z) + \gamma z f''(z) - 1 \right] \prec H(z, t) := \frac{1}{1 - 2tz + z^2} \tag{5}$$

and

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{g(w)}{w} + \alpha g'(w) + \gamma z g''(w) - 1 \right] \prec H(w, t) := \frac{1}{1 - 2wt + w^2}, \tag{6}$$

where the function  $g = f^{-1}$  is given by (2).

It is remarkable that if  $t = \cos \alpha$ , where  $\alpha \in (-\pi/3, \pi/3)$ , then

$$H(z, t) = \frac{1}{1 - 2\cos \alpha z + z^2} = 1 + \sum_{n=1}^{\infty} \frac{\sin(n+1)\alpha}{\sin \alpha} z^n \quad (z \in \Delta).$$

Thus

$$H(z, t) = 1 + 2\cos \alpha z + (3\cos^2 \alpha - \sin^2 \alpha)z^2 + \dots \quad (z \in \mathbb{U}).$$

$H(z, t)$  can be written as from reference [10],

$$H(z, t) = 1 + U_1(t)z + U_2(t)z^2 \dots \quad (z \in \Delta), t \in (-1, 1)$$

where

$$U_{n-1} = \frac{\sin(n \arccos t)}{\sqrt{1-t^2}} \quad (n \in \mathbb{N})$$

indicates the second kind of Chebyshev polynomials and we have some initial coefficients as follows:

$$U_n(t) = 2tU_{n-1}(t) - U_{n-2}(t),$$

and

$$U_1(t) = 2t, \quad U_2(t) = 4t^2 - 1, \quad U_3(t) = 8t^3 - 4t, \quad U_4(t) = 16t^4 - 12t^3 + 1 \dots \tag{7}$$

The first kind of Chebyshev polynomial  $T_n(t), t \in [-1, 1]$ , is indicated by

$$\sum_{n=0}^{\infty} T_n(t)z^n = \frac{1-tz}{1-2tz+z^2} \quad (z \in \Delta).$$

The first kind of Chebyshev polynomial  $T_n(t)$  is related to second kind of Chebyshev polynomial  $U_n(t)$  which is given below:

$$\frac{dT_n(t)}{dt} = nU_{n-1}(t); \quad T_n(t) = U_n(t) - tU_{n-1}(t); \quad 2T_n(t) = U_n(t) - U_{n-2}(t).$$

**Remark 1.2. (i)** For  $\tau = 1$ ,  $\alpha = \lambda$  and  $\gamma = 0$ , we get the class  $R_{\Sigma}(1, \lambda, 0; t) = B_{\Sigma}(\lambda, t)$  consists of functions  $f \in \Sigma$  satisfying the condition

$$(1 - \lambda) \frac{f(z)}{z} + \lambda f'(z) < H(z, t) := \frac{1}{1 - 2tz + z^2}$$

and

$$(1 - \lambda) \frac{g(w)}{w} + \lambda g'(w) < H(w, t) := \frac{1}{1 - 2tw + w^2},$$

where the function  $g = f^{-1}$  is defined by (2) (Bulut et al. [3]).

**(ii)** For  $\tau = 1$ ,  $\alpha = 1$  and  $\gamma = 0$ , we have the class  $R_{\Sigma}(1, 1, 0; t) = B_{\Sigma}(t)$  consists of functions  $f$  satisfying the condition

$$f'(z) < H(z, t) := \frac{1}{1 - 2tz + z^2}$$

and

$$g'(w) < H(w, t) := \frac{1}{1 - 2tw + w^2},$$

where the function  $g = f^{-1}$  is defined by (2).

In the current investigation, we present a subclass  $R_{\Sigma}(\tau, \alpha, \gamma; t)$  of analytic and bi-univalent functions. Additionally, we derived the initial coefficient bounds and Fekete-Szegö inequality by means of Chebyshev polynomials expansions. Furthermore, we introduce some corollaries associated with our main results.

## 2. Main Results

In this section, we propose to find the estimates on the Taylor-Maclaurin coefficients  $|a_2|$ ,  $|a_3|$  and Fekete-Szegö inequality for functions in the class  $R_{\Sigma}(\tau, \alpha, \gamma; t)$ , which is introduced by Definition 1.1. These inequalities are asserted by Theorem 2.1.

**Theorem 2.1.** For  $\alpha \geq 1$ ,  $\gamma \geq 0$ ,  $\tau \in \mathbb{C} \setminus \{0\}$  and  $t \in (\frac{1}{2}, 1]$ , let  $f \in R_{\Sigma}(\tau, \alpha, \gamma; t)$ . Then

$$|a_2| \leq \frac{2|\tau|t\sqrt{2t}}{\sqrt{|4[\tau(1+2\alpha+6\gamma) - (1+\alpha+2\gamma)^2]t^2 + (1+\alpha+2\gamma)^2|}}, \quad (8)$$

$$|a_3| \leq \frac{4|\tau|^2 t^2}{(1+\alpha+2\gamma)^2} + \frac{2|\tau|t}{1+2\alpha+6\gamma} \quad (9)$$

and for some  $\eta \in \mathbb{R}$ ,

$$|a_3 - \eta a_2^2| \leq \begin{cases} \frac{2|\tau|t}{1+2\alpha+6\gamma}, & |\eta - 1| \leq \frac{|(1+\alpha+2\gamma)^2 - 4[(1+\alpha+2\gamma)^2 - \tau(1+2\alpha+6\gamma)]t^2|}{4(1+2\alpha+6\gamma)t^2} \\ \frac{8|\tau|^2|\eta - 1|t^3}{|(1+\alpha+2\gamma)^2 - 4[(1+\alpha+2\gamma)^2 - \tau(1+2\alpha+6\gamma)]t^2|}, & |\eta - 1| \geq \frac{|(1+\alpha+2\gamma)^2 - 4[(1+\alpha+2\gamma)^2 - \tau(1+2\alpha+6\gamma)]t^2|}{4(1+2\alpha+6\gamma)t^2}. \end{cases} \quad (10)$$

*Proof.* Let  $f \in R_{\Sigma}(\tau, \alpha, \gamma; \beta)$ . Then from the equations (5) and (6), we obtain

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{f(z)}{z} + \alpha f'(z) + \gamma z f''(z) - 1 \right] = 1 + U_1(t)p(z) + U_2(t)p^2(z) + \dots \tag{11}$$

and

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{g(w)}{w} + \alpha g'(w) + \gamma w g''(w) - 1 \right] = 1 + U_1(t)q(w) + U_2(t)q^2(w) + \dots \tag{12}$$

for some analytic functions

$$p(z) = c_1z + c_2z^2 + c_3z^3 + \dots, \quad (z \in \Delta) \tag{13}$$

$$q(w) = d_1w + d_2w^2 + d_3w^3 + \dots, \quad (w \in \Delta) \tag{14}$$

such that  $p(0) = q(0) = 0$ ,  $|p(z)| < 1$  ( $z \in \Delta$ ) and  $|q(w)| < 1$  ( $w \in \Delta$ ).

Obviously, if  $|p(z)| < 1$  and  $|q(w)| < 1$ , then

$$|c_j| \leq 1 \text{ and } |d_j| \leq 1 \quad \text{for } j \in N. \tag{15}$$

From (11), (12), (13) and (14), we have

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{f(z)}{z} + \alpha f'(z) + \gamma z f''(z) - 1 \right] = 1 + U_1(t)c_1z + [U_1(t)c_2 + U_2(t)c_1^2]z^2 + \dots \tag{16}$$

and

$$1 + \frac{1}{\tau} \left[ (1 - \alpha) \frac{g(w)}{w} + \alpha g'(w) + \gamma w g''(w) - 1 \right] = 1 + U_1(t)d_1w + [U_1(t)d_2 + U_2(t)d_1^2]w^2 + \dots \tag{17}$$

When we equate the coefficients (16) and (17), we have

$$\frac{1}{\tau}(1 + \alpha + 2\gamma)a_2 = U_1(t)c_1, \tag{18}$$

$$\frac{1}{\tau}(1 + 2\alpha + 6\gamma)a_3 = U_1(t)c_2 + U_2(t)c_1^2, \tag{19}$$

$$-\frac{1}{\tau}(1 + \alpha + 2\gamma)a_2 = U_1(t)d_1, \tag{20}$$

$$\frac{1}{\tau}(1 + 2\alpha + 6\gamma)(2a_2^2 - a_3) = U_1(t)d_2 + U_2(t)d_1^2. \tag{21}$$

From (18) and (20), we obtain

$$c_1 = -d_1 \tag{22}$$

and

$$\frac{2}{\tau^2}(1 + \alpha + 2\gamma)^2 a_2^2 = U_1^2(t)(c_1^2 + d_1^2). \tag{23}$$

Also, by using (19) and (21), we obtain

$$\frac{2}{\tau}(1 + 2\alpha + 6\gamma)a_2^2 = U_1(t)(c_2 + d_2) + U_2(t)(c_1^2 + d_1^2). \tag{24}$$

By using (23) in (24), we get

$$\left[ \frac{2}{\tau}(1 + 2\alpha + 6\gamma) - \frac{2U_2(t)}{\tau^2 U_1^2(t)}(1 + \alpha + 2\gamma)^2 \right] a_2^2 = U_1(t)(c_2 + d_2). \tag{25}$$

From the eqs. (7), (15) and (25), we obtain the inequality (8).

Thereafter, we get the following equality, by subtracting (21) from (19)

$$\frac{2}{\tau}(1+2\alpha+6\gamma)(a_3-a_2^2) = U_1(t)(c_2-d_2) + U_2(c_1^2-d_1^2). \quad (26)$$

In addition, in the light of (22), we get

$$a_3 = a_2^2 + \frac{\tau U_1(t)(c_2-d_2)}{2(1+2\alpha+6\gamma)}. \quad (27)$$

Thus, we get convenient inequality (9) by using (23) and applying (7).

Now, we get the following equalities by using (25) and (27) for some  $\eta \in \mathbb{R}$

$$(a_3 - \eta a_2^2) = (1 - \eta) \left[ \frac{\tau^2 U_1^3(t)(c_2 + d_2)}{2\tau U_1^2(t)(1 + 2\alpha + 6\gamma) - 2U_2(t)(1 + \alpha + 2\gamma)^2} \right] + \frac{\tau U_1(t)(c_2 - d_2)}{2(1 + 2\alpha + 6\gamma)} \quad (28)$$

and

$$a_3 - \eta a_2^2 = \tau U_1(t) \left[ \left( h(\eta) + \frac{1}{2(1 + 2\alpha + 6\gamma)} \right) c_2 + \left( h(\eta) - \frac{1}{2(1 + 2\alpha + 6\gamma)} \right) d_2 \right],$$

where

$$h(\eta) = \frac{\tau U_1^2(t)(1 - \eta)}{(2\tau U_1^2(t)(1 + 2\alpha + 6\gamma) - 2U_2(t)(1 + \alpha + 2\gamma)^2)}.$$

So, we conclude that

$$|a_3 - \eta a_2^2| \leq \begin{cases} \frac{2|\tau|t}{1 + 2\alpha + 6\gamma}, & 0 \leq |h(\eta)| \leq \frac{1}{2(1 + 2\alpha + 6\gamma)} \\ 4|\tau||h(\eta)|t, & |h(\eta)| \geq \frac{1}{2(1 + 2\alpha + 6\gamma)}. \end{cases}$$

So, this is the proof which confirms Theorem 2.1. In the special case, we can obtain the Corollary 2.2 for the parameters  $\tau = 1$ ,  $\alpha = \lambda$  and  $\gamma = 0$  in Theorem 2.1.  $\square$

**Corollary 2.2** ([3]). *Let the function  $f \in \mathcal{B}_\Sigma(\lambda, t)$ , for  $\lambda \geq 1$  and  $t \in (1/2, 1]$ . Then*

$$|a_2| \leq \frac{2t\sqrt{2t}}{\sqrt{|(1 + \lambda)^2 - 4\lambda^2 t^2|}},$$

$$|a_3| \leq \frac{4t^2}{(1 + \lambda)^2} + \frac{2t}{(1 + 2\lambda)}$$

and for some  $\eta \in \mathbb{R}$ ,

$$|a_2 - \eta a_3| \leq \begin{cases} \frac{2t}{1 + 2\lambda}, & |\eta - 1| \leq \frac{|(1 + \lambda)^2 - 4\lambda^2 t^2|}{4(1 + 2\lambda)t^2} \\ \frac{8|\eta - 1|t^3}{|(1 + \lambda)^2 - 4\lambda^2 t^2|}, & |\eta - 1| \geq \frac{|(1 + \lambda)^2 - 4\lambda^2 t^2|}{4(1 + 2\lambda)t^2}. \end{cases}$$

In the special case, we can obtain the following Corollary 2.3 for the parameters  $\tau = 1$ ,  $\alpha = 1$  and  $\gamma = 0$  in Theorem 2.1.

**Corollary 2.3.** *Let the function  $f \in \mathcal{B}_\Sigma(t)$ , for  $t \in (\frac{1}{2}, 1]$ . Then*

$$|a_2| \leq \frac{t\sqrt{2t}}{\sqrt{|1 - t^2|}},$$

$$|a_3| \leq t^2 + \frac{2t}{3},$$

and for some  $\eta \in \mathbb{R}$ ,

$$|a_3 - \eta a_2^2| \leq \begin{cases} \frac{2t}{3}, & |\eta - 1| \leq \frac{1-t^2}{3t^2} \\ \frac{2|\eta - 1|t^3}{1-t^2}, & |\eta - 1| \geq \frac{1-t^2}{3t^2}. \end{cases}$$

### 3. Conclusion

Using the concept of Chebyshev polynomials, we have introduced a new subclass in the unit disc associated with subordination. We have then derived the initial coefficient estimations using by Chebyshev polynomials expansions, and also Fekete-Szegő inequalities for functions belonging to this subclass. Our main result is stated and proved as Theorem 2.1. Additionally, by specializing the some parameters, some relevant interesting consequences of these results which were studied in previous works, are obtained. So, these general results presented in this paper, are motivated essentially by the earlier works which are pointed out.

### Competing Interests

The authors declare that they have no competing interests.

### Authors' Contributions

All the authors contributed significantly in writing this article. The authors read and approved the final manuscript.

### References

- [1] Ş. Altınkaya and S. Yalçın, Fekete-Szegő inequalities for certain classes of bi-univalent functions, *International Scholarly Research Notices* **2014** (2014), Article ID 327962, 6 pages, DOI: 10.1155/2014/327962.
- [2] S. Aytaş, F. M. Sakar and H. Ö. Güney, Fekete-Szegő problem for  $p$ -valently starlike functions with complex order, *ARPJ Journal of Science and Technology* **5**(10) (2015), 514 – 519.
- [3] S. Bulut, N. Magesh and V. K. Balaji, Initial bounds for analytic and bi-univalent functions by means of Chebyshev polynomials, *Journal of Classical Analysis* **11**(1) (2017), 83 – 89, DOI: 10.7153/jca-11-06.
- [4] E. H. Doha, The first and second kind Chebyshev coefficients of the moments of the general-order derivative of an infinitely differentiable function, *International Journal of Computer Mathematics* **51** (1994), 21 – 35, DOI: 10.1080/00207169408804263.
- [5] J. Dziok, R. K. Raina and J. Sokół, Application of Chebyshev polynomials to classes of analytic functions, *Comptes Rendus Mathématique* **353** (2015), 433 – 438, DOI: 10.1016/j.crma.2015.02.001.
- [6] M. Fekete and G. Szegő, Eine Bemerkung über ungerade schlichte Funktionen, *Journal of the London Mathematical Society* **s1-8**(2) (1933), 85 – 89 (in German), DOI: 10.1112/jlms/s1-8.2.85.

- [7] J. C. Mason, Chebyshev polynomial approximations for the L-membrane eigenvalue problem, *SIAM Journal on Applied Mathematics* **15** (1967), 172 – 186, DOI: 10.1137/0115014.
- [8] H. Orhan, N. Magesh and V. K. Balaji, Fekete-Szegö problem for certain classes of Ma-Minda bi-univalent functions, *Afrika Matematika* **27** (2016), 889 – 897, DOI: 10.1007/s13370-015-0383-y.
- [9] F. M. Sakar, S. Aytaş and H. Ö. Güney, On the Fekete Szegö problem for generalized  $M_{\alpha,\gamma}(\beta)$  defined by differential operator, *Süleyman Demirel Üniversitesi Fen Bilimleri Enstitüsü Dergisi* **20**(3) (2016), 456 – 459, DOI: 10.19113/sdufbed.12069.
- [10] T. Whittaker and G. N. Watson, *A Course of Modern Analysis*, Cambridge University Press, Cambridge (1996), DOI: 10.1017/CBO9780511608759.
- [11] P. Zaprawa, On Fekete-Szegö problem for classes of bi-univalent functions, *Bulletin of the Belgian Mathematical Society - Simon Stevin* **21** (2014), 169 – 178, DOI: 10.36045/bbms/1394544302.