

ON INTRINSIC STRUCTURES SIMILAR TO THOSE INDUCED ON S^{2n}

BY DAVID E. BLAIR AND GERALD D. LUDDEN

1. In [1] Yano and the authors studied submanifolds of codimension 2 of almost complex manifolds and hypersurfaces of almost contact manifolds. In both cases the structure on the ambient space induced the same structure on the submanifold. The induced structure consists of a tensor field f of type $(1, 1)$, vector fields E, A , 1-forms η, α and a function λ satisfying

$$\begin{aligned}
 f^2 &= -I + \eta \otimes E + \alpha \otimes A, \\
 \eta \circ f &= \lambda \alpha, & \alpha \circ f &= -\lambda \eta, \\
 fE &= -\lambda A, & fA &= \lambda E, \\
 \eta(E) &= 1 - \lambda^2, & \alpha(E) &= 0, \\
 \eta(A) &= 0, & \alpha(A) &= 1 - \lambda^2.
 \end{aligned}
 \tag{1}$$

Moreover the metric g induced from a metric compatible with the structure on the ambient space satisfies

$$\begin{aligned}
 g(X, E) &= \eta(X), & g(X, A) &= \alpha(X), \\
 g(fX, fY) &= g(X, Y) - \eta(X)\eta(Y) - \alpha(X)\alpha(Y).
 \end{aligned}
 \tag{2}$$

It is well known that on an almost complex manifold or an almost contact manifold there exists a metric compatible with the given structure, i.e. we have an almost Hermitian structure or an almost contact metric structure. However given a $2n$ -dimensional manifold M^{2n} with tensors $(f, E, A, \eta, \alpha, \lambda)$ satisfying equations (1), we show in section 2 that there does not in general exist a Riemannian metric on M^{2n} satisfying equations (2). Thus to study manifolds with an intrinsically defined $(f, E, A, \eta, \alpha, \lambda)$ -structure from the standpoint of Riemannian geometry it is necessary to assume the existence of a Riemannian metric satisfying equations (2).

The even-dimensional spheres are clearly examples of manifolds with an $(f, E, A, \eta, \alpha, \lambda)$ -structure and a compatible metric g , the structure being induced from the natural structure on the ambient Euclidean space. If ∇ denotes the Riemannian connexion of g , then for the sphere example the structure tensors satisfy

Received September 26, 1970.

$$(3) \quad \begin{aligned} \nabla_X E &= -fX, & \nabla_X A &= -\lambda X, \\ (\nabla_X f)Y &= -\eta(Y)X + g(X, Y)E \end{aligned}$$

for any vector fields X, Y on the sphere [1]. Note also that from equations (1) and (2), $g(A, A) = g(E, E) = 1 - \lambda^2$.

In [3] Yano and Okumura obtained some characterizations of even-dimensional spheres by imposing some conditions on the tensors $f, E, A, \eta, \alpha, \lambda$. Here in section 3 we study the role that the equations (3) play in characterizing spheres.

THEOREM 3.2. *Let M^n be a compact Riemannian manifold (of any dimension $n \geq 2$) admitting a vector field A and a non-constant function λ satisfying*

$$\nabla_X A = -\lambda X, \quad g(A, A) = 1 - \lambda^2$$

for every vector field X . Then M^n is globally isometric to the unit sphere in R^{n+1} .

THEOREM 3.3. *Let M^{2n} be an even-dimensional manifold with an $(f, E, A, \eta, \alpha, \lambda)$ -structure and compatible metric g satisfying*

$$\begin{aligned} \lambda \text{ non-constant, } \nabla_X E &= -fX, \\ (\nabla_X f)Y &= -\eta(Y)X + g(X, Y)E. \end{aligned}$$

Then $\nabla_X A = -\lambda X$; in particular if M^{2n} is compact it is globally isometric to the unit sphere in R^{2n+1} .

2. Let M be an almost complex manifold with almost complex structure J and let Z be a vector field on M that is not the zero vector field. Let $\tilde{M} = M \times R^2$, where R is the real line. Define a tensor f of type $(1, 1)$, vector fields E and A , and 1-forms η and α on \tilde{M} in the following way:

$$(4) \quad \begin{aligned} f(X, t, s) &= (-JX - sZ, s, -t), \\ E &= (Z, 0, 0), \\ A &= (JZ, 0, 0), \\ \eta(X, t, s) &= t, \\ \alpha(X, t, s) &= s \end{aligned}$$

where X is any vector field on M and $t, s \in R$. Then we have that

$$\begin{aligned} f^2(X, t, s) &= f(-JX - sZ, s, -t) \\ &= (J^2X + sJZ + tZ, -t, -s) \\ &= -(X, t, s) + s(JZ, 0, 0) + t(Z, 0, 0) \\ &= -(X, t, s) + \eta(X, t, s)E + \alpha(X, t, s)A \end{aligned}$$

and hence $f^2 = -I + \eta \otimes E + \alpha \otimes A$. Also, we see that $\eta(E) = \eta(A) = \alpha(E) = \alpha(A) = 0$, $f(E) = f(Z, 0, 0) = (-JZ, 0, 0) = -A$, $f(A) = f(JZ, 0, 0) = (-J^2Z, 0, 0) = E$, $\eta \circ f(X, t, s) = \eta(-X - sZ, s, -t) = s = \alpha(X, t, s)$ and $\alpha \circ f(X, t, s) = \alpha(-X - sZ, s, -t) = -t = -\eta(X, t, s)$. Thus, (4) gives an $(f, E, A, \eta, \alpha, \lambda)$ -structure on \tilde{M} with $\lambda = 1$. If there exists a Riemannian metric \tilde{g} on \tilde{M} satisfying (2), then we have that $\tilde{g}(E, E) = \eta(E) = 0$, contradicting the fact that Z is *not* the zero vector and hence E is not the zero vector.

3. The proof of Theorem 3.2 is by means of a well known result of Obata [2] which states that a compact Riemannian manifold M^n admits a non-trivial solution λ of $(D_s d\lambda)(X, Y) = -k\lambda g(X, Y)$ for some real number $k > 0$ if and only if M^n is globally isometric to a Euclidean sphere of radius $1/\sqrt{k}$. Here D_s denotes the symmetric covariant derivative, for example for a 1-form θ ,

$$(D_s \theta)(X, Y) = \frac{1}{2} ((\nabla_X \theta)(Y) + (\nabla_Y \theta)(X)).$$

LEMMA 3.1. *Let M^n be a Riemannian manifold admitting a vector field A and a non-constant function λ satisfying $\nabla_X A = -\lambda X$, $g(A, A) = 1 - \lambda^2$. Let $\alpha(X) = g(X, A)$, then $\alpha(X) = X\lambda$.*

Proof. $\nabla_X \lambda A = (X\lambda)A - \lambda^2 X$, therefore

$$g(\nabla_X \lambda A, A) = (X\lambda)(1 - \lambda^2) - \lambda^2 \alpha(X).$$

On the other hand

$$\begin{aligned} g(\nabla_X \lambda A, A) &= -g(\lambda A, \nabla_X A) + Xg(\lambda A, A) \\ &= \lambda^2 \alpha(X) + (X\lambda)(1 - \lambda^2) + \lambda(-2\lambda X\lambda). \end{aligned}$$

Comparing we have $2\lambda^2 \alpha(X) = 2\lambda^2 X\lambda$ and hence $\alpha(X) = X\lambda$ for $\lambda \neq 0$. Let $\varphi(m) = (\alpha(X) - X\lambda)(m)$, $m \in M^n$ and suppose $\varphi(m) \neq 0$. Then there exists a neighborhood of m on which φ is non-zero. Therefore $\lambda = 0$ on this neighborhood contradicting the non-constancy of λ .

THEOREM 3.2. *Let M^n be a compact Riemannian manifold admitting a vector field A and a non-constant function λ satisfying*

$$\nabla_X A = -\lambda X, \quad g(A, A) = 1 - \lambda^2.$$

Then M^n is globally isometric to the unit sphere in R^{n+1} .

Proof. Using the Lemma and the result of Obata the proof is a short computation, the first equality holding since $d\lambda$ is an exact form.

$$\begin{aligned}
 (D_s d\lambda)(X, Y) &= X(Y\lambda) - (\nabla_X Y)\lambda \\
 &= X\alpha(Y) - \alpha(\nabla_X Y) \\
 &= Xg(Y, A) - g(\nabla_X Y, A) \\
 &= g(Y, \nabla_X A) \\
 &= -\lambda g(X, Y).
 \end{aligned}$$

THEOREM 3.3. *Let M^{2n} be a manifold with an $(f, E, A, \eta, \alpha, \lambda)$ -structure and compatible metric g satisfying*

$$\begin{aligned}
 \lambda \text{ non-constant, } \nabla_X E &= -fX, \\
 (\nabla_X f)Y &= -\eta(Y)X + g(X, Y)E.
 \end{aligned}$$

Then $\nabla_X A = -\lambda X$; in particular if M^{2n} is compact it is globally isometric to the unit sphere in \mathbb{R}^{2n+1} .

Proof. We first show that $\alpha(X) = X\lambda$. Since $g(E, E) = 1 - \lambda^2$ we have $2g(\nabla_X E, E) = -2\lambda X\lambda$ and hence $\eta(-fX) = -\lambda X\lambda$ so that by equations (1) $-\lambda\alpha(X) = -\lambda X\lambda$. Now proceeding as in the proof of Lemma 3.1 we have $\alpha(X) = X\lambda$. Thus, $\nabla_X \lambda A = \alpha(X)A + \lambda \nabla_X A$, while on the other hand

$$\begin{aligned}
 \nabla_X \lambda A &= -\nabla_X f E = -f \nabla_X E - (\nabla_X f)E \\
 &= f^2 X + \eta(E)X - g(X, E)E \\
 &= -X + \alpha(X)A + (1 - \lambda^2)X.
 \end{aligned}$$

Therefore $\lambda \nabla_X A = -\lambda^2 X$ and $\nabla_X A = -\lambda X$ for $\lambda \neq 0$. Now set $V = \nabla_X A + \lambda X$ and suppose $V(m) \neq 0$ for some $m \in M^{2n}$. Then there exists a neighborhood of m on which $V \neq 0$ and hence $\lambda = 0$, contradicting the non-constancy of λ . Thus $\nabla_X A = -\lambda X$ and the second statement follows from Theorem 3.2.

REMARK. The normality of an $(f, E, A, \eta, \alpha, \lambda)$ -structure has been defined and studied in [1] and [3]. In particular, Yano and Okumura [3] have shown that if M is a complete manifold with a normal $(f, E, A, \eta, \alpha, \lambda)$ metric structure such that $\lambda(1 - \lambda^2)$ is almost everywhere non-zero and $\nabla_X E = fX$ then M is isometric to a sphere. It can easily be seen that Theorem 3.3 implies this theorem.

REFERENCES

- [1] BLAIR, D. E., G. D. LUDDEN, AND K. YANO, Induced structures on submanifolds. *Kōdai Math. Sem. Rep.* **22** (1970), 188-198.

- [2] OBATA, M., Certain conditions for a Riemannian manifold to be isometric with a sphere. *J. Math. Soc. Japan* **14** (1962), 333-340.
- [3] YANO, K., AND M. OKUMURA, On (f, g, u, v, λ) -structures. *Kōdai Math. Sem. Rep.* **22** (1970), 401-423.

MICHIGAN STATE UNIVERSITY,
EAST LANSING, U.S.A.