

References

- [1] E. Z. Andalafte and L. M. Blumenthal, *Metric characterizations of Banach and Euclidean spaces*, Fund. Math. 55 (1964), pp. 23–55.
- [2] L. M. Blumenthal, *Theory and applications of distance geometry*, Oxford 1953.
- [3] — *An extension of a theorem of Jordan and von Neumann*, Pacific J. Math. 5 (1955), pp. 161–167.
- [4] L. D. Loveland and J. E. Valentine, *Metric criteria for Banach and Euclidean spaces*, Fund. Math. 100 (1978), pp. 75–81.
- [5] J. E. Valentine and S. G. Wayment, *Criteria for Banach spaces*, Pacific J. Math. 43 (1972), pp. 251–252.
- [6] W. H. Young, *On the analytic basis of noneuclidean geometry*, American J. Math. 33 (1911), pp. 249–286.

Accepté par la Rédaction le 17. 8. 1977

On k -regular embeddings of spaces in Euclidean space

by

David Handel (Detroit, Mich.) and Jack Segal (Seattle, Wash.)

Abstract. If $k \leq n$ are positive integers, a continuous map $f: X \rightarrow R^n$ is k -regular if whenever x_1, \dots, x_k are distinct points of X , then $f(x_1), \dots, f(x_k)$ are linearly independent. Such maps are of relevance in the theory of Čebyšev approximation. In this paper the question of existence of k -regular maps from a given X into R^n is considered. After discussing some elementary properties of k -regularity, an algebraic-topological method is introduced to obtain negative results. This method yields the fact that there does not exist a 3-regular map of the real projective plane into R^3 , and this result is best possible. Finally, it is shown how to construct explicit 2- and 3-regular maps on real projective spaces which, in terms of homogeneous coordinates, are given by quadratic functions.

1. Introduction. If $k \leq n$ are positive integers, a continuous map f of a space X into Euclidean n -space R^n is k -regular if whenever x_1, \dots, x_k are distinct points of X , then $f(x_1), \dots, f(x_k)$ are linearly independent. Closely related to this is the concept of an *affinely k -regular map* $f: X \rightarrow R^n$, where it is required that whenever x_0, \dots, x_k are distinct points of X , then $f(x_0), \dots, f(x_k)$ are affinely independent (i.e. they are the vertices of a non-degenerate k -simplex in R^n). The latter concept has been considered in [2], [1], and [9]. Clearly, a k -regular map is affinely $(k-1)$ -regular, and $f: X \rightarrow R^n$ is affinely $(k-1)$ -regular if and only if the map $g: X \rightarrow R^{n+1} = R \times R^n$ given by $g(x) = (1, f(x))$ is k -regular.

k -regular maps are of relevance in the theory of Čebyšev approximation. A set of n real-valued continuous functions on X is called a *k -Čebyšev set of length n* if these functions are the components of a k -regular map of X into R^n . The reader is referred to [10], pp. 237–242 for the significance of this concept.

The present paper is concerned with existence and non-existence of k -regular maps. The following results are obtained:

THEOREM 2.1. *X admits a 2-regular map into R^n if and only if X admits an affinely 1-regular map into R^{n-1} . (Thus if X is compact, existence of a 2-regular map of X into R^n is equivalent to X being topologically embeddable in R^{n-1}).*

THEOREM 2.2. *If X admits a k -regular map into R^n , then each $0 \leq i \leq k-1$, and S any subset of X with i points, $X-S$ admits a $(k-i)$ -regular map into R^{n-1} .*

2.2. is a slight sharpening of a result of Borsuk [2], p. 355 (proved there for the affinely k -regular case).

In Section 3 we note that existence of a k -regular map of X into \mathbb{R}^n implies existence of an equivariant map (with respect to permutation of factors) of the k th configuration space of X into $V_k(\mathbb{R}^n)$, the Stiefel manifold of k -frames in \mathbb{R}^n . Algebraic topology can be used to prove non-existence of such equivariant maps, and hence non-existence of k -regular maps. As an example, we use this technique to prove that the real projective plane P^2 does not admit a 3-regular map into \mathbb{R}^5 . (P^2 does admit a 3-regular map into \mathbb{R}^6). Since P^2 embeds in \mathbb{R}^4 and the complement of any point in P^2 embeds in \mathbb{R}^3 , this shows that the converse of 2.2 is false when $k = 3$.

In Section 4, linear algebra is used to produce quadratic 2 and 3-regular maps on real projective spaces.

2. Some properties of k -regularity.

THEOREM 2.1. *X admits a 2-regular map into \mathbb{R}^n if and only if X admits an affinely 1-regular map into \mathbb{R}^{n-1} .*

Proof. If $f: X \rightarrow \mathbb{R}^{n-1}$ is affinely 1-regular, then $g: X \rightarrow \mathbb{R}^n$ given by $g(x) = (1, f(x))$ is 2-regular, as noted in Section 1.

Conversely, suppose $g: X \rightarrow \mathbb{R}^n$ is 2-regular. Define $h: X \rightarrow S^{n-1}$ by $h(x) = g(x)/\|g(x)\|$. h is injective, and the image of h does not contain an antipodal pair since no two distinct points of X are mapped by g into the same line through the origin. In particular h maps X injectively into a proper subset of S^{n-1} , and hence X can be mapped injectively, i.e. affinely 1-regularly, into \mathbb{R}^{n-1} .

THEOREM 2.2. *If X admits a k -regular map into \mathbb{R}^n , then for each $0 \leq i \leq k-1$ and S any subset of X with i points, $X-S$ admits a $(k-i)$ -regular map into \mathbb{R}^{n-1} .*

Proof. Let $f: X \rightarrow \mathbb{R}^n$ be k -regular. Let x be any point of X , V the orthogonal complement of $f(x)$ in \mathbb{R}^n , and $\pi: \mathbb{R}^n \rightarrow V$ orthogonal projection. If x_1, \dots, x_{k-1} are distinct points of $X - \{x\}$, then $f(x), f(x_1), \dots, f(x_{k-1})$ are linearly independent, and hence $\pi f(x_1), \dots, \pi f(x_{k-1})$ are linearly independent in V . Thus πf , followed by a linear isomorphism of V onto \mathbb{R}^{n-1} , yields a $(k-1)$ -regular map of $X - \{x\}$ into \mathbb{R}^{n-1} . The general result follows by iteration.

THEOREM 2.3. *If X embeds in the n -sphere S^n , then there exists a 3-regular embedding of X in \mathbb{R}^{n+2} .*

Proof. The standard embedding of S^n in \mathbb{R}^{n+1} is affinely 2-regular (no line in \mathbb{R}^{n+1} meets S^n in more than 2 points), and hence by Section 1, S^n embeds in \mathbb{R}^{n+2} in a 3-regular fashion.

3. Equivariant maps. Let $F_k(X)$ denote the k th configuration space of X , i.e. the subspace of the k -fold cartesian product of X consisting of k -tuples of distinct points of X . Let $V_k(\mathbb{R}^n)$ denote the space of linearly independent k -frames in \mathbb{R}^n . The symmetric group on k letters, \mathcal{S}_k , acts freely on both $F_k(X)$ and $V_k(\mathbb{R}^n)$ by permutation of factors. (It is more convenient for us to use $V_k(\mathbb{R}^n)$ rather than the

Stiefel manifold of orthonormal k -frames in \mathbb{R}^n . The two are \mathcal{S}_k -equivariantly homotopy equivalent.)

THEOREM 3.1. *If X admits a k -regular map into \mathbb{R}^n , then there exists an \mathcal{S}_k -equivariant map $g: F_k(X) \rightarrow V_k(\mathbb{R}^n)$.*

Proof. If $f: X \rightarrow \mathbb{R}^n$ is k -regular, define $g: F_k(X) \rightarrow V_k(\mathbb{R}^n)$ by $g(x_1, \dots, x_k) = (f(x_1), \dots, f(x_k))$.

Let \mathcal{A}_k denote the alternating group on k letters. If X is Hausdorff, we have a double covering $F_k(X)/\mathcal{A}_k \rightarrow F_k(X)/\mathcal{S}_k$, where $F_k(X)/\mathcal{A}_k, F_k(X)/\mathcal{S}_k$ denote the orbit spaces of $F_k(X)$ with respect to the actions of $\mathcal{A}_k, \mathcal{S}_k$, respectively. Similarly we have a double covering $V_k(\mathbb{R}^n)/\mathcal{A}_k \rightarrow V_k(\mathbb{R}^n)/\mathcal{S}_k$. An \mathcal{S}_k -equivariant map $g: F_k(X) \rightarrow V_k(\mathbb{R}^n)$ induces a map of double coverings of the former to the latter. Thus, combining with 3.1 we obtain:

THEOREM 3.2. *If X admits a k -regular map into \mathbb{R}^n , then there exists a map of double coverings*

$$\begin{array}{ccc} F_k(X)/\mathcal{A}_k & \longrightarrow & V_k(\mathbb{R}^n)/\mathcal{A}_k \\ \downarrow & & \downarrow \\ F_k(X)/\mathcal{S}_k & \longrightarrow & V_k(\mathbb{R}^n)/\mathcal{S}_k \end{array}$$

If \mathcal{S}_k (resp. \mathcal{A}_k) acts on Y , for each point $y \in Y$ write $y_{\mathcal{S}}$ (resp. $y_{\mathcal{A}}$) for the point in Y/\mathcal{S}_k (resp. Y/\mathcal{A}_k) determined by y . Let $T: V_k(\mathbb{R}^n)/\mathcal{A}_k \rightarrow V_k(\mathbb{R}^n)/\mathcal{A}_k$ be the involution which interchanges the two points of each fibre in the double covering $V_k(\mathbb{R}^n)/\mathcal{A}_k \rightarrow V_k(\mathbb{R}^n)/\mathcal{S}_k$. T is given by $T(v_1, \dots, v_k)_{\mathcal{A}} = (v_2, v_1, \dots, v_k)_{\mathcal{A}}$. Let λ denote the real line bundle over $V_k(\mathbb{R}^n)/\mathcal{S}_k$ associated with the above double covering. For each positive integer m , the total space of $m\lambda$, the Whitney sum of m copies of λ , is $E(m\lambda) = V_k(\mathbb{R}^n)/\mathcal{A}_k \times_2 \mathbb{R}^m =$ quotient space of $V_k(\mathbb{R}^n)/\mathcal{A}_k \times \mathbb{R}^m$ obtained by identifying $(v_{\mathcal{A}}, x) \sim (Tv_{\mathcal{A}}, -x)$ for all $v \in V_k(\mathbb{R}^n)$ and $x \in \mathbb{R}^m$. Write $(v_{\mathcal{A}}, x)_2$ for the point in $E(m\lambda)$ determined by $(v_{\mathcal{A}}, x) \in V_k(\mathbb{R}^n)/\mathcal{A}_k \times \mathbb{R}^m$.

LEMMA 3.3. *Let $k \geq 3$ be odd, and λ the line bundle over $V_k(\mathbb{R}^n)/\mathcal{S}_k$ as above. Then $n\lambda$ is isomorphic to $\lambda \oplus \alpha \oplus \gamma$, where α is a $(k-1)$ -plane bundle with $\lambda \oplus \alpha$ orientable.*

Proof. Let β denote the k -plane subbundle of $n\lambda$ given as follows: $E(\beta) = \{(v_1, \dots, v_k)_{\mathcal{A}}, x\}_2 \mid x$ is a linear combination of $v_1, \dots, v_k\}$. If $v = (v_1, \dots, v_k) \in V_k(\mathbb{R}^n)$, the ordered basis $(v_{\mathcal{A}}, v_1)_2, \dots, (v_{\mathcal{A}}, v_k)_2$ of the fibre over $v_{\mathcal{A}}$ in β gives a well-defined orientation of this fibre since an even permutation does not change the orientation, and the above orientation coincides with that given by $(v_{\mathcal{A}}, -v_2)_2, (v_{\mathcal{A}}, -v_1)_2, (v_{\mathcal{A}}, -v_3)_2, \dots, (v_{\mathcal{A}}, -v_k)_2$ since k is odd. Hence β is orientable.

For v as above, write $\sum v = v_1 + \dots + v_k$, and let λ' denote the line subbundle of β whose fibre over $v_{\mathcal{A}}$ is spanned by $(v_{\mathcal{A}}, \sum v)_2$. We have an isomorphism of line bundles $\lambda \rightarrow \lambda'$ given by $(v_{\mathcal{A}}, i)_2 \mapsto (v_{\mathcal{A}}, i \sum v)_2$. Thus $\beta \cong \lambda' \oplus \alpha$ where α is a complementary subbundle to λ' in β . Take γ to be any complement to β in $n\lambda$.

LEMMA 3.4. *Let u denote the first Stiefel-Whitney class of the double covering $V_3(\mathbb{R}^5)/\mathcal{A}_3 \rightarrow V_3(\mathbb{R}^5)/\mathcal{S}_3$. Then $u^3 = 0$.*

Proof. We have $5\lambda \cong \lambda \oplus \alpha \oplus \gamma$ where α and γ are 2-plane bundles over $V_3(\mathbb{R}^5)/\mathcal{S}_3$ as in 3.3. We have $w(\lambda) = 1+u$. By the Whitney product formula [8], p. 37,

$$(1) \quad (1+u)^4 = w(\alpha)w(\gamma).$$

Since $\lambda \oplus \alpha$ is orientable, $w_1(\lambda \oplus \alpha) = 0$ [8], p. 146 and so it follows from the Whitney product formula that $w_1(\alpha) = u$. Thus (1) yields

$$(2) \quad 1+u^4 = (1+u+w_2(\alpha))(1+w_1(\gamma)+w_2(\gamma)).$$

Hence the 1, 2 and 3-dimensional components of the right hand side of (2) are 0, which yields

$$(3) \quad \begin{aligned} u &= w_1(\gamma), \\ uw_1(\gamma) + w_2(\alpha) + w_2(\gamma) &= 0, \\ w_1(\gamma)w_2(\alpha) + uw_2(\gamma) &= 0. \end{aligned}$$

Multiplying the second equation in (3) by u , together with the first and third equations yields $u^3 = 0$.

THEOREM 3.5. *There does not exist a 3-regular map of P^2 into \mathbb{R}^5 .*

Proof. Regard P^2 as the space of lines through the origin in \mathbb{R}^3 . If x and y are distinct points of P^2 , let $p(x, y)$ denote the unique line through 0 in \mathbb{R}^3 which is perpendicular to the plane of x and y . We have a map of double coverings

$$\begin{array}{ccc} F_2(P^2) & \xrightarrow{f} & F_3(P^2)/\mathcal{S}_3 \\ \downarrow & & \downarrow \\ F_2(P^2)/\mathcal{S}_2 & \xrightarrow{\quad} & F_3(P^2)/\mathcal{S}_3 \end{array}$$

where $f(x, y) = (x, y, p(x, y))_{\mathcal{S}}$. Let v denote the first Stiefel-Whitney class of the double covering $F_2(P^2) \rightarrow F_2(P^2)/\mathcal{S}_2$. Since $w_1(P^2) \neq 0$, it follows from [11], p. 380 (also [5], Theorem 3.7) that $v^3 \neq 0$. Thus by 3.4, there does not exist a map of double coverings

$$\begin{array}{ccc} F_3(P^2)/\mathcal{S}_3 & \xrightarrow{\quad} & V_3(\mathbb{R}^5)/\mathcal{S}_3 \\ \downarrow & & \downarrow \\ F_3(P^2)/\mathcal{S}_3 & \xrightarrow{\quad} & V_3(\mathbb{R}^5)/\mathcal{S}_3 \end{array}$$

and so we are done by 3.2.

Note that since P^2 embeds in S^4 , there does exist a 3-regular embedding of P^2 into \mathbb{R}^6 by 2.3. In fact, in the next section we will show that a quadratic (defined below) 3-regular embedding of P^2 into \mathbb{R}^6 exists.

4. Quadratic 2 and 3-regular embeddings of projective spaces. Regard real projective m -space P^m as the quotient space obtained from S^m by identifying antipodal points. Write $[x]$ for the point in P^m determined by $x \in S^m$.

DEFINITION 4.1. A quadratic map $f: P^m \rightarrow \mathbb{R}^n$ is one of the form $f[x] = g(x \otimes x)$ where $g: \mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1} \rightarrow \mathbb{R}^n$ is a linear map.

Equivalently, f is quadratic if the coordinates of $f[x]$ are homogeneous quadratic polynomials in the coordinates of x .

If $u \in \mathbb{R}^n \otimes \mathbb{R}^n$, $u \neq 0$, the rank of u is the smallest integer r such that u is expressible in the form $u = x_1 \otimes y_1 + \dots + x_r \otimes y_r$. If e_1, \dots, e_n is a basis of \mathbb{R}^n and we identify $\mathbb{R}^n \otimes \mathbb{R}^n$ with $M_n(\mathbb{R})$, the space of real $n \times n$ matrices, under the identification $\sum a_{ij} e_i \otimes e_j \mapsto (a_{ij})$, then the above notion of rank coincides with the usual matrix rank.

THEOREM 4.2. *Let $k = 2$ or 3 . If $f: P^m \rightarrow \mathbb{R}^n$ is the quadratic map given by $f[x] = g(x \otimes x)$ where $g: \mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1} \rightarrow \mathbb{R}^n$, then f is k -regular if the kernel of g contains no non-zero symmetric elements of rank $\leq k$.*

Proof. We give the proof for $k = 3$. (The case $k = 2$ is simpler.)

If $[x], [y], [z]$ are distinct points of P^m such that $f[x], f[y], f[z]$ are linearly dependent, then there exist real numbers a, b, c , not all 0, such that

$$0 = af[x] + bf[y] + cf[z] = g(ax \otimes x + by \otimes y + cz \otimes z).$$

Hence, by the hypothesis on the kernel of g ,

$$(1) \quad ax \otimes x + by \otimes y + cz \otimes z = 0.$$

Since $[x], [y], [z]$ are distinct, no two of x, y, z are linearly dependent, and hence no two of $x \otimes x, y \otimes y, z \otimes z$ are linearly dependent. Hence by (1), a, b, c must all be non-zero.

x, y, z cannot be linearly independent, for otherwise $x \otimes x, y \otimes y, z \otimes z$ would be, contradicting (1). Say $z = sx + ty$, $s, t \in \mathbb{R}$. Then both s and t are non-zero, for otherwise $[z]$ would not be distinct from $[x]$ and $[y]$. Substituting into (1) yields

$$(2) \quad (a + cs^2)x \otimes x + (b + ct^2)y \otimes y + cst(x \otimes y + y \otimes x) = 0.$$

But $x \otimes x, y \otimes y, x \otimes y + y \otimes x$ are linearly independent, and so $cst = 0$, a contradiction.

Note that if $k \geq 4$ and $m \geq 1$, a quadratic map on P^m cannot be k -regular, for if x, y are orthogonal points on S^m , then the lines $[x], [y], \left[\frac{x+y}{\sqrt{2}}\right], \left[\frac{x-y}{\sqrt{2}}\right]$ are distinct, but $x \otimes x, y \otimes y, \left(\frac{x+y}{\sqrt{2}}\right) \otimes \left(\frac{x+y}{\sqrt{2}}\right), \left(\frac{x-y}{\sqrt{2}}\right) \otimes \left(\frac{x-y}{\sqrt{2}}\right)$ are linearly dependent.

DEFINITION 4.3. A linear subspace of $M_n(\mathbb{R})$ is k -regular if it contains no non-zero symmetric matrices of rank $\leq k$.

COROLLARY 4.4. *Let $k = 2$ or 3 . If $M_{m+1}(\mathbb{R})$ contains a k -regular subspace of dimension $(m+1)^2 - n$, then P^m admits a quadratic k -regular map into \mathbb{R}^n .*

Proof. Such a subspace corresponds to a subspace K of $\mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1}$ with no non-zero symmetric elements of rank $\leq k$, and so the composition

$$\mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1} \rightarrow (\mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1})/K \cong \mathbb{R}^n$$

yields the desired quadratic k -regular map by 4.2.

EXAMPLE 4.5. Let S be the subspace of $M_{m+1}(\mathbb{R})$ consisting of all matrices $A = (a_{ij})$ satisfying $\sum_{i+j=q} a_{ij} = 0$ for $2 \leq q \leq 2m+2$. The dimension of S is $(m+1)^2 - (2m+1)$. S is 2-regular, for if (a_{ij}) is a non-zero symmetric matrix in S and q_0 is the smallest integer for which the $a_{ij}, i+j = q_0$, are not all 0, then at least 3 of these a_{ij} are non-zero and so $\text{rank}(a_{ij}) \geq 3$. Hence, by 4.4, there is a quadratic 2-regular embedding of P^m in \mathbb{R}^{2m+1} for all m .

This result is best possible when m is a power of 2, for then P^m does not embed in \mathbb{R}^{2m-1} ([4], p. 34 or [11]) and hence, by 2.1, does not 2-regularly embed in \mathbb{R}^{2m} .

EXAMPLE 4.6. Let S be the space of all skew-symmetric matrices in $M_3(\mathbb{R})$. S is k -regular for all k , S is 3-dimensional, and so by 4.4 there exists a quadratic 3-regular embedding of P^2 in \mathbb{R}^6 . By 3.5, this result is best possible.

EXAMPLE 4.7. Let S_0 consist of all matrices in $M_4(\mathbb{R})$ of the form

$$\begin{bmatrix} a & b & c & 0 \\ b & -a & 0 & c \\ c & 0 & -a & -b \\ 0 & c & -b & a \end{bmatrix}$$

and let S be the direct sum of S_0 with the space of all skew-symmetric matrices in $M_4(\mathbb{R})$. Since every non-zero matrix in S_0 has rank 4, S is 3-regular. S is 9-dimensional, and so by 4.4 there exists a quadratic 3-regular embedding of P^3 in \mathbb{R}^7 .

We conjecture that P^3 does not 3-regularly embed in \mathbb{R}^6 . (This would follow from 2.2 if we knew that the complement of a point in P^3 does not topologically embed in \mathbb{R}^4 . It is known [3], Corollary 1 that it does not differentiably embed in \mathbb{R}^4 .)

A linear map $\mathbb{R}^m \otimes \mathbb{R}^m \rightarrow \mathbb{R}^n$ is non-singular if its kernel contains no elements of rank 1. Extensive results on the existence of such maps have been obtained in [7] and [6].

THEOREM 4.8. Suppose there exists a quadratic 3-regular embedding of P^m in \mathbb{R}^n , and a non-singular map $\mathbb{R}^{m+1} \otimes \mathbb{R}^{m+1} \rightarrow \mathbb{R}^q$. Then there exists a quadratic 3-regular embedding of P^{2m+1} in \mathbb{R}^{2n+q} .

Proof. $M_{m+1}(\mathbb{R})$ contains a 3-regular subspace S_1 of dimension $(m+1)^2 - n$. The kernel of the non-singular map yields a $(m+1)^2 - q$ -dimensional subspace T of $M_{m+1}(\mathbb{R})$ which contains no elements of rank 1. Let

$$S = \left\{ \begin{bmatrix} A & B \\ C & D \end{bmatrix} \mid A \in S_1, B \in T, C \in M_{m+1}(\mathbb{R}), D \in S_1 \right\}.$$

Then S is a 3-regular subspace of $M_{2m+2}(\mathbb{R})$ of dimension $(2m+2)^2 - (2n+q)$, so we are done by 4.4.

4.8 is crude, and can undoubtedly be improved.

EXAMPLE 4.9. Quaternionic multiplication yields a non-singular map $\mathbb{R}^4 \otimes \mathbb{R}^4 \rightarrow \mathbb{R}^4$. Thus 4.7 and 4.8 yield a quadratic 3-regular embedding of P^7 in \mathbb{R}^{18} . Cayley multiplication yields a non-singular map $\mathbb{R}^8 \otimes \mathbb{R}^8 \rightarrow \mathbb{R}^8$, and hence there is a quadratic 3-regular embedding of P^{15} in \mathbb{R}^{44} .

References

- [1] V. G. Boltjanskiĭ, S. S. Ryškov, and Ju. A. Šaškin, *On k -regular imbeddings and their application to the theory of approximation of functions*, Uspehi Mat. Nauk 15 (1960), no. 6 (96), pp. 125-132 (Russian); Amer. Math. Soc. Transl. 28 (2) (1963), pp. 211-219.
- [2] K. Borsuk, *On the k -independent subsets of the Euclidean space and of the Hilbert space*, Bull. Acad. Polon. Sci. 5 (1957), pp. 351-356.
- [3] D. B. A. Epstein, *Embedding punctured manifolds*, Proc. Amer. Math. Soc. 16 (1965), pp. 175-176.
- [4] — and N. E. Steenrod, *Cohomology Operations*, Ann. of Math. Studies No. 50, Princeton Univ. Press 1962.
- [5] D. Handel, *An embedding theorem for real projective spaces*, Topology 7 (1968), pp. 125-130.
- [6] K. Y. Lam, *Construction of some nonsingular bilinear maps*, Bol. Soc. Mat. Mexicana 13 (1968), pp. 88-94.
- [7] R. J. Milgram, *Immersing projective spaces*, Ann. of Math. 85 (1967), pp. 473-482.
- [8] J. W. Milnor and J. D. Stasheff, *Characteristic Classes*, Ann. of Math. Studies No. 76, Princeton Univ. Press 1974.
- [9] Ju. A. Šaškin, *Topological properties of sets connected with approximation theory*, Izv. Akad. Nauk SSSR Ser. Mat. 29 (1965), pp. 1085-1094 (Russian).
- [10] I. Singer, *Best Approximation in Normed Linear Spaces by Elements of Linear Subspaces*, Springer-Verlag 1970.
- [11] W-T. Wu, *On the realization of complexes in Euclidean space II*, Scientia Sinica 7 (1958), pp. 365-387.

WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN
UNIVERSITY OF WASHINGTON, SEATTLE, WASHINGTON

Accepté par la Rédaction le 22. 8. 1977.