

ON MODULES INDUCED OR COINDUCED FROM HOPF SUBALGEBRAS

K.-H. ULBRICH

Let A be a Hopf algebra over a commutative ring k , $B \subset A$ a Hopf subalgebra, and W a left A -module. Koppinen and Neuvonen [2] showed that W is induced from B , that is $W \cong A \otimes_B V$ for a left B -module V , if and only if W admits a system of imprimitivity based on B ; their result assumes however that the antipode of A is bijective and that A as a right B -module is a finitely generated and projective generator. Essentially the same result holds for coinduced modules, i.e. modules of the form $W \cong \text{Hom}_B(A, V)$. The rather strong assumptions in [2] were made in order to apply the Morita theorems, and Koppinen and Neuvonen asked whether these assumptions can be weakened ([2], Remark). The present paper gives proofs for the above results which do not use Morita theory, and which only assume A to be finitely generated and projective over B . In the induced case a more general result is given which only needs A to be flat over B ; this is closely related to [1] and [4].

In the following A denotes a Hopf algebra over a commutative ring k , and $B \subset A$ a Hopf subalgebra. The antipode and counit are denoted by λ and ε , respectively, and the coproduct by δ . “ A -module” will mean left A -module.

1. Induced Modules.

Let F denote the k -algebra considered in [2]; as a k -module F consists of all right B -linear maps $f: A \rightarrow k$, i.e. $f(ab) = f(a)\varepsilon(b)$ for $a \in A, b \in B$, and the product is given by

$$(f \cdot f')(a) = \sum f(a_{(2)})f'(a_{(1)}), \quad f, f' \in F.$$

F is an A -module with $(af)(a') = f(\lambda(a)a')$ for $a, a' \in A$. By definition, an A -module W admits a system of imprimitivity based on B if it is a left F -module satisfying

$$(1) \quad a(fw) = \sum (a_{(1)}f)(a_{(2)}w), \quad a \in A, f \in F, w \in W.$$

In the following let $C = A \otimes_B k$. This is naturally an A -module coalgebra with coproduct $C \rightarrow C \otimes C$, $a \otimes 1 \mapsto \sum (a_{(1)} \otimes 1) \otimes (a_{(2)} \otimes 1)$, $a \in A$. (We note that $C \cong A/AB^+$ in the notation of [4], and that C represents the kernel of $\text{Sp}(A) \rightarrow \text{Sp}(B)$ if A is commutative, [5], p. 14). Let ${}^C_A\mathfrak{M}$ be the category of A -modules W which are supplied with a left C -coaction $\alpha: W \rightarrow C \otimes W$ such that

$$(2) \quad \alpha(aw) = \delta(a)\alpha(w), \quad a \in A, w \in W.$$

A morphism in ${}^C_A\mathfrak{M}$ is a map which is both A -linear and C -colinear.

LEMMA 1.1. *Assume A_B is finitely generated and projective. Then an A -module W admits a system of imprimitivity based on B if and only if W is an object of ${}^C_A\mathfrak{M}$.*

PROOF. First note that F is essentially the opposite of the dual algebra C^* of C . For

$$\zeta: \text{Hom}_k(C, k) \rightarrow F, \quad \zeta(g)(a) = g(a \otimes 1),$$

is an anti-isomorphism of k -algebras with $\zeta^{-1}(f)(a \otimes 1) = f(a)$; ζ is an algebra antimorphism since the product of F is defined by the transpose coproduct of A . Since A_B is finitely generated and projective, $C = A \otimes_B k$ is so over k . Therefore, W is a left F -module if and only if W is a left C -comodule, the actions being determined by each other through the formula

$$f \cdot w = \sum \langle \zeta^{-1}(f), w_{(-1)} \rangle w_{(0)}, \quad f \in F, w \in W.$$

Since $C \otimes W \cong \text{Hom}_k(F, W)$, one sees that (1) is equivalent to

$$(3) \quad (1 \otimes a)\alpha(w) = \sum (\lambda(a_{(1)}) \otimes 1)\alpha(a_{(2)}w), \quad a \in A, w \in W.$$

The latter is evidently satisfied if (2) holds. Conversely, applying (3) with a replaced by $a_{(2)}$ one obtains $\delta(a)\alpha(w) = \sum (a_{(1)}\lambda(a_{(2)}) \otimes 1)\alpha(a_{(3)}w) = \alpha(aw)$. Hence (2) is equivalent to (1), and this completes the proof.

For any left B -module V , $W = A \otimes_B V$ is naturally an object of ${}^C_A\mathfrak{M}$ with coaction $W \rightarrow C \otimes W$, $a \otimes v \mapsto \sum (a_{(1)} \otimes 1) \otimes (a_{(2)} \otimes v)$. Conversely, we want to show that any $W \in {}^C_A\mathfrak{M}$ is induced from B if A_B is flat. This is closely related to [1], Thm. 2.11, and [4], Thm. 2, and essentially the same proof as in [1] can be employed. It is based on the following lemma.

LEMMA 1.2. *For any A -module X the map*

$$\eta_X: A \otimes_B X \rightarrow C \otimes X, \quad a \otimes x \mapsto \sum (a_{(1)} \otimes 1) \otimes a_{(2)}x,$$

is an isomorphism.

PROOF. For $b \in B$ we have $\eta_X(ab \otimes x) = \sum (a_{(1)} \otimes \varepsilon(b_{(1)})) \otimes (a_{(2)}b_{(2)}x) = \eta_X(a \otimes bx)$; hence η_X is well-defined, and $(a \otimes 1) \otimes x \mapsto \sum a_{(1)} \otimes \lambda(a_{(2)})x$ gives a (well-defined) inverse.

THEOREM 1.3. *Let A be a Hopf k -algebra, $B \subset A$ a Hopf subalgebra, and $C = A \otimes_B k$. Assume A is flat as a right B -module. Then an A -module W is induced from B if and only if W is an object of ${}^C_A\mathfrak{M}$ (i.e. a left C -comodule satisfying (2)).*

PROOF. Let $\alpha: W \rightarrow C \otimes W$ be the coaction of W . Regard $C \otimes W$ as an A -module by $a \cdot (c \otimes w) = \delta(a)(c \otimes w)$. Then both α and $i: W \rightarrow C \otimes W, w \mapsto (1 \otimes 1) \otimes w$, are B -linear. Set $W_0 = \{w \in W | \alpha(w) = i(w)\}$. Since A is B -flat, the sequence

$$(4) \quad A \otimes_B W_0 \rightarrow A \otimes_B W \begin{array}{c} \xrightarrow{1 \otimes \alpha} \\ \xrightarrow{1 \otimes i} \end{array} A \otimes_B (C \otimes W)$$

is exact. Consider $\mu_W: A \otimes_B W_0 \rightarrow W, a \otimes w \mapsto aw$, which is a morphism in ${}^C_A\mathfrak{M}$. It is easy to see that $(\mu_W, \eta_W, \eta_{C \otimes W})$ transforms (4) into the sequence

$$W \xrightarrow{\alpha} C \otimes W \begin{array}{c} \xrightarrow{1 \otimes \alpha} \\ \xrightarrow{\delta \otimes 1} \end{array} C \otimes C \otimes W.$$

But this sequence is exact for any comodule. Hence Lemma 1.2 implies that μ_W is an isomorphism.

Let ${}_B\mathfrak{M}$ denote the category of left B -modules.

COROLLARY 1.4. (cf. [4], Thm. 2). *Assume A_B is faithfully flat. Then the functor ${}_B\mathfrak{M} \rightarrow {}^C_A\mathfrak{M}, V \mapsto A \otimes_B V$, is an equivalence.*

PROOF. For $V \in {}_B\mathfrak{M}$ consider the B -linear map $v: V \rightarrow (A \otimes_B V)_0, v \mapsto 1 \otimes v$. Since $\mu(1 \otimes v): A \otimes_B V \rightarrow A \otimes_B V$ is the identity, we have that $1 \otimes v$, hence v , is an isomorphism. It follows that $V \mapsto W_0$ is a quasi-inverse for $V \mapsto A \otimes_B V$.

REMARK 1.5. For $B = k$ theorem 1*3 gives the descent theorem for Hopf modules [3], Thm. 4.1.1. It should be noted however that the proof given in [3] works also for A not k -flat (and k not a field).

2. Coinduced Modules.

In the following we work with the k -algebra $E = \text{Hom}_B(A, k)$ of all left B -linear maps $f: A \rightarrow k$. The product is defined by

$$(f \cdot f')(a) = \sum f(a_{(1)})f'(a_{(2)}), \quad f, f' \in E.$$

E is an A -module algebra with $(af)(a') = f(a'a)$. We shall consider the category ${}_A\mathfrak{M}_E$ of left A -modules and right E -modules W satisfying

$$(5) \quad a(wf) = \sum (a_{(1)}w)(a_{(2)}f), \quad a \in A, w \in W, f \in E.$$

EXAMPLE. Let $W = \text{Hom}_B(A, V)$ for a left B -module V . For $g \in W$ and $f \in E$ define $gf \in W$ by $(gf)(a) = \sum g(a_{(1)})f(a_{(2)})$, $a \in A$. Then W is an object of ${}_A\mathfrak{M}_E$ with natural A -action $(ag)(a') = g(a'a)$.

REMARK 2.1. The condition for an A -module W to be an object of ${}_A\mathfrak{M}_E$ is slightly different from that of admitting a system of imprimitivity. There is no difference if A is cocommutative, for then E is commutative and $f \mapsto f\lambda$ gives an algebra isomorphism $F \simeq E$. There appears however to be a gap at the end of the proof in [2] for the coinduced case. The proof provides an action on W by the right B -endomorphisms of A , but for $W \cong \text{Hom}_B(A, V)$ a (right) action by the left B -endomorphisms of A is needed.

Let $W \in {}_A\mathfrak{M}_E$. We define a left B -module W_0 by the exact sequence

$$(6) \quad W \otimes E \begin{array}{c} \xrightarrow{m} \\ \xrightarrow{m'} \end{array} W \xrightarrow{p} W_0$$

where $m(w \otimes f) = wf$, and $m'(w \otimes f) = wf(1)$. Note that m and m' are B -linear if we regard $W \otimes E$ as a B -module by $b \cdot (w \otimes f) = \delta(b)(w \otimes f)$. We want to show that the left A - and right E -linear map

$$(7) \quad \mu_W: W \rightarrow \text{Hom}_B(A, W_0), \quad \mu_W(w)(a) = p(aw),$$

is an isomorphism if ${}_B A$ is finitely generated and projective. The proof is in some sense dual to that in section 1, and uses the following lemma (cf. the lemma in [2]).

LEMMA 2.2. Let X be an A -module, and suppose ${}_B A$ is finitely generated and projective. Then

$$\vartheta_X: X \otimes E \rightarrow \text{Hom}_B(A, X), \quad \vartheta_X(x \otimes f)(a) = \sum a_{(1)}f(a_{(2)})x,$$

is an isomorphism.

PROOF. Let ${}^e X = X$ with left B -action $b \cdot x = \varepsilon(b)x$. Then

$$\beta: \text{Hom}_B(A, {}^e X) \rightarrow \text{Hom}_B(A, X), \quad \beta(\varphi)(a) = \sum a_{(1)}\varphi(a_{(2)}),$$

is an isomorphism with $\beta^{-1}(\psi)(a) = \sum \lambda(a_{(1)})\psi(a_{(2)})$ for $\psi \in \text{Hom}_B(A, X)$. Set ${}^e \vartheta = \beta^{-1}\vartheta$. Then for $f \in E$, $a \in A$, and $x \in X$

$${}^e \vartheta(x \otimes f)(a) = \sum \lambda(a_{(1)})\vartheta(x \otimes f)(a_{(2)}) = \sum \lambda(a_{(1)})a_{(2)}f(a_{(3)})x = f(a)x.$$

Now choose a projective coordinate system $f_i \in \text{Hom}_B(A, B)$, $a_i \in A$, $1 \leq i \leq n$; then $\text{Hom}_B(A, {}^e X) \rightarrow X \otimes E$, $\varphi \mapsto \sum \varphi(a_i) \otimes \varepsilon \circ f_i$, is an inverse for ${}^e \vartheta$ as follows from $\sum f_i(a)a_i = a$ for $a \in A$.

THEOREM 2.3. Let A be a Hopf k -algebra, $B \subset A$ a Hopf subalgebra, $E = \text{Hom}_B(A, k)$, and assume that A is finitely generated and projective as a left

B-module. Then an *A*-module *W* is coinduced from *B* if and only if *W* is an object of ${}_A\mathfrak{M}_E$ (i.e. a right *E*-module satisfying (5)).

PROOF. For any right *E*-module *W* there is a canonical exact sequence

$$(8) \quad W \otimes E \otimes E \rightrightarrows W \otimes E \rightarrow W$$

defined by the action of *E* on *W*. Suppose $W \in {}_A\mathfrak{M}_E$ and consider $W \otimes E$ as a left *A*-module by $a \cdot (w \otimes f) = \delta(a)(w \otimes f)$. Then $(\mathfrak{D}_{W \otimes E}, \mathfrak{D}_W, \mu_W)$, with μ_W defined in (7), transforms (8) into the exact sequence obtained from (6) by applying $\text{Hom}_B(A, -)$. Observe that

$$\mathfrak{D}_{W \otimes E}(w \otimes f \otimes f')(a) = \sum a_{(1)}f'(a_{(2)})(w \otimes f) = \sum f'(a_{(3)})(a_{(1)}w \otimes a_{(2)}f),$$

and $p(wf) = p(wf(1))$. In particular, $(\mu_W m)(w \otimes f)(a) = p(a(wf)) = p(\sum a_{(1)}w(a_{(2)}f)) = p(\sum a_{(1)}wf(a_{(2)})) = p\mathfrak{D}_W(w \otimes f)(a)$. It follows therefore from lemma 2.2 that μ_W is an isomorphism.

COROLLARY 2.4. Suppose ${}_B A$ is finitely generated and projective. Then the functor ${}_A\mathfrak{M}_E \rightarrow {}_B\mathfrak{M}, W \mapsto W_0$, is an equivalence if and only if $B \subset A$ is a left *B*-direct summand of *A*.

PROOF. First observe that $V \mapsto \text{Hom}_B(A, V)$ is a right adjoint for $W \mapsto W_0$, the adjunction morphisms being μ , and $\nu: \text{Hom}_B(A, V)_0 \rightarrow V, p(g) \mapsto g(1)$, for $g \in \text{Hom}_B(A, V)$. Furthermore, the composite

$$\text{Hom}_B(A, V) \xrightarrow{\mu} \text{Hom}_B(A, \text{Hom}_B(A, V)_0) \xrightarrow{\text{Hom}(A, \nu)} \text{Hom}_B(A, V)$$

is the identity, since $p(ag) = (ag)(1) = g(a)$ for $a \in A$. Hence $\text{Hom}(A, \nu)$ is bijective. Now, if $A = B \oplus X$, we may conclude that ν is bijective by decomposing $\text{Hom}(A, \nu) = \text{Hom}(B, \nu) \oplus \text{Hom}(X, \nu)$. Conversely, suppose that ν is bijective for $V = B$. Then there exists $g \in \text{Hom}_B(A, B)$ with $g(1) = 1$, and therefore $A = B \oplus \text{Ker}(g)$. \square

REMARK 2.5. $B \subset A$ is a left *B*-direct summand of *A* iff *A* is a left *B*-generator; for suppose there exists a *B*-epimorphism $A^{(I)} \rightarrow B$. Pick $u \in A^{(I)}$ such that $u \mapsto 1$. Then $g: A \rightarrow Au \rightarrow B$ is a *B*-epimorphism with $g(1) = 1$, and hence with $A = B \oplus \text{Ker}(g)$.

REMARK 2.6. For $B = k$ the assumption of thm. 2.3 can not be omitted. This can be seen as follows. Let *X* be an *A*-module. Then $W = X \otimes E$ is naturally a right *E*-module and an object of ${}_A\mathfrak{M}_E$ with *A*-action defined by δ . It is not difficult to see that

$$W \otimes E \begin{array}{c} \xrightarrow{m} \\ \xrightarrow{m'} \end{array} W \xrightarrow{p} X$$

is exact where $p(x \otimes f) = f(1)x$ for $x \in X$, and $f \in E$. Hence in this case $W_0 = X$.

Furthermore, for $\mu_W: X \otimes E \rightarrow \text{Hom}_B(A, X)$ we obtain

$$\mu_W(x \otimes f)(a) = p(a \cdot (x \otimes f)) = p(\delta(a)(x \otimes f)) = \sum a_{(1)} f(a_{(2)})x.$$

Thus $\mu_W = \mathcal{G}_X$. Suppose now that μ_W is bijective for $X = A$. Then also ${}^e\mathcal{G} = \beta^{-1}\mathcal{G}$ is an isomorphism, and in case $B = k$ this means that A is finitely generated and projective over k .

ACKNOWLEDGMENT. I would like to thank the referee for some valuable remarks concerning section 2, and the Japan Soc. Prom. Sci. for financial support.

REFERENCES

1. Y. Doi and M. Takeuchi, *Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and Azumaya algebras*, J. Algebra 121 (1989), 488–516.
2. M. Koppinen and T. Neuvonen, *An imprimitivity theorem for Hopf algebras*, Math. Scand. 41 (1977), 193–198.
3. M. E. Sweedler, *Hopf Algebras*, W. A. Benjamin, Inc., New York, 1969.
4. M. Takeuchi, *Relative Hopf modules – equivalences and freeness criteria*, J. Algebra 60 (1979), 452–471.
5. W. C. Waterhouse, *Introduction to Affine Group Schemes*, Springer Verlag, Berlin, 1979.

INSTITUTE OF MATHEMATICS
UNIVERSITY OF TSUKUBA,
TSUKUBA-SHI, IBARAKI 305
JAPAN