

ON ORBIFOLD EMBEDDINGS

CHEOL-HYUN CHO, HANSOL HONG, AND HYUNG-SEOK SHIN

ABSTRACT. The concept of "orbifold embedding" is introduced. This is more general than sub-orbifolds. Some properties of orbifold embeddings are studied, and in the case of translation groupoids, orbifold embedding is shown to be equivalent to a strong equivariant immersion.

1. Introduction

Orbifolds arise naturally in many areas such as topology, geometric group theory, symplectic geometry and so on. In the last decade, they have been actively studied after Chen and Ruan introduced a new cohomology ring structure on orbifold cohomology [2]. Orbifolds also naturally appears when there is a symmetry, such as in symplectic reductions or in the presence of group actions. A very natural and basic question is to find the sub-objects for a given orbifold. A suborbifold (a subset which is also an orbifold with the induced topology) turns out to be a very restrictive notion. For example, given a product of two orbifolds the correct notion of diagonal Δ (see Definition 3.12) does not become a suborbifold but it is what is called an *orbifold embedding* into the product orbifold. Hence, it is clear that one should enlarge the class of sub-objects of an orbifold by including orbifold embeddings. Although the definition of orbifold embeddings appear in [1], such a notion has not been studied further. We became interested in this question of sub-objects in order to consider a proper notion of a Fukaya category of an orbifold. In the Fukaya category of a symplectic manifold, its objects are Lagrangian submanifolds decorated with additional data, and we believe that in the case of symplectic orbifolds, Lagrangian orbifold embeddings should become an important object in its Fukaya category (In this paper, we do not consider the Fukaya category of an orbifold, which is left for future research).

In this paper we give a slightly modified definition of orbifold embedding and explore several properties of orbifold embeddings. Also we prove that given an

Received February 22, 2013; Revised May 26, 2013.

2010 *Mathematics Subject Classification.* 57R18.

Key words and phrases. orbifold, groupoid, equivariant immersion.

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea Government (MEST) (No. 2012R1A1A2003117).

abelian orbifold embedding, the induced map between inertia orbifolds again becomes an orbifold embedding (Theorem 4.3).

One drawback of the definition of orbifold embedding is that it is rather cumbersome to work with as it is defined using local data. Also we observe that this notion of orbifold embedding is not Morita invariant. The second half of this paper is to remedy this in the case of translation groupoids. Namely, we construct an equivariant immersion in Sections 7 and 8, from the data of an orbifold embedding to an orbifold groupoid which is Morita equivalent to a translation groupoid $[M/G]$. Equivariant immersions are much easier to work with than orbifold embeddings, hence this construction should be very useful in applications. Pronk and Scull [8] showed that for translation groupoids, Morita equivalence can be defined only using translation groupoids, and our result is also in a similar point of view. To construct such an equivariant immersion, we use the Hilsum-Scandalis map, which is reviewed in Section 6.

Not all equivariant immersions give rise to orbifold embeddings, and we define what we call a *strong* equivariant immersion, which is shown to give an orbifold embedding. Also the equivariant immersion obtained from orbifold embeddings are also strong. Hence, in the case of translation groupoids, one can work with strong equivariant immersions, instead of orbifold embeddings.

2. Orbifold groupoids

In this section, we briefly recall well-known notions related to orbifold groupoids. We refer readers to [7] or [1] for details. One can define orbifolds in terms of local uniformizing charts (due to Satake).

Definition 2.1. An orbifold is a Hausdorff, second countable topological space X with a collection of uniformizing charts $(V_\alpha, G_\alpha, \phi_\alpha : V_\alpha \rightarrow X)$ of X , where the finite group G_α acts effectively on the manifold V_α , and continuous maps ϕ_α which descend to a homeomorphism $\overline{\phi}_\alpha$ of V_α/G_α onto an open subset $U_\alpha \subset X$. This data is required to satisfy following conditions:

- (1) $\{U_\alpha\}$ is a covering of X .
- (2) (Local compatibility) For $x \in U_\alpha \cap U_\beta$, there exist an open neighborhood $U \subset U_\alpha \cap U_\beta$ of x and a chart $(V, G, \phi : V \rightarrow X)$ of U which embeds to $(V_\alpha, G_\alpha, \phi_\alpha)$ and $(V_\beta, G_\beta, \phi_\beta)$.

In the modern approach of orbifolds, one usually uses the language of groupoids in the definition of orbifolds. This generalizes the notion of classical orbifolds allowing noneffective orbifolds. Recall that a groupoid is a (small) category whose morphisms are all invertible. Giving a topological structure and smooth structure on groupoids, we get the notion of Lie groupoids.

Definition 2.2. A topological groupoid \mathcal{G} is a pair of topological spaces $G_0 := \text{Obj}(\mathcal{G})$ and $G_1 := \text{Mor}(\mathcal{G})$ together with continuous structure maps:

- (1) The source and target map $s, t : G_1 \rightrightarrows G_0$, which assigns to each arrow $g \in G_1$ its source object and target object, respectively.

- (2) The multiplication map $m : G_{1s} \times_t G_1 \rightarrow G_1$, which compose two arrows.
- (3) The unit map $u : G_0 \rightarrow G_1$, which is a two-sided unit for the multiplication.
- (4) The inverse $i : G_1 \rightarrow G_1$, which assigns to each arrow its inverse arrow. This map is well-defined since all morphisms are invertible.

If all of the above maps are smooth and s (or t) is a surjective submersion (so that the domain $G_{1s} \times_t G_1$ of m is a smooth manifold), then \mathcal{G} is called a Lie groupoid.

Definition 2.3. Let \mathcal{G} be a Lie groupoid.

- (1) \mathcal{G} is proper if $(s, t) : G_1 \rightarrow G_0 \times G_0$ is a proper map.
- (2) \mathcal{G} is called a foliation groupoid if each isotropy group G_x is discrete.
- (3) \mathcal{G} is étale if s and t are local diffeomorphisms.

Note the every étale groupoid is a foliation groupoid. It can be easily checked that a proper foliation groupoid \mathcal{G} has only finite isotropy groups $G_x := (s, t)^{-1}(x, x)$ for each $x \in G_0$.

Definition 2.4. We define an orbifold groupoid to be a proper étale Lie groupoid.

Let us recall morphisms and Morita equivalence of orbifolds.

Definition 2.5. Let \mathcal{G} and \mathcal{H} be Lie groupoids. A homomorphism $\phi : \mathcal{H} \rightarrow \mathcal{G}$ consists of two smooth maps $\phi_0 : H_0 \rightarrow G_0$ and $\phi_1 : H_1 \rightarrow G_1$, that together commute with all the structure maps for the two groupoids \mathcal{G} and \mathcal{H} . It means that Lie groupoid morphisms are smooth functors between categories.

The following notion of equivalence is restrictive (it does not define equivalence relation), and later we will recall Morita equivalence which is indeed the correct notion of equivalences between orbifold groupoids.

Definition 2.6. A homomorphism between $\phi : \mathcal{H} \rightarrow \mathcal{G}$ between Lie groupoids is called equivalence if

- (i) (essentially surjective) the map

$$t\pi_1 : G_{1s} \times_{\phi_0} H_0 \rightarrow G_0$$

defined on the fibered product of manifolds

$$\{(g, y) \mid g \in G_1, y \in H_0, s(g) = \phi(y)\}$$

is a surjective submersion where $\pi_1 : G_{1s} \times_{\phi_0} H_0 \rightarrow G_1$ is the projections to the first factor;

(ii) the square

$$\begin{array}{ccc} H_1 & \xrightarrow{\phi_1} & G_1 \\ \downarrow (s,t) & & \downarrow (s,t) \\ H_0 \times H_0 & \xrightarrow{\phi_0 \times \phi_0} & G_0 \times G_0 \end{array}$$

is a fibered product of manifolds.

An equivalence in Definition 2.6 may not have an inverse. The notion of Morita equivalence is obtained by formally inverting equivalences in Definition 2.6.

Definition 2.7. \mathcal{G} and \mathcal{G}' are said to be Morita equivalent if there exists a groupoid \mathcal{H} and two equivalences

$$\mathcal{G} \xleftarrow{\phi} \mathcal{H} \xrightarrow{\phi'} \mathcal{G}.$$

It is well known that “Morita equivalence” defines an equivalence relation (see the discussion after Definition 1.43 in [1]). It is clear from the definition that equivalence is a special case of Morita equivalence.

We give an example of Morita equivalent groupoid which are not equivalent. For example, if \mathcal{G} can be made by tearing off some part \mathcal{G}' and adding arrows which contains the original gluing information, then we have an equivalence from \mathcal{G} to \mathcal{G}' . However, since “tearing off” process is not continuous, there is no map in the opposite direction in general.

Example 2.8. Consider two orbifold groupoids, \mathcal{G} and \mathcal{G}' , which are equivalent to the closed interval. From Figure 1, it is clear that they are Morita equivalent, but there are no maps neither from \mathcal{G} to \mathcal{G}' nor from \mathcal{G}' to \mathcal{G} .




FIGURE 1. Morita equivalence

Definition 2.9. Let $\phi : \mathcal{H} \rightarrow \mathcal{G}$ and $\psi : \mathcal{K} \rightarrow \mathcal{G}$ be homomorphisms of Lie groupoids. The fibered product $\mathcal{H} \times_{\mathcal{G}} \mathcal{K}$ is the Lie groupoid that makes the

following diagram a fibered product.

$$\begin{array}{ccc} \mathcal{H} \times_{\mathcal{G}} \mathcal{K} & \xrightarrow{pr_2} & \mathcal{K} \\ \downarrow pr_1 & & \downarrow \psi \\ \mathcal{H} & \xrightarrow{\phi} & \mathcal{G} \end{array}$$

which commutes up to a natural transformation. More explicitly,

$$(2.1) \quad (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 := H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0,$$

$$(2.2) \quad (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1 := H_1 \times_{s\phi_1, G_0, s} G_1 \times_{t, G_0, s\psi_1} K_1$$

with following source and target maps

$$\begin{aligned} s(h, g, k) &= (s(h), g, s(k)), \\ t(h, g, k) &= (t(h), \psi(k)g\phi(h)^{-1}, t(k)). \end{aligned}$$

We will also write $\mathcal{H} \times_{\mathcal{G}} \mathcal{K}$ as $\phi^* \mathcal{K}$ occasionally.

To be more precise, an element of $(\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0$ is a triple (x, g, z) such that

$$(2.3) \quad x \quad \phi_0(x) \xrightarrow{g} \psi_0(z) \quad z$$

and a morphism between (x, g, z) and (x', g', z') is a triple (h, g, k) which makes the following diagram commutative:

$$(2.4) \quad \begin{array}{ccccc} x & \phi_0(x) & \xrightarrow{g} & \psi_0(z) & z \\ \downarrow h & \phi_1(h) \downarrow & & \downarrow \psi_1(k) & \downarrow k \\ x' & \phi_0(x') & \xrightarrow{g'} & \psi_0(z') & z' \end{array}$$

i.e., for $(h, g, k) \in (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1$ which satisfies $s(g) = \phi_0(s(h))$ and $t(g) = \psi_0(s(k))$ by definition,

$$s(h, g, k) = s(h) \quad \phi_0(s(h)) \xrightarrow{g} \psi_0(s(k)) \quad s(k)$$

and

$$t(h, g, k) = t(h) \quad \phi_0(t(h)) \xrightarrow{g'} \psi_0(t(k)) \quad t(k)$$

where $g' = \psi_1(k)g\phi_1(h)^{-1}$.

Remark 2.10. The fibered product $\mathcal{H} \times_{\mathcal{G}} \mathcal{K}$ may not be a Lie groupoid, since $(\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0$ or $(\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1$ may not be manifolds.

The following lemma is well-known.

Lemma 2.11. *If $\psi : \mathcal{K} \rightarrow \mathcal{G}$ is an equivalence, then $\mathcal{H} \times_{\mathcal{G}} \mathcal{K}$ is a Lie groupoid and the projection $\mathcal{H} \times_{\mathcal{G}} \mathcal{K} \rightarrow \mathcal{H}$ is an equivalence*

$$(2.5) \quad \begin{array}{ccc} \mathcal{H} \times_{\mathcal{G}} \mathcal{K} & \xrightarrow{pr_2} & \mathcal{K} \\ pr_1 \downarrow & & \downarrow \psi \cong \\ \mathcal{H} & \xrightarrow{\phi} & \mathcal{G}. \end{array}$$

Proof. From (2.1), one can see that if $s \circ pr_1 : G_1 \times_{t, G_0, \psi_0} K_0 \rightarrow G_0$ is a submersion, then $(\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0$ is a manifold. This happens when ψ is an equivalence. Since $s : K_1 \rightarrow K_0$ is a submersion, a similar argument shows that $(\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1$ is a manifold for the equivalence ψ .

Recall that $\mathcal{H} \times_{\mathcal{G}} \mathcal{K}$ is a Lie groupoid whose set of objects and arrows are

$$\begin{aligned} (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 &= H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0, \\ (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1 &= H_1 \times_{s\phi_1, G_0, s} G_1 \times_{t, G_0, s\psi_1} K_1, \end{aligned}$$

respectively. We first check the condition (i) of Definition 2.6. We have to show that the following map

$$t\pi_1 : H_1 \times_{s, H_0, pr_1} (H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0) \rightarrow H_0$$

is a surjective submersion where π_1 is the projection to the first factor H_1 . Consider the following diagrams of fiber products.

$$\begin{array}{ccccc} H_1 \times_{s, H_0, pr_1} (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 & \longrightarrow & (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 & \longrightarrow & G_1 \times_{t, G_0, \psi_0} K_0 \\ \pi_1 \downarrow & & \downarrow & & \downarrow \\ H_1 & \longrightarrow & H_0 & \longrightarrow & G_0 \end{array}$$

The rightmost vertical map $G_1 \times_{t, G_0, \psi_0} K_0 \rightarrow G_0$ is a surjective submersion, since $\psi : \mathcal{K} \rightarrow \mathcal{G}$ is an equivalence. Then, it follows from a general property of fiber product diagrams that the middle vertical map $H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0 \rightarrow H_0$ is also a surjective submersion, and hence so is π_1 . Finally, $t\pi_1$ is a surjective submersion since it is given by a composition of two such kinds of maps.

To show the second condition of equivalence, we consider the following diagram:

$$\begin{array}{ccc} (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_1 & \xrightarrow{pr_1} & H_1 \\ \downarrow (s, t) & & \downarrow (s, t) \\ (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 \times (\mathcal{H} \times_{\mathcal{G}} \mathcal{K})_0 & \xrightarrow{pr_1 \times pr_1} & H_0 \times H_0 \end{array}$$

Since $(s, t) : H_1 \rightarrow H_0 \times H_0$ is a submersion, we only need to check that the above diagram is a fibered product of sets. Suppose $h \in H_1$, and denote $x = s(h)$ and $x' = t(h)$. Since $pr_1 : H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0 \rightarrow H_0$ is surjective, there exist (x, g, y) and (x', g', y') in $H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} K_0$. Since ψ is equivalence, there exists a unique $k \in K_1$ satisfying $\psi_1(k) = g' \phi_1(h) g^{-1}$. Since $h \in H_1$ determines a unique element (h, g, k) in the fiber over $((x, g, y), (x', g', y'))$, the above diagram is a fiber product as sets. \square

3. Orbifold embeddings

In this section we recall the main definition of this paper, an orbifold embedding, and explore its properties. The following notion is a slight modification from the one defined by Adem, Leida, and Ruan in their book [1].

Definition 3.1. A homomorphism of orbifold groupoids $\phi : \mathcal{H} \rightarrow \mathcal{G}$ is an embedding satisfying the following conditions:

- (1) $\phi_0 : H_0 \rightarrow G_0$ is an immersion
- (2) Let $x \in \text{im}(\phi_0) \subset G_0$ and let U_x be a neighborhood such that $\mathcal{G}|_{U_x} \cong G_x \ltimes U_x$. Then, the \mathcal{H} -action on $\phi_0^{-1}(x)$ is transitive, and there exists an open neighborhood $V_y \subset H_0$ for each $y \in \phi_0^{-1}(x)$ such that $\mathcal{H}|_{V_y} \cong H_y \ltimes V_y$ and

$$(3.1) \quad \mathcal{H}|_{\phi_0^{-1}(U_x)} \cong G_x \ltimes (G_x \times_{H_y} V_y)$$

- (3) $|\phi| : |\mathcal{H}| \rightarrow |\mathcal{G}|$ is proper and injective.

A Lie groupoid \mathcal{H} together with ϕ is called an orbifold embedding of \mathcal{G} .




FIGURE 2. Local shape of an orbifold embedding

In (2) of the above definition, the action of G_x is defined by

$$G_x \times (G_x \times_{H_y} V_y) \rightarrow G_x \times_{H_y} V_y, \quad (g, [k, z]) \mapsto [gk, z],$$

where $(k\phi_1(h), z) \sim (k, h \cdot z)$ is the equivalence relation defined by the action of H_y and $[k, z]$ denotes a class in the quotient $G_x \times_{H_y} V_y$.

There are two modifications in the definition from that of Adem, Leida, and Ruan (Definition 2.3 in [1]).

- (1) We use the local model $G_x \times_{H_y} V_y$ instead of $G_x/H_y \times V_y$.
- (2) We require that $|\phi| : |\mathcal{H}| \rightarrow |\mathcal{G}|$ is injective (which was not present in [1]).

Let us explain why we have made such modifications.

Firstly, in [1] $G_x/\phi_1(H_y) \times V_y$ was used instead of $G_x \times_{H_y} V_y$. But $\phi_1(H_y)$ may not be a normal subgroup of G_x (see Example 3.2). Also, it is not easy to find a natural G_x action on $G_x/\phi_1(H_y) \times V_y$ which reflects the H_y action on V_y . The only plausible action of G_x that may exist on $G_x/\phi_1(H_y) \times V_y$ is by the left multiplication on the first component. Now, any reasonable definition of an embedding should include the identity map, and therefore in this case we would have that $G_x \ltimes U_x \cong G_x \ltimes (G_x/G_x \times V_y)$ where $x = y$ and $U_x = V_y$ but, on U_x the group G_x acts and on $G_x/G_x \times V_y$ the action is trivial. Hence, $G_x \times_{H_y} V_y$ in (3.1) should be the correct local model.

Example 3.2. Let S_3 act on \mathbb{C}^3 as permutations on three coordinates where S_3 is the permutation group on 3 letters. Consider $V := \mathbb{C} \times \mathbb{C} \times \{0\} \subset \mathbb{C}^3$ and the subgroup H of S_3 generated by the transposition $(1, 2)$. Then, H acts on V and the natural map

$$S_3 \times_H V \rightarrow \mathbb{C}^3$$

induces an orbifold embedding $S_3 \ltimes (S_3 \times_H V) \rightarrow S_3 \ltimes \mathbb{C}^3$. Note that H is not a normal subgroup of S_3 .

Secondly, in [1], an orbifold embedding $\phi : \mathcal{H} \rightarrow \mathcal{G}$ does not necessarily induce an injective map $|\phi| : |\mathcal{H}| \rightarrow |\mathcal{G}|$. We first provide an example where $|\phi|$ is not injective but satisfies the other conditions of embedding. We will call a morphism ϕ of Lie groupoids *essentially injective* if $|\phi|$ is injective.

Example 3.3. Let \mathcal{G} be given by $G_0 = \mathbb{R} \amalg \mathbb{R}$ and $\mathbb{Z}/2\mathbb{Z}$ identifying two copies of \mathbb{R} . Suppose \mathcal{H} is the disjoint union of two copies of \mathbb{R} with only trivial arrows.

Immerse (embed) H_0 to G_0 by $id_{\mathbb{R}} \amalg id_{\mathbb{R}}$. One can easily check that ϕ satisfies the other axioms of orbifold embedding, except that $|\phi|$ is injective. The induced map between quotient space is rather a covering map from trivial double cover of \mathbb{R} to \mathbb{R} .

Remark 3.4. A morphism of groupoids $\phi : \mathcal{H} \rightarrow \mathcal{G}$ is essentially injective if the \mathcal{H} -action on $\phi_0^{-1}(t(s^{-1}(y)))$ (ϕ_0 inverse image of H_1 -orbit) is transitive for every $y \in G_0$, i.e. if one can find an arrow in H_1 from x to x' whenever there exists an arrow in G_1 from $\phi_0(x)$ to $\phi_0(x')$. If this happens, one can find an arrow in H_1 from x to x' .

Compare it with the notion of essential surjectivity: $\phi : \mathcal{H} \rightarrow \mathcal{G}$ is called essentially surjective if for any point x in G_0 , there is an arrow $g : \phi(y) \rightarrow x$ from a point in the image of ϕ to x .

Remark 3.5. The essential injectivity is Morita-invariant since it is a property of the induced map between quotient spaces.

The necessity and importance of the concept of orbifold embedding come from the fact that suborbifolds lack some essential properties. For example, objects such as the diagonal do not become suborbifolds. Therefore, we need to enlarge the definition of suborbifolds to include orbifold embeddings. See Example 3.12 to note that the diagonal homomorphism is indeed an orbifold embedding.

Now we investigate the properties of orbifold embeddings.

Lemma 3.6. *If $\phi : \mathcal{H} \rightarrow \mathcal{G}$ is an orbifold embedding, then the restriction of ϕ_1 on local isotropy groups is injective.*

Proof. Note that the point y corresponds to $[e, y]$ in this model, where e is the identity element in G_x . Since equivalence between orbifolds preserves local isotropy groups, the local group $\phi_1(H_y)$ at $[e, y]$ of $G_x \ltimes (G_x \times_{H_y} V_y)$ has to be isomorphic to H_y , and it proves the lemma. \square

Remark 3.7. For the case of an effective orbifold \mathcal{H} , Lemma 3.6 follows directly from the 0-level immersion ϕ_0 . Assume that there is a nontrivial element $h \in \ker(\phi_1|_{H_y})$. Fix a tangent vector $v \in T_y V_y$. Since the action of \mathcal{H} is effective, the difference of two vectors $v - h_* v$ is not a zero vector. By the assumption on h ,

$$(\phi_0)_*(v - h_* v) = 0,$$

and it contradicts that ϕ_0 is an immersion.

Remark 3.8. In the above Lemma 3.6, ϕ_1 may not be globally injective.

We remark that the orbifold embedding is not Morita invariant. Indeed, the following two examples illustrate this phenomenon.




FIGURE 3. An orbifold embedding and equivalence 1

Example 3.9. Let \mathcal{H} be a circle with the trivial orbifold groupoid structure and \mathcal{G} be a teardrop whose local group at the unique singular point x is $\mathbb{Z}/3$ as in Figure 3. The orbifold morphism $\phi : \mathcal{H} \rightarrow \mathcal{G}$ is not an orbifold embedding since it does not satisfy the second condition at $x \in \mathcal{G}$.

However, we can change the orbifold structure of \mathcal{H} as follows. Let $\phi(y) = x$ and U be a open neighborhood of y as in the figure. We add two more copies of U to get new objects U' and add additional arrows identifying three copies of U . Denote the resulting orbifold by \mathcal{H}' . Note that there is an equivalence from \mathcal{H}' to \mathcal{H} . The obvious modification $\phi' : \mathcal{H}' \rightarrow \mathcal{G}$ of ϕ is now an orbifold embedding.




FIGURE 4. An orbifold embedding and equivalence 2

Example 3.10. Let \mathcal{H} be the disjoint union of three copies of real lines and \mathcal{G} be \mathbb{R}^2 equipped with a $\mathbb{Z}/3$ action generated by $2\pi/3$ -rotation. Consider an orbifold embedding $\phi : \mathcal{H} \rightarrow \mathcal{G}$ shown in Figure 4. We similarly change the orbifold structure of \mathcal{H} by adding three more copies of \mathbb{R} to \mathcal{H} to get a new orbifold groupoid \mathcal{H}' , i.e., $H'_0 = \mathbb{R} \times \mathbb{Z}_6$ and $((h, g), k) \in H'_1 = H'_0 \times \mathbb{Z}_6$ sends $(h, g) \rightarrow (h, kg)$. It is clear from Figure 4 that there is an equivalence $\mathcal{H}' \rightarrow \mathcal{H}$, which is induced by the projection $\mathbb{Z}_6 \rightarrow \mathbb{Z}_3$. The morphism $\phi' : \mathcal{H}' \rightarrow \mathcal{G}$ is defined by the composition of ϕ and this equivalence. Then, we see that ϕ' is no longer an orbifold morphism because there is no transitive G_x action on $\phi'^{-1}(x)$ where x is the unique singular point in \mathcal{G} .

Example 3.11 (Orbifold diagonal). As an example of an orbifold embedding, we introduce a diagonal suborbifold of product orbifolds.

Definition 3.12. The diagonal suborbifold Δ is defined as $\mathcal{G} \times_{\mathcal{G}} \mathcal{G}$.

Lemma 3.13. The natural map $\Delta = \mathcal{G} \times_{\mathcal{G}} \mathcal{G} \rightarrow \mathcal{G} \times \mathcal{G}$ is an orbifold embedding.

Proof. We verify that Δ above is a subgroupoid in the sense of Definition 3.1. It suffices to prove this when \mathcal{G} is a global quotient orbifold $G \ltimes M$. In this case, Δ is given by $(G \times G) \ltimes (\sqcup_g \Delta_g)$, where $\Delta_g = \{(x, gx) : x \in M\}$ and $(h, k) \in G \times G$ takes (x, g, gx) to (hx, kgh^{-1}, kgx) (The second terms in the triples are used to distinguish (x, gx) from (x, ghx) for $h \in G_x$). The most natural choice of orbifold morphisms will be $\phi_0 : (x, g, gx) \mapsto (x, gx) \in M \times M$ and $\phi_1 : (h, k) \mapsto (h, k) \in G \times G$. ϕ_0 is clearly an immersion. Note that ϕ_1 is injective.

Choose a point $p = (x, y)$ in $\text{im}(\phi_0)$. Let U be a small connected neighborhood of p in $M \times M$, which is preserved under the $(G \times G)_p$ -action. Since (x, y) is in the image of ϕ_0 , there is some g in G satisfying $y = gx$. So, in particular $\phi_0^{-1}(x, y) = \{(x, g', g'x) \mid g'x = gx = y\}$. $(G \times G)_p$ acts on $\phi_0^{-1}(p)$ transitively since $(g^{-1}g', e)$ sends $(x, g', g'x) \in \phi_0^{-1}(p)$ to (x, g, gx) and $g^{-1}g' \in G_x$ when $g'x = gx$.

Let q denote $(x, g, gx) \in \phi_0^{-1}(p)$. Note that $(G \times G)_q = \{(h, k) : h \in G_x, k \in G_{gx}, kgh^{-1} = g\}$. Let V_g be the connected component of $\phi_0^{-1}(U)$ which contains q (V_g is given by $\Delta_g \cap \phi_0^{-1}(U)$). We define a smooth map ψ from $(G \times G)_p \times V_g$ to $\phi_0^{-1}(U)$ by

$$(3.2) \quad \psi : ((h, k), (x', g, gx')) \mapsto (hx', kgh^{-1}, kgx').$$

Then, ψ is $(G \times G)_p$ -equivariant by the definition of the $G \times G$ -action on Δ . Since U is preserved under the $(G \times G)_p$ and V_g is a connected component, ψ should be surjective.

Suppose two different points

$$q_1 = ((h_1, k_1), (x_1, g, gx_1)) \quad \text{and} \quad q_2 = ((h_2, k_2), (x_2, g, gx_2))$$

in $(G \times G)_p \times V_g$ are mapped to the same point in $\phi_0^{-1}(U)$ by ψ . This happens precisely when $(h_2^{-1}h_1, k_2^{-1}k_1)$ sends (x_1, g, gx_1) to (x_2, g, gx_2) . In particular, we have $(h_2^{-1}h_1, k_2^{-1}k_1) \in (G \times G)_q$. Therefore, ψ descends to a map

$$\bar{\psi} : (G \times G)_p \times_{(G \times G)_q} V_g \rightarrow \phi^{-1}(U)$$

which is bijective (Here, $(a, b) \in (G \times G)_q$ acts on the first factor of $(G \times G)_p \times V_g$ by $(h, k) \mapsto (ha^{-1}, kb^{-1})$). Since the $(G \times G)_q$ -action and the $(G \times G)_p$ -action on $(G \times G)_p \times V_g$ commute, the $(G \times G)_p$ -equivariance of ψ implies that of $\bar{\psi}$. \square

4. Inertia orbifolds and orbifold embeddings

In this section we show that given an orbifold embedding, there is an induced orbifold embedding between their inertia orbifolds under abelian assumption.

First, let us recall inertia orbifolds. The following diagram defines a smooth manifold $\mathcal{S}_{\mathcal{G}}$, which can be interpreted intuitively as a set of loops (i.e., elements

of local groups) in \mathcal{G} :

$$(4.1) \quad \begin{array}{ccc} \mathcal{S}_{\mathcal{G}} & \longrightarrow & G_1 \\ \downarrow \beta & & \downarrow (s,t) \\ G_0 & \xrightarrow{\text{diag}} & G_0 \times G_0. \end{array}$$

Then, the inertia orbifold $\Lambda\mathcal{G}$ will be an action groupoid $\mathcal{G} \ltimes \mathcal{S}_{\mathcal{G}}$, i.e.,

$$\begin{aligned} (\Lambda\mathcal{G})_0 &= \mathcal{S}_{\mathcal{G}}, \\ (\Lambda\mathcal{G})_1 &= G_1 \times_{G_0} \mathcal{S}_{\mathcal{G}}, \end{aligned}$$

where for $h \in G_1$ the induced map $h : \beta^{-1}(s(h)) \rightarrow \beta^{-1}(t(h))$ is given by the conjugation. More precisely, for any $g \in \beta^{-1}(s(h))$, set $h(g) = hgh^{-1}$. This gives a target map from $(\Lambda\mathcal{G})_1$ to $(\Lambda\mathcal{G})_0$ whereas the source map is simply the projection to the second factor of $(\Lambda\mathcal{G})_1$. Note that $\beta^{-1}(s(h))$ and $\beta^{-1}(t(h))$ are the sets of loops in \mathcal{G} based at $s(h)$ and $t(h)$, respectively. Similarly, one can define $\mathcal{S}_{\mathcal{H}}$ and $\Lambda\mathcal{H}$ for a suborbifold \mathcal{H} of \mathcal{G} .

Now, let us see how $\phi : \mathcal{H} \rightarrow \mathcal{G}$ induces a morphism $\Lambda\phi$ between inertia orbifolds. $\Lambda\phi_0$ should be a map from $(\Lambda\mathcal{H})_0 = \mathcal{S}_{\mathcal{H}}$ to $(\Lambda\mathcal{G})_0 = \mathcal{S}_{\mathcal{G}}$. Suppose (h, y) , $y = \beta(h) \in H_0$ is a loop $h : y \rightarrow y$ in \mathcal{H} . Then, the image of this loop is $(\phi_1(h), \phi_0(y))$ or, $\phi_1(h) : \phi_0(y) \rightarrow \phi_0(y)$, i.e.,

$$\Lambda\phi_0 : (h, y) \mapsto (\phi_1(h), \phi_0(y)).$$

$\Lambda\phi_1$ maps $(h', h) \in (\Lambda\mathcal{H})_1 = H_1 \times_{H_0} \mathcal{S}_{\mathcal{H}}$ as follows:

$$\Lambda\phi_1 : (h', h) \mapsto (\phi_1(h'), \phi_1(h)).$$

If $h : y \rightarrow y$, then $\phi_1(h) : \phi_0(y) \rightarrow \phi_0(y)$.

Lemma 4.1. *If \mathcal{G} is abelian, i.e., G_x is an abelian group for each $x \in \mathcal{G}_0$, then $\Lambda\mathcal{H}$ -action on $\Lambda\phi_0^{-1}(g, x)$ is transitive.*

Proof. To observe the local behavior of $\Lambda\phi$, we use the local model of embeddings. Near $y \in H_0$, the local model and the morphism, again denoted by ϕ , is given as follows:

$$\phi : G_x \ltimes (G_x \times_{H_y} V_y) \rightarrow G_x \ltimes U_x,$$

where V_y and U_x are suitable neighborhoods of y and x , respectively and $x = \phi_0(y)$. Note that $\phi_0 : G_x \times_{H_y} V_y \rightarrow U_x$ is given as $\phi_0[g, y'] = g \cdot \phi_0(y')$ and $\phi_1 = (id, \phi_0) : G_x \times (G_x \times_{H_y} V_y) \rightarrow G_x \times U_x$. One can easily check that ϕ is well-defined.

Recall that we assumed ϕ_1 to be injective and identify H_y as a subgroup of G_x . We observe the fiber $\Lambda\phi_0^{-1}(g, x)$ for a loop $g : x \rightarrow x$ in \mathcal{G} in these local models.

In our local model, any objects in $\Lambda\phi_0^{-1}(g, x)$ can be written as $(g, [g', y])$ for some $g' \in G_x$. Suppose that $(g, [g_1, y])$ and $(g, [g_2, y])$ are distinct objects in

$\Lambda\phi_0^{-1}(g, x)$. Now we want to find $k \in \Lambda\mathcal{H}_1$ which sends $(g, [g_1, y])$ to $(g, [g_2, y])$, i.e., k such that $k \cdot (g, [g_1, y]) = (g, [g_2, y])$ or, equivalently $(kgk^{-1}, [kg_1, y]) = (g, [g_2, y])$. This can be achieved simply by choosing $k = g_2g_1^{-1}$. \square

For general \mathcal{G} , $\Lambda\mathcal{H}$ -action on $\Lambda\phi_0^{-1}(g, x)$ is not necessarily transitive. In the last paragraph of the proof of the lemma, the abelian assumption is crucial to find $k \in \Lambda\mathcal{H}_1$ satisfying $(kgk^{-1}, [kg_1, y]) = (g, [g_2, y])$. If G_x is not abelian, such k may not exist. One may try with $k = g_2g_1^{-1}$ which sends $[g_1, y]$ to $[g_2, y]$, but the loop kgk^{-1} is different from g if k does not commute with g . See the following example.

Example 4.2. Let G be the subgroup of $\mathrm{SL}(2, \mathbb{C})$ generated by

$$(4.2) \quad a = \begin{pmatrix} \rho & 0 \\ 0 & \rho^{-1} \end{pmatrix}, \quad b = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

where $\rho = e^{\pi i/3}$. Then G is the binary dihedral group of order 12. Consider its fundamental representation on \mathbb{C}^2 . Relations on generators a and b are given by

$$a^6 = b^4 = 1, \quad bab^{-1} = a^{-1}, \quad a^3 = b^2.$$

Let V be the first coordinate axis in \mathbb{C}^2 . Then, the subgroup H of G generated by a acts on V . Now,

$$G \ltimes (G \times_H V)$$

gives rise to an orbifold embedding into $[\mathbb{C}^2/G]$ whose image is the union of two coordinate axes in \mathbb{C}^2 . Note that on the level of inertia, $(a, [e, 0])$ and $(a, [b, 0])$ in $\Lambda(G \ltimes (G \times_H V))$ are both mapped to $(a, (0, 0))$ in $\Lambda[\mathbb{C}^2/G]$ by the induced map between inertias.

We claim that there is no arrow between $(a, [e, 0])$ and $(a, [b, 0])$ in the inertia $\Lambda(G \ltimes (G \times_H V))$ and therefore, the induced map is not an orbifold embedding. Such an arrow would first send $[e, 0]$ to $[b, 0]$ and hence, it would be of the form bh for some $h \in H$. This arrow sends the loop a at $[e, 0]$ to the loop $(bh)a(bh)^{-1}$ at $[b, 0]$. However, for any $h \in H$, $(bh)a(bh)^{-1} = bab^{-1} = a^{-1}$ since H is abelian.

Finally we prove that $\Lambda\phi : \Lambda\mathcal{H} \rightarrow \Lambda\mathcal{G}$ satisfies the condition (2) of the orbifold embedding (3.1) under the abelian assumption.

Proposition 4.3. *Given an orbifold embedding $\phi : \mathcal{H} \rightarrow \mathcal{G}$, consider the induced map between inertia orbifolds $\Lambda\phi : \Lambda\mathcal{H} \rightarrow \Lambda\mathcal{G}$. If \mathcal{G} is an abelian orbifold, then $\Lambda\phi$ is again an orbifold embedding.*

Proof. The sector fixed by the loop $h \in H_1$ should be mapped through ϕ_0 to the sector fixed by $\phi_1(h) \in G$. So $\Lambda\phi_{0*}$ is essentially the same as ϕ_{1*} , which sends tangent vector to the sector fixed by $h \in H_1$ to the one determined by $\phi_1(h)$. Therefore $\Lambda\phi_0$ is an immersion.

The only non-trivial part is the second condition of the orbifold embedding. For this, we can work on local charts. Suppose $\phi_0(y) = x$ for $y \in \mathcal{H}_0, x \in \mathcal{G}_0$,

and we fix a neighborhood U_x of x in G_0 from the embedding property of ϕ , so that $H|_{\phi^{-1}(U_x)}$ can be identified with the action groupoid $G_x \ltimes (G_x \times_{H_y} V_y)$. We may identify H_y as a subgroup of G_x via the embedding map.

We fix an element $g \in H_y \subset G_x$. In general, the local chart of inertia orbifold $\Lambda\mathcal{G}$ near $(g, x) \in \Lambda\mathcal{G}_0$ can be written as

$$(4.3) \quad C_G(g) \ltimes (U_x^g \times \{g\}),$$

where U_x^g is the set of g -fixed points in U_x and $C_G(g) = \{h \in G_x \mid hg = gh\}$ acts on U_x^g by the left multiplication. We put $\{g\}$ in (4.3) to indicate the sector in the inertia orbifold $\Lambda\mathcal{G}$, and we will suppress it for notational simplicity in the following.

In our case, $C_G(g) = G_x$ since \mathcal{G} is abelian. We rewrite the local chart (4.3) as

$$(4.4) \quad G_x \ltimes U_x^g.$$

Choose an element $(g, [e, y])$ in $\Lambda\phi_0^{-1}(g, x)$. Then V_y^g is an open neighborhood of $(g, [e, y])$ and

$$\Lambda\mathcal{H}|_{V_y^g} \cong H_y \ltimes V_y^g.$$

Note that $H_y = C_H(g) := \{h \in H_y \mid hg = gh\}$, since H_y is a subgroup of the abelian group G_x .

We claim that the inverse image of the local chart (4.4) of (g, x) by $\Lambda\phi^{-1}$ can be written as

$$G_x \ltimes (G_x \times_{H_y} V_y^g).$$

To see this, we only need to show that the twisted g -sector in the local chart of $\mathcal{H}|_{\phi^{-1}(G_x \ltimes U_x)}$ is isomorphic to $G_x \ltimes (G_x \times_{H_y} V_y^g)$. It can be checked as follows:

If $[g', z] \in G_x \times_{H_y} V_y$ is a fixed point of g , then $[gg', z] = [g', z]$. By definition, this happens if $gg' = g'h$ and $hz = z$ for some $h \in H_y$. This is equivalent to $(g')^{-1}gg'z = z$, and by the abelian assumption on G_x , $gz = z$. Hence, objects in g -twist sector of $\mathcal{H}|_{\phi^{-1}(G_x \ltimes U_x)}$ are contained in $G_x \times_{H_y} V_y^g$. Conversely, using the condition that G_x is abelian and $g \in H_y$, it follows that any element $[g', z] \in G_x \times_{H_y} V_y^g$ is fixed by g .

By the definition of arrows in an inertia groupoid, $G_x \times (G_x \times_{H_y} V_y^g)$ is the arrow space of the g -twisted sector of $\mathcal{H}|_{\phi^{-1}(G_x \ltimes U_x)}$ with an obvious action map, and this proves the proposition. \square

5. Orbifold embeddings and equivariant immersions

In this section we show that equivariant immersions which are *strong* (which will be defined later) give rise to orbifold embeddings between orbifold quotients.

First, let us review orbifold quotients and its groupoid analogue, translation groupoids. Let G be a compact Lie group which acts on M smoothly. The quotient $[M/G]$ naturally has a structure of a translation groupoid.

Definition 5.1. Suppose a Lie group G acts smoothly on a manifold M from the left. The translation groupoid $[G \ltimes M]$ associated to this group action is defined as follows. Let $(G \ltimes M)_0 := M$ and $(G \ltimes M)_1 := G \times M$, with $s : G \times M \rightarrow M$ the projection and $t : G \times M \rightarrow M$ the action. The other structure maps are defined in the natural way.

In particular, we are interested in group actions which give rise to an orbifold groupoid structure.

Definition 5.2. A G -action on M is said to be locally free if the isotropy groups G_p are discrete for all $p \in M$.

Now we assume that the G -action on M is locally free. The compactness of G implies that G_x is finite for all $x \in M$. Since G acts on M locally freely, we have a representation of $[M/G]$ as an orbifold groupoids in the following manner which is called the slice representation in [7].

Proposition 5.3. *For any translation groupoid $[M/G]$, there is an orbifold groupoid \mathcal{G} with an equivalence groupoid homomorphism $p : \mathcal{G} \rightarrow [M/G]$.*

Proof. By the slice theorem, we can cover M by a collection of G -invariant open sets $\{U_i\}$ with G -equivariant diffeomorphisms

$$\psi_i : G \times_{G_i} V_i \rightarrow U_i,$$

where V_i is a normal slice with local action of $G_i \leq G$. Define \mathcal{G} as follows. Let $G_0 := \sqcup_i V_i$ be the disjoint union of all the V_i , and define a map $p : G_0 \rightarrow M$ as $p(i, v) := \psi_i([1, v])$. Define G_1 as the pullback bundle of following diagram:

$$\begin{array}{ccc} G_1 & \longrightarrow & G \times M \\ (s,t) \downarrow & & \downarrow (s,t) \\ G_0 \times G_0 & \xrightarrow{(p,p)} & M \times M \end{array}$$

Then groupoid homomorphism $p : \mathcal{G} \rightarrow [M/G]$ is an equivalence. See the proof of Theorem 4.1 in [7] for more details. \square

The converse in general still remains as a conjecture. The conjecture was partially proven in the case of effective orbifold groupoids (Theorem 1.23 of [1]).

Conjecture 5.4. Every orbifold groupoid can be represented by translation groupoid with locally free group action.

Now, let us recall the definition of an equivariant immersion and introduce what we call strong equivariant immersion.

Definition 5.5. Let N, M be G -manifolds. A G -equivariant immersion from N into M is a smooth map $\iota : N \rightarrow M$ such that

- (1) the derivative $d\iota : T_x N \rightarrow T_{\iota(x)} M$ is injective at every point in N ;

$$(2) \quad \iota(g \cdot x) = g \cdot \iota(x).$$

When ι is an equivariant immersion, the inverse image of $p \in \iota(N) \subset M$ admits a natural G_p action. If $q \in N$ is a point in $\iota^{-1}(p)$, then for $g \in G_p$

$$(5.1) \quad \iota(g \cdot q) = g \cdot \iota(q) = g \cdot p = p.$$

Definition 5.6. Suppose the G -action on N is locally free and $\iota : N \rightarrow M$ be a G -equivariant immersion. We call ι a *strong* G -equivariant immersion if for every $p \in M$, G_p action on $\iota^{-1}(p)$ is transitive.

Here is an example. Let N be a submanifold of M , which may not be necessarily preserved by G -action. We take G copies of N , and denote it by \tilde{N} , i.e., $\tilde{N} = G \times N$. \tilde{N} admits a natural G -action

$$(5.2) \quad g : (h, x) \mapsto (gh, x)$$

for $g, h \in G$ and $x \in N$. An immersion $\iota : \tilde{N} \rightarrow M$ defined by $\iota(g, x) = g \cdot x$ is then G -equivariant.

Lemma 5.7. *The G -equivariant immersion $\tilde{\iota} : \tilde{N} \rightarrow M$ obtained above is strong if and only if*

$$(5.3) \quad N \cap g \cdot N = N^g$$

for all $g \in G$.

Proof. From the definition of ι , only the image under ι of a point in $h \cdot N \cap g \cdot N$ can have a multiple fiber. It suffices to consider the case when $h = 1$ and other cases can be handled by a parallel argument. Let y be a point in $N \cap g \cdot N$. Then, there exists $x \in N$ such that $g \cdot x = y$. Observe that $(1, y)$ and (g, x) in \tilde{N} maps to the same point $y \in M$. For ι to be strong, there should be a group element mapping $(1, y)$ to (g, x) , and from (5.2), this implies $x = y$. Therefore, $g \cdot x = x$ and, hence $x \in N^g$. \square

When a nontrivial subgroup G_N of G preserves N but does not fix N , then condition (5.3) cannot be satisfied in general. However, we may try to use the minimal number of copies of N . Define G_N so that we have the property, $g \cdot N = h \cdot N$ if and only if $g^{-1}h \in G_N$. Thus, for an element α of the coset space G/G_N , αN is well defined. Let

$$\tilde{N} = \bigcup_{\alpha \in G/G_N} \alpha N \times \{\alpha\}.$$

\tilde{N} is a G -space by letting

$$g : (x, \alpha) \mapsto (g \cdot x, g \cdot \alpha).$$

Obviously, the natural immersion $\iota : \tilde{N} \rightarrow M$, $\iota(x, \alpha) = x$, is G -equivariant.

With this construction, we can interpret the orbifold diagonal for a global quotient orbifolds (cf. 3.12) as a strong equivariant immersion: Suppose a finite group G acts on M and let N be the diagonal submanifold of $M \times M$. Then,

$G \times G/\Delta_G$ parametrizes sheets of the domain of the immersion where $\Delta_G = \{(g, g) \mid g \in G\}$, i.e.,

$$\tilde{N} = \bigcup_{\alpha \in G \times G/\Delta_G} \alpha N \times \{\alpha\}.$$

To see that $\tilde{N} \rightarrow M \times M$ is strong, assume $(x, gx, [1, g])$ and $(x, hx, [1, h])$ are mapped to the same point z in $M \times M$ ($[1, g], [1, h] \in G \times G/\Delta_G$). Then, $(h^{-1}g, 1)$ belongs to the local isotropy of $z = (x, gx) = (x, hx) \in M \times M$ and it sends $(x, gx, [1, g])$ to $(x, hx, [1, h])$ since $h^{-1}gx = x$ and

$$(h^{-1}g, 1)[1, g] = [h^{-1}g, g] = [h^{-1}, 1] = [1, h].$$

One nice property which follows from the strong condition is that the strong equivariant immersions always induce injective maps between the quotient spaces.

Lemma 5.8. *If $\iota : N \rightarrow M$ is a strong G -equivariant immersion between two G -spaces N and M , then,*

$$|\iota| : |N/G| \rightarrow |M/G|$$

is injective.

Proof. Let $|\iota|(\overline{q_1}) = |\iota|(\overline{q_2})$ in $|M/G|$ for $\overline{q_i} \in |N/G|$. Then,

$$\iota(q_1) = g \cdot \iota(q_2)$$

for some $g \in G$. Denote $\iota(q_1)$ by p . We have to find $h \in G$ such that $h \cdot q_1 = q_2$. Observe that

$$\iota(g \cdot q_2) = g \cdot \iota(q_2) = \iota(q_1) = p,$$

which implies that $g \cdot q_2$ and q_1 lie over the same fiber $\iota^{-1}(p)$ of ι . Since ι is strong, there is $h' \in G_p$ such that $h' \cdot q_1 = g \cdot q_2$. By letting $h = g^{-1}h'$, we prove the claim. \square

Next, we use the local model of strong G -equivariant immersion to construct an orbifold embedding.

Proposition 5.9. *Let $\iota : N \rightarrow M$ be a strong G -equivariant immersion between two G -manifolds with locally free G -actions. Then, there exist orbifold groupoid representations \mathcal{H} and \mathcal{G} of $|N/G|$ and $|M/G|$ respectively so that ι induces an orbifold embedding $\phi_\iota : \mathcal{H} \rightarrow \mathcal{G}$ whose underlying map between quotient spaces is injective.*

We will give a proof at the end of this section, after we discuss local models. The following lemma is an analogue of standard slice theorem.

Lemma 5.10. *Let M be a manifold on which a compact Lie group G acts locally freely. Suppose $\iota : N \rightarrow M$ is a G -equivariant immersion. For $q \in N$ and $p = \iota(q) \in M$, we can find a G -invariant neighborhood \tilde{U}_p of $G \cdot p$ in M and \tilde{V}_q of $G \cdot q$ in N with the following properties:*

- (i) *There are normal slices U_p and V_q to $G \cdot p$ and $G \cdot q$ at p and q respectively such that*

$$(5.4) \quad \tilde{U}_p \cong G \times_{G_p} U_p, \quad \tilde{V}_q \cong G \times_{G_q} V_q.$$

- (ii) *There is an G_q -equivariant embedding $e : V_q \rightarrow U_p$ such that the diagram*

$$\begin{array}{ccc} \tilde{V}_q & \xrightarrow{\iota} & \tilde{U}_p \\ \cong \downarrow & & \downarrow \cong \\ G \times_{G_q} V_q & \xrightarrow{[id, e]} & G \times_{G_p} U_p \end{array}$$

commutes where the map on the second row is given by $(g, v) \mapsto (g, e(v))$.

Proof. This is a relative version of the slice theorem (see for example Theorem B.24 of [3]). We briefly sketch the construction of the slice here. Fix any G -invariant metric ξ on M . The exponential map E identifies a neighborhood of p in M with a neighborhood of 0 in $T_p M$. Moreover, ξ induces a decomposition

$$(5.5) \quad T_p M \cong T_p(G \cdot p) \oplus W,$$

where W is normal to the orbit and hence, it is equipped with the linear G_p action (coming from the one on $T_p M / T_p(G \cdot p)$). Let $U_p \subset W$ be a G_p -invariant small disk in W around the origin on which E is a diffeomorphism. Now,

$$\psi : G \times_{G_p} U_p \rightarrow M, \quad [g, u] \mapsto g \cdot E(u)$$

is well defined and G -equivariant. Since ψ is a local diffeomorphism at the point $[e, 0]$, G -equivariance implies that it is a local diffeomorphism at all points of the form $[g, 0]$. One can check that ψ is indeed injective if U_p is sufficiently small (see the proof of Theorem B.24 in [3] for details).

To get the relative version, we pull back ξ to N by ι . Since ι is an immersion, $\iota^* \xi$ gives a metric on N . From the G -equivariant injection $T_q N \xrightarrow{\iota_*} T_p M$, we can choose a decomposition compatible with (5.5):

$$T_q N \cong T_p(G \cdot q) \oplus W',$$

i.e., ι_* is decomposed as

$$\iota_* = (\iota_*^O, \iota_*^N) : T_q(G \cdot q) \oplus W' \rightarrow T_p(G \cdot p) \oplus W.$$

Note that $G_q \subset G_p$ and ι_*^N is G_q -equivariant. Let V_q be the inverse image of U_p by ι_*^N . We may assume that $G \times_{G_q} V_q$ is diffeomorphic to a neighborhood of $G \cdot q$ by shrinking U_p if necessary. Thus, we proved (i).

Finally, by letting e the restriction of ι_*^N to V_q , we get (ii). \square

The following lemma provides the local model which is needed for the orbifold embedding.




FIGURE 5. Slices

Lemma 5.11. *Under the setting of Lemma 5.10, assume further that ι is strong. Then, we can find a G_p -invariant neighborhood $\bar{U}_p(\subset \tilde{U}_p)$ of p in M and a G_q -invariant neighborhood $\bar{V}_q(\subset \tilde{V}_q)$ of q in N such that there is a G_p -equivariant isomorphism*

$$(5.6) \quad \iota^{-1}(\bar{U}_p) \cong G_p \times_{G_q} \bar{V}_q,$$

where the G_q -action on $G_p \times \bar{V}_q$ is given by

$$h \cdot (g, x) = (gh^{-1}, h \cdot x).$$

Proof. We take a *product type* neighborhood $\bar{V}_q(\subset \tilde{V}_q)$ and $\bar{U}_p(\subset \tilde{U}_p)$ of q and p as follows. We first identify the orbit $G \cdot q$ with G/G_q where q corresponds to the image of identity $[e]$ in G/G_q and $G \cdot p$ with G/G_p in a similar way. Then the tubular neighborhood $G \times_{G_q} V_q$ can be regarded as a fiber bundle over G/G_q . Take an open neighborhood O_q of $[e]$ in G/G_q which is invariant under the left G_q -action on G/G_q . As $[e]$ is fixed by this G_q -action, one can for instance choose left G_q -invariant metric on G/G_q and then, take O to be a small open ball around $[e]$. We may assume O is small enough so that

$$(5.7) \quad g \cdot O_q \cap O_q = \emptyset$$

for nontrivial $g \in G_p \setminus G_q$ (Note that G_p also acts on G/G_q from the left). This is possible since G_p is finite. Let O_p be the image of O_q by the map $G/G_q \rightarrow G/G_p$. (5.7) implies that the map $O_q \rightarrow O_p$ is an embedding.

Finally, we define \bar{V}_q and \bar{U}_p to be open neighborhoods of q and p , respectively, such that the following diagrams are cartesian (see Figure 5).

$$(5.8) \quad \begin{array}{ccc} \bar{V}_q & \hookrightarrow & G \times_{G_q} V_q \\ \downarrow & & \downarrow \\ O_q & \hookrightarrow & G/G_q \end{array} \quad \begin{array}{ccc} \bar{U}_p & \hookrightarrow & G \times_{G_p} V_p \\ \downarrow & & \downarrow \\ O_p & \hookrightarrow & G/G_p \end{array}$$

Then, by (5.7) we have

$$(5.9) \quad g \cdot \bar{V}_q = \begin{cases} \bar{V}_q & \text{if } g \in G_q, \\ \text{disjoint from } \bar{V}_q & \text{if } g \in G_p \setminus G_q \end{cases}$$

(More precisely, \bar{V}_q and \bar{U}_p are image of these fiber products under the isomorphisms shown in (i) of the previous lemma). Observe that $\iota|_{\bar{V}_q} : \bar{V}_q \rightarrow \bar{U}_p$ is an embedding since both $O_q \rightarrow O_p$ and $V_q \rightarrow U_p$ are embeddings.

Since the G_p -action on $\iota^{-1}(p)$ is transitive, there is $|G_p|/|G_q|$ -open subsets of N (isomorphic to \bar{V}_q) which are mapped to \bar{U}_p . By (5.9), $\iota^{-1}(\bar{U}_p)$ is the disjoint union of these open subsets of N .

Now, define $\tilde{\phi} : G_p \times \bar{V}_q \rightarrow \iota^{-1}(\bar{U}_p)$ by

$$\begin{aligned} \tilde{\phi} : G_p \times \bar{V}_q &\longrightarrow \iota^{-1}(\bar{U}_p) \\ (g, x) &\longmapsto g \cdot x \end{aligned}$$

This map is well defined because ι is G_p -equivariant and \bar{U}_p is G_p -invariant subset of M . Furthermore, $\tilde{\phi}$ is surjective (and hence a submersion) by the strong condition of ι . It remains to show that it is injective up to G_q -action.

Suppose $\tilde{\phi}$ sends (g, x) and (g', x') in $G_p \times \bar{V}_q$ to the same point in $\iota^{-1}(\bar{U}_p)$. Then $g \cdot x = g' \cdot x'$, equivalently $(g')^{-1} g \cdot x = x'$. Note that both x and x' belong to \bar{V}_q and $(g')^{-1} g \in G_p$. From the dichotomy (5.9), we have $(g')^{-1} g = h$ for some $h \in G_q$. Therefore, $g' = gh^{-1}$ and $x' = h \cdot x$ for $h \in G_q$. We conclude that $\tilde{\phi}$ is indeed a principal G_q -bundle and the isomorphism (5.6) follows. \square

Remark 5.12. Note that the induced map $\phi : G_p \times_{G_q} \bar{V}_q \rightarrow \iota^{-1}(\bar{U}_p)$ is G_p -equivariant by definition.

Proof of Proposition 5.9. Suppose we have a strong G -equivariant immersion $\iota : N \rightarrow M$. By equivariance, ι induces a map $\phi'_\iota : [N/G] \rightarrow [M/G]$ and $|\phi'_\iota| : N/G \rightarrow M/G$ is clearly injective.

Consider a point \bar{p} in $[M/G]$ and let $\pi_M(p) = \bar{p}$ for the quotient map $\pi_M : M \rightarrow [M/G]$. From the definition of strong equivariant immersion, the group action on $\iota^{-1}(p)$ is transitive. If $q \in N$ maps to p , then there exists a G_p -invariant product type neighborhood \bar{U}_p of p in M and G_q -invariant \bar{V}_q of q in N which satisfies (5.6) from the lemma 5.11. We add all such slices U_p and V_q into the slice representations of $[N/G]$ and $[M/G]$ to get orbifold groupoids \mathcal{H} and \mathcal{G} .

Now the previous remark implies that

$$\mathcal{H}|_{\phi_0^{-1}(U_p)} \cong G_p \ltimes (G_p \times_{G_q} V_q).$$

Essential injectivity of the resulting orbifold morphism follows directly from Lemma 5.8. Since the other conditions in Definition 3.1 are automatic, we get an orbifold embedding $\phi : \mathcal{H} \rightarrow \mathcal{G}$. \square

Section 7 will be devoted to prove the converse of this proposition.

6. Hilsum-Scandalis maps

We give brief review on Hilsum-Scandalis map, and we refer readers to [8] and [5] for further details. We first recall the definition of the action of a orbifold groupoid on manifolds

Definition 6.1. Let \mathcal{G} be an orbifold groupoid. A left \mathcal{G} -space is a manifold E equipped with an action by \mathcal{G} . Such an action is given by two maps:

- an anchor $\pi : E \rightarrow G_0$;
- an action $\mu : G_1 \times_{G_0} E \rightarrow E$.

The latter map is defined on pairs (g, e) with $\pi(e) = s(g)$, and written $\mu(g, e) = g \cdot e$. It satisfies the usual identities for an action:

- $\pi(g \cdot e) = t(g)$;
- $1_x \cdot e = e$;
- $g \cdot (h \cdot e) = (gh) \cdot e$

for $x \xrightarrow{h} y \xrightarrow{g} z$ in G_1 with $\pi(e) = x$.

A right \mathcal{G} -space is the same thing as a left \mathcal{G}^{op} -space, where \mathcal{G}^{op} is the opposite groupoid obtained by exchanging the roles of the target and source maps.

Definition 6.2. A left \mathcal{G} -bundle over a manifold M is a manifold R with smooth maps

$$\begin{array}{ccc} R & \xrightarrow{\rho} & M \\ \downarrow r & & \\ G_0 & & \end{array}$$

and a left \mathcal{G} -action μ on R , with anchor map $r : R \rightarrow G_0$, such that $\rho(gx) = \rho(x)$ for any $x \in R$ and any $g \in G_1$ with $r(x) = s(g)$.

Such a bundle R is principal if

- (1) ρ is a surjective submersion,
- (2) the map $(\pi_1, \mu) : R \times_{r, G_0, s} G_1 \rightarrow R \times_M R$, sending (x, g) to (x, gx) , is a diffeomorphism.

Definition 6.3. A Hilsum-Scandalis map $\mathcal{G} \rightarrow \mathcal{H}$ is represented by a principal left \mathcal{H} -bundle R over G_0

$$\begin{array}{ccc} R & \xrightarrow{\rho} & G_0 \\ \downarrow r & & \\ H_0 & & \end{array}$$

which is also a right \mathcal{G} -bundle (over H_0), and the right \mathcal{G} -action commutes with the \mathcal{H} -action. R is called the Hilsum-Scandalis bibundle.

Definition 6.4. For two bibundles $R : \mathcal{G} \rightarrow \mathcal{H}$ and $Q : \mathcal{H} \rightarrow \mathcal{K}$, their composition is defined by the quotient of the fiber product $Q \times_{H_0} R$ by the action of \mathcal{H} :

$$(6.1) \quad Q \circ R := (Q \times_{H_0} R) / H_1,$$

where the action of H_1 on $Q \times_{H_0} R$ is given by $h \cdot (q, r) := (qh, h^{-1}r)$. Since the left action of \mathcal{H} on R is principal, the action of \mathcal{H} on $Q \times_{H_0} R$ is free and proper; hence, the $Q \circ R$ is a smooth manifold. It also admits a principal \mathcal{K} -bundle structure with a right \mathcal{G} -action, because \mathcal{H} -action commutes with \mathcal{G} - and \mathcal{K} -actions on R and Q , respectively.

One can compose two Hilsum-Scandalis maps as follows:

Definition 6.5. Two Hilsum-Scandalis maps $P, R : \mathcal{G} \rightarrow \mathcal{H}$ are isomorphic if they are diffeomorphic as left \mathcal{H} - and right \mathcal{G} -bundles: i.e, there is a diffeomorphism $\alpha : P \rightarrow R$ satisfying $\alpha(h \cdot p \cdot g) = h \cdot \alpha(p) \cdot g$ for all $(h, p, g) \in H_1 \times_{H_0} P \times_{G_0} G_1$.

For example, any Lie groupoid homomorphism $\phi : \mathcal{G} \rightarrow \mathcal{H}$ defines a Hilsum-Scandalis map

$$\begin{array}{ccc} R_\phi := H_1 \times_{s, H_0, \phi_0} G_0 & \xrightarrow{\pi_2} & G_0 \\ \downarrow t \circ \pi_1 & & \\ H_0 & & \end{array}$$

where π_1 and π_2 are the projection maps. One can easily check that π_2 is principal in this case. We will use this construction crucially in the next section to construct an equivariant immersion from an orbifold embedding.

Remark 6.6.

- (1) Not every Hilsum-Scandalis map is induced from Lie groupoid homomorphisms. In fact, a Hilsum-Scandalis map $R : \mathcal{G} \rightarrow \mathcal{H}$ is isomorphic to some R_ϕ for some Lie groupoid homomorphism $\phi : \mathcal{G} \rightarrow \mathcal{H}$ if and only if the map $\rho : R \rightarrow G_0$ has a global section. See Lemma 3.36 in [5].
- (2) We use slightly different notion of the Hilsum-Skandalis map from [8]. In [8],

$$R_\phi^{ps} = H_0 \times_{\phi_0, G_0, t} G_1$$

is used to construct a Hilsum-Skandalis map from ϕ . Here, we use R_ϕ^* to make it a left G -space. See the following diagrams.

$$\begin{array}{ccc} R_\phi & \xrightarrow{\pi_1} & G_1 \\ \pi_2 \downarrow & & \downarrow s \\ H_0 & \xrightarrow{\phi_0} & G_0 \end{array} \quad \begin{array}{ccc} R_\phi^{ps} & \longrightarrow & H_0 \\ \downarrow & & \downarrow \phi_0 \\ G_1 & \xrightarrow{t} & G_0 \end{array}$$

Now, we want to translate the notion of equivalence in the category of orbifold groupoids into Hilsum-Scandalis maps. We first refer to the following two lemmas from [5].

Lemma 6.7 ([5], Lemma 3.34). *A Lie groupoid homomorphism $\phi : \mathcal{G} \rightarrow \mathcal{H}$ is an equivalence of Lie groupoids if and only if the corresponding R_ϕ is \mathcal{G} -principal.*

Lemma 6.8 ([5], Lemma 3.37). *Let $P : \mathcal{G} \rightarrow \mathcal{H}$ be a Hilsum-Scandalis map. Then, there is a cover $\phi : \mathcal{U} \rightarrow G_0$ and a groupoid homomorphism $f : \phi^*\mathcal{G} \rightarrow \mathcal{H}$ so that*

$$P \circ R_{\tilde{\phi}} \xrightarrow{\simeq} R_f,$$

where $\tilde{\phi} : \phi^*\mathcal{G} \rightarrow \mathcal{G}$ is the induced functor and “ $\xrightarrow{\simeq}$ ” an isomorphism of Hilsum-Scandalis maps. Here, $\phi^*\mathcal{G}$ is the Lie groupoid with $(\phi^*\mathcal{G})_0 = \mathcal{U}$ and $(\phi^*\mathcal{G})_1$ given by

$$\begin{array}{ccc} (\phi^*\mathcal{G})_1 & \longrightarrow & G_1 \\ \downarrow (s,t) & & \downarrow (s,t) \\ \mathcal{U} \times \mathcal{U} & \xrightarrow{(\phi,\phi)} & G_0 \times G_0 \end{array}$$

From the above two lemmas, we obtain the following characterization of Morita equivalence in terms of Hilsum-Scandalis language.

Lemma 6.9. *If a Hilsum-Scandalis map $P : \mathcal{G} \rightarrow \mathcal{H}$ is also right \mathcal{G} -principal, then $f : \phi^*\mathcal{G} \rightarrow \mathcal{H}$ obtained from the above lemma is an equivalence of groupoids. Note that $\tilde{\phi} : \phi^*\mathcal{G} \rightarrow \mathcal{G}$ is trivially an equivalence of groupoids.*

Proof. Note that $R_{\tilde{\phi}}$ is biprincipal, since $\tilde{\phi}$ is an equivalence of groupoids. The composition of two biprincipal bundle $P \circ R_{\tilde{\phi}}$ is also biprincipal, and hence the isomorphic bibundle R_f also biprincipal. Therefore f is an equivalence of groupoids $\phi^*\mathcal{G}$ and \mathcal{H} . \square

The above lemma justifies the notion of the Morita equivalence in the Hilsum-Scandalis setting.

Definition 6.10. A Hilsum-Scandalis map (R, ρ, r) is a Morita equivalence when it is both a principal \mathcal{G} -bundle and a principal \mathcal{H} -bundle.

7. Construction of equivariant immersions from orbifold embeddings

Let \mathcal{H} be an orbifold groupoid, and $\phi : \mathcal{H} \rightarrow [M/G]$ be a groupoid morphism which factors through an orbifold embedding ψ and a slice representation $p :$

$$\mathcal{G} \rightarrow [M/G].$$

$$(7.1) \quad \begin{array}{ccc} \mathcal{H} & \xrightarrow{\quad \phi \quad} & [M/G] \\ & \searrow \psi: \text{orb.emb.} & \nearrow p: \cong \\ & \mathcal{G} & \end{array}$$

With this assumption in this section, we construct a G -equivariant immersion map $\iota: N \rightarrow M$ for some G -manifold N .

Proposition 7.1. *Consider $\mathcal{H}, [M/G], \phi, \psi$ as above. Then, there exists G -space N and a strong G -equivariant immersion $\iota: N \rightarrow M$ such that*

- $[N/G]$ is Morita equivalent to \mathcal{H} .
- the induced map $[N/G] \rightarrow [M/G]$, again denoted by ι , fits into the following diagram of Lie groupoid homomorphisms

$$(7.2) \quad \begin{array}{ccc} [N/G] & \xrightarrow{\quad \iota \quad} & [M/G] \\ \text{Morita } \cong \downarrow & & \nearrow \phi \\ \mathcal{H} & & \end{array}$$

and ι is a G -equivariant immersion.

Proof. From the groupoid structure of $[M/G]$, we have smooth maps

$$\phi_0: H_0 \rightarrow M, \quad \phi_1: H_1 \rightarrow G \times M$$

which are compatible with the structure maps of an orbifold groupoid. As in [8], we interpret ϕ as a Hilsum-Skandalis-type map. So, we define a bibundle R_ϕ as

$$R_\phi := (G \times M) \times_{s, M, \phi_0} H_0.$$

Note that R_ϕ is a smooth manifold since s is a submersion.

$$(7.3) \quad \begin{array}{ccc} R_\phi & \xrightarrow{\quad \pi_1 \quad} & G \times M \\ \pi_2 \downarrow & & \downarrow s \\ H_0 & \xrightarrow{\quad \phi_0 \quad} & M \end{array}$$

This space is first of all smooth and has two maps to H_0 and M ,

$$H_0 \xleftarrow{\pi_2} R_\phi \xrightarrow{\tilde{t} := t \circ \pi_1} M$$

which will be used as anchor maps below (We denote $t \circ \pi_1$ by \tilde{t}).

We define a right \mathcal{H} -action and a left G -action on R_ϕ as follows: Write an element of R_ϕ by $(g, \phi_0(y), y)$ which indicates a point y in H_0 and an arrow $g \in G \times M$ whose source is $\phi_0(y)$. Then,

- For an arrow $h \in H_1$,

$$(g, \phi_0(y), y) \cdot h := (g \circ \phi_1(h), \phi_0(s(h)), s(h));$$

- For $g \in G$,

$$g' \cdot (g, \phi_0(y), y) := (g' \circ g, \phi_0(y), y).$$

R_ϕ is a right \mathcal{H} -space and a left G -space as Figure 6 below shows. As mentioned, the corresponding anchor maps are π_2 and $\tilde{\iota}$, respectively. Indeed, π_2 is a principal left G -bundle (π_2 is a submersion since s in the diagram (7.3) is submersion).




FIGURE 6. the right \mathcal{H} -action and the left G -action on R_ϕ^*

Now, the following are clear from the definition of both actions.

Lemma 7.2. *Two actions defined above have the following properties:*

- (1) *The right \mathcal{H} -action we have defined is free;*
- (2) *The left G -action and the right \mathcal{H} -action commute;*
- (2) *$\tilde{\iota}$ is a G -equivariant map which is invariant under the \mathcal{H} -action.*

Proof. We only show (1) and the others follow from the definition. Suppose h fixes $((g, \phi_0(y)), y) \in R_\phi$. Then h should be an element of H_y and $g \cdot \phi_1(h) = g$, where H_y is a local isotropy group of $y \in V_y$ for some local chart $H_y \ltimes V_y$ of \mathcal{H} . Thus h lies in the kernel of the group homomorphism

$$\phi_1|_{H_y} : H_y \rightarrow G.$$

Note that equivalence map $p : \mathcal{G} \rightarrow [M/G]$ preserves isotropy groups. More precisely,

$$\text{Hom}_G(x, z) \cong \text{Hom}_{[M/G]}(p(x), p(z))$$

for all $x, z \in G_0$. Now it follows that $\phi_1| = p_1 \circ \psi_1|$ is injective, because $\psi_1|$ is injective from the definition of orbifold embedding and p_1 preserves isotropy groups. Hence, the h is the identity. \square

We denote by N the quotient space of R_ϕ by the right \mathcal{H} -action and by $\pi : R_\phi^* \rightarrow N$ the quotient map.

Lemma 7.3. *N is a smooth manifold.*

Proof. This follows directly from the fact that the right \mathcal{H} -action on R_ϕ is free ((2) of Lemma 7.2) and proper (because \mathcal{H} itself is étale and hence proper). \square

From (2) of Lemma 7.2, N admits a left G -action which is induced by G -action on R_ϕ . Since the G -action on M is locally free, so is it on N . Therefore, we get a global quotient orbifold $[N/G]$ from the orbifold embedding $\phi : \mathcal{H} \rightarrow [M/G]$.

Lemma 7.4. *$[N/G]$ is Morita equivalent to \mathcal{H} .*

Proof. Note that we have a Hilsum-Skandalis map $\mathcal{H} \rightarrow [N/G]$ (or, $\pi_2 : [N/G] \rightarrow \mathcal{H}$ from $\pi_2 : R_\phi \rightarrow H_0$):

$$\begin{array}{ccc} R_\phi & \xrightarrow{\pi_2} & H_0 \\ \pi \downarrow & & \\ N & & \end{array}$$

We have shown that π is principal in (2) of Lemma 7.2. It is also obvious from the Hilsum-Skandalis construction that π_2 is principal. So the Hilsum-Skandalis map from R_ϕ is a Morita equivalence. \square

From (3) of Lemma 7.2, we can observe that $\tilde{\iota}$ factors through the quotient space H . Since $\tilde{\iota}$ is a G -equivariant, we get a G -equivariant map $\iota : N \rightarrow M$. Furthermore,

Lemma 7.5. *ι is a G -equivariant immersion.*

$$(7.4) \quad \begin{array}{ccc} R_\phi & \xrightarrow{\tilde{\iota}} & M \\ \pi \downarrow & \nearrow \iota & \\ N & & \end{array}$$

Proof. Since being an immersion is a local property, it suffices to prove it locally. However, we have a nice local model of ι from (3.1). Thus, it is enough to prove it with $\mathcal{H} = G_x \ltimes (G_x \times_{H_y} V_y)$, $\mathcal{G} = G_x \ltimes U_x$, $\psi : G_x \ltimes (G_x \times_{H_y} V_y) \rightarrow G_x \ltimes U_x$ and $p : G_x \ltimes U_x \rightarrow G \ltimes \tilde{U}_x$ for $\tilde{U}_x \cong G \times_{G_x} U_x$ as in the Lemma 5.10.

Then, $R_\phi = (G \times \tilde{U}_x) \times_{s, \tilde{U}_x, \phi_0} (G_x \times_{H_y} V_y)$. We mod it out by the right G_x -action (considered as a local H_1 -action) to get the local shape of N , again denoted by N in this proof. Recall that this G_x -action is given by

$$((k_1, a), [g_1, b]) \cdot g = [(k_1 g, g^{-1} \cdot a), [g^{-1} g_1, b]]$$

for $g \in G_x$, $k_1 \in G$ and $g_1 \in G_x$ where $a = g_1 \cdot \phi_0(b)$. And $\tilde{\iota}$ on \mathcal{R}_ϕ which projects down to ι on N is defined as

$$(7.5) \quad \tilde{\iota}((k_1, a), [g_1, b]) = k_1 \cdot a = k_1 g_1 \cdot \phi_0(b) \in \tilde{\tau}_x,$$

where $\phi = p \circ \psi : \mathcal{H} \rightarrow [M/G]$.

For given $z \in \tilde{U}_x$, we check how many points in N are mapped to z . Suppose

$$\iota[(k_1, a_1), [g_1, b_1]] = z \quad \text{and} \quad \iota[(k_2, a_2), [g_2, b_2]] = z.$$

Up to the G_x -action, we may assume that $g_1 = g_2 = 1$ (recall $N = G_x \setminus R_\phi$). Therefore, we have $k_1 \phi_0(b_1) = k_2 \phi_0(b_2) = z$ by (7.5). This implies that $k_2^{-1} k_1 \in G_x$ since both $\phi_0(b_1)$ and $\phi_0(b_2)$ belong to the normal slice at x . As G_x is finite, there are finitely many k_2 with this property.

Since p_0 is an embedding and every fiber of ψ_0 is finite, $\phi_0 = p_0 \circ \psi_0$ is an immersion whose fibers are all finite as well. Finally, as b_2 lies in the fiber $\phi_0^{-1}(k_2^{-1} k_1 b_1)$, there can exist only finitely many such b_2 's. \square

Finally, we show in the following lemma that the resulting equivariant immersion is strong which will finish the proof of the proposition. \square

Lemma 7.6. $\iota : N \rightarrow M$ constructed above is strong.

Proof. Note that $|N/G| \cong |\mathcal{H}|$ and $|M/G| \cong |\mathcal{G}|$. From the construction in Section 8 (or Section 7), we have

$$\begin{array}{ccc} |N/G| & \xrightarrow{|\iota|} & |M/G| \\ \cong \downarrow & & \downarrow \cong \\ |\mathcal{H}| & \xrightarrow{|\phi|} & |\mathcal{G}| \end{array}$$

Since $|\phi|$ is injective from the definition of the orbifold embedding, $|\iota|$ is injective. \square

8. General case

So far, we have considered a translation groupoid $[M/G]$ as our target space. The construction can be generalized to the case of general orbifolds which we will discuss from now on. We state this as a theorem, first.

Theorem 8.1. Let $\phi : \mathcal{H} \rightarrow \mathcal{G}$ be an orbifold embedding, where \mathcal{G} is Morita equivalent to a translation groupoid $[M/G]$. Then, there exist a manifold N on which the Lie group G acts locally freely such that

- (i) $\mathcal{H} \simeq [N/G]$ and $\mathcal{G} \simeq [M/G]$, and

- (ii) *there exists a G -equivariant immersion $\iota : N \rightarrow M$ which makes the diagram*

$$(8.1) \quad \begin{array}{ccc} \mathcal{H} & \xrightarrow{\phi} & \mathcal{G} \\ \text{Morita} \downarrow & & \downarrow \text{Morita} \\ [N/G] & \xrightarrow{\iota} & [M/G] \end{array}$$

commute.

Remark 8.2. The diagram in (ii) of the theorem can be regarded as a diagram of morphisms in the category of Lie groupoids where we can invert equivalences (see Definition 2.7, or [1] for the precise definition of the morphisms in the category of groupoids).

We proceed the proof of theorem 8.1 as follows:

After fixing a Morita equivalence map

$$\mathcal{G} \xrightarrow{\psi} \mathcal{G}' \xrightarrow{\sigma} [M/G]$$

for some Lie groupoid \mathcal{G}' , we pull back the equivalence map $\psi : \mathcal{G}' \rightarrow \mathcal{G}$ to \mathcal{H} to get $\phi^*\mathcal{G}' = \mathcal{H} \times_{\mathcal{G}} \mathcal{G}'$. Recall that

$$\begin{aligned} (\phi^*\mathcal{G}')_0 &= H_0 \times_{\phi_0, G_0, s} G_1 \times_{t, G_0, \psi_0} G'_0, \\ (\phi^*\mathcal{G}')_1 &= H_1 \times_{s\phi_1, G_0, s} G_1 \times_{t, G_0, s\psi_1} G'_1. \end{aligned}$$

We denote the composition $\sigma \circ pr_2 : \phi^*\mathcal{G}' \rightarrow [M/G]$ by $\tilde{\phi}$. Then,

$$\tilde{\phi}_0 = \sigma_0 \circ (pr_2)_0 \quad \tilde{\phi}_1 = \sigma_1 \circ (pr_2)_1,$$

where pr_2 is the projection from $\phi^*\mathcal{G}' = \mathcal{H} \times_{\mathcal{G}} \mathcal{G}'$ to \mathcal{G}' . We will apply the construction in the previous section to $\tilde{\phi}$.

$$(8.2) \quad \begin{array}{ccc} & & [M/G] \\ & \nearrow \tilde{\phi} & \uparrow \sigma: \text{equiv.} \\ \phi^*\mathcal{G}' & \xrightarrow{pr_2} & \mathcal{G}' \\ \downarrow pr_1 & & \downarrow \psi: \text{equiv.} \\ \mathcal{H} & \xrightarrow{\phi: \text{orb.emb.}} & \mathcal{G} \end{array}$$

First of all $\phi^*\mathcal{G}'$ is equivalent to \mathcal{H} . Pull-back of any equivalence is again an equivalence as shown in Lemma 2.11.

To construct a G -equivariant immersion from $\tilde{\phi} : \phi^*\mathcal{G}' \rightarrow [M/G]$, we introduce the Hilsum-Skandalis bibundle associated to $\tilde{\phi}$ as we did in the previous section. Recall

$$R_{\tilde{\phi}} = (G \times M) \times_{s, M, \tilde{\phi}_0} (\phi^*\mathcal{G}')_0.$$

An element of $R_{\tilde{\phi}}$ consists of the following data:

$$\begin{array}{c} \begin{array}{c} m \\ \odot \\ \downarrow a \\ \downarrow \end{array} \quad \star \left(= [x, \phi_0(x) \xrightarrow{g} \psi_0(z), z] \right) \end{array}$$

where $m \in M$, $a \in G$, $x \in H_0$, $z \in G'_0$ and $\sigma_0(z) = m$. Write \mathbf{r} for this element. Then, $pr_2(\mathbf{r}) = z$ and the G -equivariant map $\tilde{\iota} : R_{\tilde{\phi}} \rightarrow M$ is given by $\tilde{\iota}(\mathbf{r}) = a \cdot \sigma_0(z)$. Recall

$$(8.3) \quad \begin{array}{ccc} R_{\tilde{\phi}} & \xrightarrow{\tilde{\iota}} & M \\ \downarrow & \nearrow \iota & \\ N & & \end{array}$$

where N is obtained from $R_{\tilde{\phi}}$ after taking a quotient by $\phi^*\mathcal{G}'$ -action. Since local groups are preserved by equivalences, restriction of 1-level maps appearing in 8.2 to any local groups are all injective. Then, similar argument as in Lemma 7.2 shows the right $\phi^*\mathcal{G}$ -action on $R_{\tilde{\phi}}$ is free and proper. Note that $\phi^*\mathcal{G}$ is proper since it is equivalent to the proper Lie (indeed, étale) groupoid \mathcal{G} . Therefore, N is a smooth manifold.

It remains to show that the induced G -equivariant map $\iota : N \rightarrow M$ is indeed an immersion. We will directly compute the kernel of $d\tilde{\iota}$. For notational simplicity, we will write τ_* for the derivative $d\tau$ of a smooth map τ between two manifolds.

A tangent vector on $R_{\tilde{\phi}}$ at \mathbf{r} is given by the tuple

$$\mathbf{v} = [(v_l = v_a \oplus v_m, v_m), v_x, v_g, v_z],$$

where $v_l \in T(G \times M)$ and $v_a \in TG$ with the relations

- $s_*(v_l) = v_m$,
- $(\sigma_0)_*(v_z) = v_m$,
- $s_*(v_g) = (\phi_0)_*(v_x)$ $t_*(v_g) = (\psi_0)_*(v_z)$.

Since \mathcal{G} is étale and hence both

$$s_* : T_g G_1 \rightarrow T_{\phi_0(x)} G_0 \quad \text{and} \quad t_* : T_g G_1 \rightarrow T_{\psi_0(z)} G_0$$

are isomorphisms, we may rewrite the third relation as

$$(8.4) \quad \bullet \quad s_*^{-1}(\phi_0)_*(v_x) = t_*^{-1}(\psi_0)_*(v_z) = v_g.$$

From the first relation, it suffices to represent \mathbf{v} as

$$\mathbf{v} = [v_a \oplus v_m, v_x, v_g, v_z].$$

Note that v_m and v_g are determined by v_x and v_z . One can easily check that

$$\tilde{t}_*(\mathbf{v}) = t_*(v_a \oplus v_m) = (v_a)^\# + (L_a)_*(v_m),$$

where $(v_a)^\#$ is a vector field on M generated by the infinitesimal action of v_a on M . For simplicity, we assume that a is the identity element of G . Then,

$$\tilde{t}_*(\mathbf{v}) = (v_a)^\# + v_m.$$

Our goal is to compute the kernel of this map. If we can show that the kernel of \tilde{t}_* lies in the tangent direction of $(\phi^*\mathcal{G}')$ -orbit, then it will imply that ι is an immersion.

To do this, we first characterize the direction of $(\phi^*\mathcal{G}')$ -orbit. By the definition of the right $\phi^*\mathcal{G}'$ action on $R_{\tilde{\phi}}$, we should consider an arrow of $\phi^*\mathcal{G}'$ given by a pair $(h, k) \in H_1 \times G'_1$ such that

$$(8.5) \quad t(h) = x \text{ and } t(k) = z.$$

Considering the infinitesimal version of $\phi^*\mathcal{G}'$ -action carefully, we have the following lemma.

Lemma 8.3. \mathbf{v} is tangent to $(\phi^*\mathcal{G}')$ -orbit if and only if there exists $(v_h, v_k) \in T_h H_1 \times T_k G'_1$ such that

$$(8.6) \quad t_*(v_h) = t_*(v_k) = 0$$

and

$$(8.7) \quad \mathbf{v} = [(\sigma_1)_*(v_k), s_*(v_h), v_g, s_*(v_k)]$$

(Here, we do not specify v_g since they are completely determined by other components (8.4)).

Remark 8.4. In the equation (8.6), $t_*(v_h) = 0$ implies $v_h = 0$ since \mathcal{H} is étale. Then the equation (8.7) can be rewritten as

$$(8.8) \quad \mathbf{v} = [(\sigma_1)_*(v_k), 0, 0, s_*(v_k)].$$

Now, we are ready to prove the desired property of ι .

Lemma 8.5. $\iota : N \rightarrow M$ above is an immersion.

Proof. Let $\mathbf{v} = [(v_a, v_m), v_x, v_g, v_z] \in \ker \tilde{t}_*$, i.e., $t_*(v_a \oplus v_m) = (v_a)^\# + v_m = 0$. We should show that $v_h = 0$ and find v_k satisfying (8.6) and (8.8). First, we find v_k as follows:

From the condition $(\sigma_0)_*(v_z) = v_m = (-v_a)^\#$, we get

$$\exp(t \cdot (-v_a)) \cdot m = m(t),$$

where $m(t)$ is a curve in M with $m'(0) = v_m$. Since $\exp(-t \cdot v_a) = \exp(t \cdot v_a)^{-1}$,

$$m \equiv \exp(t \cdot v_a) \cdot m(t).$$

Note that $m(t) = \sigma_0(z(t))$ for some curve $z(t)$ in G'_0 with $z'(0) = v_z$. Since σ is an equivalence map and $(a(t), m(t)) \in G \times M$ is an arrow from $\sigma_0(z(t))$ to $\sigma_0(z)$ for each t , there is unique $k(t) \in G'_1$ which maps to $\sigma_1(k(t)) = (a(t), m(t))$ with $s(k(t)) = z(t)$ and $t(k(t)) \equiv z$. We define $v_k := k'(0) \in T_k G'_1$, then $(\sigma_1)_*(v_k) = (v_a, v_m) \in TG \times TM$ and $s_*(v_k) = v_z$.

Let γ be a curve in the R_ϕ^* such that $\gamma'(0) = \mathbf{v}$. Consider a component of γ , $g(t) \in G_1$ such that $g'(0) = v_g$. We claim that this curve $g(t)$ is a constant curve.

Note that $\psi_0 \circ t \circ k(t) \equiv \psi_0(z)$. Since ψ is an equivalence map and \mathcal{G} is étale, $\psi_0 \circ s \circ k(t) \equiv \psi_0(z)$ (ψ_1 maps “infinitesimal action” on \mathcal{G}' whose image of target is fixed to a “constant action” on \mathcal{G}). Note that $t \circ g(t) = \psi_0 \circ z(t) = \psi_0 \circ s \circ k(t) = \psi_0(z)$. Since \mathcal{G} is étale and target points of $g(t)$ is fixed, $g(t)$ is a constant arrow in G_1 . Since $\phi_0 \circ x(t) = s \circ g(t) = \phi_0(x)$ and ϕ_0 is an immersion map, $x(t) \equiv x$.

We conclude that, if $\tilde{t}_*(\mathbf{v}) = 0$, then there exist v_k such that

$$(8.9) \quad \mathbf{v} = [(\sigma_1)_*(v_k), s_*(0) = 0, v_g = 0, s_*(v_k)],$$

it proves that $\iota : N \rightarrow M$ is an immersion. \square

Lastly, the equivariant immersion ι is strong by basically the same argument as in Lemma 7.6.

References

- [1] A. Adem, J. Leida, and Y. Ruan, *Orbifolds and Stringy Topology*, Cambridge Tracts in Mathematics, 171. Cambridge University Press, Cambridge, 2007.
- [2] W. Chen and Y. Ruan, *A new cohomology theory of orbifold*, Comm. Math. Phys. **248** (2004), no. 1, 1–31.
- [3] V. Ginzburg, V. Guillemin, and Y. Karshon, *Moment maps, cobordisms, and Hamiltonian group actions*, Mathematical Surveys and Monographs **98** American Mathematical Society, Providence, RI, 2002.
- [4] A. Haefliger, *Groupoïdes d’holonomie et classifiants*, Astérisque **116** (1984), 70–97.
- [5] E. Lerman, *Orbifolds as stacks?*, L’Enseign. Math. (2) **56** (2010), no. 3–4, 315–363.
- [6] I. Moerdijk and J. Mrčun, *Lie groupoids, sheaves and cohomology*, In: Poisson Geometry, Deformation Quantisation and Group Representations. London Math. Soc. Lecture Note Ser. 323, pp. 145–272, Cambridge University Press, Cambridge, 2005.
- [7] I. Moerdijk and D. Pronk, *Orbifolds, sheaves and groupoids*, K-Theory **12** (1997), no. 1, 3–21.
- [8] D. Pronk and L. Scull, *Translation groupoids and orbifold cohomology*, Canad. J. Math. **62** (2010), no. 3, 614–645.

CHEOL-HYUN CHO
 DEPARTMENT OF MATHEMATICAL SCIENCES
 RESEARCH INSTITUTE OF MATHEMATICS
 SEOUL NATIONAL UNIVERSITY
 SEOUL 151-747, KOREA
E-mail address: chocheol@snu.ac.kr

HANSOL HONG
DEPARTMENT OF MATHEMATICAL SCIENCES
RESEARCH INSTITUTE OF MATHEMATICS
SEOUL NATIONAL UNIVERSITY
SEOUL 151-747, KOREA
E-mail address: `hansol84@snu.ac.kr`

HYUNG-SEOK SHIN
DEPARTMENT OF MATHEMATICAL SCIENCES
RESEARCH INSTITUTE OF MATHEMATICS
SEOUL NATIONAL UNIVERSITY
SEOUL 151-747, KOREA
E-mail address: `shs83@snu.ac.kr`