

## ON QUADRATIC FUNCTIONALS

PETER ŠEMRL

In this note a general solution of the problem of the characterisation of quadratic functionals posed by Vukman is given.

**THEOREM.** *Let  $A$  be a complex  $\ast$ -algebra with identity  $e$  and let  $X$  be a vector space which is also a unitary left  $A$ -module. Suppose there exists a mapping  $Q: X \rightarrow A$  with the properties*

- (i)  $Q(x + y) + Q(x - y) = 2Q(x) + 2Q(y)$  for all pairs  $x, y \in X$ , and
- (ii)  $Q(ax) = aQ(x)a^\ast$  for all  $x \in X$  and all  $a \in A$ .

*Under these conditions the mapping  $B(\cdot, \cdot): X \times X \rightarrow A$  defined by the relation*

$$B(x, y) = (1/4)(Q(x + y) - Q(x - y)) + (i/4)(Q(x + iy) - Q(x - iy))$$

*satisfies the following:*

- (1)  $B(\cdot, \cdot)$  is additive in both arguments;
- (2)  $B(ax, y) = aB(x, y)$   
 $B(x, ay) = B(x, y)a^\ast$ , for all pairs  $x, y \in X$  and all  $a \in A$ ;
- (3)  $Q(x) = B(x, x)$  for all  $x \in X$ .

**REMARK:** A functional  $Q: X \rightarrow A$  which satisfies (i) and (ii) is called an  $A$ -quadratic functional and a mapping  $B: X \times X \rightarrow A$  for which conditions (1) and (2) are fulfilled is called an  $A$ -sesquilinear functional. If  $A$  is the complex number field then this result reduces to Kurepa's extension of the Jordan-Neumann theorem which characterises pre-Hilbert space among all normed spaces.([3])

**PROOF:** As in the proof of Kurepa's result (see [3], [5] and also [6]) one can prove that the function  $W(\cdot, \cdot)$  defined by relation  $W(x, y) = Q(x + y) - Q(x - y)$  is additive in both variables. Therefore the same is true for the functional  $B$ . A short computation shows that  $Q(x) = B(x, x)$  for all  $x \in X$ . Hence it remains to prove (2). For this purpose we define a new functional  $S: A \times A \rightarrow A$  by  $S(a, b) = aB(x, y)b^\ast - B(ax, by)$

---

Received 17 March 1987

This work was supported by the Research Council of Slovenia

where  $x$  and  $y$  are fixed vectors. From the fact that  $B$  is biadditive it follows that the functional  $S$  is also biadditive. Using (ii) one can easily obtain

$$S(ca, cb) = cS(a, b)c^*, \quad a, b, c \in A.$$

A short computation yields  $S(ia, b) = iS(a, b)$  and  $S(a, ib) = -iS(a, b)$ . For any four elements  $a, b, c, d \in A$  we have that  $S(ab, ac) + S(ab, dc) + S(db, ac) + S(db, dc) = S((a+d)b, (a+d)c) = (a+d)S(b, c)(a^* + d^*) = aS(b, c)a^* + dS(b, c)a^* + aS(b, c)d^* + dS(b, c)d^*$ . This yields  $S(ab, dc) + S(db, ac) = dS(b, c)a^* + aS(b, c)d^*$ . Replacing  $d$  and  $c$  by  $e$  we get

$$(4) \quad S(ab, e) + S(b, a) = S(b, e)a^* + aS(b, e).$$

Let us put the element  $ia$  instead of  $a$ . We obtain

$$(5) \quad iS(ab, e) - iS(b, a) = -iS(b, e)a^* + iaS(b, e).$$

Comparing (4) and (5) we see that  $S(ab, e) = aS(b, e)$  and  $S(b, a) = S(b, e)a^*$ . Replacing  $b$  by  $e$  by using the relation  $S(e, e) = 0$  we complete the proof. ■

This result was proved in [4] and [8] under the stronger assumption that  $A$  is a Banach  $*$ -algebra (see also [6] and [7]) using the fact that such algebras have enough invertible elements. It should be mentioned that in the proof of the present general result an idea similar to those of Davison [1] was used.

#### REFERENCES

- [1] T.M.K. Davison, 'Jordan derivations and quasi-bilinear forms', *Comm. Algebra* **121** (1984), 23–32.
- [2] S. Kurepa, 'The Cauchy functional equation and scalar product in vector spaces', *Glasnik Math. Fiz.-Astr.* **10** (1964), 23–36.
- [3] S. Kurepa, 'Quadratic and sesquilinear functionals', *Glas. Mat. Fiz.-Astr.* **20** (1965), 79–92.
- [4] P. Šemrl, 'On quadratic and sesquilinear functionals', *Aequationes Math.* **31** (1986), 184–190.
- [5] P. Vrbová, 'Quadratic and bilinear forms', *Časopis Pěst. Mat.* **98** (1973), 159–161.
- [6] J. Vukman, 'A result concerning additive functions in hermitian Banach  $*$ -algebras and an application', *Proc. Amer. Math. Soc.* **91** (1984), 367–372.
- [7] J. Vukman, 'Some results concerning the Cauchy functional equation in certain Banach algebras', *Bull. Austral. Math. Soc.* **31** (1985), 137–144.
- [8] J. Vukman, 'Some functional equations in Banach algebras and an application', *Proc. Amer. Math. Soc.* (to appear).

Institute of Mathematics, Physics and Mechanics  
University of Ljubljana  
P.O. Box 543  
61001 Ljubljana  
Yugoslavia