

## On semiparallel anti-invariant submanifolds of generalized Sasakian space forms

Cihan ÖZGÜR<sup>1,\*</sup>, Fatma GÜRLER<sup>1</sup>, Cengizhan MURATHAN<sup>2</sup>

<sup>1</sup>Department of Mathematics, Balıkesir University, Çağış, Balıkesir, Turkey

<sup>2</sup>Department of Mathematics, Uludağ University, Bursa, Turkey

Received: 17.09.2013 • Accepted: 22.01.2014 • Published Online: 25.04.2014 • Printed: 23.05.2014

**Abstract:** We consider minimal anti-invariant semiparallel submanifolds of generalized Sasakian space forms. We show that the submanifolds are totally geodesic under certain conditions.

**Key words:** Semiparallel submanifold, generalized Sasakian space form, Laplacian of the second fundamental form, totally geodesic submanifold

### 1. Introduction

Let  $(M, g)$  and  $(N, \tilde{g})$  be Riemannian manifolds and  $f: M \rightarrow N$  an isometric immersion. Denote by  $\sigma$  and  $\bar{\nabla}$  its second fundamental form and van der Waerden–Bortolotti connection, respectively. If  $\bar{\nabla}\sigma = 0$ , then the submanifold  $M$  is said to have a parallel second fundamental form [6]. The act of  $\bar{R}$  to the second fundamental form  $\sigma$  is defined by

$$\begin{aligned}(\bar{R}(X, Y) \cdot \sigma)(Z, W) &= R^\perp(X, Y)h(Z, W) - \sigma(R(X, Y)Z, W) - \sigma(Z, R(X, Y)W) \\ &= (\bar{\nabla}_X \bar{\nabla}_Y \sigma)(Z, W) - (\bar{\nabla}_Y \bar{\nabla}_X \sigma)(Z, W),\end{aligned}\tag{1}$$

where  $\bar{R}$  is the curvature tensor of the van der Waerden–Bortolotti connection  $\bar{\nabla}$ . Semiparallel submanifolds were introduced by Deprez in [7]. If  $\bar{R} \cdot \sigma = 0$ , then  $f$  is called semiparallel. It is clear that if  $f$  has parallel second fundamental form, then it is semiparallel. Hence, a semiparallel submanifold can be considered as a natural generalization of a submanifold with a parallel second fundamental form. Semiparallel submanifolds have been studied by various authors; see, for example [3, 7, 8, 9, 13, 16] and the references therein. Recently, in [18], Yıldız et al. studied  $C$ -totally real pseudoparallel submanifolds of Sasakian space forms, which are generalizations of semiparallel submanifolds. In [5], Brasil et al. studied  $C$ -totally real pseudoparallel submanifolds of  $\lambda$ -Sasakian space forms. In [15], Sular, et al. studied anti-invariant pseudoparallel submanifolds of Kenmotsu space forms with  $\xi$  tangent to the submanifold. In [14], Sular studied pseudoparallel submanifolds of Kenmotsu space forms with  $\xi$  normal to the submanifold.

Motivated by the studies of the above authors, in the present paper, we study anti-invariant minimal semiparallel submanifolds of generalized Sasakian space forms.

\*Correspondence: [cozgur@balikesir.edu.tr](mailto:cozgur@balikesir.edu.tr)

2010 AMS Mathematics Subject Classification: 53C40, 53C42, 53C25.

**2. Generalized Sasakian space forms**

Let  $M^{2n+1} = M(\varphi, \xi, \eta, g)$  be an almost contact metric manifold. If  $[\varphi, \varphi](X, Y) = -2d\eta(X, Y)\xi$  for all vector fields  $X, Y$  on  $M^{2n+1}$  then the almost contact metric structure is called *normal*, where  $[\varphi, \varphi]$  denotes the Nijenhuis torsion. If  $d\eta(X, Y) = g(X, \varphi Y)$  for all vector fields  $X, Y$  on  $M$ , then the almost contact metric structure  $(\varphi, \xi, \eta, g)$  is a *contact metric structure*. In this case, the manifold  $M^{2n+1}$  with the contact metric structure  $(\varphi, \xi, \eta, g)$  is called a *contact metric manifold*. A normal contact metric manifold is called a *Sasakian manifold* [4]. An almost contact metric manifold  $M$  is called a *Kenmotsu manifold* [11] if

$$(\nabla_X \varphi)Y = g(\varphi X, Y)\xi - \eta(Y)\varphi X,$$

where  $\nabla$  is the Levi-Civita connection. A Kenmotsu manifold is normal but not a contact manifold.

An almost contact metric manifold  $M$  is called a *cosymplectic manifold* [12] if  $\nabla\varphi = 0$ , which implies that  $\nabla\xi = 0$ . Hence,  $\xi$  is a Killing vector field for a cosymplectic manifold.

An almost contact metric manifold is called a  $\lambda$ -*Sasakian manifold* [10] if

$$(\nabla_X \varphi)Y = \lambda [g(X, Y)\xi - \eta(Y)X].$$

If  $\lambda = 1$ , a  $\lambda$ -*Sasakian manifold* is a Sasakian manifold.

The sectional curvature of a  $\varphi$ -section is called a  $\varphi$ -*sectional curvature*. A Sasakian (resp. Kenmotsu, cosymplectic,  $\lambda$ -Sasakian) manifold with constant  $\varphi$ -sectional curvature  $c$  is called a *Sasakian (resp. Kenmotsu, cosymplectic,  $\lambda$ -Sasakian) space form*; see [4, 11, 12, 10], respectively.

The notion of a generalized Sasakian space form was introduced by Alegre et al. in [1]. An almost contact metric manifold  $M^{2n+1} = M(\varphi, \xi, \eta, g)$  whose curvature tensor satisfies

$$\begin{aligned} R(X, Y)Z &= f_1\{g(Y, Z)X - g(X, Z)Y\} \\ &+ f_2\{g(X, \varphi Z)\varphi Y - g(Y, \varphi Z)\varphi X + 2g(X, \varphi Y)\varphi Z\} \\ &+ f_3\{\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + g(X, Z)\eta(Y)\xi - g(Y, Z)\eta(X)\xi\} \end{aligned} \tag{2}$$

for certain differentiable functions  $f_1, f_2$ , and  $f_3$  on  $M^{2n+1}$  is called a generalized Sasakian space form [1]. The natural examples of generalized Sasakian space forms with constant functions are a Sasakian space form ( $f_1 = \frac{c+3}{4}, f_2 = f_3 = \frac{c-1}{4}$ ) [4], a Kenmotsu space form ( $f_1 = \frac{c-3}{4}, f_2 = f_3 = \frac{c+1}{4}$ ) [11], and a cosymplectic space form ( $f_1 = f_2 = f_3 = \frac{c}{4}$ ) [12]. If  $M$  is a  $\lambda$ -Sasakian space form then  $f_1 = \frac{c+3\lambda}{4}, f_2 = f_3 = \frac{c-\lambda}{4}$  [10].

Let  $M$  be an  $n$ -dimensional submanifold of a Riemannian manifold  $\widetilde{M}$ . We denote by  $\widetilde{\nabla}, \nabla$  the Riemannian and induced Riemannian connections in  $\widetilde{M}$  and  $M$ , respectively, and let  $\sigma$  be the second fundamental form of the submanifold. The equation of Gauss is given by

$$\begin{aligned} \widetilde{R}(X, Y, Z, W) &= R(X, Y, Z, W) \\ &-g(\sigma(X, W), \sigma(Y, Z)) + g(\sigma(X, Z), \sigma(Y, W)) \end{aligned} \tag{3}$$

for all vector fields  $X, Y, Z, W$  tangent to  $M$ , where  $\widetilde{R}$  and  $R$  denote the curvature tensors of the connections  $\widetilde{\nabla}, \nabla$ , respectively. The mean curvature vector field  $H$  is given by  $H = \frac{1}{n} \text{trace}(\sigma)$ . The submanifold  $M$  is *totally geodesic* in  $\widetilde{M}$  if  $\sigma = 0$ , and *minimal* if  $H = 0$  [6].

Using (3), the Gauss equation for the submanifold  $M^n$  of a generalized Sasakian space form  $\widetilde{M}^{2m+1}$  is

$$\begin{aligned} \widetilde{R}(X, Y, Z, W) = & \\ & f_1\{g(Y, Z)X - g(X, Z)Y\} \\ & + f_2\{g(X, \varphi Z)\varphi Y - g(Y, \varphi Z)\varphi X + 2g(X, \varphi Y)\varphi Z\} \\ & + f_3\{\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X + g(X, Z)\eta(Y)\xi - g(Y, Z)\eta(X)\xi\}, \\ & + g(\sigma(X, W), \sigma(Y, Z)) - g(\sigma(X, Z), \sigma(Y, W)). \end{aligned} \tag{4}$$

A submanifold  $M$  of a generalized Sasakian space form  $\widetilde{M}^{2m+1}$  is called *anti-invariant* if and only if  $\varphi(T_x M) \subset T_x^\perp M$  for all  $x \in M$  [2]. For more information about anti-invariant submanifolds we refer to [17].

### 3. Semiparallel anti-invariant submanifolds of a generalized Sasakian space form

In this section, we give the main results of the paper.

For an  $n$ -dimensional submanifold  $M$  of a  $(2n + 1)$ -dimensional Riemannian manifold  $\widetilde{M}^{2n+1}$ , it is known that the Laplacian  $\Delta\sigma_{ij}^\alpha$  of  $\sigma_{ij}^\alpha$  is defined by

$$\Delta\sigma_{ij}^\alpha = \sum_{i,j,k=1}^n \sigma_{ijkk}^\alpha. \tag{5}$$

Then

$$\frac{1}{2}\Delta(\|\sigma\|^2) = \sum_{i,j,k=1}^n \sum_{\alpha=n+1}^{2n+1} \sigma_{ij}^\alpha \sigma_{ijkk}^\alpha + \|\overline{\nabla}\sigma\|^2, \tag{6}$$

(see [17]), where

$$\|\sigma\|^2 = \sum_{i,j,k=1}^n \sum_{\alpha=n+1}^{2n+1} (\sigma_{ij}^\alpha)^2, \tag{7}$$

and

$$\|\overline{\nabla}\sigma\|^2 = \sum_{i,j,k=1}^n \sum_{\alpha=n+1}^{2n+1} (\sigma_{ijkk}^\alpha)^2 \tag{8}$$

are the square of the length of second and the third fundamental forms of  $M$ , respectively.

A simple calculation gives us the following proposition:

**Proposition 1** *Let  $M$  be an  $n$ -dimensional minimal anti-invariant submanifold of a  $(2n + 1)$ -dimensional generalized Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  normal to  $M$ . Then we have*

$$\begin{aligned} \frac{1}{2}\Delta(\|\sigma\|^2) = & \|\overline{\nabla}\sigma\|^2 + (f_2 + nf_1)\|\sigma\|^2 \\ & - \left[ \sum_{\alpha,\beta=n+1}^{2n+1} tr(A_\alpha \circ A_\beta)^2 + \|[A_\alpha, A_\beta]\|^2 \right]. \end{aligned} \tag{9}$$

**Theorem 2** Let  $M$  be an  $n$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional generalized Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  normal to  $M$ . If

$$f_2 + nf_1 \leq 0,$$

then  $M$  is totally geodesic.

**Proof** Let  $\{e_1, e_2, \dots, e_n, \xi, \varphi e_1, \varphi e_2, \dots, \varphi e_n\}$  be an orthonormal frame in  $\widetilde{M}^{2n+1}$  such that  $e_1, e_2, \dots, e_n$  are tangent to  $M$ . By definition, the semiparallelity of  $M$ , for  $1 \leq k, l \leq n$ , gives us

$$\overline{R}(e_l, e_k) \cdot \sigma = 0. \tag{10}$$

By (1), we can write

$$(\overline{R}(e_l, e_k) \cdot \sigma)(e_i, e_j) = (\overline{\nabla}_{e_l} \overline{\nabla}_{e_k} \sigma)(e_i, e_j) - (\overline{\nabla}_{e_k} \overline{\nabla}_{e_l} \sigma)(e_i, e_j) = 0, \tag{11}$$

where  $1 \leq i, j, k, l \leq n$ .

Hence, equation (6) turns into

$$\frac{1}{2} \Delta(\|\sigma\|^2) = \sum_{i,j,k=1}^n g((\overline{\nabla}_{e_k} \overline{\nabla}_{e_k} \sigma)(e_i, e_j), \sigma(e_i, e_j)) + \|\overline{\nabla} \sigma\|^2. \tag{12}$$

Furthermore, using equations (5) and (6), we have

$$\frac{1}{2} \Delta(\|\sigma\|^2) = \sum_{i,j,k=1}^n \sum_{\alpha=n+1}^{2n+1} \sigma_{ij}^\alpha (\overline{\nabla}_{e_i} \overline{\nabla}_{e_j} H^\alpha) + \|\overline{\nabla} \sigma\|^2. \tag{13}$$

Since  $M$  is minimal, equation (13) can be written as

$$\frac{1}{2} \Delta(\|\sigma\|^2) = \|\overline{\nabla} \sigma\|^2 \tag{14}$$

(see [18]). Comparing (9) and (14), we find

$$\begin{aligned} & - (f_2 + nf_1) \|\sigma\|^2 \\ & + \sum_{\alpha,\beta=n+1}^{2n+1} tr(A_\alpha \circ A_\beta)^2 + \|[A_\alpha, A_\beta]\|^2 = 0. \end{aligned}$$

From the assumption, if

$$f_2 + nf_1 \leq 0,$$

then  $tr(A_\alpha \circ A_\beta) = 0$ . In particular,  $\|A_\alpha\|^2 = tr(A_\alpha \circ A_\alpha) = 0$ , and thus  $A_\alpha = 0$ , which means that  $\sigma = 0$ . Then  $M$  is totally geodesic.  $\square$

Using Theorem 2, we have the following corollaries:

**Corollary 3** [18] *Let  $M$  be an  $n$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  normal to  $M$ . If*

$$n(c + 3) + c - 1 \leq 0,$$

*then  $M$  is totally geodesic.*

**Corollary 4** *Let  $M$  be an  $n$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional cosymplectic space form  $\widetilde{M}^{2n+1}$  with  $\xi$  normal to  $M$ . If*

$$c \leq 0,$$

*then  $M$  is totally geodesic.*

**Corollary 5** [5] *Let  $M$  be an  $n$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional  $\lambda$ -Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  normal to  $M$ . If*

$$c - \lambda + n(c + 3\lambda) \leq 0,$$

*then  $M$  is totally geodesic.*

If  $M$  is an  $(n + 1)$ -dimensional minimal anti-invariant submanifold of a  $(2n + 1)$ -dimensional generalized Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ , then we have the following proposition:

**Proposition 6** *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant submanifold of a  $(2n + 1)$ -dimensional generalized Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ . Then we have*

$$\begin{aligned} \frac{1}{2}\Delta(\|\sigma\|^2) &= \|\overline{\nabla}\sigma\|^2 + (f_2 + (n + 1)f_1 - f_3)\|\sigma\|^2 \\ &- f_3 \sum_{i=1}^{n+1} \|\sigma(e_i, \xi)\|^2 - \left[ \sum_{\alpha, \beta=n+2}^{2n+1} \text{tr}(A_\alpha \circ A_\beta)^2 + \|[A_\alpha, A_\beta]\|^2 \right]. \end{aligned} \tag{15}$$

**Theorem 7** *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant semiparallel submanifold of  $(2n + 1)$ -dimensional generalized Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ . If*

$$f_2 + (n + 1)f_1 - f_3 \leq 0$$

*and*

$$f_3 \geq 0,$$

*then  $M$  is totally geodesic.*

**Proof** Let  $\{e_1, e_2, \dots, e_n, \xi, \varphi e_1, \varphi e_2, \dots, \varphi e_n\}$  be an orthonormal frame in  $\widetilde{M}^{2n+1}$  such that  $e_1, e_2, \dots, e_n, \xi$  are tangent to  $M$ . Then for  $1 \leq i, j \leq n + 1$  and  $n + 2 \leq \alpha \leq 2n + 1$ . Similar to the proof of Theorem 2, using the minimality condition, we obtain

$$\frac{1}{2}\Delta(\|\sigma\|^2) = \|\overline{\nabla}\sigma\|^2. \tag{16}$$

Comparing (15) and (16) we find

$$\begin{aligned}
 & - (f_2 + (n + 1) f_1 - f_3) \|\sigma\|^2 + f_3 \sum_{i=1}^{n+1} \|\sigma(e_i, \xi)\|^2 \\
 & + \sum_{\alpha, \beta=n+2}^{2n+1} \text{tr}(A_\alpha \circ A_\beta)^2 + \|[A_\alpha, A_\beta]\|^2 = 0.
 \end{aligned}$$

From the assumption, if

$$f_2 + (n + 1) f_1 - f_3 \leq 0$$

and

$$f_3 \geq 0,$$

then  $\text{tr}(A_\alpha \circ A_\beta) = 0$ . Similar to the proof of Theorem 2, this gives us  $\sigma = 0$ . Then  $M$  is totally geodesic.  $\square$

Using Theorem 7, we have the following corollaries:

**Corollary 8** *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ . If*

$$c \in (-\infty, -3] \cup [1, \infty),$$

*then  $M$  is totally geodesic.*

**Corollary 9** *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional cosymplectic space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ . If  $c = 0$ , then  $M$  is totally geodesic.*

**Corollary 10** [15] *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional Kenmotsu space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ . If  $c \in [-1, 3]$ , then  $M$  is totally geodesic.*

**Corollary 11** *Let  $M$  be an  $(n + 1)$ -dimensional minimal anti-invariant semiparallel submanifold of a  $(2n + 1)$ -dimensional  $\lambda$ -Sasakian space form  $\widetilde{M}^{2n+1}$  with  $\xi$  tangent to  $M$ .*

*i) If  $\lambda$  is a positive function on  $M$  and*

$$c \in (-\infty, -3\lambda] \cup [\lambda, \infty)$$

*or*

*ii) If  $\lambda$  is a negative function on  $M$  and*

$$c \in [\lambda, -3\lambda],$$

*then  $M$  is totally geodesic.*

### References

- [1] Alegre P, Blair DE, Carriazo A. Generalized Sasakian space-forms. *Israel J Math* 2004; 141: 157–183.
- [2] Alegre P, Carriazo A. Submanifolds of generalized Sasakian space forms. *Taiwanese J Math* 2009; 13: 923–941.
- [3] Asperti AC, Mercuri F. Semi-parallel immersions into space forms. *Boll Un Mat Ital* 1994; 8B: 883–895.
- [4] Blair DE. *Riemannian Geometry of Contact and Symplectic Manifolds*. Boston: Birkhauser, 2002.
- [5] Brasil A, Lobos GA, Mariano M.  $C$ -totally real submanifolds with parallel mean curvature in  $\lambda$ -Sasakian space forms. *Mat Contemp* 2008; 34: 83–102.
- [6] Chen BY. *Geometry of Submanifolds*. Pure and Applied Mathematics. New York: Marcel Dekker, 1973.
- [7] Deprez J. Semiparallel hypersurfaces. *Rend Sem Mat Univ Politec Torino* 1986; 44: 303–316.
- [8] Deprez J. Semi-parallel immersions. In: Morvan JM, Verstraelen L, editors. *Geometry and Topology of Submanifolds*. Singapore: World Scientific, 1989, pp. 73–88.
- [9] Dillen F. Semi-parallel hypersurfaces of a real space form. *Israel J Math* 1991; 75: 193–202.
- [10] Janssens D, Vanhecke L. Almost contact structures and curvature tensors. *Kodai Math J* 1981; 4: 1–27.
- [11] Kenmotsu K. A class of almost contact Riemannian manifolds. *Tohoku Math J* 1972; 24: 93–103.
- [12] Ludden GD. Submanifolds of cosymplectic manifolds. *J Differential Geometry* 1970; 4: 237–244.
- [13] Özgür C, Murathan C. On invariant submanifolds of Lorentzian para-Sasakian manifolds. *Arab J Sci Eng Sect A Sci* 2009; 34: 177–185.
- [14] Sular S. Kenmotsu manifolds and their some submanifolds. PhD, Balıkesir University, Balıkesir, Turkey, 2009.
- [15] Sular S, Özgür C, Murathan C. Pseudoparallel anti-invariant submanifolds of Kenmotsu manifolds. *Hacettepe J Math Stat* 2010; 39: 535–543.
- [16] Van der Veken J, Vrancken L. Parallel and semi-parallel hypersurfaces of  $S^n \times \mathbb{R}$ . *Bull Braz Math Soc (NS)* 2008; 39: 355–370.
- [17] Yano K, Kon M. *Structures on Manifolds*. Series in Pure Mathematics. Singapore: World Scientific, 1984.
- [18] Yıldız A, Murathan C, Arslan K, Ezentaş R.  $C$ -totally real pseudo-parallel submanifolds of Sasakian space forms. *Monatsh- Math* 2007; 151: 247–256.