

Research Article

On Soft Separation Axioms and Their Applications on Decision-Making Problem

T. M. Al-shami 

Department of Mathematics, Sana'a University, Sana'a, Yemen

Correspondence should be addressed to T. M. Al-shami; tareqalshami83@gmail.com

Received 7 September 2020; Revised 24 December 2020; Accepted 31 December 2020; Published 30 January 2021

Academic Editor: Serdar Ulubeyli

Copyright © 2021 T. M. Al-shami. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In this work, we introduce new types of soft separation axioms called pt -soft α regular and pt -soft αT_i -spaces ($i = 0, 1, 2, 3, 4$) using partial belong and total nonbelong relations between ordinary points and soft α -open sets. These soft separation axioms enable us to initiate new families of soft spaces and then obtain new interesting properties. We provide several examples to elucidate the relationships between them as well as their relationships with e -soft T_i , soft αT_i , and tt -soft αT_i -spaces. Also, we determine the conditions under which they are equivalent and link them with their counterparts on topological spaces. Furthermore, we prove that pt -soft αT_i -spaces ($i = 0, 1, 2, 3, 4$) are additive and topological properties and demonstrate that pt -soft αT_i -spaces ($i = 0, 1, 2$) are preserved under finite product of soft spaces. Finally, we discuss an application of optimal choices using the idea of pt -soft T_i -spaces ($i = 0, 1, 2$) on the content of soft weak structure. We provide an algorithm of this application with an example showing how this algorithm is carried out. In fact, this study represents the first investigation of real applications of soft separation axioms.

1. Introduction

The urgent need of theories dealing with uncertainties comes from daily facing complicated problems containing data which are not always crisp. The recent mathematical tool to handle these problems is soft set which was initiated by Molodtsov [1] in 1999. The rationale of soft sets is based on parameterization idea, which references that complex objects should be perceived from many aspects and each solo facet only provides a partial and approximate description of the whole entity. Molodtsov [1] in his pioneering work provided some applications of soft set in different fields and elaborated its merits compared with probability theory and fuzzy sets theory which deals with vagueness or uncertainties.

Afterwards, Maji et al. [2] started studying the operations between soft sets such as soft union and soft intersections. To overcome the shortcomings of these operations, Ali et al. [3] proposed new operations such as restricted union and intersection and a complement of a soft set and revealed some of their properties. Babitha and Sunil [4] investigated some

properties of relations and functions in soft setting. Qin and Hong [5] defined new types of soft equal relations and proved some algebraic properties with respect to them. Their novel work opened up a new direction which was a goal of many papers (for more details, see [6, 7] and the references mentioned therein). Recently, soft set theory has become very prevalent tool among researchers to overcome the problems of uncertainty in different fields such as information theory [8], computer sciences [9], engineering [10], and medical sciences [11].

In 2011, Shabir and Naz [12] and Ça man et al. [13] initiated a new research line by defining soft topology. However, they used two different techniques of defining soft topology. On the one hand, Shabir and Naz formulated soft topology on the collection of soft sets over a universal crisp set with a fixed set of parameters. On the other hand, Ça man et al. formulated soft topology on the collection of soft sets over an absolute soft set with different sets of parameters which are subsets of the universal set of parameters. In this paper, we continue studying soft topology using the definition given by Shabir and Naz. They formulated the

notions of soft interior and soft closure operators and soft subspaces and shed light on soft separation axioms. Following Shabir and Naz's work, many researchers explored the topological concepts on the domain of soft topology and examined the similarity and divergence between classical topology and soft topology.

Aygün and Aygün [14] first introduced the concept of soft compactness and then Hida [15] distinguished between two types of soft compactness depending on the total belong relation. After that, Al-shami et al. [16] studied new types of soft compact and soft Lindelöf spaces. Al-shami and Kočinac [17] defined and discussed the soft version of nearly Menger spaces. Babitha and Sunil investigated some notions on soft topological spaces in [18] and presented some techniques of generated soft topology from soft relations in [19]. The authors of [20] presented soft maps by using two crisp maps, one of them between the sets of parameters and the second one between the universal sets. However, the authors of [21] introduced soft maps by using the concept of soft points.

In 2018, the authors of [22] came up new relations of belong and nonbelong between an element and soft set, namely, partial belong and total nonbelong relations. In fact, these relations widely opened the door to study and redefine many soft topological notions. This leads to obtain many fruitful properties and changes which can be seen significantly on the study of soft separation axioms and decision-making problems as it was shown in [23, 24]. As another part of study on soft separation axioms, the authors of [25, 26] studied them with respect to the distinct soft points. Singh and Noorie [27] carried out a comparative study between soft separation axioms, and Terepeta [28] studied soft separating axioms and similarity of soft topological spaces. Alcantud [29] investigated the properties of countability axioms in soft setting. Recently, Al-shami [30, 31] has done amendments for some previous studies of soft separation axioms. For more details of conducted studies on soft set theory, we refer the reader to [32].

By decision making, we mean select the optimal alternative from the finite set of alternatives according to the multiple criteria. Decision-making theory is a very significant area, which is used mostly in human activities. In the literature, there are many studies which were conducted in this regard (see, for example, [33–35]).

The rest of this paper is organized as follows. In Section 2, we review some basic concepts about soft sets, soft topology, and soft separation axioms. Section 3 explores the concepts pt -soft αT_i ($i = 0, 1, 2, 3, 4$) and pt -soft α -regular spaces using partial belong and total nonbelong relations between ordinary points and soft α -open sets. This section shows the relationships between these concepts and discusses their main properties with the help of examples. In Section 4, we present the first investigation of real applications of soft separation axioms in decision-making problems. We initiate an algorithm of this application and show the way of carrying out this algorithm by an illustrative example. In Section 5, we discuss the advantages and limitations of the soft weak structure approach and propose some application in the engineering sciences. Finally,

conclusions and some directions for future works are given in Section 6.

2. Preliminaries

In this section, we recall some basic concepts and properties regarding soft set theory and soft topology.

2.1. Soft Sets

Definition 1 (see [1]). For a nonempty set X and a set of parameters E , a pair (G, E) is said to be a soft set over X provided that G is a map of E into the power set $P(X)$.

In this study, we use a symbol G_E to refer a soft set instead of (G, E) and we identify it as ordered pairs $G_E = \{(e, G(e)) : e \in E \text{ and } G(e) \in P(X)\}$.

Each $G(e)$ is called a component of G_E (or e -approximate), and a family of all soft sets defined over X with E is denoted by $S(X_E)$.

Definition 2 (see [36]). A soft set G_E is said to be a subset of a soft set H_E , denoted by $G_E \widetilde{\subseteq} H_E$, if $G(e) \subseteq H(e)$ for each $e \in E$.

The soft sets G_E and H_E are said to be soft equal if each one of them is a subset of the other.

In the literature, the relations between ordinary points and soft sets were described by four types of belong and nonbelong relations. Consequentially, new kinds of soft topological notions and concepts can be defined and studied.

Definition 3 (see [12, 22]). Let G_E be a soft set over X and $x \in X$. We have the following relations:

- (i) $x \in G_E$ if $x \in G(e)$ for each $e \in E$. We read it as x totally belongs to G_E or G_E totally contains x .
- (ii) $x \notin G_E$ if $x \notin G(e)$ for some $e \in E$. We read it as x does not partially belong to G_E or G_E does not partially contain x .
- (iii) $x \in G_E$ if $x \in G(e)$ for some $e \in E$. We read it as x partially belongs to G_E or G_E partially contains x .
- (iv) $x \notin G_E$ if $x \notin G(e)$ for each $e \in E$. We read it as x does not totally belong to G_E or G_E does not totally contain x .

Definition 4 (see [3]). The relative complement of a soft set G_E is a soft set G_E^c , where $G^c : E \rightarrow 2^X$ is a mapping defined by $G^c(e) = X \setminus G(e)$ for all $e \in E$.

Two special soft sets over X are the null soft set $\tilde{\emptyset}$ in which all components are the empty set and the absolute \tilde{X} soft set in which all components are the initial (universal) set X . Also, we say that a soft set is countable (resp. finite) if all components are countable (resp. finite). Otherwise, it is said to be uncountable (resp. infinite). Soft point is an important type of soft sets which was defined by making one of its approximations a singleton set and all the other approximations empty set. If we make all approximations of a soft set equal to a fixed subset S of the universal set X , then we call it a stable soft set and denote it by \tilde{S} . In particular, if

$S = \{x\}$, we write x_E instead of \tilde{S} . To see the main properties of these types of soft set, we refer to [2, 22, 37, 38].

Definition 5 (see [2, 3]). Let G_E and H_E be two soft sets over X .

- (i) Their intersection, denoted by $G_E \widetilde{\cap} H_E$, is a soft set U_E , where a mapping $U: E \rightarrow 2^X$ is given by $U(e) = G(e) \cap H(e)$.
- (ii) Their union, denoted by $G_E \widetilde{\cup} H_E$, is a soft set U_E , where a mapping $U: E \rightarrow 2^X$ is given by $U(e) = G(e) \cup H(e)$.

By using a similar method, the soft union and intersection operators were generalized for an arbitrary number of soft sets.

Definition 6 (see [4]). The Cartesian product of two soft sets G_E and H_F over X and Y , respectively, is a soft set $G \times H_{E \times F}$ over $X \times Y$ defined by $(G \times H)(e, f) = G(e) \times H(f)$ for each $(e, f) \in E \times F$.

2.2. Soft Topology

Definition 7 (see [12]). A family τ of soft sets over X under a fixed set of parameters E is said to be a soft topology on X if it satisfies the following.

- (i) \tilde{X} and $\tilde{\Phi}$ are members of τ .
- (ii) The intersection of a finite number of soft sets in τ is a member of τ .
- (iii) The union of an arbitrary number of soft sets in τ is a member of τ .

The triple (X, τ, E) is called a soft topological space. A member in τ is called soft open and its relative complement is called soft closed.

Throughout this paper, (X, τ, E) and (Y, θ, E) denote soft topological spaces unless otherwise explicitly stated.

Proposition 1 (see [12]). In (X, τ, E) , a family $\tau_e = \{G(e): G_E \in \tau\}$ is a classical topology on X for each $e \in E$. τ_e is called a parametric topology and (X, τ_e) is called a parametric topological space.

Definition 8 (see [12]). Let (X, τ, E) be a soft topological space and $\emptyset \neq Y \subseteq X$. A family $\tau_Y = \{\tilde{Y} \cap tG_E: qG_E h \in \tau\}$ is called a soft relative topology on Y and the triple (Y, τ_Y, E) is called a soft subspace of (X, τ, E) .

Definition 9 (see [39]). A subset G_E of (X, τ, E) is called soft α -open if $G_E \subseteq \text{int}(cl(\text{int}(G_E)))$.

Theorem 1 (see [39])

- (i) Every soft open set is soft α -open.
- (ii) The arbitrary union (finite intersection) of soft α -open sets is soft α -open.

The above theorem implies that the family of all soft α -open subsets of (X, τ, E) forms a new soft topology τ_α finer than τ .

Proposition 2. Let \tilde{Y} be soft open subset of (X, τ, E) . Then:

- (1) If (H, E) is soft α -open and \tilde{Y} is soft open in (X, τ, E) , then $(H, E) \widetilde{\cap} (Y, E)$ is a soft α -open subset of (Y, τ_Y, E) .
- (2) If \tilde{Y} is soft open in (X, τ, E) and (H, E) is a soft α -open in (Y, τ_Y, E) , then (H, E) is a soft α -open subset of (X, τ, E) .

Definition 10 (see [40]). (X, τ, E) is said to be

- (i) Soft αT_0 if for every $x \neq y \in X$, there is a soft α -open set U_E such that $x \in U_E$ and $y \notin U_E$ or $y \in U_E$ and $x \notin U_E$.
- (ii) Soft αT_1 if for every $x \neq y \in X$, there are two soft α -open sets U_E and V_E such that $x \in U_E$ and $y \notin U_E$ and $y \in V_E$ and $x \notin V_E$.
- (iii) Soft αT_2 if for every $x \neq y \in X$, there are two disjoint soft α -open sets U_E and V_E such that $x \in G_E$ and $y \in F_E$.
- (iv) Soft α -regular if for every soft α -closed set H_E and $x \in X$ such that $x \notin H_E$, there are two disjoint soft α -open sets U_E and V_E such that $H_E \subseteq U_E$ and $x \in V_E$.
- (v) Soft α -normal if for every two disjoint soft α -closed sets H_E and F_E , there are two disjoint soft α -open sets U_E and V_E such that $H_E \subseteq U_E$ and $F_E \subseteq V_E$.
- (vi) Soft αT_3 (resp. soft αT_4) if it is both soft α -regular (resp. soft α -normal) and soft αT_1 -space.

Remark 1

- (i) The definitions of e -soft T_i -spaces ($i = 0, 1, 2, 3, 4$) of [23] were given by replacing soft α -open and soft α -closed sets of the above definition by soft open and soft closed sets with respect to partial belong and total nonbelong relations.
- (ii) The definitions of tt -soft αT_i -spaces ($i = 0, 1, 2, 3, 4$) of [41] were given by replacing a partial nonbelong relation of the above definition by a total nonbelong relation.

Definition 11. A soft topology τ on X is said to be

- (i) An enriched soft topology [14] if all soft sets G_E such that $G(e) = \emptyset$ or X are members of τ .
- (ii) An extended soft topology [38] if $\tau = \{G_E: G(e) \in \tau_e \text{ for each } e \in E\}$, where τ_e is a parametric topology on X .

The equivalence of enriched and extended soft topologies was proved in [42]. This result helps to probe the

relationships between soft topology and its parametric topologies.

Theorem 2 (see [42]). *A subset (F, E) of an extended soft topological space (X, τ, E) is soft α -open if and only if each e -approximate element of (F, E) is α -open.*

Proposition 3 (see [43]). *Let $\{(X_i, \tau_i, E): i \in I\}$ be a family of pairwise disjoint soft topological spaces and $X = \cup_{i \in I} X_i$. Then, the collection $\tau = \{(G, E) \subseteq \widetilde{X}: (G, E) \cap \widetilde{X}_i \text{ which is a soft open set in } (X_i, \tau_i, E) \text{ for every } i \in I\}$ defines a soft topology on X with a fixed set of parameters E .*

Definition 12 (see [43]). The soft topological space (X, τ, E) given in the above proposition is said to be the sum of soft topological spaces and is denoted by $(\oplus_{i \in I} X_i, \tau, E)$.

Theorem 3 (see [43]). *A soft set $(G, E) \subseteq \widetilde{\oplus_{i \in I} X_i}$ is soft α -open (resp. soft α -closed) in $(\oplus_{i \in I} X_i, \tau, E)$ if and only if all $(G, E) \cap \widetilde{X}_i$ are soft α -open (resp. soft α -closed) in (X_i, τ_i, E) .*

Theorem 4 (see [44]). *Let (X, τ, A) and (Y, θ, B) be two soft topological spaces and $\Omega = \{G_A \times F_B: G_A \in \tau \text{ and } F_B \in \theta\}$. Then, the family of all arbitrary union of elements of Ω is a soft topology over $X \times Y$ under a fixed set of parameters $A \times B$.*

Definition 13 (see [45]). A family τ of soft sets over X under a fixed set of parameters E is said to be a soft weak structure on X if τ contains the null soft set $\widetilde{\Phi}$.

3. pt -Soft αT_0 -Spaces ($i = 0, 1, 2, 3, 4$)

In this section, we define a new class of soft separation axioms called pt -soft αT_0 -spaces ($i = 0, 1, 2, 3, 4$), where the notations p and t indicate partial belong and total nonbelong relations, respectively. The initiation of this class is based on the relationship between ordinary points and soft α -open sets with respect to partial belong and total nonbelong relations. We ascertain the relationships between them and reveal their main properties.

Definition 14. (X, τ, E) is said to be

- (i) pt -soft αT_0 if for every $x \neq y \in X$, there exists a soft α -open set U_E such that $x \in U_E$ and $y \notin U_E$ or $y \in U_E$ and $x \notin U_E$.
- (ii) pt -soft αT_1 if for every $x \neq y \in X$, there exist soft α -open sets U_E and V_E such that $x \in U_E$ and $y \notin U_E$ and $y \in V_E$ and $x \notin V_E$.
- (iii) pt -soft αT_2 if for every $x \neq y \in X$, there exist two disjoint soft α -open sets U_E and V_E such that $x \in U_E$ and $y \notin U_E$ and $y \in V_E$ and $x \notin V_E$.
- (iv) pt -soft α regular if for every soft α -closed set H_E and $x \in X$ such that $x \notin H_E$, there exist disjoint soft α -open sets U_E and V_E such that $H_E \subseteq U_E$ and $x \in V_E$.
- (v) pt -soft αT_3 (resp. pt -soft αT_4) if it is both pt -soft α regular (resp. soft α normal) and pt -soft αT_1 .

We begin this work by showing the relationships between pt -soft αT_i -spaces as well as their relationships with e -soft T_i -spaces and soft αT_i -spaces.

Proposition 4

- (i) Every pt -soft αT_i -space is pt -soft αT_{i-1} for $i = 0, 1, 2, 3$.
- (ii) Every e -soft T_i -space is pt -soft αT_i for $i = 0, 1, 2, 4$.
- (iii) Every tt -soft αT_i -space is pt -soft αT_i for $i = 0, 1, 2, 4$.
- (iv) Every soft αT_i -space is pt -soft αT_i for $i = 2, 3$.

Proof

- (i) It immediately follows from Definition 16 that pt -soft αT_2 implies pt -soft αT_1 and pt -soft αT_1 implies pt -soft αT_0 .

To prove that pt -soft αT_3 implies pt -soft αT_2 , let $x \neq y$ in a pt -soft αT_3 -space (X, τ, E) . Since it is pt -soft αT_1 , then there are two soft α -open sets U_E and V_E such that $x \in U_E$, $y \notin U_E$ and $y \in V_E$, $x \notin V_E$. Now, $x \notin U_E^c$ and $y \notin V_E^c$. By hypothesis, (X, τ, E) is pt -soft regular; then, we have the following:

- (1) There are two disjoint soft α -open sets M_E and N_E such that $U_E^c \subseteq M_E$ and $x \in N_E$. Therefore, $y \in M_E$ and $y \notin N_E$.
- (2) There are two disjoint soft α -open sets H_E and F_E such that $V_E^c \subseteq H_E$ and $y \in F_E$. Therefore, $x \in H_E$ and $x \notin F_E$.

From (1) and (2) above, we find that $x \in N_E \cap H_E$, $y \notin N_E \cap H_E$ and $y \in M_E \cap F_E$, $x \notin M_E \cap F_E$. It follows from Theorem 1 that $N_E \cap H_E$ and $M_E \cap F_E$ are soft α -open sets. The disjointness of them proves that (X, τ, E) is pt -soft αT_2 .

- (ii) It follows from the fact that every soft α -open set is soft open.
- (iii) It follows from the fact that a total belong relation implies partial belong.
- (iv) When $i = 2$, let $x \neq y$ in a soft αT_2 -space (X, τ, E) . Then, there exist two disjoint soft α -open sets U_E and V_E such that $x \in U_E$ and $y \notin U_E$ and $y \in V_E$ and $x \notin V_E$. The disjointness of U_E and V_E leads to $y \notin U_E$ and $x \notin V_E$ as well. Thus, (X, τ, E) is pt -soft αT_2 .

When $i = 3$, it is clear that a soft α regular space is pt -soft α regular. Also, we know that every soft α -open and soft α -closed subsets of a soft α regular space are stable. Then, a soft αT_1 -space is pt -soft αT_1 . Hence, a soft αT_3 -space is pt -soft αT_3 , as required.

The succeeding examples illustrate that the above proposition is not always reversible. \square

Example 1. Consider the following three soft sets defined over the universal set $X = \{x, y\}$ with a set of parameters $E = \{e_1, e_2\}$ as follows:

$$\begin{aligned} U_{1_E} &= \{(e_1, \{x\}), (e_2, \emptyset)\}, \\ U_{2_E} &= \{(e_1, \emptyset), (e_2, \{y\})\}, \\ U_{3_E} &= \{(e_1, \{x\}), (e_2, \{y\})\}. \end{aligned} \tag{1}$$

Then, $\tau = \{\tilde{\Phi}, t\tilde{X}n, qU_i: i = 1, 2, 3\}$ is a soft topology on X . Now, $x \neq y$. One can note the following cases:

- (1) (X, τ, E) is *pt*-soft αT_2 because U_{1_E} and U_{2_E} are two disjoint soft α -open sets partially containing x and y , respectively, such that $x \notin U_{2_E}$ and $y \notin U_{1_E}$. In turn, (X, τ, E) is not soft αT_2 because there does not exist a proper soft α -open subset of (X, τ, E) totally containing x or y .
- (2) (X, τ, E) is soft α normal because the absolute and null soft sets are the only disjoint soft α -closed subsets of (X, τ, E) . Then, it is *pt*-soft αT_4 .
- (3) (X, τ, E) is not a *tt*-soft αT_0 -space because there does not exist a proper soft α -open subset of (X, τ, E) totally containing x or y . This also means that (X, τ, E) is not *tt*-soft αT_4 .
- (4) For a soft α -closed set $U_{3_E}^c$, we have $x \notin U_{3_E}^c$. Since the only soft α -open set containing $U_{3_E}^c$ is the absolute soft set, (X, τ, E) is not *pt*-soft α regular space. Consequently, it is not *pt*-soft αT_3 .

Example 2. Let E be any set of parameters and $\tau = \{\tilde{\Phi}, tU_E n \subseteq qNn: \cup E^x \text{ is finite}\}$ be a soft topology on the set of natural numbers \mathbb{N} . It is clear that a soft subset of (\mathbb{N}, τ, E) is soft α -open if and only if it is soft open. For each $x \neq y \in \mathbb{N}$, $\mathbb{N} \setminus \{y\}$ and $\mathbb{N} \setminus \{x\}$ are soft α -open sets such that $x \in \mathbb{N} \setminus \{y\}$ and $y \notin \mathbb{N} \setminus \{y\}$ and $y \in \mathbb{N} \setminus \{x\}$ and $x \notin \mathbb{N} \setminus \{x\}$. Therefore, (\mathbb{N}, τ, E) is *pt*-soft αT_1 . On the other hand, there do not exist two disjoint soft α -open sets except for the null and absolute soft sets. Hence, (\mathbb{N}, τ, E) is not *pt*-soft αT_2 .

Example 3. Let $\tau = \{\tilde{\Phi}, t\tilde{X}n, q\{(e_1, \{x\}), (e_2, \emptyset)\}\}$ be a soft topology on $X = \{x, y\}$ with a set of parameters $E = \{e_1, e_2\}$. It can be easily checked that (X, τ, E) is *pt*-soft αT_0 , but not *pt*-soft αT_1 .

Example 4. It is well known that a soft topological space is a classical topological space if E is a singleton. In this case, the concepts of e -soft T_i -spaces and T_i -spaces are identical, and the concepts of *pt*-soft αT_i (soft αT_i) and αT_i -spaces are identical for $i = 0, 1, 2, 3, 4$. In order to show that item (ii) of the above proposition fails, we suffice by examples of classical topological spaces that satisfy an αT_i -space, but not T_i for $i = 0, 1, 2, 3, 4$.

Now, we proceed to show the main properties of *pt*-soft αT_i -spaces and determine the conditions under which they are equivalent.

Proposition 5. (X, τ, E) is *pt*-soft αT_1 if x_E is a soft α -closed set for every $x \in X$.

Proof. For every $x \neq y$, $y \in (x_E)^c$ and $x \in (y_E)^c$ are soft α -open sets. It is clear that $x \in (y_E)^c$ and $y \notin (y_E)^c$ and $y \in (x_E)^c$ and $x \notin (x_E)^c$. Hence, (X, τ, E) is *pt*-soft αT_1 . \square

Theorem 5. If (X, τ, E) has a soft basis consisting of soft α -clopen sets, then (X, τ, E) is *pt*-soft α regular.

Proof. Suppose that H_E is a soft α -closed subset of (X, τ, E) such that $x \notin H_E$ for some $x \in X$. Then, H_E^c is a soft α -open set such that $x \in H_E^c$. By hypothesis, there is a soft α -clopen set F_E in the basis of (X, τ, E) such that $x \in F_E \subseteq H_E^c$. Now, $H_E \subseteq F_E^c$. Obviously, F_E and F_E^c are disjoint soft α -open sets. Hence, (X, τ, E) is *pt*-soft α regular. \square

The following results determine the condition under which *pt*-soft αT_i , soft αT_i , and *tt*-soft αT_i -spaces are equivalent.

Theorem 6. The concepts of *pt*-soft αT_i and soft αT_i are equivalent for $i = 0, 1$ if (X, τ, E) is extended.

Proof. We prove the theorem in the case of $i = 0$, as the proof of $i = 1$ is analogous.

\Rightarrow : Let (X, τ, E) be a soft αT_0 -space and let $x \neq y$. Without loss of generality, there exists a soft α -open set U_E such that $x \in U_E$ and $y \notin U_E$. If $y \notin U(e)$ for each $e \in E$, then the proof is trivial. So, without loss of generality, we consider there exists $e \in E$ such that $y \notin U(e)$ and $y \in U(e')$ for each $e' \in E \setminus \{e\}$. Since (X, τ, E) is extended, then $U(e)$ is an α -open subset of (X, τ, E) . It follows from Theorem 2 that a soft set V_E given by $V(e) = U(e)$ and $V(e') = \emptyset$ for each $e' \in E \setminus \{e\}$ is a soft α -open set. Now, we have $x \in V_E$ and $y \notin V_E$. Hence, (X, τ, E) is *pt*-soft αT_0 .

\Leftarrow : Let (X, τ, E) be a *pt*-soft αT_0 -space and let $x \neq y$. Without loss of generality, there exists a soft α -open set U_E such that $x \in U_E$ and $y \notin U_E$. If $x \in U(e)$ for each $e \in E$, then the proof is trivial. So, without loss of generality, we consider there exists $e \in E$ such that $x \in U(e)$ and $x \notin U(e')$ for each $e' \in E \setminus \{e\}$. Since (X, τ, E) is extended, then there exists a soft α -open set V_E such that $V(e) = U(e)$ and $V(e') = X$ for each $e' \in E \setminus \{e\}$. Obviously, $x \in V_E$ and $y \notin V_E$. Hence, (X, τ, E) is soft αT_1 . \square

Corollary 1. The concepts of *pt*-soft αT_4 and soft αT_4 are equivalent if (X, τ, E) is extended.

Definition 15. (X, τ, E) is said to be α stable if every soft α -open set is stable.

Theorem 7. If (X, τ, E) is α stable, then the concepts of *pt*-soft αT_i , soft αT_i , and *tt*-soft αT_i -spaces are equivalent for each $i = 0, 1, 2, 3, 4$.

Proof. In the case of an α stable space, the relations of partial belong and total belong between ordinary points and soft α -open (soft α -closed) sets are identical, and the relations of partial nonbelong and total nonbelong between ordinary points and soft α -open (soft α -closed) sets are identical too. Hence, we obtain the desired equivalences. \square

Corollary 2. If (X, τ, E) is a soft α regular space, then the concepts of *pt*-soft αT_i , soft αT_i , and *tt*-soft αT_i -spaces are equivalent for each $i = 0, 1, 2, 3, 4$.

Interchangeability “transmission” of pt -soft αT_i -spaces and their corresponding spaces on classical topology (which are αT_i -spaces) are investigated in the following findings.

Theorem 8. *Let (X, τ, E) be extended. If there exists $e \in E$ such that (X, τ_e) is αT_i , then (X, τ, E) is pt -soft αT_i for each $i = 0, 1, 2$.*

Proof. We prove the theorem in the case of $i = 2$. The other cases follow similar lines.

Let (X, τ_e) be αT_2 and let $x \neq y \in X$. Then, there exist two disjoint α -open subsets M, N of (X, τ_e) containing x and y , respectively. It follows from Theorem 2 that there are two disjoint soft α -open subsets U_E and V_E of (X, τ, E) such that $U(e) = M$, $V(e) = N$ and $U(e') = V(e') = \emptyset$ for each $e' \in E \setminus \{e\}$. It can be seen that $x \in U_E$ and $y \notin U_E$ and $y \in V_E$ and $x \notin V_E$. Hence, (X, τ, E) is pt -soft αT_2 . \square

Theorem 9. *Let (X, τ, E) be extended. If all parametric topological space (X, τ_e) is αT_i , then (X, τ, E) is pt -soft αT_i for each $i = 0, 1, 2, 3, 4$.*

Proof. We prove the theorem in the cases of $i = 3, 4$. The other cases follow from the above theorem. It suffices to prove the property of pt -soft regular and soft normal.

First, we prove that (X, τ, E) is pt -soft regular. Let H_E be a soft α -closed set such that $x \notin H_E$. Then, there exists $e \in E$ such that $x \notin H(e)$. Since (X, τ_e) is α regular, then there exist α -open subsets M, N of (X, τ_e) such that $x \in M$ and $H(e) \subseteq N$. It follows from Theorem 2 that there exist soft α -open subsets U_E and V_E of (X, τ, E) which are defined as follows:

$$\begin{aligned} U(e) = M, U(e') = \emptyset, \quad & \text{for each } e' \in E \setminus \{e\}, \\ V(e) = N, V(e') = X, \quad & \text{for each } e' \in E \setminus \{e\}. \end{aligned} \quad (2)$$

This shows that $x \in U_E$, $H_E \subseteq V_E$. Obviously, U_E and V_E are disjoint. So, (X, τ, E) is pt -soft α regular. Hence, it is pt -soft αT_3 .

Second, we prove that (X, τ, E) is soft α normal. Let H_E and L_E be two disjoint soft α -closed sets. Then, $H(e)$ and $L(e)$ are two disjoint α -closed sets for each $e \in E$. Since (X, τ_e) is α normal, then there exists two disjoint α -open sets M and N such that $H(e) \subseteq M$ and $L(e) \subseteq N$. It follows from Theorem 2 that there exist soft α -open subsets U_E and V_E of (X, τ, E) which are defined as follows:

$$\begin{aligned} U(e) = M, U(e') = \emptyset, \quad & \text{for each } e' \in E \setminus \{e\}, \\ V(e) = N, V(e') = \emptyset, \quad & \text{for each } e' \in E \setminus \{e\}. \end{aligned} \quad (3)$$

Now, $\widetilde{U}_{\forall e \in E} U_E$ and $\widetilde{U}_{\forall e \in E} V_E$ are disjoint soft α -open sets such that $H_E \subseteq \widetilde{U}_{\forall e \in E} U_E$ and $L_E \subseteq \widetilde{U}_{\forall e \in E} V_E$. Thus, (X, τ, E) is soft α normal. Hence, it is pt -soft αT_4 . \square

For the sake of brevity, we present the following two theorems without proof.

Theorem 10. *If (X, τ, E) is α stable (soft α regular), then (X, τ_e) is αT_i iff (X, τ, E) is pt -soft αT_i for each $i = 0, 1, 2, 3, 4$.*

Theorem 11. *The property of being a pt -soft αT_i -space is hereditary for $i = 0, 1, 2, 3$.*

Now, we proceed to discuss the behaviour of pt -soft αT_i -spaces in relation with additive and topological properties and finite product spaces.

Theorem 12. *The property of being a pt -soft αT_i -space is an additive property for $i = 0, 1, 2, 3, 4$.*

Proof. We only prove the theorem in the case of $i = 4$. First, we prove that a property of pt -soft αT_i is additive. Let $x \neq y \in \bigoplus_{i \in I} X_i$. Then, the proof is trivial if x and y belong to the same X_{i_0} . Therefore, we consider there exist $i_0 \neq j_0 \in I$ such that $x \in X_{i_0}$ and $y \in X_{j_0}$. According to the definition of sum of soft topological spaces, we obtain that \widetilde{X}_{i_0} and \widetilde{X}_{j_0} are soft α -open subsets of $(\bigoplus_{i \in I} X_i, \tau, E)$. Hence, $(\bigoplus_{i \in I} X_i, \tau, E)$ is pt -soft αT_1 .

Second, we prove a property of soft α -normality is additive. Suppose that F_E and H_E are two disjoint soft α -closed subsets of $(\bigoplus_{i \in I} X_i, \tau, E)$. Then, $F_E \cap \widetilde{X}_i$ and $H_E \cap \widetilde{X}_i$ are soft α -closed in (X_i, τ_i, E) for each $i \in I$. By hypothesis, there are two disjoint soft α -open subsets U_{i_E} and V_{i_E} of (X_i, τ_i, E) such that $F_E \cap \widetilde{X}_i \subseteq U_{i_E}$ and $H_E \cap \widetilde{X}_i \subseteq V_{i_E}$. This implies that $F_E \subseteq \bigcup_{i \in I} U_{i_E}$, $H_E \subseteq \bigcup_{i \in I} V_{i_E}$, and $[\bigcup_{i \in I} U_{i_E}] \cap \bigcup_{i \in I} V_{i_E} = \emptyset$. Hence, $(\bigoplus_{i \in I} X_i, \tau, E)$ is a soft α -normal space. \square

Theorem 13. *The finite product of pt -soft αT_i -spaces is a pt -soft αT_i -space for $i = 0, 1, 2$.*

Proof. We prove the theorem in case of $i = 2$. The other cases follow similar lines.

Let (X, τ, E) and (Y, θ, E) be two pt -soft αT_2 -spaces and let $(x_1, y_1) \neq (x_2, y_2)$ in $X \times Y$. Then, $x_1 \neq x_2$ or $y_1 \neq y_2$. Without loss of generality, let $x_1 \neq x_2$. Then, there exist two disjoint soft α -open subsets G_E and H_E of (X, τ, E) such that $x_1 \in G_E$ and $x_2 \notin G_E$ and $x_2 \in H_E$ and $x_1 \notin H_E$. Obviously, $G_E \times \widetilde{Y}$ and $H_E \times \widetilde{X}$ are two disjoint soft α -open subsets $X \times Y$ such that $(x_1, y_1) \in G_E \times \widetilde{Y}$ and $(x_2, y_2) \notin G_E \times \widetilde{Y}$ and $(x_2, y_2) \in H_E \times \widetilde{Y}$ and $(x_1, y_1) \notin H_E \times \widetilde{Y}$. Hence, $X \times Y$ is pt -soft αT_2 . \square

4. An Application of Optimization via Soft Weak Structure Using pt -Soft T_i -Spaces ($i = 0, 1, 2$)

In this section, we present an application of optimal choices using the idea of pt -soft T_i -spaces ($i = 0, 1, 2$) on the content of soft weak structure. The idea of this application is based on personality characteristics of the applicants. We construct an algorithm of this application and provide an example to demonstrate how this algorithm is carried out.

First of all, we define pt -soft T_i -spaces ($i = 0, 1, 2$) on soft weak structure in a similar way of their counterparts on topological spaces.

Definition 16. A soft weak structure (X, τ, E) is said to be

- (i) pt -soft $\mathcal{W}T_0$ if for every two distinct points $x, y \in X$, there exists a \mathcal{W} -soft open set G_E such that $x \in G_E$ and $y \notin G_E$ or $y \in G_E$ and $x \notin G_E$.

- (ii) pt -soft $\mathscr{W}T_1$ if for every two distinct points $x, y \in X$, there exist \mathscr{W} -soft open sets G_E and F_E such that $x \in G_E$ and $y \notin G_E$ and $y \in F_E$ and $x \notin F_E$.
- (iii) pt -soft $\mathscr{W}T_2$ if for every two distinct points $x, y \in X$, there exist two disjoint \mathscr{W} -soft open sets G_E and F_E such that $x \in G_E$ and $y \notin G_E$ and $y \in F_E$ and $x \notin F_E$.

Tourism companies compete with each other to attract the customers by offering different programmes of their trip activities. One of the most important methods of promotion is the choice of places of carrying out the trip programmes. Herein, we will propose a method to help the customers to select the suitable tourism company that satisfies their options.

To this end, we consider some tourism companies will carry out their trips on the same region, and their trip programmes are distributed in some places of this region for a week. We consider the places are $X = \{h_i: i = 1, 2, \dots, n\}$ and the trip programmes are $E = \{e_i: i = 1, 2, \dots, m\}$, where e_1 are available places of accommodation, e_2 are available places of eating, e_3 are available places of watching cinema, e_4 are available places of watching cinema, e_5 are available places of celebrations, and so on.

The idea of this application is based on three factors: the first one is the classification of the places of trip activities as a soft set for each day. For example, let the places of trip activities on Tuesday be given as follows.

Then, we describe these activities by a soft set as follows:

$$G_E = \{(e_1, \{h_4, h_5, h_6\}), (e_2, \{h_1, h_5\}), (e_3, \emptyset), (e_4, \{h_2, h_5\}), (e_5, \{h_3, h_7\})\}. \tag{4}$$

Note that $G(e_3) = \emptyset$ does not imply any shortcoming. This case means that an activity of watching cinema is unavailable on Tuesday. This matter is reasonable because the participants of the trips need not carry out all the activities every day.

The second factor is the combination of a soft weak structure from the soft sets that represent the activities of the seven days (the whole period of the trip is a week). Then, we classify this soft weak structure in four categories: non- pt -soft $\mathscr{W}T_0$, pt -soft $\mathscr{W}T_0$, pt -soft $\mathscr{W}T_1$, and pt -soft $\mathscr{W}T_2$.

The third factor is based on the personality characteristics of the customers. In this application, we classify the customers into two groups: group of many visited places without repetition as much as possible and group of few visited places with repetition. According to this classification, if the soft weak structure is non- pt -soft $\mathscr{W}T_0$, then the customer will visit many places per day. Therefore, the customer who belongs to the group of many visited places will prefer tourism company that has trip programmes satisfying this condition. On the other hand, the customer who belongs to the group of few visited places will prefer tourism company that has trip programmes not satisfying this condition. He will prefer tourism company that has trip

programmes with few reiterated places. In other words, his or her optimal choice will be the weaker form of available pt -soft $\mathscr{W}T_i$ -spaces.

To illustrate this method, we give the succeeding interesting example.

Example 5. Consider that four tourism companies A, B, C, and D will carry out their trips on the same region and their trip programmes are distributed in seven places of this region for a week. We consider the places are $X = \{h_i: i = 1, 2, \dots, 7\}$ and we consider trip programmes are $E = \{e_i: i = 1, 2, \dots, 5\}$, where e_1, e_2, e_3, e_4 , and e_5 stand for the places of accommodation, eating, watching cinema, swimming, and celebrations, respectively.

In the succeeding four tables, we outline their trip programmes as given in their brochures.

Now, we transfer the four programmes of trips given in the above four tables to four soft weak structures as follows.

- (1) The soft weak structure of programmes of trip proposed by tourism company A (given in Table 1) is $\tau_A = \{\tilde{\Phi}, tG_{iE}n: qih =_1, 2x, 7 \dots C, ; 7\}$, where G_{iE} is given by

$$\begin{aligned} G_{1E} &= \{(e_1, \{h_1, h_2, h_3\}), (e_2, \{h_3, h_7\}), (e_3, \emptyset), (e_4, \{h_2, h_5\}), (e_5, \emptyset)\}, \\ G_{2E} &= \{(e_1, \{h_6\}), (e_2, \{h_6, h_7\}), (e_3, \{h_2\}), (e_4, \emptyset), (e_5, \{h_3, h_4, h_5\})\}, \\ G_{3E} &= \{(e_1, \{h_4\}), (e_2, \{h_4\}), (e_3, \{h_1, h_6\}), (e_4, \{h_3, h_6\}), (e_5, \{h_7\})\}, \\ G_{4E} &= \{(e_1, \{h_3, h_5, h_7\}), (e_2, \{h_2, h_3, h_6\}), (e_3, \{h_2, h_3\}), (e_4, \{h_4, h_5\}), (e_5, \{h_3, h_5\})\}, \\ G_{5E} &= \{(e_1, \{h_2, h_5\}), (e_2, \{h_2, h_5\}), (e_3, \emptyset), (e_4, \{h_1, h_3, h_6\}), (e_5, \emptyset)\}, \\ G_{6E} &= \{(e_1, \{h_4, h_7\}), (e_2, \{h_2, h_5\}), (e_3, \emptyset), (e_4, \emptyset), (e_5, \{h_6, h_7\})\}, \\ G_{7E} &= \{(e_1, \{h_6, h_7\}), (e_2, \{h_4\}), (e_3, \{h_3, h_4, h_6\}), (e_4, \emptyset), (e_5, \emptyset)\}. \end{aligned} \tag{5}$$

TABLE 1: Trip programmes of tourism company A.

X	E				
	e_1 : accommodation	e_2 : eating	e_3 : cinema	e_4 : swimming	e_5 : celebrations
Saturday	$\{h_1, h_2, h_3\}$	$\{h_3, h_7\}$	\emptyset	$\{h_2, h_5\}$	\emptyset
Sunday	$\{h_6\}$	$\{h_6, h_7\}$	$\{h_2\}$	\emptyset	$\{h_3, h_4, h_5\}$
Monday	$\{h_4\}$	$\{h_4\}$	$\{h_1, h_6\}$	$\{h_3, h_6\}$	$\{h_7\}$
Tuesday	$\{h_3, h_5, h_7\}$	$\{h_2, h_3, h_6\}$	$\{h_2, h_3\}$	$\{h_4, h_5\}$	$\{h_3, h_5\}$
Wednesday	$\{h_2, h_5\}$	$\{h_2, h_5\}$	\emptyset	$\{h_1, h_3, h_6\}$	\emptyset
Thursday	$\{h_4, h_7\}$	$\{h_2, h_5\}$	\emptyset	\emptyset	$\{h_6, h_7\}$
Friday	$\{h_6, h_7\}$	$\{h_4\}$	$\{h_3, h_4, h_6\}$	\emptyset	\emptyset

(2) The soft weak structure of programmes of trip proposed by tourism company B (given in Table 2) is

$\tau_B = \{\tilde{\Phi}, tG_{iE}n: qih =_1, 2x, 7 \dots C, ; 7\}$, where G_{iE} is given by

$$\begin{aligned}
 G_{1E} &= \{(e_1, \{h_1, h_2\}), (e_2, \{h_3\}), (e_3, \{h_1, h_2\}), (e_4, \emptyset), (e_5, \emptyset)\}, \\
 G_{2E} &= \{(e_1, \{h_7\}), (e_2, \{h_7\}), (e_3, \emptyset), (e_4, \{h_1, h_4, h_6\}), (e_5, \{h_1, h_4, h_6\})\}, \\
 G_{3E} &= \{(e_1, \{h_3, h_4\}), (e_2, \{h_1, h_2\}), (e_3, \{h_1, h_3\}), (e_4, \{h_3, h_4, h_6\}), (e_5, \emptyset)\}, \\
 G_{4E} &= \{(e_1, \{h_2, h_3\}), (e_2, \{h_4\}), (e_3, \{h_1, h_4, h_5\}), (e_4, \{h_2\}), (e_5, \emptyset)\}, \\
 G_{5E} &= \{(e_1, \{h_2, h_4, h_6\}), (e_2, \{h_1\}), (e_3, \{h_7\}), (e_4, \{h_3, h_5\}), (e_5, \emptyset)\}, \\
 G_{6E} &= \{(e_1, \{h_3\}), (e_2, \{h_7\}), (e_3, \emptyset), (e_4, \{h_1, h_6\}), (e_5, \{h_4, h_5, h_7\})\}, \\
 G_{7E} &= \{(e_1, \{h_1, h_4, h_5\}), (e_2, \{h_1, h_4, h_5\}), (e_3, \emptyset), (e_4, \{h_7\}), (e_5, \emptyset)\}.
 \end{aligned} \tag{6}$$

(3) The soft weak structure of programmes of trip proposed by tourism company C (given in Table 3) is

$\tau_C = \{\tilde{\Phi}, tG_{iE}n: qih =_1, 2x, 7 \dots C, ; 7\}$, where G_{iE} is given by

$$\begin{aligned}
 G_{1E} &= \{(e_1, \{h_1\}), (e_2, \{h_1, h_3\}), (e_3, \{h_2, h_3\}), (e_4, \emptyset), (e_5, \emptyset)\}, \\
 G_{2E} &= \{(e_1, \{h_6, h_7\}), (e_2, \{h_4, h_7\}), (e_3, \emptyset), (e_4, \{h_4, h_5\}), (e_5, \{h_4, h_5\})\}, \\
 G_{3E} &= \{(e_1, \{h_1, h_2, h_5\}), (e_2, \{h_3\}), (e_3, \{h_1, h_3\}), (e_4, \emptyset), (e_5, \{h_6\})\}, \\
 G_{4E} &= \{(e_1, \{h_2\}), (e_2, \{h_4\}), (e_3, \{h_1, h_2, h_4\}), (e_4, \emptyset), (e_5, \emptyset)\}, \\
 G_{5E} &= \{(e_1, \{h_4, h_5\}), (e_2, \{h_1\}), (e_3, \{h_4\}), (e_4, \{h_5\}), (e_5, \{h_4, h_5\})\}, \\
 G_{6E} &= \{(e_1, \{h_6, h_7\}), (e_2, \{h_2, h_3\}), (e_3, \{h_2, h_3, h_7\}), (e_4, \{h_6, h_7\}), (e_5, \emptyset)\}, \\
 G_{7E} &= \{(e_1, \{h_2, h_4, h_7\}), (e_2, \{h_4, h_7\}), (e_3, \{h_1, h_5\}), (e_4, \emptyset), (e_5, \emptyset)\}.
 \end{aligned} \tag{7}$$

(4) The soft weak structure of programmes of trip proposed by tourism company D (given in Table 4) is

$\tau_D = \{\tilde{\Phi}, tG_{iE}n: qih =_1, 2x, 7 \dots C, ; 7\}$, where G_{iE} is given by

TABLE 2: Trip programmes of tourism company B.

X	E				
	e_1 : accommodation	e_2 : eating	e_3 : cinema	e_4 : swimming	e_5 : celebrations
Saturday	$\{h_1, h_2\}$	$\{h_3\}$	$\{h_1, h_2\}$	\emptyset	\emptyset
Sunday	$\{h_7\}$	$\{h_7\}$	\emptyset	$\{h_1, h_4, h_6\}$	$\{h_1, h_4, h_6\}$
Monday	$\{h_3, h_4\}$	$\{h_1, h_2\}$	$\{h_1, h_3\}$	$\{h_3, h_4, h_6\}$	\emptyset
Tuesday	$\{h_2, h_3\}$	$\{h_4\}$	$\{h_1, h_4, h_5\}$	$\{h_2\}$	\emptyset
Wednesday	$\{h_2, h_4, h_6\}$	$\{h_1\}$	$\{h_7\}$	$\{h_3, h_5\}$	$\{h_5, h_6, h_7\}$
Thursday	$\{h_3\}$	$\{h_7\}$	\emptyset	$\{h_1, h_6\}$	$\{h_4, h_5, h_7\}$
Friday	$\{h_1, h_4, h_5\}$	$\{h_1, h_4, h_5\}$	\emptyset	$\{h_7\}$	\emptyset

TABLE 3: Trip programmes of tourism company C.

X	E				
	e_1 : accommodation	e_2 : eating	e_3 : cinema	e_4 : swimming	e_5 : celebrations
Saturday	$\{h_1\}$	$\{h_1, h_3\}$	$\{h_2, h_3\}$	\emptyset	\emptyset
Sunday	$\{h_6, h_7\}$	$\{h_4, h_7\}$	\emptyset	$\{h_4, h_5\}$	$\{h_4, h_5\}$
Monday	$\{h_1, h_2, h_5\}$	$\{h_3\}$	$\{h_1, h_3\}$	\emptyset	$\{h_6\}$
Tuesday	$\{h_2\}$	$\{h_4\}$	$\{h_1, h_2, h_4\}$	\emptyset	\emptyset
Wednesday	$\{h_4, h_5\}$	$\{h_1\}$	$\{h_4\}$	$\{h_5\}$	$\{h_4, h_5\}$
Thursday	$\{h_6, h_7\}$	$\{h_2, h_3\}$	$\{h_2, h_3, h_7\}$	$\{h_6, h_7\}$	\emptyset
Friday	$\{h_2, h_4, h_7\}$	$\{h_4, h_7\}$	$\{h_1, h_5\}$	\emptyset	\emptyset

TABLE 4: Trip programmes of tourism company D.

X	E				
	e_1 : accommodation	e_2 : eating	e_3 : cinema	e_4 : swimming	e_5 : celebrations
Saturday	$\{h_4, h_5, h_7\}$	$\{h_4, h_5, h_7\}$	$\{h_4, h_5, h_7\}$	\emptyset	\emptyset
Sunday	$\{h_1, h_6\}$	$\{h_3\}$	$\{h_2\}$	$\{h_1, h_2, h_6\}$	\emptyset
Monday	$\{h_6\}$	$\{h_3\}$	$\{h_1, h_6\}$	$\{h_3, h_5\}$	$\{h_7\}$
Tuesday	$\{h_3, h_5, h_7\}$	$\{h_3, h_6\}$	$\{h_3\}$	$\{h_4, h_5\}$	$\{h_3, h_5\}$
Wednesday	$\{h_2, h_5\}$	$\{h_2, h_5\}$	\emptyset	$\{h_1, h_2, h_4\}$	\emptyset
Thursday	$\{h_4, h_7\}$	$\{h_2, h_5\}$	\emptyset	\emptyset	$\{h_6, h_7\}$
Friday	$\{h_1, h_2\}$	$\{h_3\}$	$\{h_4\}$	$\{h_3, h_7\}$	\emptyset

$$\begin{aligned}
 G_{1E} &= \{(e_1, \{h_4, h_5, h_7\}), (e_2, \{h_4, h_5, h_7\}), (e_3, \{h_4, h_5, h_7\}), (e_4, \emptyset), (e_5, \emptyset)\}, \\
 G_{2E} &= \{(e_1, \{h_1, h_6\}), (e_2, \{h_3\}), (e_3, \{h_2\}), (e_4, \{h_1, h_2, h_6\}), (e_5, \emptyset)\}, \\
 G_{3E} &= \{(e_1, \{h_6\}), (e_2, \{h_3\}), (e_3, \{h_1, h_6\}), (e_4, \{h_3, h_5\}), (e_5, \{h_7\})\}, \\
 G_{4E} &= \{(e_1, \{h_3, h_5, h_7\}), (e_2, \{h_3, h_6\}), (e_3, \{h_3\}), (e_4, \{h_4, h_5\}), (e_5, \{h_3, h_5\})\}, \\
 G_{5E} &= \{(e_1, \{h_2, h_5\}), (e_2, \{h_2, h_5\}), (e_3, \emptyset), (e_4, \{h_1, h_2, h_4\}), (e_5, \emptyset)\}, \\
 G_{6E} &= \{(e_1, \{h_4, h_7\}), (e_2, \{h_2, h_5\}), (e_3, \emptyset), (e_4, \emptyset), (e_5, \{h_6, h_7\})\}, \\
 G_{7E} &= \{(e_1, \{h_1, h_2\}), (e_2, \{h_3\}), (e_3, \{h_4\}), (e_4, \{h_3, h_7\}), (e_5, \emptyset)\}.
 \end{aligned} \tag{8}$$

Now, we analyze the four programmes of trips with respect to soft separation axioms as follows.

- (1) A soft weak structure (X, τ_A, E) of tourism company A is non- pt -soft $\mathscr{W}T_0$ because $h_2 \neq h_5$ and there does not exist a \mathscr{W} -soft open subset G_E of (X, τ_A, E) such that h_2 partially belongs to G_E and h_5 does not totally belong to it or h_5 partially belongs to G_E and h_2 does not totally belong to it.

- (2) It can be checked that a soft weak structure (X, τ_B, E) of tourism company B is pt -soft $\mathscr{W}T_0$. On the other hand, it is not pt -soft $\mathscr{W}T_1$ because $h_2 \neq h_3$ and there does not exist a \mathscr{W} -soft open subset G_E of (X, τ_A, E) such that h_2 partially belongs to G_E and h_3 does not totally belong to it.
- (3) It can be checked that a soft weak structure (X, τ_C, E) of tourism company C is pt -soft $\mathscr{W}T_0$. On the other

hand, it is not pt -soft $\mathscr{W}T_1$ because $h_2 \neq h_3$ and there does not exist a \mathscr{W} -soft open subset G_E of (X, τ_A, E) such that h_3 partially belongs to G_E and h_2 does not totally belong to it.

- (4) It can be checked that a soft weak structure (X, τ_D, E) of tourism company D is pt -soft $\mathscr{W}T_1$. On the other hand, it is not pt -soft $\mathscr{W}T_2$ because $h_1 \neq h_2$ and there do not exist disjoint \mathscr{W} -soft open subsets G_{1E} and G_{2E} of (X, τ_A, E) partially containing h_1 and h_2 , respectively, such that h_2 does not totally belong to G_{1E} and h_1 does not totally belong to G_{2E} .

According to the data given above and their analysis, we infer that the offer presented by tourism company D is more convenient (optimal choice) for the customers who belong to the group of many visited places without repetition as much as possible. However, we infer that the optimal offers presented by tourism company B and tourism company C are more convenient (optimal choices) for the customers who belong to the group of few visited places with repetition as much as possible.

In what follows, we present an algorithm showing the method of selecting the optimal offers.

- (1) Select the desired location of trip.
- (2) Take the offering brochures given from some tourism companies.
- (3) Determine the number of visited places $X = \{h_i; i = 1, 2, \dots, n\}$ and available activities $E = \{e_i; i = 1, 2, \dots, m\}$.
- (4) Write every trip programmes in a table (as given in Tables 1–4).
- (5) Transfer each table in the previous step to its corresponding soft weak structure.
- (6) Classify the obtained soft weak structures with respect to soft pt -soft $\mathscr{W}T_i$ -spaces. In other words, determine which one is non- pt -soft $\mathscr{W}T_0$, pt -soft $\mathscr{W}T_0$, pt -soft $\mathscr{W}T_1$, or pt -soft $\mathscr{W}T_2$.
- (7) Determine an optimal choice according to your personality characteristic: if you belong to the group of many visited places without iteration, then you will prefer tourism company which has trip programmes satisfying strong form of available pt -soft $\mathscr{W}T_i$ -space. In contrast, If you belong to the group of few visited places, then you will prefer tourism company which has trip programmes satisfying weak form of available pt -soft $\mathscr{W}T_i$ -space.
- (8) If there is more than one optimal choice, then you can select any one of them satisfying his or her option.

Last but not least, we recommend the tourism companies to take into consideration the criteria proposed in this study when they prepare tourism programmes as they can enlist experts in this field and benefit from their experience to raise the standard of turnout.

5. Discussion

The method followed in the previous application is based on pt -soft $\mathscr{W}T_1$ -spaces ($i = 0, 1, 2$) which are defined in this study. This method relies on two factors, the first one comes from the classification induced from soft separation axioms and the second one comes from the personality characteristics of the customers.

One of the advantages of this technique is the relaxation of conditions of some structures such as soft topology, supra soft topology, and generalized soft topology. In other words, we do not need to check the finite soft union. We write a finite case instead of arbitrary case for a soft union because we deal with a system consisting of finite elements and attributes and intersection which gives us freedom and ease to model the phenomena under study. Another merit of this technique is the nature (type) of the belong and nonbelong relations that are utilized to define those types of soft separation axioms. These relations (partial belong and total nonbelong relations which are the core of our approach in this manuscript) offer multiple options to “transfer” the real-life problem to a mathematical model compared with their counterparts of soft separation axioms using total belong and total nonbelong relations introduced in [22].

On the other hand, there are some limitations of our method with the number of variables (which in this study are the days, places, and activities). According to our application, we should examine $\binom{7}{2} = 21$ different cases of places h_i . In this case, the total relation whether it is belonging or nonbelonging requires more soft sets to satisfy the different cases. This implies that modeling under total belong or total nonbelong hampers the description of the phenomenon; consequentially, the flexibility (completeness and accuracy) of this method to model a phenomenon is less than that of a method induced from partial belong and partial nonbelong relations given in [24]. We conclude that we can represent the phenomena using partial belong and partial nonbelong relations more easier than using partial belong and total nonbelong relations.

It noteworthy that the conditions regarding the number of variables in our approach will be similar to the required conditions in the case of total belong and partial nonbelong relations given in [23]. The differences are induced by replacing the total nonbelong relation \notin by the total belong relation \in .

Finally, we can apply this method, taking into account the nature of each phenomenon, in the engineering sciences as follows.

- (1) Decision making and topology have a long joint tradition since the modern statement of the classical Weierstrass extreme value theorem. It combines two topological concepts called continuity of a real-valued function and compactness of the domain (both with respect to a given topology). They represent a necessary and sufficient condition to guarantee the existence of the maximum and minimum values of the function. The success of this

technique was amplified by its adoption in fields like engineering sciences, computer sciences, and mathematical economics. This matter can be adopted on the version of soft setting by replacing the classical notions (compactness, function, and real numbers) by their soft counterparts (soft compactness, soft function, and soft real numbers).

- (2) Some practical experiments in the civil engineering require classification of the materials according to their characteristics (attribute set or parameter set E) which can be expressed using the concept of soft sets. Then, we study the separation of them with respect to the group of soft sets which are constructed from the practical experiments. In this group of soft sets, we add the absolute and null soft sets to initiate a soft weak structure. Finally, we determine the type of this soft weak structure with respect to pt -soft $\mathscr{W}T_i$ ($i = 0, 1, 2$) or non- pt -soft $\mathscr{W}T_0$.
- (3) The researchers in the communication engineering endeavor to select the best protocol to solve the noisy problems in wireless networks. They evaluate the performance of these protocols according to the proposed scenarios. We plan with some engineers to propose some protocols using the appropriate soft structure and compare with those proposed [46] to select the optimal protocol to solve the interference problems in wireless networks.

6. Conclusion

In this study, we have obtained a new class of soft topological spaces by defining the concepts of pt -soft αT_i -spaces ($i = 0, 1, 2, 3, 4$). They are formulated with respect to partial belong and total nonbelong relations between ordinary points and soft α -open sets. We have investigated the interrelations between these concepts and their parametric topological spaces. Some illustrative examples are given to clarify the obtained relationships and results. In the end, we have defined pt -soft T_i -spaces ($i = 0, 1, 2$) on a soft weak structure and applied them in solving a decision-making problem. In this regard, we have proposed an algorithm of an optimal selection and provided a real example to explain how this algorithm works. It is worthily noted that this paper is the first emergence of real applications of soft separation axioms in decision-making problems.

As future works, we shall study these concepts with respect to another generalizations of soft open sets such as soft preopen and soft semiopen sets. Also, we shall redefine these concepts using partial belong and partial nonbelong relations and investigate their characterizations. Moreover, we will study them on the contents of supra soft topological spaces, minimal soft topological spaces, and soft weak structures. In addition, we attempt to apply these concepts in the areas of engineering sciences and computer sciences (as we explained in Discussion section).

In conclusion, we hope that the initiated notions will be beneficial for researchers and scholars to promote and progress the study in soft topology and decision-making problems.

Data Availability

No data were used to support this study.

Conflicts of Interest

The author declares that there are no conflicts of interest.

References

- [1] D. Molodtsov, "Soft set theory—first results," *Computers & Mathematics with Applications*, vol. 37, no. 4-5, pp. 19–31, 1999.
- [2] P. K. Maji, R. Biswas, and A. R. Roy, "Soft set theory," *Computers & Mathematics with Applications*, vol. 45, no. 4-5, pp. 555–562, 2003.
- [3] M. I. Ali, F. Feng, X. Liu, W. K. Min, and M. Shabir, "On some new operations in soft set theory," *Computers & Mathematics with Applications*, vol. 57, no. 9, pp. 1547–1553, 2009.
- [4] K. V. Babitha and J. J. Sunil, "Soft set relations and functions," *Computers & Mathematics with Applications*, vol. 60, no. 7, pp. 1840–1849, 2010.
- [5] K. Qin and Z. Hong, "On soft equality," *Journal of Computational and Applied Mathematics*, vol. 234, no. 5, pp. 1347–1355, 2010.
- [6] T. M. Al-shami, "Investigation and corrigendum to some results related to g -soft equality and gf -soft equality relations," *Filomat*, vol. 33, no. 11, pp. 3375–3383, 2019.
- [7] T. M. Al-shami and M. E. El-Shafei, " T -soft equality relation," *Turkish Journal of Mathematics*, vol. 44, no. 4, pp. 1427–1441, 2020.
- [8] M. Ali, H. Khan, L. H. Son, F. Smarandache, and W. B. V. Kandasamy, "New soft sets based class of linear algebraic codes," *Symmetry*, vol. 10, pp. 1–10, 2018.
- [9] N. Çağman and S. Enginoglu, "Soft matrix theory and its decision making," *Computers and Mathematics with Applications*, vol. 59, pp. 3308–3314, 2010.
- [10] F. Karaaslan, "Soft classes and soft rough classes with applications in decision making," *Mathematical Problems in Engineering*, vol. 2016, p. 11, Article ID 1584528, 2016.
- [11] S. Yuksel, T. Dizman, G. Yildizdan, and U. Sert, "Application of soft sets to diagnose the prostate cancer risk," *Journal of Inequalities and Applications*, vol. 2013, p. 229, 2013.
- [12] M. Shabir and M. Naz, "On soft topological spaces," *Computers & Mathematics with Applications*, vol. 61, no. 7, pp. 1786–1799, 2011.
- [13] N. Çağman, S. Karataş, and S. Enginoglu, "Soft topology," *Computers and Mathematics with Applications*, vol. 62, pp. 351–358, 2011.
- [14] A. Aygüno lu and H. Aygün, "Some notes on soft topological spaces," *Neural Computing and Applications*, vol. 21, pp. 113–119, 2012.
- [15] T. Hida, "A comprasion of two formulations of soft compactness," *Annals of Fuzzy Mathematics and Informatics*, vol. 8, no. 4, pp. 511–524, 2014.
- [16] T. M. Al-shami, M. E. El-Shafei, and M. Abo-Elhamayel, "Almost soft compact and approximately soft Lindelöf spaces," *Journal of Taibah University for Science*, vol. 12, no. 5, pp. 620–630, 2018.
- [17] T. M. Al-shami and L. D. R. Koçinac, "Nearly soft Menger spaces," *Journal of Mathematics*, vol. 2020, p. 9, Article ID 3807418, 2020.

- [18] K. V. Babitha and S. J. John, "Studies on soft topological spaces," *Journal of Intelligent & Fuzzy Systems*, vol. 28, no. 4, pp. 1713–1722, 2015.
- [19] K. V. Babitha and J. J. Sunil, "Soft topologies generated by soft set relations," in *Handbook of Research on Generalized and Hybrid Set Structures and Applications for Soft Computing*, pp. 118–126, IGI Global Pub, Hershey, PA, USA, 2015.
- [20] A. Kharal and B. Ahmad, "Mappings on soft classes," *New Mathematics and Natural Computation*, vol. 7, no. 3, pp. 471–481, 2011.
- [21] D. Wardowski, "On a soft mapping and its fixed points," *Fixed Point Theory Application*, vol. 2013, p. 182, 2013.
- [22] M. E. El-Shafei, M. Abo-Elhamayel, and T. M. Al-shami, "Partial soft separation axioms and soft compact spaces," *Filomat*, vol. 32, no. 13, pp. 4755–4771, 2018.
- [23] T. M. Al-shami and M. E. El-Shafei, "Partial belong relation on soft separation axioms and decision-making problem, two birds with one stone," *Soft Computing*, vol. 24, no. 7, pp. 5377–5387, 2020.
- [24] M. E. El-Shafei and T. M. Al-shami, "Applications of partial belong and total non-belong relations on soft separation axioms and decision-making problem," *Journal of Computational and Applied Mathematics*, vol. 39, p. 138, 2020.
- [25] S. Bayramov and C. G. Aras, "A new approach to separability and compactness in soft topological spaces," *TWMS Journal of Applied and Engineering Mathematics*, vol. 9, no. 1, pp. 82–93, 2018.
- [26] O. Tantawy, S. A. El-Sheikh, and S. Hamde, "Separation axioms on soft topological spaces," *Annals of Fuzzy Mathematics and Informatics*, vol. 11, no. 4, pp. 511–525, 2016.
- [27] A. Singh and N. S. Noorie, "Remarks on soft axioms," *Annals of Fuzzy Mathematics and Informatics*, vol. 14, no. 5, pp. 503–513, 2017.
- [28] M. Terepeta, "On separating axioms and similarity of soft topological spaces," *Soft Computing*, vol. 23, no. 3, pp. 1049–1057, 2019.
- [29] J. C. R. Alcantud, "Soft open bases and a novel construction of soft topologies from bases for topologies," *Mathematics*, vol. 8, no. 5, p. 672, 2020.
- [30] T. M. Al-shami, "Comments on "Soft mappings spaces," *The Scientific World Journal*, vol. 2019, p. 2, Article ID 6903809, 2019.
- [31] T. M. Al-shami, "Comments on some results related to soft separation axioms," *Afrika Matematika*, vol. 31, no. 7-8, pp. 1105–1119, 2020.
- [32] S. J. John, *Soft Sets: Theory and Applications*, Springer, Berlin, Germany, 2021.
- [33] S. Ashraf, S. Abdullah, and L. Abdullah, "Child development influence environmental factors determined using spherical fuzzy distance measures," *Mathematics*, vol. 7, no. 8, p. 661, 2019.
- [34] S. Ashraf, S. Abdullah, T. Mahmood, F. Ghani, and T. Mahmood, "Spherical fuzzy sets and their applications in multi-attribute decision making problems," *Journal of Intelligent & Fuzzy Systems*, vol. 36, no. 3, p. 2829, 2019.
- [35] S. Ashraf, T. Mahmood, S. Abdullah, and Q. Khan, "Different approaches to multi-criteria group decision making problems for picture fuzzy environment," *Bulletin of the Brazilian Mathematical Society, New Series*, vol. 50, no. 2, pp. 373–397, 2019.
- [36] F. Feng, C. Li, B. Davvaz, and M. I. Ali, "Soft sets combined with fuzzy sets and rough sets: a tentative approach," *Soft Computing*, vol. 14, no. 9, pp. 899–911, 2010.
- [37] S. Das and S. K. Samanta, "Soft metric," *Annals of Fuzzy Mathematics and Informatics*, vol. 6, no. 1, pp. 77–94, 2013.
- [38] S. Nazmul and S. K. Samanta, "Neighbourhood properties of soft topological spaces," *Annals of Fuzzy Mathematics and Informatics*, vol. 6, no. 1, pp. 1–15, 2013.
- [39] M. Akdag and A. Ozkan, "Soft α -open sets and soft α -continuous functions α -open sets and soft α -continuous functions," *Abstract and Applied Analysis*, vol. 2014, Article ID 891341, 7 pages, 2014.
- [40] M. Akdag and A. Ozkan, "On soft α -separation axioms," *Journal of Advanced Studies in Topology*, vol. 5, no. 4, pp. 16–24, 2014.
- [41] T. M. Al-shami and E. A. Abo-Tabl, *Soft α -separation Axioms and α -Fixed Soft Points*.
- [42] T. M. Al-shami and L. D. R. Kočinac, "The equivalence between the enriched and extended soft topologies," *Journal of Computational and Applied Mathematics*, vol. 18, no. 2, pp. 149–162, 2019.
- [43] T. M. Al-shami, L. D. R. Kočinac, and B. A. Asaad, "Sum of soft topological spaces," *Mathematics*, vol. 8, no. 6, p. 990, 2020.
- [44] I. Zorlutuna, M. Akdag, W. K. Min, and S. Atmaca, "Remarks on soft topological spaces," *Annals of Fuzzy Mathematics and Informatics*, vol. 2, pp. 171–185, 2012.
- [45] A. H. Zakari, A. Ghareeb, and S. Omran, "On soft weak structures," *Soft Computing*, vol. 21, no. 10, pp. 2553–2559, 2017.
- [46] W. M. Bahgat, M. M. Al-dabaa, and H. H. Soliman, "Performance evaluation of cooperative relaying for MAC protocols," *International Journal of Computer Applications*, vol. 158, no. 1, pp. 35–49, 2017.