



Applied Mathematics and Nonlinear Sciences

<https://www.sciendo.com>

On Solutions of Fractional order Telegraph Partial Differential Equation by Crank-Nicholson Finite Difference Method

Mahmut Modanlı¹, Ali Akgül^{2†}

¹Harran University, Faculty of Arts and Sciences, Department of Mathematics, 63300-Şanlıurfa/Turkey

² Siirt University, Art and Science Faculty, Department Of Mathematics, 56100-Siirt/Turkey

Submission Info

Communicated by Hacı Mehmet Baskonus

Received March 24th 2019

Accepted May 15th 2019

Available online March 31st 2020

Abstract

The exact solution is calculated for fractional telegraph partial differential equation depend on initial boundary value problem. Stability estimates are obtained for this equation. Crank-Nicholson difference schemes are constructed for this problem. The stability of difference schemes for this problem is presented. This technique has been applied to deal with fractional telegraph differential equation defined by Caputo fractional derivative for fractional orders $\alpha = 1.1, 1.5, 1.9$. Numerical results confirm the accuracy and effectiveness of the technique.

Keywords: Fractional order Telegraph Partial Differential equations, Finite Difference Method, Stability

AMS 2010 codes: 26A33

Introduction and Preliminaries

Fractional differential equations have many implementations in finance, engineering, physics and seismology [1–3]. These type equations are solvable with respect to variables time and space. Some difference schemes are given for the space-fractional heat equations in [4–7, 18–22]. A new difference scheme for time fractional heat equation based on the Crank-Nicholson method has been presented in [5]. Orsingher and Beghin [14] have presented the Fourier transform of the fundamental solutions to time-fractional telegraph equations of order 2α . In [15], the time-fractional advection dispersion equations have been presented. In [16], Liu has studied fractional difference approximations for time-fractional telegraph equation. Modanlı and Akgül [12] have worked the second-order partial differential equations by two accurate methods. Finally, Modanlı and Akgül [13] have solved the fractional telegraph differential equations by theta-method. For more details see [23–27].

In this study, the Crank-Nicholson difference schemes method has been applied to fractional derivatives to get numerical results.

[†]Corresponding author.

Email address: aliakgul00727@gmail.com

Now, we examine the following fractional telegraph equations

$$\begin{cases} \frac{\partial^2 u(t,x)}{\partial t^2} + \frac{\partial^{\alpha-1} u(t,x)}{\partial t^{\alpha-1}} - \frac{\partial^2 u(t,x)}{\partial x^2} + u(t,x) = f(t,x), \\ 0 < x < L, 0 < t < T, 1 < \alpha < 2, \\ u(0,x) = r_1(x), u_t(0,x) = r_2(x), 0 \leq t \leq T, \\ u(t, X_L) = u(t, X_R) = 0, X_L < x < X_R. \end{cases} \quad (1)$$

Here, $r_1(x)$, $r_2(x)$ are smooth function defined with the space $[0, T]$, $f(t,x)$ is smooth function defined with the space $(0, L) \times (0, t)$ and $u(t,x)$ is unknown function with the domain $[0, L] \times [0, T]$. For the equation (1), the Crank-Nicholson finite difference scheme method is applied. With using this method, obtained numerical results are very good and efficient for given examples.

Definition 1. The Caputo fractional derivative $D_t^\alpha u(t,x)$ of order α with respect to time is defined as:

$$\begin{aligned} \frac{\partial^\alpha u(t,x)}{\partial t^\alpha} &= D_t^\alpha u(t,x) \\ &= \frac{1}{\Gamma(n-\alpha)} \int_0^t \frac{1}{(t-p)^{\alpha-n+1}} \frac{\partial^\alpha u(p,x)}{\partial p^\alpha} dp, \quad (n-1 < \alpha < n) \end{aligned} \quad (2)$$

and for $\alpha = n \in N$ defined as:

$$D_t^\alpha u(t,x) = \frac{\partial^\alpha u(t,x)}{\partial t^\alpha} = \frac{\partial^n u(t,x)}{\partial t^n}.$$

Definition 2. First-order approach difference method for the computation of the problem (1) has been presented as:

$$D_t^\alpha U_n^k \cong g_{\alpha,\tau} \sum_{j=0}^{k-1} b_j^{(\alpha)} (U_n^{k-j} - U_n^{k-j-1}), \quad (3)$$

where $g_{\alpha,\tau} = \frac{\tau^{2-\alpha}}{\Gamma(3-\alpha)}$ and $b_j^{(\alpha)} = (j+1)^{2-\alpha} - j^{2-\alpha}$.

Next section, we shall give Crank-Nicholson difference scheme for fractional order telegraph differential equation.

1 Crank-Nicolson Difference Scheme and its Stability

Using the formula (3) and definition of Crank-Nicholson first order difference schemes, we can construct the following difference scheme formula for (1) as:

$$\left\{ \begin{aligned} & \frac{u_n^{k+1} - 2u_n^k + u_n^{k-1}}{\tau^2} + g_{\alpha, \tau} \sum_{j=0}^{k-1} b_j^{(\alpha)} (u_n^{k-j} - u_n^{k-j-1}) + \frac{1}{2} (u_n^{k+1} + u_n^k) \\ & - \frac{1}{h^2} ((u_{n+1}^{k+1} - 2u_n^{k+1} + u_{n-1}^{k+1}) + (u_{n+1}^k - 2u_n^k + u_{n-1}^k)) = f_n^k, \\ & f_n^k = f(t_k, x_n), \quad 1 < \alpha < 2, \\ & u_n^0 = r_1(x_n), \quad \frac{u_n^1 - u_n^0}{\tau} = r_2(x_n), \quad 0 \leq n \leq M, \\ & u_0^k = u_M^k = 0, \quad 0 \leq k \leq N. \end{aligned} \right. \tag{4}$$

The formula (4) can be rewritten as:

$$\left\{ \begin{aligned} & (-\frac{1}{2h^2})u_{n+1}^k + (-\frac{1}{2h^2})u_{n+1}^{k+1} + (\frac{1}{\tau^2})u_n^{k-1} + (-\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2})u_n^k \\ & + (\frac{1}{\tau^2} + \frac{1}{h^2} + \frac{1}{2})u_n^{k+1} + (-\frac{1}{2h^2})u_{n-1}^k + (-\frac{1}{2h^2})u_{n-1}^{k+1} \\ & + \frac{1}{2}(u_n^{k+1} + u_n^k) + g_{\alpha, \tau} \sum_{j=0}^k b_j^{(\alpha)} (u_n^{k-j} - u_n^{k-j-1}) \\ & - \tau g_{\alpha, \tau} b_k r_2(x_n) = f_n^k, \quad 1 < \alpha < 2, \\ & f_n^k = f(t_k, x_n), \\ & u_n^0 = r_1(x_n), \quad \frac{u_n^1 - u_n^0}{\tau} = r_2(x_n), \quad 0 \leq n \leq M, \\ & u_0^k = u_M^k = 0, \quad 0 \leq k \leq N. \end{aligned} \right. \tag{5}$$

We can write the above system in matrix form as

$$Au_{n+1} + Bu_n + Cu_{n-1} = \varphi_n, \tag{6}$$

where $u_n = [u_n^0, u_n^1, \dots, u_n^N]$, $\varphi_n = [\varphi_n^0, \varphi_n^1, \dots, \varphi_n^N]^T$ and $\varphi_n^k = f(t_k, x_n) + \tau g_{\alpha, \tau} b_k r_2(x_n)$. Here $A_{(N+1) \times (N+1)}$ and $B_{(N+1) \times (N+1)}$ are the matrices of the following form,

$$A = -\frac{1}{2h^2} \begin{bmatrix} 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 & 0 \\ \cdot & \cdot & \cdot & \dots & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot & \cdot & \cdot \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 & 0 \end{bmatrix},$$

$$B = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ a - b_0 g_{\alpha, \tau} & b + g_{\alpha, \tau} & c & \dots & 0 \\ -b_1 g_{\alpha, \tau} & a + g_{\alpha, \tau}(b_1 - b_0) & b + g_{\alpha, \tau} & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ \vdots & \vdots & \vdots & \dots & \vdots \\ -b_{k-1} g_{\alpha, \tau} & (b_{k-2} - b_{k-3}) g_{\alpha, \tau} & (b_{k-3} - b_{k-4}) g_{\alpha, \tau} & \dots & c \\ -b_k g_{\alpha, \tau} & (b_k - b_{k-1}) g_{\alpha, \tau} & (b_{k-1} - b_{k-2}) g_{\alpha, \tau} & \dots & b + b_0 g_{\alpha, \tau} \end{bmatrix},$$

where $a = \frac{1}{\tau^2}$, $b = -\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2}$ and $c = \frac{1}{\tau^2} + \frac{1}{h^2} + \frac{1}{2}$.

Then we have

$$u_n = \alpha_{n+1} u_{n+1} + \beta_{n+1}, \quad 1 \leq k \leq M. \tag{7}$$

Next we should determine the matrices α_{n+1} and β_{n+1} above. Using the Dirichlet boundary condition

$$u(0, t) = u(0, L) = 0, \quad 0 \leq t \leq T,$$

we obtain $u_0 = \alpha_1 u_1 + \beta_1$. From that, we can choose $\alpha_1 = O_{(N+1) \times (N+1)}$ and $\beta_1 = O_{N+1}$. Substitute $u_n = \alpha_{n+1} u_{n+1} + \beta_{n+1}$ and $u_{n-1} = \alpha_n u_n + \beta_n$ into the equation (7), then

$$(A + B\alpha_{n+1} + A\alpha_n \alpha_{n+1})u_{n+1} + (B\beta_{n+1} + A\alpha_n \beta_{n+1} + A\beta_n) = \varphi_n.$$

Thus, we get

$$\begin{aligned} A + B\alpha_{n+1} + A\alpha_n \alpha_{n+1} &= 0, \\ B\beta_{n+1} + A\alpha_n \beta_{n+1} + A\beta_n &= \varphi_n. \end{aligned}$$

Thus, we obtain the following equalities

$$\begin{aligned} \alpha_{n+1} &= -(B + A\alpha_n)^{-1} A, \\ \beta_{n+1} &= (B + A\alpha_n)^{-1} (\varphi_n - A\beta_n), \end{aligned}$$

where $1 \leq n \leq M$.

For the stability, implementing the technique of analyzing the eigenvalues of the iteration matrices of the schemes.

Let $\rho(A)$ be the spectral radius of a matrix A , which indicates the maximum of the absolute value of the eigenvalues of the matrix A . We can write the following results.

Theorem 1. *The difference scheme (5) is stable.*

Proof. From the method [15], we should prove that $\rho(\alpha_n) < 1$, $1 \leq n \leq M$.

$\rho(\alpha_1) = 0 < 1$ is clearly.

$$\begin{aligned} \rho(\alpha_2) &= \|-B^{-1}A\| \leq \|-B^{-1}\| \|A\| = \frac{1}{\min_{1 \leq k \leq N-1} \left\{ |a_{kk}| - \sum_{\substack{m \neq k, \\ m=1}}^{N-1} |a_{km}| \right\}} \|A\| \\ &= \frac{\left| -\frac{1}{2h^2} \right| + \left| -\frac{1}{2h^2} \right|}{\left| -\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2} + \frac{\tau^{1-\alpha}}{\Gamma(3-\alpha)} \right|} = \frac{\frac{1}{h^2}}{-\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2} + \frac{\tau^{1-\alpha}}{\Gamma(3-\alpha)}} \leq 1, \text{ for } \frac{1}{2} + \frac{1}{h^2} \frac{\tau^{2-\alpha}}{\Gamma(3-\alpha)} - \frac{2}{\tau^2} \geq 0. \end{aligned}$$

If $\rho(\alpha_n) < 1$, let us calculate $\rho(\alpha_{n+1})$.

$$A = - \begin{bmatrix} 0 & 0 & 0 \dots 0 & 0 & 0 \\ 0 & \frac{\frac{1}{h^2}}{-\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2} + \frac{\tau^{1-\alpha}}{\Gamma(3-\alpha)} - \frac{1}{2h^2} \alpha_{n2,2}} & 1 \dots 0 & 0 & 0 \\ \cdot & \cdot & \cdot \dots \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \dots \cdot & \cdot & \cdot \\ 0 & 0 & 0 \dots 0 & 0 & 0 \\ 0 & 0 & 0 \dots 0 & 0 & \frac{\frac{1}{h^2}}{-\frac{2}{\tau^2} + \frac{1}{h^2} + \frac{1}{2} + \frac{\tau^{1-\alpha}}{\Gamma(3-\alpha)} - \frac{1}{2h^2} \alpha_{nN+1,N+1}} \end{bmatrix}.$$

We know that $\alpha_{ni} = \rho(\alpha_n)$ and $0 \leq \rho(\alpha_n) < 1$ for $2 \leq i \leq N + 1$. Then, we can obtain that $\rho(\alpha_{n+1}) < 1$. As a result, we obtain the desired result with induction. □

Remark 2. Using Matlab programming for $N = M = 10, \alpha = 1.5, 0 \leq t \leq 1, 0 \leq x \leq \pi$ and $h = \frac{\pi}{M}, \tau = \frac{1}{N}$, we obtain the following spectral radius of a matrix as:

$\rho(\alpha_1) = 0, \rho(\alpha_2) = 0.0482, \rho(\alpha_3) = 0.0484, \rho(\alpha_4) = 0.0485, \rho(\alpha_5) = 0.0482, \rho(\alpha_6) = 0.0487, \rho(\alpha_7) = 0.0487, \rho(\alpha_8) = 0.0475, \rho(\alpha_9) = 0.0487$ and $\rho(\alpha_{10}) = 0.0486$.

These final results prove the stability estimation of the Theorem 1.

Remark 3. Applying the method in [16, 17], we can get the convergence of the method from stability and consistency of the proposed method.

Now, we give numerical applications for the fractional telegraph partial differential equation by Crank-Nicholson method.

2 Numerical implementation

Example . We take into consideration the following fractional telegraph partial differential equation:

$$\begin{cases} \frac{\partial^2 u(t,x)}{\partial t^2} + \frac{\partial^{\alpha-1} u(t,x)}{\partial t^{\alpha-1}} - \frac{\partial^2 u(t,x)}{\partial x^2} + u(t,x) = \sin x(6t + 6\frac{t^{4-\alpha}}{\Gamma(5-\alpha)} + 2(t^3 + 1)), \\ 0 < x < \pi, 0 < t < 1, 1 < \alpha < 2, \\ u(0,x) = \sin x, u_t(0,x) = 0, 0 \leq t \leq 1, \\ u(t,0) = u(t,\pi) = 0, 0 \leq x \leq \pi. \end{cases} \tag{8}$$

The exact solution is given as $u(t,x) = (t^3 + 1) \sin x$. We implement difference schemes method to solve the problem. We utilize a procedure of modified Gauss elimination method for difference equation (8). We obtain the maximum norm of the error of the numerical solution by:

$$\varepsilon = \max_{\substack{n = 0, 1, \dots, M \\ k = 0, 1, 2, \dots, N}} |u(t,x) - u(t_k, x_n)|,$$

where $u_n^k = u(t_k, x_n)$ is the approximate solution. The error analysis in Table 1 gives our error analysis for difference schemes method.

Table 1. Error Analysis

$\tau = \frac{1}{N}, h = \frac{p^i}{M}$			
The difference scheme (8)			In method [13]
	α		
$N = M = 40$	1.5	0.0040	0.0242
$N = M = 80$	1.5	5.4707×10^{-4}	0.0118
$N = M = 160$	1.5	0.0022	0.0058
$N = 100, M = 10$	1.1	7.0178×10^{-4}	
	1.5	0.0045	
	1.9	0.0093	
$N = 225, M = 15$	1.1	3.4496×10^{-4}	
	1.5	0.0040	
	1.9	0.0083	
$N = 400, M = 20$	1.1	2.0762×10^{-4}	
	1.5	0.0034	
	1.9	0.0079	

We have compared Crank-Nicholson finite difference scheme method by the theta method [13] for the variable values $N = M = 40, 80, 160$. From these comparisons, we see that this method is more effective than the method used in [13].

3 Conclusion

In this work, stability estimates were presented for fractional telegraph differential equations. Stability inequalities were given for the difference schemes method. We applied the difference schemes-method for investigating fractional telegraph partial differential equations. Approximate solutions were obtained by this method. MATLAB software program was utilized for all results.

References

- [1] C. Celik, M. Duman. Crank-Nicholson method for the fractional equation with the Riesz fractional derivative. *Journal of computational physics*, 231:1743-1750, 2012.
- [2] I. I. Gorial. Numerical methods for fractional reaction-dispersion equation with Riesz space fractional derivative. *Eng. and Tech. Journal*, 29:709-715, 2011.
- [3] H. Jafari, V. D. Gejii. Solving linear and nonlinear fractional diffusion and wave equations by adomian decomposition. *Appl. Math. and Comput.*, 180:488-497, 2006.
- [4] I. Karatay, S. R. Bayramoglu, A. Sahin. Implicit difference approximation for the time fractional heat equation with the nonlocal condition. *Applied Numerical Mathematics*, 61:1281-1288, 2011.
- [5] I. Karatay, S. R. Bayramoglu, A. Sahin. A new difference scheme for time fractional heat equation based on the Crank-Nicholson method. *Fractional Calculus and Applied Analysis*, 16:892-910, 2013.
- [6] L. Su, W. Wang, Z. Yang. Finite difference approximations for the fractional advection-diffusion equation. *Physics Letters A.*, 373:4405-4408.2009.
- [7] C. Tadjeran, M. M. Meerschaert, H. P. Scheffler. A Second-order Accurate Numerical Approximation for the Fractional Diffusion Equation. *Journal of Computational Physics*, 213:205-213, 2006.
- [8] Li, C. and Cao, J. (2012) A Finite Difference Method for Time-Fractional Telegraph Equation. *Mechatronics and Embedded Systems and Applications (MESA)*, 2012.IEEE/ASME International Conference on. IEEE, 314-318.
- [9] Zhao, Z. and Li, C. (2012) Fractional Difference/Finite Element Approximations for the Time-Space Fractional Telegraph Equation. *Applied Mathematics and Computation*, 219, 2975-2988.
- [10] Ford, N.J., Rodrigues, M.M., Xiao, J. and Yan, Y. (2013) Numerical Analysis of a Two-Parameter Fractional Telegraph Equation. *Journal of Computational and Applied Mathematics*, 249, 95-106.

- [11] Sevimlican, A. (2010) An Approximation to Solution of Space and Time Fractional Telegraph Equations by He's Variational Iteration Method. *Mathematical Problems in Engineering*, 1-11 2010.
- [12] Modanli, Mahmut, and Ali Akgül. "On solutions to the second-order partial differential equations by two accurate methods." *Numerical Methods for Partial Differential Equations* 34.5 1678-1692 2018.
- [13] Modanli, Mahmut, and Ali Akgül. "Numerical solution of fractional telegraph differential equations by theta-method." *The European Physical Journal Special Topics* 226.16-18 3693-3703. 2017
- [14] Orsingher, Enzo, and Luisa Beghin. "Time-fractional telegraph equations and telegraph processes with Brownian time." *Probability Theory and Related Fields* 128.1 (2004): 141-160.
- [15] Smith, Gordon D. *Numerical solution of partial differential equations: finite difference methods*. Oxford university press, 1985.
- [16] Liu, Ru. "Fractional Difference Approximations for Time-Fractional Telegraph Equation." *Journal of Applied Mathematics and Physics* 6.01: 301 2018.
- [17] Richtmyer, Robert D., and Keith W. Morton. "Difference methods for initial-value problems." Malabar, Fla.: Krieger Publishing Co., c1994, 2nd ed. 1994.
- [18] A. Ashyralyev, M. Modanli. Nonlocal boundary value problem for telegraph equations. *AIP Conference Proceedings*, 1676:020078-1-020078-4, 2015.
- [19] S. Samko, A. Kibas, O. Marichev. *Fractional Integrals and derivatives: Theory and Applications*. Gordon and Breach, London, 1993.
- [20] I. Podlubny. *Fractional Differential Equations*, "Mathematics in Science and Engineering V198", Academic Press, San Diego 1999.
- [21] K. Nishimoto; *An essence of Nishimoto's Fractional Calculus*, Descartes Press Co. 1991.
- [22] K.M. Kowankar, A.D Gangal; *Fractional Differentiability of nowhere differentiable functions and dimensions*, CHAOS V.6, No. 4, 1996, American Institute of Physics.
- [23] A. Yokus, Comparison of Caputo and conformable derivatives for time-fractional Korteweg-de Vries equation via the finite difference method, *International Journal of Modern Physics B* Vol. 32, No. 29, 1850365 2018.
- [24] A. Yokus. D. Kaya, Numerical and exact solutions for time fractional Burgers' equation, *J. Nonlinear Sci. Appl.*, 10 3419-3428 2017.
- [25] A. Yokus. D. Kaya, Conservation laws and a new expansion method for sixth order Boussinesq equation, *AIP Conference Proceedings* 1676, 020062 2015.
- [26] A. Yokus., Numerical solution for space and time fractional order Burger type equation, *Alexandria Engineering Journal*, 2085-2091 2018.
- [27] A. Yokus., H. Bulut, On the numerical investigations to the Cahn-Allen equation by using finite difference method. *An International Journal of Optimization and Control: Theories & Applications (IJOCTA)*, 9(1), 18-23, 2018.

This page is intentionally left blank