

Mumtaz Ahmad Khan

On some operational representations of q -polynomials

Czechoslovak Mathematical Journal, Vol. 45 (1995), No. 3, 457–464

Persistent URL: <http://dml.cz/dmlcz/128538>

Terms of use:

© Institute of Mathematics AS CR, 1995

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON SOME OPERATIONAL REPRESENTATIONS
OF q -POLYNOMIALS

MUMTAZ AHMAD KHAN, Aligarh

(Received August 9, 1993)

1. INTRODUCTION

In an earlier paper [16] the present author defined the $T_{k,q,x}$ -operator by the relation

$$(1) \quad T_{k,q,x} \equiv x(1-q)\{[k] + q^k x D_{q,x}\},$$

where k is a constant, $|q| < 1$, $[k]$ is a q -number and $D_{q,x}$ is the q -derivative with respect to x .

The present paper gives applications of the $T_{k,q,x}$ -operator in finding operational representations for certain q -polynomials. In a separate communication it has been demonstrated how successfully this operator can be used to obtain generating functions and recurrence relations for q -Laguerre and other polynomials.

Some of the results obtained in this paper are q -analogues of those obtained by Al-Salam [5], Mittal [19] and Rainville [20] while the rest are believed to be new.

2. DEFINITIONS AND NOTATION

For most of the definitions and the notation needed in this paper, the reader is referred to the papers by Agarwal and Verma [2], Hahn [9], Khan [13–18] and to the books by Exton [8] and Slater [21]. However, definitions of some q -polynomials are given below:

The q -Jacobi polynomials are defined by

$$(1) \quad J_n(q, \gamma, \beta; x) = \frac{(-1)^n (q^\gamma)_n q^{n\gamma + n(n-1)/2}}{(q^{\beta+n-1})_n} {}_2\varphi_1[q^{-n}, q^{\beta+n-1}; q^\gamma; q^{1-\gamma}x]$$

and

$$(2) \quad P_{n,q}^{(\alpha,\beta)}(x) = \frac{(q^{1+\alpha})_n}{(q)_n} {}_2\varphi_1[q^{-n}, q^{1+\alpha+\beta+n}; q^{1+\alpha}; x].$$

Here the q -polynomial (2.1) is due to Hahn [10].

The q -Rice and generalized q -Rice polynomials are given by the relations

$$(3) \quad H_{n,q}(\xi, p, x) = {}_3\varphi_2[q^{-n}, q^{1+n}, q^\xi; q, q^p; x]$$

and

$$(4) \quad H_{n,q}^{(\alpha,\beta)}(\xi, p, x) = \frac{(q^{1+\alpha})_n}{(q)_n} {}_3\varphi_2[q^{-n}, q^{1+\alpha+\beta+n}, q^\xi; q^{1+\alpha}, q^p; x].$$

Further, the q -polynomial due to Al-Salam and Carlitz [7] is defined by

$$(5) \quad U_n^{(a)}(x) = x^n \left(\frac{1}{x}\right)_n {}_1\varphi_1 \left[\begin{matrix} q^{-n}; & -a \\ xq^{1-n}; & q \end{matrix} \right].$$

For $a = -1$ this polynomial gives the q -analogue of the Hermite polynomial.

Besides, the reader is referred to the papers by Jackson [12] and Khan [14] for q -Laguerre polynomials and Abdi [1] and Ismail [11] for q -Bessel polynomials.

3. RESULTS USED

Some of the results of Khan [16] required in this paper are listed below:

$$(1) \quad T_{k,q}^n {}_r\varphi_s^{(q)}[(a_r); (b_s); x] = x^n (q^k)_n {}_{r+1}\varphi_{s+1}^{(q)}[(a_r), n+k; (b_s), k; x],$$

$$(2) \quad \begin{aligned} T_{k,q}^n &= x^n (1-q)^n \prod_{j=0}^{n-1} ([k+j] + q^{k+j} x D_q) \\ &= x^n (1-q)^n \prod_{j=0}^{n-1} x^{-1} (1-q)^{-1} T_{k+j,q}, \end{aligned}$$

$$(3) \quad F(T_{k,q})\{x^\alpha f(x)\} = x^\alpha F(T_{k+\alpha,q})f(x),$$

$$(4) \quad T_{k,q}^n \{u(x)v(x)\} = \sum_{r=0}^n \binom{n}{r}_q q^{kr} T_{k,q}^{n-r} v(q^r x) T_{0,q}^r u(x).$$

4. OPERATIONAL REPRESENTATIONS

Here we give certain operational formulae and derive certain results for q -Laguerre polynomials. Besides, certain operational representations of some other q -polynomials will also be obtained.

Using (3.2) the following equivalent forms are obtained.

$$\begin{aligned}
 (1) \quad & \{x(1 - q^\alpha) + q^\alpha T_{k,q}\}^n f(x) = T_{k+\alpha,q}^n f(x) \\
 & = x^n (1 - q)^n \prod_{j=0}^{n-1} x^{-1} (1 - q)^{-1} T_{k+\alpha+j,q} f(x), \\
 (2) \quad & \{q^\alpha(1 + x)T_{k,q} + x(1 - q^\alpha) - x^2 q^\alpha\}^n f(x) \\
 & = x^n (1 - q)^n \prod_{j=0}^{n-1} \left\{ x(1 + x)q^{k+\alpha+j} D_q - \frac{xq^{k+\alpha+j}}{1 - q} + [k + \alpha + j] \right\} f(x),
 \end{aligned}$$

and

$$\begin{aligned}
 (3) \quad & \prod_{j=0}^{n-1} \left\{ q^\alpha T_{k,q} + x(1 - q^\alpha) - \frac{x^2 q^j}{1 - q} \right\} f(x) \\
 & = x^n (1 - q)^n \prod_{j=0}^{n-1} \left\{ xq^{k+\alpha+j} D_q - \frac{xq^j}{1 - q} + [k + \alpha + j] \right\} f(x).
 \end{aligned}$$

Formulae (4.2) and (4.3) are obtained by applying (4.1) to $e_q(-x)f(x)$ and $E_q(x)f(x)$, respectively.

Now the left hand side of (4.2) can also be written as

$$E_q(-x)T_{k+\alpha,q}^n \{e_q(-x)f(x)\} = x^{-\alpha} E_q(-x)T_{k,q}^n \{x^\alpha e_q(-x)f(x)\}.$$

Thus, we get the identity

$$\begin{aligned}
 (4) \quad & T_{k,q}^n \{x^\alpha e_q(-x)f(x)\} \\
 & = x^{\alpha+n} (q)_n e_q(-x) \sum_{r=0}^n \frac{(1+x)_r}{(q)_r x^r} {}_q L_{n-r}^{(\alpha+r)}(xq^{n+\alpha+k-1}, 1) T_{0,q}^r f(x).
 \end{aligned}$$

Similarly, we have

$$\begin{aligned}
 (5) \quad & T_{k,q}^n \{x^\alpha E_q(x)f(x)\} \\
 & = x^{\alpha+n} (q)_n E_q(xq^n) \sum_{r=0}^n \frac{q^{r(k+r)}}{(q)_r x^r} {}_q L_{n-r}^{(\alpha+k-1)}(xq^n) T_{0,q}^r f(x).
 \end{aligned}$$

Next, considering the operator ${}_0\varphi_1[-, q^{\alpha+k}; -tT_{k,q}]$, we obtain

$$(6) \quad {}_0\varphi_1[-, q^{\alpha+k}; -tT_{k,q}]x^{\alpha+n} = x^{\alpha+n} {}_1\varphi_1[q^{k+\alpha+n}, q^{k+\alpha}; -xt].$$

One can also easily obtain the operational formulae

$$(7) \quad {}_0\varphi_1[-; q^{\alpha+k}; T_{k,q}] \left\{ \frac{x^\alpha}{(1-xt)_{k+\alpha}} \right\} = \frac{x^\alpha}{(1-xt)_{k+\alpha}} e_q \left(\frac{x}{[1-xtq^{k+\alpha}]} \right)$$

and

$$(8) \quad {}_0\varphi_1 \left[\begin{matrix} ; & T_{k,q} \\ q^{k+\alpha}; & q \end{matrix} \right] \left\{ \frac{x^\alpha}{(1-xt)_{k+\alpha}} \right\} = \frac{x^\alpha}{(1-xt)_{k+\alpha}} E_q \left(\frac{-xq}{[1-xtq^{k+\alpha}]} \right).$$

As this stage we consider the following q -polynomials:

(A) *q-Laguerre Polynomials.* We shall obtain certain formulae and operational representations of q -Laguerre polynomials. Putting $f(x) = 1$ in (4.4) and (4.5) and taking different values of α and k , we get a number of operational representations for the q -Laguerre polynomials ${}_qL_n^{(\alpha)}(x, 1)$ and ${}_qL_n^{(\alpha)}(x)$, e.g.,

$$(9) \quad T_{k,q}^n \{x^\alpha e_q(-x)\} = x^{\alpha+n} (q)_n e_q(-x) {}_qL_n^{(\alpha+k-1)}(xq^{n+\alpha+k-1}, 1),$$

$$(10) \quad T_{k,q}^n \{x^\alpha E_q(x)\} = x^{\alpha+n} (q)_n E_q(xq^n) {}_qL_n^{(\alpha+k-1)}(xq^n)$$

are obtained by taking $f(x) = 1$ in (4.4) and (4.5).

By a simple change of variable, we also note that

$$(11) \quad T_{k,q}^n \{x^\alpha e_q(-\lambda x)\} = x^{\alpha+n} (q)_n e_q(-\lambda x) {}_qL_n^{(\alpha+k-1)}(\lambda xq^{n+\alpha+k-1}, 1)$$

and

$$(12) \quad T_{k,q}^n \{x^\alpha E_q(\lambda x)\} = x^{\alpha+n} (q)_n E_q(\lambda xq^n) {}_qL_n^{(\alpha+k-1)}(\lambda xq^n).$$

Now (4.11) and (4.12) can also be written as

$$(13) \quad \{q^\alpha(1+\lambda x)T_{k,q} + x(1-q^\alpha) - \lambda x^2 q^\alpha\}^n \cdot 1 = x^n (q)_n {}_qL_n^{(\alpha+k-1)}(\lambda xq^{n+\alpha+k-1}, 1)$$

and

$$(14) \quad \{q^\alpha T_{k,q} + x(1-q^\alpha) - \lambda x^2\}^n \cdot 1 = x^n (q)_n {}_qL_n^{(\alpha+k-1)}(\lambda xq^n).$$

Further, (4.9) gives

$$T_{k,q}^m \{x^{\alpha+n} e_q(-x) {}_qL_n^{(\alpha+k-1)}(xq^{n+\alpha+k-1}, 1)\} = T_{k,q}^m \left[\frac{1}{(q)_n} T_{k,q}^n \{x^\alpha e_q(-x)\} \right].$$

Hence

$$(15) \quad \begin{aligned} T_{k,q}^m \{x^{\alpha+n} e_q(-x)_q L_n^{(\alpha+k-1)}(xq^{n+\alpha+k-1}, 1)\} \\ = \frac{(q)_{m+n}}{(q)_n} x^{\alpha+m+n} e_q(-x)_q L_{m+n}^{(\alpha+k-1)}(xq^{m+n+\alpha+k-1}, 1). \end{aligned}$$

Similarly, (4.10) gives

$$(16) \quad \begin{aligned} T_{k,q}^m \{x^{\alpha+n} E_q(xq^n)_q L_n^{(\alpha+k-1)}(xq^n)\} \\ = \frac{(q)_{m+n}}{(q)_n} x^{\alpha+m+n} E_q(xq^{m+n})_q L_{m+n}^{(\alpha+k-1)}(xq^{m+n}). \end{aligned}$$

Using the q -analogue of Kummer's transform (4.6) yields

$${}_0\varphi_1[-; q^{\alpha+k}; -tT_{k,q}]x^{\alpha+n} = x^{\alpha+n} e_q(-xt)_1 \varphi_1 \left[\begin{matrix} q^{-n}; & xtq^{n+\alpha+k-1} \\ q^{k+\alpha}; & q \end{matrix} \right]$$

which can alternatively be written as

$$(17) \quad \begin{aligned} {}_0\varphi_1[-; q^{\alpha+k}; -tT_{k,q}]x^{\alpha+n} \\ = \frac{(q)_n}{(q^{k+\alpha})_n} x^{\alpha+n} e_q(-xt)_q L_n^{(\alpha+k-1)}(xtq^{n+\alpha+k-1}, 1). \end{aligned}$$

Similarly,

$$(18) \quad {}_0\varphi_1 \left[\begin{matrix} ; & -tT_{k,q} \\ q^{k+\alpha}; & q \end{matrix} \right] x^{\alpha+n} = \frac{(q)_n x^{\alpha+n}}{(q^{k+\alpha})_n} E_q(xtq^{1+n})_q L_n^{(\alpha+k-1)}(xtq^{1+n}).$$

Also, we have

$$(19) \quad \left(1 + \frac{t}{T_{k,q}}\right)_n x^{-\alpha-k} = \frac{x^{-\alpha-k} (q)_n}{(q^{1+\alpha})_n} {}_qL_n^{(\alpha)}(tq^{\alpha+n}/x, 1).$$

As an immediate consequence of the Leibniz formula (3.4) and the formula (4.9) we get

$$(20) \quad \begin{aligned} {}_qL_n^{(\alpha+\beta+k)}(xq^{n+\alpha+\beta+k}, 1) \\ = \sum_{r=0}^n \binom{\beta+r}{r}_q q^{r(k+\alpha)} (1+x)_r {}_qL_{n-r}^{(\alpha+k-1)}(xq^{\alpha+n+k-1}, 1), \end{aligned}$$

and using (3.4) and (4.10) we obtain

$$(21) \quad {}_qL_n^{(\alpha+\beta+k)}(xq^n) = \sum_{r=0}^n \binom{\beta+r}{r}_q q^{r(k+\alpha)} {}_qL_{n-r}^{(\alpha+k-1)}(xq^n).$$

Formula (4.20) is obtained by putting $u = x^{1+\beta}$ and $v = x^\alpha e_q(-x)$ in (3.4), while (4.21) is obtained by putting $u = x^{1+\beta}$ and $v = x^\alpha E_q(x)$ in (3.4). On the other hand, if we put $u = x^\beta E_q(\mu, x)$, $v = x^\alpha e_q(-\lambda x)$ in (3.4) and then employ (4.11) and (4.12), we get the following addition-like theorem, involving the q -Laguerre polynomials ${}_qL_n^{(\alpha)}(x, 1)$ and ${}_qL_n^{(\alpha)}(x)$:

$$(22) \quad {}_qL_n^{(\alpha+\beta+k-1)}([\lambda + \mu]xq^{n+\alpha+\beta+k-1}, 1) \\ = \sum_{r=0}^n \frac{q^{r(k+\alpha)}(1 + \lambda x)_r}{(1 - \mu x)_r} {}_qL_{n-r}^{(\alpha+k-1)}(\lambda xq^{n+\alpha+k-1}, 1) {}_qL_r^{(\beta-1)}(\mu xq^r).$$

From (4.13) and the shift rule (3.3) we have the following formula:

$$(23) \quad \frac{(q)_{m+n}}{(q^{k+\alpha})_n (q)_m} {}_qL_{m+n}^{(\alpha+k-1)}(xq^{m+n+\alpha+k-1}, 1) \\ = \sum_{r=0}^n \binom{n}{r}_q \frac{(-x)^r q^{r(r+\alpha+k-1)}}{(q^{k+\alpha})_r} {}_qL_m^{(n+r+\alpha+k-1)}(xq^{m+n+r+\alpha+k-1}, 1).$$

Similarly, we obtain

$$(24) \quad \frac{(q)_{m+n}}{(q)_m (q^{k+\alpha})_n} {}_qL_{m+n}^{(\alpha+k-1)}(xq^{m+n}) = \sum_{r=0}^n \frac{(q^{-n})_r x^r q^{rn}}{(q)_r (q^{k+\alpha})_r} {}_qL_m^{(n+r+\alpha+k-1)}(xq^m).$$

(B) *q-Bessel Polynomials.* Here we shall give three operational representations for q -Bessel Polynomials. One can obtain many others

$$(25) \quad T_{c+n,q}^n e_q(q^{n+1}/x) = \frac{(q)_n}{(q^c)_n} (-1)^n q^{\frac{1}{2}n(n+1)} e_q(q/x) J(q; c, n; x).$$

To obtain (4.25), $e_q(q^{n+1}/x)$ is replaced by its equivalent infinite series and $T_{c+n,q}^n$ is operated on the variable x of the series. We then use the q -analogue of Kummer's transform and finally the resulting finite ${}_1\varphi_1$ series is written in reverse order.

Similarly, we also have

$$(26) \quad T_{c,q}^n e_q\left(\frac{1}{x}\right) = \frac{(q)_n (-x)^n}{(xq)_n (q^{1-c})_n} q^{\frac{1}{2}n(n+1)-nc} e_q\left(\frac{1}{x}\right) J(q; c-n, n; xq^{n+1})$$

and

$$(27) \quad T_{c-n,q}^n e_q\left(\frac{1}{x}\right) = \frac{(q)_n (q^{1-c})_n (-x)^n q^{\frac{1}{2}n(3n+1)-nc}}{(xq)_n (q^{1-c})_{2n}} e_q\left(\frac{1}{x}\right) J(q; c-2n, n; xq^{n+1}).$$

(C) *q*-Jacobi Polynomials. We give here the following operational representations for the *q*-Jacobi polynomials $J_n(q, \alpha, \beta; x)$ due to Hahn [10] and the *q*-Jacobi polynomials $P_{n,q}^{(\alpha,\beta)}(x)$:

$$(28) \quad T_{a,q}^n(1 - xq^{1-a-n})_{n+b-a-1} = \frac{(xq^{1-a})_\infty (q^{b+n-1})_n (-x)^n}{(xq^{b-2a})_\infty q^{(1/2)n(n-1)+na}} J_n(q, a, b; x)$$

and

$$(29) \quad T_{a+1,q}^n(1 - xq^{-n})_{b+n} = x^n(1-x)_b (q)_n P_{n,q}^{(a,b)}(x).$$

Also, we have

$$(30) \quad T_{a+1,q}^n(1-x)_b = \frac{(q)_n(1-xq^n)_\infty x^n}{(1-xq^b)_\infty} P_{n,q}^{(a,b-n)}(xq^n)$$

and

$$(31) \quad T_{1+a-n,q}^n(1-x)_b = \frac{(q)_n(1-xq^n)_\infty x^n}{(1-xq^b)_\infty} P_{n,q}^{(a-n,b)}(xq^n).$$

Relations (4.30) and (4.31) can alternatively be written as follows:

$$(32) \quad T_{a,q}^n(1-x)_b = \frac{(q^{a+b})_n(1-xq^b)_\infty (-x)^n}{q^{\frac{1}{2}n(n-1)+na}(1-xq^b)_\infty} J_n(q, a, 1+a+b-n; xq^{n+a-1}),$$

$$(33) \quad T_{a-n,q}^n(1-x)_b = \frac{(q^{a+b})_n(1-xq^n)_\infty (-x)^n q^{-na+n(n+1)/2}}{(1-xq^b)_\infty} J_n(q, a-n, 1+a+b-n; xq^{a-1}).$$

(D) *Generalized q-Rice Polynomials*. Using (3.1) we have the following operational representation for the generalized *q*-Rice polynomials $H_{n,q}^{(\alpha,\beta)}(\xi, p, x)$:

$$(34) \quad T_{1+\alpha,q}^{n+\beta} {}_2\varphi_1[q^{-n}, q^\xi; q^p; x] = x^{n+\beta} (q^{1+\alpha+n})_\beta (q)_n H_{n,q}^{(\alpha,\beta)}(\xi, p, x).$$

If we put $\alpha = 0 = \beta$, (4.34) reduces to

$$(35) \quad T_{1,q}^n {}_2\varphi_1[q^{-n}, q^\xi; q^p; x] = x^n (q)_n H_{n,q}(\xi, p, x).$$

(E) *A q-polynomial of Al-Salam and Carlitz*. One can easily obtain the following operational representation for $U_n^{(a)}(x)$:

$$(37) \quad (1-x)(-1)^n q^{n(n-1)/2} e_q\left(\frac{1}{a} T_{1-n,q,xa}\right) G_n(aq^{-1}, q) = U_n^{(a)}(x)$$

where $G_n(x, q)$ is the Szegő polynomial defined by

$$(38) \quad G_n(x, q) = \sum_{r=0}^n \binom{n}{r}_q q^{r(r-n)} x^r.$$

References

- [1] *W.H. Abidi*: A basic analogue of the Bessel polynomials. *Math. Nachr.* 30 (1965), 209–219.
- [2] *R.P. Agarwal and A. Verma*: Generalized basic hypergeometric series with unconnected bases. *Proc. Camb. Philos. Soc.* 63 (1967), 727–734.
- [3] *N.A. Al-Salam*: Some operational formulas for the q -Laguerre polynomials. *Fibonacci Quarterly* 22 (1984), 166–170.
- [4] *N.A. Al-Salam*: On some q -operators with applications. *Proc. Konen. Nederl. Akademiev. Wetenschappen, Ser. A* 92 (1989), 1–13.
- [5] *W.A. Al-Salam*: Operational representation for the Laguerre and Hermite polynomials. *Duke Math. Journal* 31 (1964), 127–142.
- [6] *W.A. Al-Salam and A. Verma*: Orthogonality Preserving Operators. I. *Atti Della Accademia Nazionale Dei Lincei, ser VIII Vol. 2 VIII* (1975), 833–838.
- [7] *W.A. Al-Salam and L. Carlitz*: Some orthogonal q -polynomials. *Math. Nachr.* 30 (1965), 47–61.
- [8] *H. Exton*: q -Hypergeometric Functions and Applications. John Wiley and Sons (Halsted Press), New York — Ellis Horwood, Chichester, 1983.
- [9] *W. Hahn*: Beiträge zur Theorie der Heinschen Reihen, Die 24 Integrale der hypergeometrischen q -Differenzgleichung, Das q -Analogen der Laplace Transformation. *Math. Nachr.*
- [10] *W. Hahn*: q -Differenzgleichung, Das q -Analogen der Laplace Transformation. *Math. Nachr.* 2 (1949).
- [11] *M.E.H. Ismail*: The zeros of basic Bessel functions, the functions $J_{\nu+ax}(x)$ and associated orthogonal polynomials. *J. Math. Anal. Appl.* 86 (1982), no. 1, 1–19.
- [12] *F.H. Jackson*: Basic double hypergeometric functions. *Quart. J. Math. (Oxford)* 15 (1944), 49–61.
- [13] *M.A. Khan*: Certain fractional q -integrals and q -derivatives. *Nanta Mathematica* 7 (1974), no. 1, 52–60.
- [14] *M.A. Khan*: On q -Laguerre polynomials. *Ganita* 34 (1983), no. 1 and 2, 111–123.
- [15] *M.A. Khan*: q -Analogue of certain operational formulae. *Houston J. Math.* 13 (1987), no. 1, 75–82.
- [16] *M.A. Khan*: On a calculus for the $T_{k,q,x}$ -operator. *Mathematica Balkanica, New Series* 6 (1992), no. 1, 83–90.
- [17] *M.A. Khan and A.H. Khan*: Fractional q -integration and integral representations of ‘Bibasic’ double hypergeometric series of higher order. *Acta Mathematica Vietnamica* 11 (1986), no. 2, 234–240.
- [18] *M.A. Khan and A.H. Khan*: On some characterizations of q -Bessel polynomials. *Acta Math. Viet.* 15 (1990), no. 1, 55–59.
- [19] *H.B. Mittal*: Operational representation for the generalized Laguerre polynomials. *Glasnik Matematički* 6(26) (1977), no. 1, 45–53.
- [20] *E.D. Rainville*: Special Functions. The MacMillan Co., New York, 1960.
- [21] *L.J. Slater*: Generalized Hypergeometric Functions. Cambridge University Press, 1966.
- [22] *H.M. Srivastava and H.L. Manocha*: A Treatise on Generating Functions. John Wiley and Sons (Halsted Press), New York; Ellis Horwood, Chichester, 1984.

Author's address: Department of Applied Mathematics, Z.H. College of Engg. and Technology, Faculty of Engineering, A.M.U., Aligarh-202002, U.P., India.