

Work supported by the National Institute of Health, Public Health
Service, Grant No. GM-12868-04.

ON SOME SELECTION PROCEDURES IN TWO-WAY LAYOUTS

by

PRANAB KUMAR SEN AND MADAN LAL PURI

University of North Carolina, Chapel Hill
and Indiana University, Bloomington

Institute of Statistics Mimeo Series No. 630

June, 1969

ON SOME SELECTION PROCEDURES IN TWO-WAY LAYOUTS*

BY PRANAB KUMAR SEN AND MADAN LAL PURI

University of North Carolina, Chapel Hill

and Indiana University, Bloomington

0. Summary. In the one-way layout problem, Puri and Puri (1969) have considered some selection procedures which are nonparametric analogues of some earlier procedures by Bechhofer (1954), Gupta and Sobel (1958), and Paulson (1952), among others. In the present paper, some of these procedures are extended to the two-way layout problem in the parametric as well as nonparametric setup. Allied efficiency results are also studied.

1. Introduction. For a two-factor complete block design with one observation per cell, we express the observable random variables $X_{i\alpha}$ ($i=1, \dots, c; \alpha=1, \dots, n$) as

$$(1.1) \quad X_{i\alpha} = \mu + \beta_{\alpha} + \tau_i + \epsilon_{i\alpha}, \quad \sum_{i=1}^c \tau_i = 0,$$

where μ is the mean-effect, β_1, \dots, β_n are the block effects (nuisance parameters for the fixed effects model or random variables for the mixed effects model), τ_1, \dots, τ_c are the treatment effects, and the $\epsilon_{i\alpha}$ are the error components. It is assumed that $\underline{\epsilon}_{\alpha} = (\epsilon_{1\alpha}, \dots, \epsilon_{c\alpha})$, $\alpha=1, \dots, n$ are independent and identically distributed stochastic vectors with a continuous cumulative distribution function (cdf) $F(\underline{\epsilon})$, $\underline{\epsilon} \in \mathbb{R}^c$ (the real c space), where $F(\underline{\epsilon})$ is symmetric in its c arguments, that is, for any $\underline{\epsilon} \in \mathbb{R}^c$ and any permutation (i_1, \dots, i_c) of $(1, \dots, c)$

$$(1.2) \quad F(\epsilon_1, \dots, \epsilon_c) = F(\epsilon_{i_1}, \dots, \epsilon_{i_c}).$$

[Note that if all the nc errors are independent and identically distributed, (1.2) holds, but the converse is not necessarily true.] Our purpose is to study some (parametric as well as nonparametric) multiple decision procedures for the following three problems: (i) selection of the best t treatments without regard to order, (ii) selection of the best t treatments with regard to order, and (iii) selection

* Work supported by the National Institute of Health, Public Health Service, Grant No. GM-12868-04.

of all the treatments which are as good as or better than a standard treatment. The bestness of the treatments is judged by the largeness of the τ_i .

In the parametric case, Bechhofer (1954) has studied the first problem under the assumption that the errors are independent and normally distributed. It is shown here that if the errors are jointly (within each block) normally distributed and are equally correlated then his procedure remains valid. This covers the situation (1.2) which may arise often in mixed effects model. It is also shown that if (1.2) holds and F admits of the existence of second order moments, then the Bechhofer procedure remains valid for large samples, even if F is not normal.

In the nonparametric setup, Puri and Puri (1969) have considered these selection procedures for the one-way layout problems, and their procedures are based on a class of rank order statistics. Here, instead of these statistics, we consider the rank order estimators of $\{\tau_i\}$ by Puri and Sen (1967) to provide asymptotically distribution-free selection procedures for the two-way layout problems. The asymptotic relative performances and efficiencies of the parametric as well as nonparametric procedures are studied. The cases of paired comparisons designs as well as one-way layout problems are also briefly presented.

2. Preliminary notions. Let $\tau_{[1]} \leq \dots \leq \tau_{[c]}$ denote the (actual) ranked τ 's (which are not known). Let then

$$(2.1) \quad Z_{ni} = \bar{X}_{ni} - \bar{X}_n, \quad \bar{X}_{ni} = n^{-1} \sum_{\alpha=1}^n X_{i\alpha}, \quad i=1, \dots, c; \quad \bar{X}_n = c^{-1} \sum_{i=1}^c \bar{X}_{ni}.$$

We denote the ordered values of the Z_{ni} by $Z_{n[1]} < \dots < Z_{n[c]}$, and let $Z_{n(i)}$ be the statistic associated with $\tau_{[i]}$, $i=1, \dots, c$. The parametric procedures involve the statistics $Z_{n[1]}, \dots, Z_{n[c]}$.

For the nonparametric procedures, to be considered later on, we let

$$(2.2) \quad X_{ij,\alpha}^* = X_{i\alpha} - X_{j\alpha}, \quad e_{ij,\alpha} = e_{i\alpha} - e_{j\alpha}, \quad \alpha=1, \dots, n; \quad \Delta_{ij} = \tau_i - \tau_j, \quad \text{for } 1 \leq i < j \leq c.$$

We denote the marginal cdf of $e_{ij,\alpha}$ by $G(e)$, so that the marginal cdf of $X_{ij,\alpha}^*$ is $G(x - \Delta_{ij})$. It may be noted that by virtue of (1.2), $G(e)$ is symmetric about 0, so

that the distribution of $X_{ij,\alpha}^*$ is symmetric about Δ_{ij} for all $1 \leq i < j \leq c$ and $\alpha=1, \dots, n$. Also, note that even if the block effects β_1, \dots, β_n are random, the $X_{ij,\alpha}^*$ are independent of these variables. We write $\underline{X}_{ij} = (X_{ij,1}^*, \dots, X_{ij,n}^*)$, and consider the statistic

$$(2.3) \quad h_n(\underline{X}_{ij}^*) = n^{-1} \sum_{\alpha=1}^n E_{n,\alpha} Z_{n,\alpha},$$

where $Z_{n,\alpha} = 1$ if the α -th smallest observation among $|X_{ij,1}^*|, \dots, |X_{ij,n}^*|$ is from a positive $X_{ij,\beta}^*$, and $Z_{n,\alpha} = 0$, otherwise, for $\alpha=1, \dots, n$; $E_{n,\alpha}$ is the expected value of the α -th smallest observation of a sample of size n from the distribution $\Psi^*(x)$, given by

$$(2.4) \quad \Psi^*(x) = \begin{cases} \Psi(x) - \Psi(-x), & x \geq 0, \\ 0, & x < 0. \end{cases}$$

We assume that $\Psi(x)$ satisfies the following assumptions:

Assumption I. $\Psi(x)$ is symmetric about zero, i.e., $\Psi(x) + \Psi(-x) = 1$, for all x .

Assumption II. $n^{-1} \sum_{\alpha=1}^n [E_{n,\alpha} - \Psi^{*-1}(\alpha/n+1)] Z_{n,\alpha} = o_p(n^{-\frac{1}{2}})$.

Assumption III. $J(u) = \Psi^{-1}(u)$ ($0 < u < 1$) is absolutely continuous, and

$$(2.5) \quad |J^{(i)}(u)| = |d^i J(u)/du^i| \leq \kappa [u(1-u)]^{-i-\frac{1}{2}+\delta}, \quad i=0,1$$

where $\kappa (> 0)$ is finite and $\delta > 0$.

(It may be noted that the Wilcoxon signed rank statistic and the one sample normal scores statistic are special cases of (2.2), see for example [5,6]). Let $\underline{1}_n = (1, \dots, 1)$, and note that by definition $h_n(\underline{X}_{ij} - t\underline{1}_n)$ is non-increasing in t ($-\infty < t < \infty$). As in Puri and Sen (1967), we consider as an estimate of Δ_{ij} , the statistic

$$(2.6) \quad \hat{\Delta}_{ij}^{(n)} = [\hat{\Delta}_{ij,1}^{(n)} + \hat{\Delta}_{ij,2}^{(n)}]/2, \quad (1 \leq i < j \leq c)$$

where

$$(2.7) \quad \hat{\Delta}_{ij,1}^{(n)} = \text{Sup}\{t: h_n(\underline{X}_{ij} - t\underline{1}_n) > \mu_n\},$$

$$(2.8) \quad \hat{\Delta}_{ij,2}^{(n)} = \text{Inf}\{t: h_n(\underline{X}_{ij} - t\underline{1}_n) < \mu_n\},$$

and $\mu_n = (2n)^{-1} \sum_{\alpha=1}^n E_{n,\alpha} = \frac{1}{2} E_{\Psi}\{|X_{ij,\alpha}^*|\}$. Conventionally, we let $\hat{\Delta}_{ii}^{(n)} = \Delta_{ii} = 0$, $i=1, \dots, c$.

Also, as in Puri and Sen (1967), we define

$$(2.9) \quad Y_i^{(n)} = c^{-1} \sum_{j=1}^c \hat{\Delta}_{ij}^{(n)}, \quad i=1, \dots, c.$$

Let then $Y_{[1]}^{(n)} < \dots < Y_{[c]}^{(n)}$ be the ordered values of the $Y_i^{(n)}$, and let $Y_{(i)}^{(n)}$ be the statistic associated with $\tau_{[i]}$, $i=1, \dots, c$. The nonparametric procedures are based on $Y_{[1]}^{(n)}, \dots, Y_{[c]}^{(n)}$.

3. Exact as well as large sample (parametric) solutions to problem 1. In the case of independent and normally distributed errors, Bechhofer (1954) considered the solution based on the order statistics associated with $\bar{X}_{n1}, \dots, \bar{X}_{nc}$. His technique is not directly applicable when the errors are not independent (as in (1.2)). For this reason, we consider the following modification of his procedure:

Extended Bechhofer procedure (to be referred in the sequel as the B*-procedure).

Select the t best treatments which are associated with

$$(3.1) \quad Z_{n[c-t+1]}, \dots, Z_{n[c]}.$$

Our basic problem is to determine the sample size n is such a way that for any pre-assigned γ ($0 < \gamma < 1$), the probability of correct selection of the t best population is $\geq \gamma$, where $\tau_{[c-t]}$ and $\tau_{[c-t+1]}$ are subject to the condition that

$$(3.2) \quad \tau_{[c-t+1]} - \tau_{[c-t]} \geq \zeta,$$

ζ being the smallest worth detecting difference. Note that the choice of ζ is left to the practical considerations. We denote the condition in (3.2) by a sequence $\{\zeta_m\}$ where m is a positive integer and $\zeta_m \rightarrow 0$ as $m \rightarrow \infty$, we denote the corresponding sequence of conditions by $\{A_m\}$. In the context of one-way layout (which also extends to two-way layouts for independent errors), Bechhofer (1954) has shown that for the parent distribution being normal,

$$(3.3) \quad P\{\text{correct selection of the } t \text{ best treatments}\} = \gamma$$

when the following least favorable configuration holds:

$$(3.4) \quad \begin{cases} \tau_{[1]} = \dots = \tau_{[c-t]} = \tau_{[c-t+1]} - \zeta, \\ \tau_{[c-t+1]} = \dots = \tau_{[c]}. \end{cases}$$

We denote by $L(c,t;\zeta)$ the configuration in (3.4), and note that if ζ be replaced by $\{\zeta_m\}$, the corresponding sequence will be denoted by $\{L(c,t;\zeta_m)\}$.

We first show that $L(c,t;\zeta)$ is the least favorable configuration for the entire class of symmetric dependent multinormal distributions. This basic result will be used throughout the paper.

Theorem 3.1. Let (W_1, \dots, W_c) have jointly a multinormal distribution with mean vector $(\tau_{[1]}, \dots, \tau_{[c]})$ and dispersion matrix $\Sigma = \sigma^2[(1-\rho)\underline{I} + \rho\underline{J}]$, ($\sigma^2 > 0$, $-1/(c-1) \leq \rho < 1$, \underline{I} is the identity matrix of order c and $\underline{J} = \underline{1}'\underline{1}_c$). Then for given γ , under (A), (3.3) holds when (3.4) holds.

Proof. Denote by

$$(3.5) \quad U_{ij} = (W_i - W_j - [\tau_{[i]} - \tau_{[j]}]) / [2\sigma^2(1-\rho)]^{\frac{1}{2}}, \quad 1 \leq i \leq c-t, \quad c-t+1 \leq j \leq c,$$

and note that $\{U_{ij}, 1 \leq i \leq c-t, c-t+1 \leq j \leq c\}$ have jointly a (singular) multinormal distribution with null mean vector and dispersion matrix with the elements

$$(3.6) \quad \text{Cov}(U_{ij}, U_{i'j'}) = \begin{cases} 1, & i=i', j=j', \\ \frac{1}{2}, & i=i', j \neq j' \text{ or } i \neq i', j=j', \\ 0, & i \neq i', j \neq j'. \end{cases}$$

Now, by the procedure in (3.1), the probability of a correct selection is

$$(3.7) \quad \begin{aligned} & P\{\text{Max}[W_1, \dots, W_{c-t}] < \text{Min}[W_{c-t+1}, \dots, W_c]\} \\ &= P\{W_i - W_j < 0, i=1, \dots, c-t, j=c-t+1, \dots, c\} \\ &= P\{U_{ij} \leq (\tau_{[j]} - \tau_{[i]}) / [2\sigma^2(1-\rho)]^{\frac{1}{2}}, i=1, \dots, c-t, j=c-t+1, \dots, c\}. \end{aligned}$$

Now, in view of (3.2), the $\tau_{[i]}$ satisfy

$$(3.8) \quad \tau_{[1]} \leq \dots \leq \tau_{[c-t]} \leq \tau_{[c-t+1]} - \zeta \leq \tau_{[c-t+2]} \leq \dots \leq \tau_{[c]}.$$

Since the right hand side of (3.7) agrees with the corresponding expression in Bechhofer (1954, pp. 20-21), [with the only change that his σ^2 is replaced by our $\sigma^2(1-\rho)$], the least favorable configuration for which (3.3) holds can easily be shown (as in [1, pp. 20-21]) to be (3.4). Hence the theorem.

It may be noted that for the Z_{ni} , defined by (2.1), σ^2 in theorem 3.1 has to be replaced by σ^2/n , and as in Sen (1968), it follows that the mean square due to error consistently estimates $\sigma^2(1-\rho)$, for all parent distribution having finite

second moments. While proceeding to the large sample solutions to the problem, we note that by virtue of the consistency of the estimates Z_{ni} , $i=1, \dots, c$, for any fixed ζ in (3.2), the probability of correct decision will tend to unity as $n \rightarrow \infty$. Hence, to avoid this limiting degeneracy, we conceive of a sequence of worth-detecting differences which also converge to zero as $n \rightarrow \infty$, in such a way that (3.3) holds with a γ , strictly less than 1. The justification of the use of such a sequence can be made in the same light as the use of the well-known Pitman-efficiency is justified in nonparametric inference procedures. Theoretically speaking, we replace the single sequence $\{X_{i\alpha}, \alpha=1, \dots, n, i=1, \dots, c\}$ by a double sequence $\{[X_{i\alpha}^{(n)}, \alpha=1, \dots, n, i=1, \dots, c], n \geq 1\}$, where for the $X_{i\alpha}^{(n)}$, the corresponding $\tau_{[i]}$ (say $\tau_{[i]}^{(n)}$) satisfy (3.2) with ζ replaced by ζ_n , and $\zeta_n \rightarrow 0$ as $n \rightarrow \infty$. On the other hand, to practicing people, the solution can be made justified for small values of ζ ; small enough to make the applied approximations to be valid and reasonable.

We now drop the assumption of multinormality of the errors, and, as in (1.2), we let F to be arbitrary. For the parametric procedures based on sample means, we require that F possesses second order moments.

Theorem 3.2. Suppose in (3.3) γ is fixed, and in (3.2), ζ is replaced by ζ_n . Then for a γ - probability of correct selection, we have

$$(3.9) \quad |n^{\frac{1}{2}}\zeta_n - \delta\sigma\sqrt{1-\rho}| \rightarrow 0 \text{ as } n \rightarrow \infty,$$

where σ^2 is the variance of $e_{i\alpha}$, ρ is the common correlation of $e_{i\alpha}$, $e_{i\beta}$ ($\beta \neq \alpha$), and δ is determined by the condition:

$$(3.10) \quad \gamma = tQ_{c-1}(\underbrace{\delta//2, \dots, \delta//2}_{c-t \text{ times}}, \underbrace{0, \dots, 0}_{t-1 \text{ times}}),$$

Q_{c-1} being the cumulative distribution function of a normally distributed $(c-1)$ -vector $(U_1, \dots, U_{c-t}, W_{c-t+2}, \dots, W_c)$, with $EU_i = EW_j = 0$, $\text{Cov}[U_i, U_{i'}] = \frac{1}{2}(1 + \delta_{ii'})$, $\text{Cov}[W_j, W_{j'}] = \frac{1}{2}(1 + \delta_{jj'})$ and $\text{Cov}[U_i, W_{j'}] = -\frac{1}{2}$ for $i, i' = 1, \dots, c-t$, $j, j' = c-t+2, \dots, c$; $\delta_{rs} = 0, 1$ according as $r \neq s$ or $r = s$.

The proof follows as a special case of (5.4) [with simplified (5.1)], and hence is omitted.

Suppose now we are given a small ζ^* and we wish to determine n such that (3.3) holds [subject to (3.2), with ζ replaced by ζ^*]. Then theorem 3.2 provides the following large sample solution:

$$(3.11) \quad n \approx \delta^2 \sigma^2 (1-\rho) / (\zeta^*)^2,$$

and as s^2 , the mean square due to error estimates $\sigma^2(1-\rho)$ consistently, we have also asymptotically

$$(3.12) \quad n \approx \delta^2 s^2 / (\zeta^*)^2; \quad s^2 = \frac{1}{(n-1)(c-1)} \sum_{\alpha=1}^n \sum_{i=1}^c (X_{i\alpha} - \bar{X}_{i.} - \bar{X}_{.\alpha} + \bar{X}_{..})^2,$$

$$X_{i.} = n^{-1} \sum_{\alpha=1}^n X_{i\alpha}, \quad i = 1, \dots, e, \quad \bar{X}_{.\alpha} = c^{-1} \sum_{i=1}^c X_{i\alpha}, \quad \alpha = 1, \dots, n, \text{ and}$$

$$X_{..} = (nc)^{-1} \sum_{\alpha=1}^n \sum_{i=1}^c X_{i\alpha}.$$

4. Procedures based on rank order estimates. Here, we select the t populations associated with

$$(4.1) \quad Y_{[c-t+1]}^{(n)}, \dots, Y_{[c]}^{(n)}.$$

The basic theorem of this section is the following.

Theorem 4.1. Under (1.1) and (1.2) and the assumptions I, II and III of Section 2, the limiting distribution of $n^{1/2}[Y_{(i)}^{(n)} - Y_{(j)}^{(n)} - \Delta_{ij}; 1 \leq i \leq c-t, c-t+1 \leq j \leq c]$ is multivariate normal with null means, and

$$(4.2) \quad \text{cov}\{n^{1/2}[Y_{(i)}^{(n)} - Y_{(j)}^{(n)}], n^{1/2}[Y_{(i')}^{(n)} - Y_{(j')}^{(n)}]\} = \begin{cases} \sigma_0^2, & \text{if } i=i', j=j' \\ \frac{1}{2} \sigma_0^2, & \text{if } i=i', j \neq j' \text{ or } i \neq i', j=j', \\ 0, & \text{if } i \neq i', j \neq j', \end{cases}$$

where

$$(4.3) \quad \sigma_0^2 = [A^2 + (c-2)\lambda_J(F)] / (cB^2),$$

$$(4.4) \quad A^2 = \int_0^1 J^2(u) du, \quad B = \int_{-\infty}^{\infty} (d/dx) J[G(x)] dG(x); \quad J(u) = \Psi^{-1}(u),$$

$$(4.5) \quad \lambda_J(F) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} J[G(x)] J[G(x)] J[G(y)] dG^*(x, y),$$

and $G^*(x,y)$ is the joint cdf of $(e_{ij,\alpha}^*, e_{ij',\alpha}^*)$ ($j \neq j'$), whose marginals are both $G(x)$.

The proof of the theorem when the errors $e_{i\alpha}$ are mutually independent follows from theorem 3.2 of Puri and Sen (1967). However, the multivariate approach of Sen and Puri (1967) along with lemma 2.1 of Sen (1968) extends the proof to the case of symmetric dependent errors. For intended brevity, the details are omitted.

Now, the probability of correct selection of t best populations is given by

$$(4.6) \quad P\{\max[Y_{(1)}^{(n)}, \dots, Y_{(c-t)}^{(n)}] < \min[Y_{(c-t+1)}^{(n)}, \dots, Y_{(c)}^{(n)}]\} \\ = P\{(\frac{n}{2\sigma_0^2})^{1/2} [Y_{(i)}^{(n)} - Y_{(j)}^{(n)} - \Delta_{ij}] < (\frac{n}{2\sigma_0^2})^{1/2} \Delta_{ji}, i=1, \dots, c-t, j=c-t+1, \dots, c\} .$$

Thus, as in the asymptotic parametric case, we replace Δ_{ji} by a sequence $\{\Delta_{ji}^{(n)}\}$ such that $n^{1/2} \Delta_{ji}^{(n)} \rightarrow \lambda_{ji}$ (real and finite) as $n \rightarrow \infty$. Then, by using theorem 4.1, we conclude that the right hand side of (4.6) is asymptotically equal to

$$(4.7) \quad P\{U_{ij} < (n/2\sigma_0^2)^{1/2} \Delta_{ij}^{(n)}, i=1, \dots, c-t, j=c-t+1, \dots, c\} ,$$

where the U_{ij} have a multinormal distribution with null mean vector and covariance matrix, given by (4.2). Since, this multinormal distribution satisfies the condition of theorem 3.1, we can again check easily that the least favorable configuration turns out to be (3.4) with ζ replaced by ζ_n . Moreover, by the same technique as in theorem 3.2, it follows that

$$(4.8) \quad |n^{1/2} \zeta_n - \delta \sigma_0| \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

where σ_0 is defined by (4.3) and δ by (3.10). Thus, a large sample solution to the sample size needed for the probability of correct selection being equal to γ is given by

$$(4.9) \quad n \approx \delta^2 \sigma_0^2 / (\zeta^*)^2,$$

where ζ^* is a given small worth-detecting difference. Now, σ_0^2 involves the two unknown parameters B and $\lambda_J(F)$. B can be estimated (by \hat{B}) as in Sen (1966), while $\lambda_J(F)$ can be estimated (by $L_J(F)$) as in Puri and Sen (1967). Hence, we have

$$(4.10) \quad n \approx \delta^2 [A^2 + (c-2)L_J(F)] / c\hat{B}^2.$$

Now, using the same notion of asymptotic relative efficiency (ARE) as in Puri and Puri (1969), it follows from (3.11) and (4.10) that the ARE of the rank order procedure (based on ψ -scores) with respect to the B^* -procedure is

$$(4.11) \quad e(\psi, B^*) = \sigma^2(1-\rho)/\sigma_0^2 \\ = \{[2\sigma^2(1-\rho)]B^2\} \{c/2[A^2+(c-2)\lambda_J(F)]\} .$$

Now, B relates to the cdf G [cf. (4.4)] whose variance is $2\sigma^2(1-\rho)$. Hence, the first factor on the right hand side of (4.11) is the A.R.E. of the one-sample rank order tests (for location) with respect to the Student t -test when the parent distribution is $G(x)$ [cf. Puri and Sen (1969)], and we denote it by $e_{\psi, B^*}(G)$. Further, it has been shown in Puri and Sen (1967) that $\lambda_J(F) \leq \frac{1}{2} A^2$, where the equality sign holds iff $J[F(x)]$ is linear in x , with probability one. As such, we have

$$(4.12) \quad e(\psi, B^*) \geq e_{\psi, B^*}(G),$$

where the equality sign holds iff $J[F(x)] = a+bx$, with probability one. Now, various known bounds for $e_{\psi, B^*}(G)$ can be used to provide bounds to $e(\psi, B^*)$. For example, if we use Ψ as the standard normal distribution, $e_{\psi, B^*}(G)$ is bounded below by 1, where the lower bound is attained iff G is normal. Thus, the procedure based on the normal Scores estimators is asymptotically at least as efficient as the extended Bechhofer procedure. If we use the Wilcoxon-Scores estimator, it follows that $e(\psi, B^*)$ is bounded below by 0.864 (though not attainable) for all $F(G)$, while the same can be greater than unity for many non-normal F . For normal F , it is bounded below by $3/\pi$ for all $c(\geq 2)$, while it can be as high as 0.98.

5. Relative performance characteristics when (3.4) is not necessarily true.

To study the relative performance of the Ψ -score procedure with respect to the B^* -procedure, we consider any sequence of parameter points satisfying

$$(5.1) \quad \tau_{[c-t+1]}^{(n)} - \tau_{[i]}^{(n)} = \delta_i^{(n)} = n^{-1/2} \theta_i + o(n^{-1/2}), \quad i=1, \dots, c-t, c-t+2, \dots, c,$$

where not all the $\theta_1, \dots, \theta_{c-t}$ are equal to θ , and/or not all the $\theta_{c-t+2}, \dots, \theta_c$ are equal to zero, i.e. the least favorable configuration does not hold, but

$\tau_{[i]}^{(n)} \leq \tau_{[j]}^{(n)}$ whenever $1 \leq i \leq j \leq c$. Then, for the B*-procedure, we have

$$(5.2) \quad \begin{aligned} & \lim_{n \rightarrow \infty} P\{\text{correct selection of } t \text{ best treatments}\} \\ &= \lim_{n \rightarrow \infty} P\{\max[Z_{n(1)}, \dots, Z_{n(c-t)}] < \min[Z_{n(c-t+1)}, \dots, Z_{n(c)}]\} \\ &= \lim_{n \rightarrow \infty} \left[\sum_{\ell=1}^t P\{\text{all the } Z_{n(1)}, \dots, Z_{n(c-t)} < Z_{(c-t+\ell)} < \min[Z_{n(i)}, i=c-t+1, \dots, c(\neq c-t+\ell)]\} \right] \\ &= \sum_{\ell=1}^t \left[\lim_{n \rightarrow \infty} P\{Z_{n(s)} - Z_{n(\ell)} < 0, s=1, \dots, c-t, \right. \\ & \quad \left. Z_{n(\ell)} - Z_{n(r)} < 0, r=c-t+1, \dots, c(\neq c-t+\ell)\} \right] \\ &= \sum_{\ell=1}^t \left[\lim_{n \rightarrow \infty} P\{U_s < \xi_{\ell, s}^{(n)}, W_r < \xi_{r, \ell}^{(n)}, s=1, \dots, c-t, \right. \\ & \quad \left. r=c-t+1, \dots, c(\neq c-t+\ell)\} \right], \end{aligned}$$

where $[U_1, \dots, U_{c-t}, W_{c-t+1}, \dots, W_{-1}, W_{i+1}, \dots, W_c]$ has the multivariate normal distribution Q_{c-1} , defined in theorem 3.2, and

$$(5.3) \quad \xi_{r, s}^{(n)} = [n/2\sigma^2(1-\rho)]^{1/2} [\tau_{[r]}^{(n)} - \tau_{[s]}^{(n)}], \quad 1 \leq r \leq s \leq c;$$

the last identify in (5.2) is a direct consequence of the central limit theorem as applied on the $Z_{n(r)}$. Thus, from (5.1), (5.2) and (5.3), we obtain that (5.2) is equal to

$$(5.4) \quad \sum_{\ell=c-t+1}^c Q_{c-1} \left(\frac{1}{\sqrt{2(1-\rho)}} [\theta_{1, \ell}, \dots, \theta_{c-t, \ell}, \theta_{c-t+1, \ell}, \dots, \theta_{\ell-1, \ell}, \theta_{\ell+1, \ell}, \dots, \theta_{c, \ell}] \right),$$

where $\theta_{i, j} = \theta_i - \theta_j$.

Consider now the rank procedures, where the assumptions I, II and III of section 2 hold and the sequence in (5.1) also holds. Then, by theorem 4.1, it follows as in (5.2) that under (5.1), for the ψ -scores procedure

$$(5.5) \quad \begin{aligned} & \lim_{n \rightarrow \infty} P\{\text{correct solution of } t \text{ best treatments}\} \\ &= \sum_{\ell=c-t+1}^c Q_{c-1} \left(\frac{\sigma}{\sqrt{2} \sigma_0} \theta_{1, \ell}, \dots, \theta_{c-t, \ell}, \theta_{c-t+1, \ell}, \dots, \theta_{\ell-1, \ell}, \theta_{\ell+1, \ell}, \dots, \theta_{c, \ell} \right), \end{aligned}$$

where σ_0^2 is defined by (4.3). Thus, comparing (5.4) and (5.5) (in the Pitman sense), we may conclude that the A.R.E. remains the same for (5.1), even when the least favorable configuration may not hold.

6. Selection of best treatments with regard to order. Here the Bechhofer procedure consists in selecting the t best treatments associated with $Z_{n[c-t+1]}, \dots, Z_{n[c]}$ respectively. By virtue of our theorem 3.1, we can readily extend the original procedure by Bechhofer (1954), and derive the following results. [The details are omitted for intended brevity.]

For a fixed $\gamma(0 < \gamma < 1)$ and under the condition that

$$(6.1) \quad \tau_{[i+1]}^{-\tau_{[i]}} \geq \zeta_n \text{ (worth detecting distance), } i=c-t, \dots, c-1,$$

let n be determined so that (a) the following (least favorable) configuration holds:

$$(6.2) \quad \tau_{[1]} = \dots = \tau_{[c-t]} = \tau_{[c-t+1]}^{-\zeta_n}, \tau_{[i+1]}^{-\tau_{[i]}} = \zeta_n, i=c-t, \dots, c-1,$$

and (b)

$$(6.3) \quad P\{\max_{1 \leq i \leq c-t} Z_{n(i)} < Z_{n(c-t+1)} < \dots < Z_{n(c)}\} = \gamma.$$

Then asymptotically,

$$(6.4) \quad |n^{1/2} \zeta_n - \delta \sigma \sqrt{1-\rho}| \rightarrow 0, \text{ (as } n \rightarrow \infty),$$

where σ and ρ are defined as in theorem 3.2 and δ is determined by the condition

$$(6.5) \quad (c-t)Q_{c-1} \underset{c-t-1 \text{ times}}{(0, \dots, 0, \delta/\sqrt{2}, \dots, \delta/\sqrt{2})} \underset{t \text{ times}}{=} \gamma,$$

and Q_{c-1} is the cdf of a normally distributed vector $(U_1, \dots, U_{c-t-1}, W_{c-t}, \dots, W_{c-1})$ satisfying $EU_i = EW_j = 0, i=1, \dots, c-t-1, j=c-t, \dots, c-1$ and (i) $\text{Cov}(U_i, U_{i'}) = \frac{1}{2}(1 + \delta_{ii'})$, (ii) $\text{Cov}(U_i, W_j) = -\frac{1}{2}$ if $j=c-t$ and 0, otherwise, and (iii) $\text{Cov}(W_j, W_{j'}) = 1, -\frac{1}{2}$ or 0 according as $j=j', |j-j'|=1$ or $|j-j'| > 1$, for $i, i'=1, \dots, c-t-1, j, j'=c-t, \dots, c-1$; $\delta_{ii'}$ being the usual Kronecker delta.

Again by virtue of our theorems 3.1 and 4.1, it follows along the same line as in [4] that for the rank scores procedure based on (4.1) (with regard to the

order), the least favorable configuration turns out to be (6.2) (for small ζ_n), and hence, for this procedure

$$(6.6) \quad |n^{1/2} \zeta_n - \delta\sigma_0| \rightarrow 0, \text{ as } n \rightarrow \infty,$$

where σ_0 is defined by (4.3) and δ by (6.5).

Hence, the A.R.E. is the same as in (4.11) and (4.12). Also, the results of section 5 can readily be extended in this situation and similar conclusions be derived; for brevity the details are omitted.

7. Selection of a subset of treatments better than a standard one. Instead of (1.1), we consider the model

$$(7.1) \quad X_{i\alpha} = \mu + \beta_\alpha + \tau_i + \epsilon_{i\alpha}, \quad i = 0, 1, \dots, c; \alpha = 1, \dots, n,$$

where the notations are all explained after (1.1) and τ_0 is the standard treatment effect. We say that the i th treatment is better than the standard if

$$(7.2) \quad \tau_i \geq \tau_0 + \zeta; \quad \zeta = \text{worth detecting difference.}$$

For the one-way layout case with normally distributed errors, Gupta and Sobel (1958) proposed the procedure: Select the subset of treatments for which $X_{ni} - X_{n0} > 0$, where $X_{ni} = n^{-1} \sum_{\alpha=1}^n X_{i\alpha}$, $i = 0, 1, \dots, c$. An elegant solution for the sample size (n) needed to achieve a γ ($0 < \gamma < 1$) probability of correct selection has also been provided by them. Noting that for (within-block) symmetric dependent normally distributed errors, the distribution of $[n^{1/2}(\bar{X}_{ni} - \bar{X}_{n0} - \tau_i + \tau_0)]$, $i=1, \dots, c$ is also multinormal with null means and covariance matrix $\sigma^2(1-\rho)[I_c + J_c]$, (where σ^2 and ρ are defined in theorem 3.2), it turns out that the only change needed in Gupta-Sobel solution for the two-way layout problem is to replace their σ^2 by $\sigma^2(1-\rho)$. By virtue of the central limit theorem, the same solution holds asymptotically for the entire class of cdf's with finite second moments.

For the rank scores procedure, we define the estimators $\hat{\Delta}_{i0}^{(n)}$, $i=1, \dots, c$ as in (2.5) and (2.6). Then, we select the subset of treatments for which $\hat{\Delta}_{i0}^{(n)} > 0$.

Along the same line as in theorem 4.1, it follows that $[n^{1/2}(\hat{\Delta}_{i0}^{(n)} - \tau_i + \tau_0), i=1, \dots, c]$ have asymptotically a multinormal distribution with null means and covariance matrix $\sigma_0^2 [I_c + J_c]$, where σ_0^2 is defined by (4.3). Consequently, the Gupta-Sobel solution also asymptotically holds for the rank scores procedure provided we replace σ^2 by σ_0^2 . Hence, the A.R.E. of the rank scores procedure with respect to the Gupta-Sobel procedure can again be measured by $\sigma^2(1-\rho)/\sigma_0^2$, and it agrees with (4.11). The details are therefore omitted. However, the results clearly indicate the superiority of the normal scores procedure over the normal theory procedure (for large samples).

8. Few additional comments. Unlike Puri and Puri (1969), we have considered here the procedures based on the rank order estimates, not statistics. The same procedure can be suggested for the one-way layout problem as an alternative to the procedures by Puri and Puri (1969). Also, we note that the procedure considered in this paper can readily be extended to incomplete block designs (such as the paired comparisons designs, etc.), whereas the original procedures by Puri and Puri face considerable difficulties.

REFERENCES

- [1] BECHHOFFER, R. E. (1954). A single sample multiple decision procedure for ranking means of normal populations with known variances. Ann. Math. Statist. 25, 16-39.
- [2] GUPTA, S. S. and SOBEL, M. (1958). On selecting a subset which contains all populations better than a standard. Ann. Math. Statist. 29, 235-244.
- [3] PAULSON, E. (1952). On the comparison of several experimental categories with a control. Ann. Math. Statist. 23, 239-246.
- [4] PURI, M. L. and PURI, P. S. (1969). Multiple decision procedures based on ranks for certain problems in analysis of variance. Ann. Math. Statist. 40, 619-632.
- [5] PURI, M. L. and SEN, P. K. (1967). On some optimum nonparametric procedures in two-way layouts. J. Amer. Statist. Asso. 62, 1214-1229.
- [6] PURI, M. L. and SEN, P. K. (1969). On the asymptotic normality of one-sample rank order test statistics. Teor. Veroi. Prim. 14, 172-177.
- [7] SEN, P. K. (1966). On a distribution-free method of estimating asymptotic efficiency of a class of nonparametric tests. Ann. Math. Statist. 37, 1759-1770.
- [8] SEN, P. K. (1968). Robustness of some nonparametric procedures in linear models. Ann. Math. Statist. 39, 1913-1922.
- [9] SEN, P. K. and PURI, M. L. (1967). On the theory of rank order tests for location in the multivariate one sample problem. Ann. Math. Statist. 38, 1216-1228.